

Expression of homologous RND efflux pump genes is dependent upon AcrB expression

Blair, Jessica M A; Smith, Helen E; Ricci, Vito; Lawler, Amelia J; Thompson, Louisa J; Piddock, Laura J V

DOI:

[10.1093/jac/dku380](https://doi.org/10.1093/jac/dku380)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Blair, JMA, Smith, HE, Ricci, V, Lawler, AJ, Thompson, LJ & Piddock, LJV 2015, 'Expression of homologous RND efflux pump genes is dependent upon AcrB expression: implications for efflux and virulence inhibitor design', *Journal of Antimicrobial Chemotherapy*, vol. 70, no. 2, pp. 424-431. <https://doi.org/10.1093/jac/dku380>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Eligibility for repository: Checked on 9/2/2016

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Expression of homologous RND efflux pump genes is dependent upon AcrB expression:
Implications for efflux and virulence inhibitor design

Jessica M. A. Blair, Helen E. Smith, Vito Ricci, Amelia J. Lawler, Louisa J. Thompson and Laura
J.V. Piddock*

*Antimicrobials Research Group, School of Immunity and Infection, College of Medical and
Dental Sciences, and Institute of Microbiology and Infection, The University of Birmingham,
B15 2TT, UK*

Running title: Efflux pump redundancy

Keywords: redundancy, AcrB, AcrD, AcrF

*Corresponding author. Mailing address: Antimicrobials Research Group, School of Immunity
and Infection and Institute of Microbiology & Infection, College of Medical and Dental
Sciences. University of Birmingham, Birmingham, B15 2TT, United Kingdom. Telephone:
(+44) (0)121-414-6966; Fax (+44) (0)121-414-6819. E-mail: l.j.v.piddock@bham.ac.uk

20

Synopsis

21 **Objectives** Enterobacteriaceae have multiple efflux pumps which confer intrinsic resistance
22 to antibiotics. AcrB mediates clinically relevant multi-drug resistance and is required for
23 virulence and biofilm formation making it an attractive target for the design of inhibitors.
24 The aim of this study was to assess the viability of single transporters as a target for efflux
25 inhibition using *S. Typhimurium* as the model pathogen.

26 **Methods** The expression of RND efflux pump genes, in response to inactivation of single or
27 multiple homologues was measured using real-time RT-PCR. Phenotypes of mutants were
28 characterised by measuring antimicrobial susceptibility, dye accumulation and ability to
29 cause infection *in vitro*.

30 **Results** Expression of all RND efflux pump genes was increased when single or multiple *acr*
31 genes were inactivated, suggesting a feedback mechanism which activates transcription of
32 homologous efflux pump genes. When two or three *acr* genes were inactivated, the
33 mutants had further reduced efflux, altered susceptibility to antimicrobials (including
34 increased susceptibility to some, but conversely and counter-intuitively, decreased
35 susceptibility to some others) and were more attenuated in the tissue culture model than
36 mutants lacking single pumps.

37 **Conclusions** These data indicate that it is critical to understand which pumps an inhibitor is
38 active against and the effect of this on expression of homologous systems. For some
39 antimicrobials, an inhibitor with activity against multiple pumps will have a greater impact
40 on susceptibility but, an unintended consequence of this may be decreased susceptibility to
41 other drugs, such as aminoglycosides.

42

Introduction

Efflux is an important mechanism of multidrug resistance in bacteria, conferring decreased susceptibility to a wide range of substrates including antibiotics, dyes, detergents and biocides.¹ This makes them an attractive target for the design of inhibitors which could be used to potentiate the use of existing antimicrobials.

Resistance nodulation division (RND) efflux transporters are found in the inner membrane of Gram-negative bacteria and form a complex with an outer membrane channel and a periplasmic adaptor protein (PAP) to form a tri-partite efflux pump system spanning both the inner and outer membrane.^{2, 3} Substrates of these multi-protein complexes are structurally diverse and include antibiotics, biocides, dyes, detergents and host-derived molecules. Active efflux of substrates by RND systems is responsible for the intrinsic resistance of Gram negative bacteria to multiple classes of structurally distinct antimicrobials.¹

Salmonella have five RND efflux pump systems: AcrAB, AcrD, AcrEF, MdtABC and MdsABC. Further RND efflux pumps are found in some other members of the Enterobacteriaceae including MdtF in *Escherichia coli*.⁴ The transporter protein, AcrB, and its homologues in other Gram-negative bacteria, is considered the most clinically relevant RND system because it has the broadest substrate profile and is more abundant within the cell than other efflux systems.¹ Inactivation of *acrB* increases susceptibility of laboratory mutants of *E. coli*, *Salmonella enterica* and other Enterobacteriaceae to many antimicrobials, whereas overexpression confers multi-drug resistance (MDR), including to clinically efficacious agents. Such mutants have been selected *in vitro* and *in vivo*.⁵⁻⁸ Efflux via AcrB is driven by the proton-motive force and forms a tri-partite complex with the PAP, AcrA, and the outer

66 membrane channel, TolC.⁹ The recent elucidation of the structure of AcrB in complex with
 67 different substrates of varying molecular weights has revealed two large, discrete, multisite
 68 binding pockets within AcrB, which may explain how AcrB can transport such structurally
 69 varied substrates.^{10, 11}

70 Single deletions of RND efflux pump genes other than *acrB* have little or no effect on the
 71 susceptibility of Enterobacteriaceae to most antimicrobial agents.¹² Antimicrobial
 72 susceptibility of deletion mutants and strains with increased expression of certain RND
 73 efflux pumps indicates that there is overlap, or redundancy between the antimicrobials,
 74 biocides, dyes and detergents that can be transported by the different RND pumps of
 75 *Salmonella*.^{12, 13} AcrF is the closest homologue of AcrB in *Salmonella* (80% identity) and
 76 AcrEF overexpression can suppress antibiotic hyper-susceptibility in AcrB deficient strains.^{12,}
 77 ¹³ AcrD (64% identity to AcrB) and MdtABC have similar substrate profiles including SDS,
 78 novobiocin, deoxycholate, some β -lactams, copper and zinc.^{12, 14} In *E. coli* and *Salmonella*,
 79 AcrD also transports aminoglycoside antibiotics.^{15, 16} MdsABC is found only in *Salmonella*
 80 and in LB medium is expressed at lower levels than the other four RND efflux pumps.¹⁷
 81 However, overexpression of *mdsAB*, encoding the pump and periplasmic adaptor protein,
 82 or *MdsABC*, encoding all three components, were overexpressed in a strain lacking AcrB,
 83 susceptibility to novobiocin, acriflavine, crystal violet, methylene blue, rhodamine 6G,
 84 benzalkonium chloride and SDS was decreased compared to the mutant suggesting that
 85 MdsB is capable of exporting these compounds.¹⁸

86 In addition to the role in resistance to antimicrobials, some RND efflux pumps are also
 87 required for virulence of many Gram-negative pathogens.¹⁹ In *S. Typhimurium*, inactivation
 88 of *acrB* attenuated invasion of tissue culture cells *in vitro* and colonisation in poultry.^{20, 21}

Inactivation of *acrAB* or *acrEF* has also been shown to attenuate the virulence of the organism in mice.¹²

Regulation of RND efflux pumps is complex; transcriptional control is multi-layered and some regulators control expression of more than one pump. In *Salmonella* and *E. coli*, the regulation of *acrAB* is the best studied. The *acrA* and *acrB* genes are encoded in a single operon and are co-regulated. At a local level *acrAB* is repressed by AcrR, which is encoded alongside and divergently transcribed from *acrAB*. At a global level members of the AraC/XylS family of DNA transcriptional activators, such as MarA, SoxS, Rob and RamA, all influence expression of *acrAB-tolC* and have overlapping recognition sites.²²⁻²⁷ AcrD and MdtABC are both under the control of the two-component regulatory systems BaeSR and CpxAR which induce expression of AcrD and MdtABC in response to high levels of indole, copper, zinc or envelope stress.^{28, 29} Expression of *acrEF* in *E. coli* is generally low due to repression by the global regulator H-NS.³⁰ However, *acrEF* is also encoded alongside a gene encoding a local repressor, AcrS (previously EnvR) which inhibits expression of *acrAB* and acts as a regulatory switch between expression of *acrAB* and *acrEF*.³¹

Due to the functional redundancy of RND pumps, potential exists for the loss of certain pump components to be compensated by increased expression of a homologous component that could fulfil, at least to some extent, the same function. For example, Eaves and colleagues³² showed that when *acrB* or *acrF* of *Salmonella* was inactivated, expression of *acrD* increased (3.6 and 4.9 fold, respectively). To allow compensatory changes in expression levels of efflux systems upon inactivation of homologous systems, there must be a tightly controlled and integrated regulatory network that can respond to loss of efflux function. Whilst the literature contains multiple examples of regulation of single efflux

112 systems or a small number of efflux systems, an integrated network of regulation is yet to
113 be elucidated.

114 The role of RND systems in both antimicrobial resistance and virulence makes them
115 attractive targets for the design of inhibitors. Using *S. Typhimurium* as a model, the aim of
116 the present study was to investigate the viability of single transporters such as AcrB as a
117 target for efflux inhibition by investigating the expression and roles of structurally similar
118 efflux pumps in antimicrobial resistance and virulence, and the extent of the redundancy
119 between RND efflux pumps. This information is crucial for the rational design of inhibitors
120 which inhibit all pumps so preventing resistance by compensatory over-production of
121 homologous RND efflux systems.

Materials and Methods

Strains and growth conditions. All strains were derived from *Salmonella enterica* serovar Typhimurium SL1344.³³ *Salmonella* was used as a model organism in this study as it is an important human pathogen that causes a significant number of infections annually. SL1344 is a widely studied pathogenic strain of *Salmonella* for which there are well validated models of infection including an *in vitro* tissue culture model. Single gene inactivated mutants were constructed as described previously.^{32, 34} Mutants with multiple efflux pumps inactivated were created by P22 transduction between mutants in which single genes were inactivated or deleted. All mutants were verified by PCR and DNA sequencing. All experiments including MICs reveal that the phenotype of the marked and unmarked mutants for the same gene is indistinguishable. LB broth (Sigma-Aldrich, UK) and MOPS minimal medium (Teknova Inc., USA) were used throughout this study.

RNA extraction and Real Time qRT-PCR. Overnight cultures of *S. Typhimurium* SL1344 and the test strains were grown in MOPS minimal medium at 37°C. From each strain three biological replicate RNA preparations were made and quantified as described previously.^{22, 35} Primers (Supplementary Table 1) were designed with an annealing temperature of 57.3°C using Beacon Designer 4.0 (Premier Biosoft, USA). cDNA was synthesised from 2 µg total RNA using superscript III cDNA synthesis kit (Invitrogen). Validation experiments were carried out using 5 cDNA standards of different concentrations (10, 1, 0.1, 0.01, 0.001 ng/µl) to determine PCR efficiency for the housekeeping gene 16S and each test gene. qRT-PCRs were set up in biological triplicate and technical duplicate in a BIORAD PCR tray using 1 µl neat cDNA for test genes and 1 µl of a 1:1000 dilution cDNA for 16S in a 25 µl reaction containing 12.5 µl iQ SYBR green supermix (BIORAD, UK), 1 µl primers (500nm) and 9.5 µl

sterile water. qRT-PCR was carried out in a CFX-96 Real-time machine (BIORAD, UK) using the following protocol; 95°C for 5 minutes followed by 40 plate read cycles of 95°C for 30 s, 57.3°C for 30 s and 72°C for 30 s. Data were analysed using CFX Manager (BIORAD, UK) and expression ratios were calculated using the $\Delta\Delta\text{Ct}$ method and normalised to expression of 16S.³⁶

Determination of susceptibility to antimicrobials. Biolog data was confirmed by measuring growth in the presence of representative AcrB substrates. Briefly, overnight bacterial cultures were diluted to 10⁴ CFU/ml and grown in a 96 well plate in iso-sensitest broth in the presence of selected drugs at 0.25 x the wild-type MIC.

The minimum inhibitory concentrations (MIC) of antibiotics, dyes and detergents were determined for each strain according to the standardised agar doubling dilution method procedure of the British Society of Antimicrobial Chemotherapy (BSAC) using iso-sensitest agar.³⁷ The MIC was determined as the lowest concentration of antimicrobial that caused no visible growth. Values stated are the mode value from at least three biological replicates performed on at least three independent occasions. All antimicrobials tested were obtained from Sigma, UK.

Accumulation of Hoechst 33342 and norfloxacin. The efflux activity of the mutants was assessed by determining the accumulation of the fluorescent dye Hoechst 33342 and norfloxacin (Sigma, UK) as described previously.^{38, 39} Differences in steady state accumulation values between mutants and parental strains were analysed for statistical significance using a two-tailed Student's t-test, where $P < 0.05$ was considered significant. Data presented are the mean of three independent biological replicates \pm standard error of the mean.

168 **Adhesion and invasion assays.** The ability of the strains to adhere to, and invade, INT-407
169 (human embryonic intestine cells) was measured as previously described.²¹ Each assay was
170 repeated a minimum of three times, with each experiment including four technical
171 replicates per bacterial strain. The results were analysed using Student's *t*-test and P values
172 of ≤ 0.05 were considered significant.

173

Results

***In silico* analysis**

The *Salmonella* genome encodes five efflux systems of the RND family. The three most similar pump proteins are AcrB, AcrD and AcrF (Table 1). The two remaining RND pumps, MdsB and MdtB/C, which is a heteromultimer of MdtB and MdtC, are 46.7% and 48.6% identical to AcrB, respectively. ClustalW alignments of all six RND type proteins revealed that five residues (Asp407, Asp408, Lys940, Arg971 and Thr978) vital to proton transport within AcrB of *E. coli*^{40, 41} are identical in all six *Salmonella* proteins indicating that energy transduction is conserved in this family of proteins. However, greater variation was seen in residues involved in substrate recognition or binding. For example, of six important residues in the hydrophobic, phenylalanine rich binding pocket of AcrB (Phe178, 615, 610, 136, 617 and 628)⁴² all six were conserved in AcrF, two in AcrD, one in MdtC, one in MdsB and none of these residues were found in MdtB.

Expression of efflux pump genes is altered upon inactivation of homologous pumps.

To determine whether the expression of each gene was altered upon inactivation of one or more homologous genes, real-time quantitative RT-PCR was used to measure the level of *acrB*, *acrD* and *acrF* transcription in the single, double and triple efflux mutants compared to SL1344 (Table 2). As shown previously, in the *acrB* mutant both *acrD* and *acrF* had increased expression.³² In the *acrD* mutant, expression of *acrB* was increased while *acrF* expression was unchanged. In the *acrF* mutant both *acrB* and *acrD* had increased expression. The expression of *mdsB* and *mdtB*, was also affected by loss of single RND pumps. The expression of *mdtB* was increased in the *acrF* mutant while expression of the *Salmonella* specific efflux pump *mdsB* was decreased upon inactivation of *acrB*, *acrD* or *acrF* (Table 2).

When two genes were inactivated (e.g. *acrB* and *acrF* or *acrB* and *acrD*) expression of *acrD* or *acrF* was increased, although expression was lower than in the strain lacking only *acrB* (L110). When *acrF* and *acrD* were inactivated expression of *acrB* was increased, this was greater than that seen upon inactivation of *acrD* and similar to that in the *acrF* mutant. Expression of the *mdtB* and *mdsB* efflux genes was increased in all *acr* gene double mutants (L646, L1297 and L1395) but in the triple *acrBDF* mutant only *mdtB* expression was increased. Expression of both *mdtB* and *mdsB* was highest in the *acrDF* mutant (L1395) (Table 2).

The expression level of known regulators of RND efflux was also measured. The expression of *ramA* and *marA* was increased when *acrB* was inactivated but not changed in the *acrD* (L132) or *acrF* (L131) mutants (Table 2). Expression of *soxS* was increased in the *acrBF* mutant (6.3 fold), in the *acrDF* mutant (2.2 fold) and in the strain lacking all three RND pump genes. Expression of *rob* was not significantly altered in any of the mutants. The expression of the genes encoding the repressor proteins AcrR and EnvR/AcrS was also measured. Transcription of *acrR* was decreased in the *acrB* mutant and transcription of *envR* was increased in the *acrBDF* mutant (Table 2).

Inactivation of two or more RND efflux systems altered antimicrobial susceptibility.

As previously described, inactivation of *acrB* led to multi-drug hyper-susceptibility while single inactivation of either *acrD* or *acrF* did not significantly alter MICs of antibiotics, dyes and detergents compared to the wildtype strain. Re-interrogation of previously published data⁴³ from the Biolog Phenotype Microarray showed that the *acrD* and *acrF* mutants grew better than SL1344 when exposed to four β -lactams, five macrolides, and five quinolones

(Supplementary Table S2). This observation was confirmed by measuring the growth kinetics of the strains in the presence of representative AcrB substrates. However, the beneficial effect of lacking *acrD* or *acrF* was lost when *acrB* was deleted in the same strain (L1297 and L646, respectively) (Figure 1).

The antimicrobial susceptibility of the double mutant lacking AcrD and AcrF (L1395) was not significantly different from that of SL1344 (Table 3). Furthermore, except for ethidium bromide (for which the MIC value was lower) and the aminoglycosides (for which the MIC values were increased), the susceptibility of the *acrBD* (L1297), *acrBF* (L646) and the triple *acrBDF* (L1405) mutants to antibiotics, dyes and detergents was not significantly different to that of the *acrB* mutant. Surprisingly, the MICs of the aminoglycoside antibiotics, streptomycin, gentamicin and amikacin, were higher for the *acrBF* (L646) mutant and the triple *acrBDF* mutant (L1405) than for the wild-type parental strain SL1344; the MIC of tobramycin was also greater for L1405 than SL1344 (Table 3B). All *acrB* mutants (L110, L646, L1297 and L1405) were more susceptible to the efflux inhibitors Phe-arg- β -naphthylamide (PA β N) and carbonyl cyanide *m*-chlorophenyl hydrazone (CCCP) than the wild-type parental strain (SL1344).

Inactivation of two or more RND efflux systems decreased efflux activity.

We previously showed that inactivation of *acrB* led to increased accumulation of the dye Hoechst 33342.²⁰ Compared to SL1344, inactivation of *acrD* (L132) or *acrF* (L131) or inactivation of both *acrD* and *acrF* (L1395) did not significantly alter the accumulation of Hoechst 33342 (Figure 2). However, mutants with *acrB* and another *acr* gene inactivated (*acrBD* (L1297) or *acrBF* (L646)) accumulated less Hoechst 33342 than the *acrB* mutant. The *acrBDF* mutant (L1405) accumulated the highest level of Hoechst 33342 indicating the

lowest level of efflux. Accumulation of the fluoroquinolone antibiotic, norfloxacin, showed a similar pattern although the *acrB* mutant accumulated a higher concentration than the *acrBDF* mutant (Figure 3).

Inactivation of two or more RND efflux systems attenuated the ability of *Salmonella* to infect tissue culture cells.

In addition to their role in antimicrobial resistance, in many Gram-negative bacterial pathogens RND efflux pumps are required for the ability to cause infection.¹⁹ Inactivation of *acrB* has been previously shown to attenuate invasion of *S. Typhimurium* into mammalian cells growing in tissue culture.^{12, 20, 21} We now show that single gene inactivation of either *acrD* (L132) or *acrF* (L131) also significantly attenuate the virulence of *Salmonella*. Adhesion by L131 (*acrF::aph*) to human intestinal cells (INT-407) was 56.0% that of SL1344 and it only invaded at 39.1% of wild-type level. L132 (*acrD::aph*) was even more attenuated; adhesion was only 29.0% that of SL1344 and invasion was only 39.1% of the SL1344 level (Figure 4).

When two efflux pump genes were inactivated or deleted, the ability of *Salmonella* to adhere to, and invade, INT-407 cells was attenuated more than that seen in single *acr* mutants (Figure 4). Invasion of the *acrBD* double mutant was the lowest among all single and double mutants at only 0.4% of wild-type level. The adhesion of the *acrDF* mutant, L1395 (*acrD::cat, acrF::aph*), was 83.7% that of SL1344 adhesion, which is significantly greater than mutants lacking only one of these systems. Despite this, the invasion level of this mutant was only 17.8% that of the parental strain showing that these two mutations have an additive effect upon invasive ability. When all three of the efflux genes were inactivated in L1405, *Salmonella* was almost completely unable to adhere to or invade INT-407 cells (adhesion = 0.16% of wild-type level, invasion = 0.004% of wild-type level).

Discussion

The role of RND efflux pumps in multi-drug resistance and virulence makes them attractive targets for the design of efflux inhibitors. We have shown that expression of all RND efflux pump genes can be altered when single or multiple *acr* genes are inactivated. These data suggest that the bacterium can sense and respond to the levels of RND transporters and due to overlapping substrate specificity this affords resilience to the bacterium to prevent intracellular accumulation of toxic metabolites,⁴⁴⁻⁴⁷ or survival in toxic environments.

Critically, we can correlate alteration in efflux level and sensitivity to antimicrobials to the compensatory changes in efflux pump gene expression in strains lacking specific RND pumps. For example, in the Biolog phenotype microarray many of the compounds in which the *acrD* and *acrF* mutant grew better are known substrates of AcrB which is over-expressed in these mutants. Furthermore, this beneficial effect is lost when *acrB* is also inactivated. Other than to the aminoglycosides, the susceptibility of the *acrBD* and *acrBF* mutants was not significantly different to that of the mutant lacking only *acrB* and it is likely that this is because the other three RND systems are overproduced in both cases.

AcrD, which is known to transport aminoglycosides, and MdtB, which has a similar substrate profile to AcrD, are over-expressed in the *acrBF* mutant. This could explain the decreased susceptibility to the aminoglycosides seen in this mutant. The *acrBDF* mutant also had decreased susceptibility to the aminoglycosides. Expression of *mdsB* is increased in this mutant but there is currently no evidence this pump can transport aminoglycosides. Aminoglycosides enter bacterial cells by self-promoted uptake and it is possible that changes in expression of genes encoding cell envelope components, including LPS, could be responsible for this effect.⁴⁸

289 These data provide proof of principle that changes in expression of pumps in response to
290 inactivation of RND efflux pumps can alter susceptibility to clinically relevant antimicrobials.
291 We postulate the same will be true when the pump proteins themselves are inhibited and
292 recent evidence showing that the efflux inhibitors PA β N and NMP altered expression of RND
293 efflux pump genes in *E. coli* supports this.⁴⁹ Additionally, this highlights that any change in
294 the phenotype of strains with single or multiple genes inactivated should be interpreted
295 with caution as the phenotype represents, the engineered inactivation and any consequent
296 transcriptional changes.

297 The role of AcrAB-TolC in virulence of *S. Typhimurium* is well established and inactivation of
298 *acrB* causes decreased expression of genes in SPI-1, which are known to be required for
299 infection.^{12, 20, 21, 35} Nishino *et al.*, showed that inactivation of *acrD* did not confer significant
300 attenuation in the BALB/C mouse model of infection and inactivation of *acrEF* (encoding the
301 RND pump protein and the periplasmic adaptor protein) increased the host survival rate
302 with 20% of mice surviving to 21 days rather than none when infected with the wild-type
303 strain.¹² In the tissue culture model lack of either AcrD or AcrF caused a significant reduction
304 in the ability of *Salmonella* to infect INT-407 cells with the *acrD* mutant (L132) being more
305 attenuated than the *acrF* mutant (L131). There are several hypotheses to explain these data.
306 First, as inactivation of *acrB* is known to alter expression of genes found in SPI-1³⁵ it is
307 possible that inactivation of other RND pump genes also affects virulence gene expression.
308 Alternative explanations include that RND efflux pumps export substrates that are required
309 for infection or that absence of some RND efflux pumps causes damage or stress to the
310 bacterial cell membrane which compromises the ability to cause infection.

311 The effect of inactivating *acrB* plus one or two other efflux pump genes upon the ability to
312 cause infection was additive. The *acrDF* (L1395) mutant was less attenuated than either of
313 the single mutants (L131 and L132). One explanation for this is that *acrB*, *mdtB* and *mdsB*
314 are all overexpressed in this mutant and so are able to partially compensate for the
315 functions of the other two systems. The triple mutant lacking AcrB, AcrD and AcrF was
316 unable to adhere to, or invade the INT-407 cells. This could suggest that no other
317 transporter could compensate for the loss of these proteins or that inactivation of multiple
318 RND efflux pump genes causes greater changes in expression of virulence genes.

319 The role of efflux pumps in antibiotic resistance makes them targets for the design of
320 inhibitors. Due to the role of efflux pumps in virulence we also postulate that efflux
321 inhibitors will inhibit virulence as well as augment the activity of antibacterial drugs. Our
322 data show that inhibitors designed to inhibit all RND efflux systems will have a greater anti-
323 virulence effect on the organism.

324 The compensatory expression of efflux pump genes was associated with changes in
325 regulatory gene expression. We hypothesise that the bacterial cell is attempting to increase
326 expression of the inactivated/deleted genes by increasing expression of factors known to
327 regulate expression of RND efflux pump genes such as *ramA*, *marA*, *soxS* and *rob*.^{22-24, 26, 27,}

328 ^{43, 50} Our data suggest that these regulators are involved in the modulation of RND efflux
329 pump expression in the absence of homologous systems. Expression of *ramA* was increased
330 when *acrB* was inactivated,⁵¹ however, expression of *soxS* was increased when two or more
331 *acr* genes were inactivated. SoxS is also a transcription factor of the AraC/XylS family
332 involved in regulating the response to oxidative stress and genes including *acrAB* and *micF*.⁵²
333 Increased expression of *soxS* could suggest that lack of efflux by Acr pump proteins leads to

334 accumulation of toxic metabolites, as proposed by Rosner and Martin when *E. coli tolC* is
335 inactivated.^{44, 47}

336 The critical role of RND systems in both antimicrobial resistance and virulence of pathogenic
337 bacteria makes them attractive targets for the design of inhibitors. These molecules could
338 be used to re-sensitise the bacterium to antimicrobials whilst simultaneously attenuating
339 virulence of the infecting organism. Critically, our data indicate that care should be taken
340 when developing efflux pump inhibitors against the RND pumps to determine which pumps
341 are inhibited and to understand the effect of this on expression of homologous systems. In
342 terms of attenuating virulence, the effect of inhibition was additive so inhibition of multiple
343 pumps is a good strategy. However, the benefit of this strategy on increasing susceptibility
344 to antimicrobials may be more complex and the impact of this will depend on which drugs
345 are used to treat infections by a particular pathogen. For some antimicrobials, an inhibitor
346 with activity against multiple pumps will have a greater impact on susceptibility but, an
347 unintended consequence of this may be decreased susceptibility to other drugs, such as the
348 aminoglycosides.

349

350

351

Acknowledgements

352 This work was funded by an MRC Programme grant (G0501415) to LJP and a research grant
353 (GA2011-04R) from BSAC to JMAB. We thank Mark Webber, Michelle Buckner, Lee Rosner
354 and Bill Shafer for reading and constructive criticism of this manuscript.

Transparency declaration

356 The authors declare no conflict of interest.

357

Figure legends

Figure 1. Growth of efflux pump mutants in the presence of AcrB substrates. Growth of the efflux in the presence of (a) ciprofloxacin and (b) doxycycline at a concentration of 0.25 x the MIC for wild-type. Data presented is the mean of three biological replicates.

Figure 2. Hoechst 33342 Accumulation in (A) single efflux mutants and (B) double and triple efflux mutants. The data presented are the mean of three separate experiments presented as fold change compared to SL1344 at the end point of the assay \pm standard error of the mean. Student's *t*-tests were performed to compare the Hoechst 33342 accumulation of each strain to that of SL1344 and those returning *P* values of less than 0.05 are indicated by *.

Figure 3. Accumulation of norfloxacin. Data shown is the mean of three biological replicates \pm the standard error of the mean.

Figure 4. Adhesion (A) and Invasion (B) of strains lacking AcrB, AcrD, AcrF and combinations thereof to invade INT-407 cells *in vitro*. Data shown is the mean of at least three independent experiments. Student's *t*-tests were used to compare values for each strain with that of the wild-type, SL1344. *P* values of ≤ 0.05 were considered significant and are indicated by asterisks.

376 Figure 1.







Figure 2. Hoechst 33342 accumulation




382 **Figure 3. Accumulation of norfloxacin**



383

384

Figure 4. Adhesion (A) and Invasion (B) of strains lacking AcrB, AcrD, AcrF and combinations thereof to invade INT-407 cells *in vitro*



389 **Table 1. Percentage nucleotide identity and amino acid similarity between RND efflux**
390 **pump genes and proteins of *Salmonella***

	<i>acrB</i> AcrB	/	<i>acrD</i> AcrD	/	<i>acrF</i> AcrF	/	<i>mdtB</i> MdtB	/	<i>mdtC</i> MdtC	/	<i>mdsB</i> MdsB	/
<i>acrB</i> /AcrB	-		70 / 79.1		74 / 90.4		55 / 46.7		55 / 48.6		59 / 63.4	
<i>acrD</i> /AcrD	-		-		68 / 78.2		54 / 49.1		54 / 48.8		59 / 61.3	
<i>acrF</i> /AcrF	-		-		-		54 / 47.4		53 / 48.5		57 / 63.1	
<i>mdtB</i> /MdtB	-		-		-		-		62 / 66.0		56 / 49.0	
<i>mdtC</i> /MdtC	-		-		-		-		-		56 / 49.2	
<i>mdsB</i> /MdsB	-		-		-		-		-		-	

391

392

Table 2. Expression of RND efflux pump genes and regulators thereof, quantified by real time RT-PCR

		Fold change in gene expression										
		RND efflux pump genes					Known regulators of efflux					
Strain		<i>acrB</i>	<i>acrD</i>	<i>acrF</i>	<i>mdtB</i>	<i>mdsB</i>	<i>marA</i>	<i>ramA</i>	<i>rob</i>	<i>soxS</i>	<i>acrR</i>	<i>envR</i>
SL1344	WT	1.0	1.00	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
L644	Δ <i>acrB</i>	-	13.4	16.0	1.0	<i>0.6</i>	2.0	2.8	1.4	2.6	0.4	1.2
L132	<i>acrD::aph</i>	1.8	-	1.0	0.9	<i>0.5</i>	1.2	0.8	1.4	1.3	0.3	0.6
L131	<i>acrF::aph</i>	3.4	3.4	-	1.8	<i>0.7</i>	0.9	1.0	1.3	1.4	0.9	1.5
L646	Δ <i>acrF</i> <i>acrB::aph</i>	-	4.6		4.5	3.0	2.5	1.6	1.2	6.3	0.4	0.8
L1297	Δ <i>acrB</i> <i>acrD::aph</i>	-	-	2.4	4.6	2.1	1.2	1.4	1.1	1.8	2.2	1.2
L1395	Δ <i>acrF</i> <i>acrD::aph</i>	3.8	-	-	6.0	5.9	1.0	1.3	1.4	2.2	0.4	1.0
L1405	Δ <i>acrB</i> Δ <i>acrF</i> <i>acrD::aph</i>	-	-	-	1.1	3.2	1.6	1.9	1.0	5.0	0.5	4.5

Bold text indicated statistically significant ($P \leq 0.05$) increased expression. Italic text indicates statistically significant decreased expression.

Table 3. Antimicrobial susceptibility of SL1344 and RND mutants thereof.

A. Minimum Inhibitory Concentration of antimicrobials to SL1344 and mutants thereof.

	Genotype	MIC (mg/L)													
		Amp	Chl	Cip	Tet	Nal	EtBr	Fus	Ami	Gent	Hyg	Strep	Tob	PAβN	CCCP
SL1344	Wild-type	2	4	0.015	1	4	>256	>256	4	0.5	32	8	2	>1024	64
L110	<i>acrB::aph</i>	0.25	1	<0.008	0.5	1	64	8	4	1	32	8	1	64	32
L644	Δ <i>acrB</i>	0.25	1	<0.008	0.5	1	64	8	4	1	32	8	1	64	32
L131	<i>acrF::aph</i>	2	4	0.015	2	4	>256	>256	4	1	32	16	2	>1024	64
L132	<i>acrD::aph</i>	2	4	0.015	1	4	>256	>256	4	1	32	16	2	>1024	64
L646	Δ <i>acrF</i> <i>acrB::aph</i>	2	1	<0.008	1	1	16	4	8	2	64	32	4	64	32
L1297	Δ <i>acrB</i> <i>acrD::aph</i>	0.25	1	<0.008	0.5	1	64	8	2	0.5	32	8	1	64	32
L1395	<i>acrF::cat</i> <i>acrD::aph</i>	2	16	0.015	2	4	>256	>256	4	1	64	16	2	>1024	64
L1405	Δ <i>acrB</i> Δ <i>acrF</i> <i>acrD::aph</i>	0.12	1	<0.008	0.5	1	16	8	16	2	64	32	8	64	32

B. Fold change in MIC compared to $\Delta acrB$ (L644)

	Genotype	Amp	Chl	Cip	Tet	Nal	EtBr	Fus	Ami	Gent	Hyg	Strep	Tob	PA β N	CCCP
L646	<i>$\Delta acrF$ $acrB::aph$</i>	4			2		-4	-2	2	2	2	4	4		
L1297	<i>$\Delta acrB$ $acrD::aph$</i>								-2	-2	-2				
L1405	<i>$\Delta acrB$ $\Delta acrF$ $acrD::aph$</i>	-2					-4		4	2	2	4	8		

Amp, ampicillin; Chl, chloramphenicol; Tet, tetracycline; Nal, nalidixic acid; EtBr, Ethidium bromide; Fus, Fusidic Acid; Ami, Amikacin, Gent, Gentamicin; Hyg, Hygromycin; Strep, Streptomycin; Tob, Tobramycin; PA β N, Phe-Arg β -naphthylamide dihydrochloride; CCCP, Carbonyl cyanide 3-chlorophenylhydrazone. The *aph* gene used is *aph(3')*-1 which gives resistance to kanamycin, neomycin and paromycin.

Table 3A: Bold red font indicates an increase in the MIC of the same compound compared to SL1344. Italic blue font indicates a decrease in the MIC of the same compound compared to SL1344. Table 3B: Bold red font indicates an increase in MIC compared to the same compound for L644. Italic blue font indicates a decrease in the MIC of the same compound for $\Delta acrB$. No value indicates no difference in MIC values.

References

1. Piddock LJV. Clinically relevant chromosomally encoded multidrug resistance efflux pumps in bacteria. *Clin Microbiol Rev* 2006; **19**: 382-402.
2. Hinchliffe P, Symmons MF, Hughes C *et al.* Structure and Operation of Bacterial Tripartite Pumps. *Annu Rev Microbiol* 2013; **67**: 221-42.
3. Ruggerone P, Murakami S, Pos KM *et al.* RND Efflux Pumps: Structural Information Translated into Function and Inhibition Mechanisms. *Curr Top Med Chem* 2013; **13**: 3079-100.
4. Zhang Y, Xiao M, Horiyama T *et al.* The Multidrug Efflux Pump MdtEF Protects against Nitrosative Damage during the Anaerobic Respiration in *Escherichia coli*. *J Biol Chem* 2011; **286**: 26576-84.
5. Unemo M, Shafer WM. Antibiotic resistance in *Neisseria gonorrhoeae*: origin, evolution, and lessons learned for the future. *Ann NY Acad Sci* 2011; **1230**: E19-E28.
6. Pumbwe L, Piddock LJV. Two Efflux Systems Expressed Simultaneously in Multidrug-Resistant *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother* 2000; **44**: 2861-4.
7. Yoon E-J, Courvalin P, Grillot-Courvalin C. RND-Type Efflux Pumps in Multidrug-Resistant Clinical Isolates of *Acinetobacter baumannii*: Major Role for AdeABC Overexpression and AdeRS Mutations. *Antimicrob Agents Chemother* 2013; **57**: 2989-95.
8. Piddock LJV, Whale K, Wise R. Quinolone resistance in *salmonella*: clinical experience. *Lancet* 1990; **335**: 1459.
9. Nishino K, Yamada J, Hirakawa H *et al.* Roles of TolC-dependent multidrug transporters of *Escherichia coli* in resistance to beta-lactams. *Antimicrob Agents Chemother* 2003; **47**: 3030-3.
10. Eicher T, Cha H-j, Seeger MA *et al.* Transport of drugs by the multidrug transporter AcrB involves an access and a deep binding pocket that are separated by a switch-loop. *Proc Natl Acad Sci* 2012.
11. Nakashima R, Sakurai K, Yamasaki S *et al.* Structures of the multidrug exporter AcrB reveal a proximal multisite drug-binding pocket. *Nature* 2011; **480**: 565-9.
12. Nishino K, Latifi T, Groisman EA. Virulence and drug resistance roles of multidrug efflux systems of *Salmonella enterica* serovar Typhimurium. *Mol Microbiol* 2006; **59**: 126-41.
13. Kobayashi K, Tsukagoshi N, Aono R. Suppression of hypersensitivity of *Escherichia coli* *acrB* mutant to organic solvents by integrational activation of the *acrEF* operon with the IS1 or IS2 element. *J Bacteriol* 2001; **183**: 2646-53.
14. Nishino K, Nikaido E, Yamaguchi A. Regulation of multidrug efflux systems involved in multidrug and metal resistance of *Salmonella enterica* serovar Typhimurium. *J Bacteriol* 2007; **189**: 9066-75.
15. Rosenberg EY, Ma D, Nikaido H. AcrD of *Escherichia coli* is an aminoglycoside efflux pump. *J Bacteriol* 2000; **182**: 1754-6.
16. Aires JR, Nikaido H. Aminoglycosides are captured from both periplasm and cytoplasm by the AcrD multidrug efflux transporter of *Escherichia coli*. *J Bacteriol* 2005; **187**: 1923-9.
17. Nishino K, Nikaido E, Yamaguchi A. Regulation and physiological function of multidrug efflux pumps in *Escherichia coli* and *Salmonella*. *Biochimica et Biophysica Acta (BBA) - Proteins & Proteomics* 2009; **1794**: 834-43.
18. Horiyama T, Yamaguchi A, Nishino K. TolC dependency of multidrug efflux systems in *Salmonella enterica* serovar Typhimurium. *J Antimicrob Chemoth* 2010; **65**: 1372-6.
19. Piddock LJ. Multidrug-resistance efflux pumps - not just for resistance. *Nat Rev Microbiol* 2006; **4**: 629-36.
20. Blair JMA, La Ragione RM, Woodward MJ *et al.* Periplasmic adaptor protein AcrA has a distinct role in the antibiotic resistance and virulence of *Salmonella enterica* serovar Typhimurium. *J Antimicrob Chemoth* 2009; **64**: 965-72.
21. Buckley AM, Webber MA, Cooles S *et al.* The AcrAB-TolC efflux system of *Salmonella enterica* serovar Typhimurium plays a role in pathogenesis. *Cell Microbiol* 2006; **8**: 847-56.

22. Bailey AM, Ivens A, Kingsley R *et al.* RamA, a member of the AraC/XylS family, influences both virulence and efflux in *Salmonella enterica* serovar Typhimurium. *J Bacteriol* 2010; **192**: 1607-16.
23. Barbosa TM, Levy SB. Differential Expression of over 60 Chromosomal Genes in *Escherichia coli* by Constitutive Expression of MarA. *J Bacteriol* 2000; **182**: 3467-74.
24. Tanaka T, Horii T, Shibayama K *et al.* RobA-induced multiple antibiotic resistance largely depends on the activation of the AcrAB efflux. *Microbiol Immunol* 1997; **41**: 697-702.
25. Zhang A, Rosner JL, Martin RG. Transcriptional activation by MarA, SoxS and Rob of two *tolC* promoters using one binding site: a complex promoter configuration for *tolC* in *Escherichia coli*. *Mol Microbiol* 2008; **69**: 1450-5.
26. White DG, Goldman JD, Demple B *et al.* Role of the *acrAB* locus in organic solvent tolerance mediated by expression of *marA*, *soxS*, or *robA* in *Escherichia coli*. *J Bacteriol* 1997; **179**: 6122-6.
27. Hachler H, Cohen SP, Levy SB. *marA*, a regulated locus which controls expression of chromosomal multiple antibiotic resistance in *Escherichia coli*. *J Bacteriol* 1991; **173**: 5532-8.
28. Nishino K, Yamasaki S, Hayashi-Nishino M *et al.* Effect of NlpE overproduction on multidrug resistance in *Escherichia coli*. *Antimicrob Agents Chemother* 2010; **54**: 2239-43.
29. De Wulf P, McGuire AM, Liu X *et al.* Genome-wide Profiling of Promoter Recognition by the Two-component Response Regulator CpxR-P in *Escherichia coli*. *J Biol Chem* 2002; **277**: 26652-61.
30. Nishino K, Yamaguchi A. Role of histone-like protein H-NS in multidrug resistance of *Escherichia coli*. *J Bacteriol* 2004; **186**: 1423-9.
31. Hirakawa H, Takumi-Kobayashi A, Theisen U *et al.* AcrS/EnvR represses expression of the *acrAB* multidrug efflux genes in *Escherichia coli*. *J Bacteriol* 2008; **190**: 6276-9.
32. Eaves DJ, Ricci V, Piddock LJV. Expression of *acrB*, *acrF*, *acrD*, *marA*, and *soxS* in *Salmonella enterica* serovar Typhimurium: role in multiple antibiotic resistance. *Antimicrob Agents Chemother* 2004; **48**: 1145-50.
33. Wray C, Sojka W. Experimental *Salmonella typhimurium* infection in calves. *Res Vet Sci* 1978; **25**: 139-43.
34. Datsenko KA, Wanner BL. One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proc Natl Acad Sci* 2000; **97**: 6640-5.
35. Webber MA, Bailey AM, Blair JMA *et al.* The global consequence of disruption of the AcrAB-TolC efflux pump in *Salmonella enterica* includes reduced expression of SPI-1 and other attributes required to infect the host. *J Bacteriol* 2009; **191**: 4276-85.
36. Pfaffl MW. A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res* 2001; **29**: e45.
37. Andrews JM. Determination of minimum inhibitory concentrations. *J Antimicrob Chemother* 2001; **48**: 5-16.
38. Coldham NG, Webber M, Woodward MJ *et al.* A 96-well plate fluorescence assay for assessment of cellular permeability and active efflux in *Salmonella enterica* serovar Typhimurium and *Escherichia coli*. *J Antimicrob Chemother* 2010; **65**: 1655-63.
39. Mortimer PGS, Piddock LJV. A comparison of methods used for measuring the accumulation of quinolones by Enterobacteriaceae, *Pseudomonas aeruginosa* and *Staphylococcus aureus*. *J Antimicrob Chemother* 1991; **28**: 639-53.
40. Su C-C, Li M, Gu R *et al.* Conformation of the AcrB multidrug efflux pump in mutants of the putative proton relay pathway. *J Bacteriol* 2006; **188**: 7290-6.
41. Takatsuka Y, Nikaido H. Threonine-978 in the transmembrane segment of the multidrug efflux pump AcrB of *Escherichia coli* is crucial for drug transport as a probable component of the proton relay network. *J Bacteriol* 2006; **188**: 7284-9.
42. Bohnert JA, Schuster S, Seeger MA *et al.* Site-directed mutagenesis reveals putative substrate binding residues in the *Escherichia coli* RND efflux pump AcrB. *J Bacteriol* 2008; **190**: 8225-9.

43. Bailey AM, Paulsen IT, Piddock LJV. RamA confers multidrug-resistance in *Salmonella enterica* via increased expression of *acrB*, which is inhibited by chlorpromazine. *Antimicrob Agents Chemother* 2008; **52**: 3604-11.
44. Rosner JL, Martin RG. An Excretory Function for the Escherichia coli Outer Membrane Pore TolC: Upregulation of *marA* and *soxS* Transcription and Rob Activity Due to Metabolites Accumulated in *tolC* Mutants. *J Bacteriol* 2009; **191**: 5283-92.
45. Rosner JL, Martin RG. Reduction of Cellular Stress by TolC-Dependent Efflux Pumps in *Escherichia coli* Indicated by BaeSR and CpxARP Activation of *spy* in Efflux Mutants. *J Bacteriol* 2013; **195**: 1042-50.
46. Biot FV, Lopez MM, Poyot T *et al.* Interplay between Three RND Efflux Pumps in Doxycycline-Selected Strains of *Burkholderia thailandensis*. *PLOS One* 2013; **8**: e84068.
47. Helling RB, Janes BK, Kimball H *et al.* Toxic Waste Disposal in *Escherichia coli*. *J Bacteriol* 2002; **184**: 3699-703.
48. Hancock R, Farmer SW, Li Z *et al.* Interaction of aminoglycosides with the outer membranes and purified lipopolysaccharide and OmpF porin of *Escherichia coli*. *Antimicrobial agents and chemotherapy* 1991; **35**: 1309-14.
49. Barrero MAO, Pietralonga PAG, Schwarz DGG *et al.* Effect of the inhibitors phenylalanine arginyl beta-naphthylamide (PABN) and 1-(1-naphthylmethyl)-piperazine (NMP) on expression of genes in multidrug efflux systems of *Escherichia coli* isolates from bovine mastitis. *Res Vet Sci* 2014; **In Press**.
50. Pomposiello PJ, Bennik MHJ, Demple B. Genome-Wide Transcriptional Profiling of the *Escherichia coli* Responses to Superoxide Stress and Sodium Salicylate. *J Bacteriol* 2001; **183**: 3890-902.
51. Lawler AJ, Ricci V, Busby SJW *et al.* Genetic inactivation of *acrAB* or inhibition of efflux induces expression of *ramA*. *J Antimicrob Chemoth* 2013; **68**: 1551-7.
52. Pomposiello PJ, Demple B. Identification of SoxS-regulated genes in *Salmonella enterica* serovar Typhimurium. *J Bacteriol* 2000; **182**: 23-9.