

# Prevalence and predictors of externalizing behavior in young adult survivors of pediatric traumatic brain injury

Ryan, Nicholas P.; Hughes, Nathan; Godfrey, Celia; Rosema, Stefanie; Catroppa, Cathy; Anderson, Vicki A.

DOI:

[10.1097/HTR.000000000000123](https://doi.org/10.1097/HTR.000000000000123)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Ryan, NP, Hughes, N, Godfrey, C, Rosema, S, Catroppa, C & Anderson, VA 2015, 'Prevalence and predictors of externalizing behavior in young adult survivors of pediatric traumatic brain injury', *Journal of Head Trauma Rehabilitation*, vol. 30, no. 2, pp. 75-85. <https://doi.org/10.1097/HTR.000000000000123>

[Link to publication on Research at Birmingham portal](#)

## Publisher Rights Statement:

Checked for eligibility 31/03/2016. Ryan, Nicholas P., et al. "Prevalence and predictors of externalizing behavior in young adult survivors of pediatric traumatic brain injury." *The Journal of head trauma rehabilitation* 30.2 (2015): 75-85. Published version can be found at doi: 10.1097/HTR.000000000000123

## General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

**Prevalence and predictors of externalising behavior in young adult survivors of pediatric traumatic brain injury.**

Nicholas P. Ryan, BA(Hons)<sup>1,2</sup>

Nathan Hughes, PhD<sup>1,2,3</sup>,

Celia Godfrey, DPsych<sup>1,2</sup>,

Stefanie Rosema, PhD<sup>1,4</sup>,

Cathy Catroppa, PhD<sup>1,2,4</sup>,

Vicki A. Anderson, PhD<sup>1,2,4</sup>

1. Murdoch Childrens Research Institute, Melbourne, Australia

2. University of Melbourne, Australia

3. University of Birmingham, UK

4. Royal Children's Hospital, Melbourne, Australia

Corresponding author:

Nicholas Ryan. Murdoch Childrens Research Institute, Level 4 West, The Royal Children's Hospital Melbourne, Flemington Road, Parkville, Melbourne, Victoria, Australia 3052.

Email: [nicholas.ryan@mcri.edu.au](mailto:nicholas.ryan@mcri.edu.au)

Conflicts of Interest and Source of Funding:

No competing financial interest exists. This work was funded through research grants awarded by the National Health and Medical Research Council, the Victorian Neurotrauma Institute and the Murdoch Childrens Research Institute, Clinical Sciences and supported by the Victorian Government's Operational Infrastructure Scheme.

**Abstract**

*Objectives:* To investigate rates of clinically significant externalizing behavior in young adult survivors of pediatric TBI, and evaluate the contribution of pre- and post-injury risk and resilience factors to externalizing behaviour outcomes 16 years after injury.

*Setting:* Melbourne, Australia

*Participants:* Fifty-five young adults ( $M$  age = 23.85; Injury Age: 1.0 - 12 years) admitted to an emergency department following TBI between 1993 and 1997.

*Design:* Longitudinal prospective study with data collected at the acute, 10-year and 16-year post-injury time points.

*Main Measures:* Severity of TBI, adaptive functioning, family functioning, full scale IQ, executive functioning, social communication, and symptoms of externalizing behavior (EB).

*Results:* One of every four young people with a history of pediatric TBI demonstrated clinical or sub-threshold levels of EB in young adulthood. More frequent externalizing behavior was associated with poorer pre-injury adaptive functioning, reduced full scale IQ and more frequent pragmatic communication difficulty.

*Conclusion:* Pediatric TBI is associated with elevated risk for externalizing disorders in the transition to adulthood. Results underscore the need for screening and assessment of TBI among young offenders, and suggest that early and long-term targeted interventions may be required to address risk factors for EB in children and young people with TBI.

**Keywords:**

Traumatic brain injury, externalising behavior (EB), crime, antisocial behavior, young adulthood

## Introduction

A growing number of research studies, undertaken within various national contexts, consistently demonstrate a disproportionately high prevalence of traumatic brain injury (TBI) amongst youth and adult offending populations. Studies estimate that between 12% and 24% of the general population have experienced a head injury resulting in loss of consciousness<sup>1-4</sup>. This compares to equivalent reported rates of between 32% and 46% amongst young people in custody<sup>5-8</sup>, with several studies suggesting over 60% of adult prisoners have experienced a head injury with loss of consciousness<sup>9-11</sup>.

Studies of adult offending populations seldom indicate whether TBI was sustained in childhood or adulthood. Nonetheless research has repeatedly demonstrated that ‘life-course persistent’ offending typically begins in childhood and is commonly associated with ‘neurocognitive impairments’<sup>12,13</sup>. This would suggest that children experiencing pediatric TBI are at elevated risk for persistent offending beyond adolescence, and that their experiences run counter to the norm of desistence from criminality during young adulthood<sup>14,15</sup>. Despite this apparent association between TBI and persistent adult offending, to date there is insufficient research regarding such behavior in the transition into young adulthood amongst those who experience pediatric TBI.

There is considerable evidence that pediatric TBI contributes to impairments in executive functions, cognitive skills and EB (such as aggression, hyperactivity, bullying and defiance) that are commonly identified as risk factors for antisocial behavior and criminality. TBI commonly involves pathology to anterior brain regions implicated in executive functioning, as well as traumatic axonal injury that may contribute to deficits in social cognition, attention, learning difficulties and pragmatic communication<sup>16-18</sup>. Impairments in executive functioning have been detected soon after injury and appear to persist or even worsen with time since injury<sup>19-21</sup>, likely reflecting a failure to develop and acquire skills at

an age appropriate rate<sup>22</sup>. It may be that such deficits contribute to antisocial or offending behavior through decreased inhibition, poor anticipation of the consequences of specific actions, or an inability to recognise when certain behavior is inappropriate in a given social context<sup>23,24</sup>.

Reduced cognitive empathy implies an inability to see the consequences of antisocial behavior or to empathize with victims<sup>25</sup>. These deficits are a common consequence of pediatric TBI, particularly among children with frontal injuries<sup>26</sup>. In addition, children and adolescents with TBI experience persisting difficulty interpreting non-verbal emotion cues from facial expressions and prosody, as well as impairments in social or pragmatic communication<sup>22,27</sup>. These deficits are likely to contribute to reduced interpersonal effectiveness, which may in turn lead to frustration and distress, reflected in EBs that are shown to persist or even worsen with time since injury<sup>28-31</sup>.

Though persisting injury-related neurocognitive impairments may elevate risk for antisocial or offending behavior after TBI, criminological research suggests that such injuries may influence offending via exposure to social and environmental experiences that may exacerbate the neurological consequences of brain injury<sup>32-34</sup>. For example, studies have linked permissive or authoritarian parenting styles, poor parental mental health and lower socio-economic status to long-term behavioral problems following TBI<sup>35-38</sup>. Persistent problems in academic performance, including specific difficulties in reading, spelling and arithmetic are commonly reported after TBI<sup>39-42</sup>, and are likely to have a cumulative impact on educational opportunities, leading to challenges in engaging in later stages of education, particularly in the transition to secondary school<sup>43-46</sup>.

The Heuristic Model of Social Competence (HMSC)<sup>47</sup> provides a useful framework for conceptualizing how such a range of injury and non-injury related risk and resilience factors may contribute to variability in social functioning after TBI, and can therefore be

usefully applied to the study of crime and antisocial behavior. Injury related factors, including injury severity and pathology location, are conceptualized as factors that increase the likelihood of deficits in social cognition and atypical social interaction, while environmental factors may further heighten risk or represent sources of resilience that buffer against the negative long term consequences of TBI. In addition, the model posits reciprocal interactions between various components of social information processing (SIP; cognitive-executive functions, social cognition, social problem solving), social interaction, and social adjustment, such that deficits in any one component in SIP may contribute to impaired social interaction and poor social adjustment, including EB.

In order to address limited understanding of the links between criminality and pediatric TBI, further research is required to investigate factors that may contribute to maladaptive or antisocial behavior among young adult survivors of pediatric TBI. In conceptualizing EB as a marker of risk for delinquency and/or criminal behaviour, it may be that injury-related neurocognitive impairments and environmental factors confer risk for criminality via their influence on EB.

The objectives of the present investigation were to (1) examine the prevalence of clinically significant EB problems in young adult survivors of pediatric TBI and (2) evaluate the respective contributions of a variety of injury and non-injury related risk and resilience factors at various time points to variability in EB outcomes 16 years after pediatric TBI. More specifically, guided by the HMSC model<sup>47</sup>, we aimed to examine relationships between long-term externalizing symptoms and a variety of individual and environmental factors including: pre-injury individual and family functioning; injury severity and acute intellectual functioning; executive functioning; and social cognitive and affective functions, including social perception and pragmatic communication.

We predicted that relative to population expectations, a significantly greater proportion of young adults with TBI would show clinically significant EB. Furthermore, we hypothesized that more frequent EBs would be related to (a) greater injury severity; (b) poorer pre-injury adaptive and family functioning; (c) poorer acute intellectual functioning; (d) poorer executive function at 10-years post injury; and (e) reduced emotional perception and more frequent pragmatic communication difficulty at 16-years post-TBI.

## Method

### Participants

This longitudinal study followed up a sample originally recruited from consecutive admissions to the emergency department at The Royal Children's Hospital, Melbourne, Australia (RCH), between 1993 and 1997 for traumatic brain injury<sup>48</sup>. Inclusion criteria for the original study were: (1) age at injury 1.0 to 12.0 years; (2) documented evidence of TBI, including a period of altered consciousness; (3) sufficiently detailed medical records for diagnosis of injury severity. Exclusion criteria were: penetrating or non-accidental head injury; history of previous closed head injury; or pre-existing physical, neurological, psychiatric or developmental disorder.

During the initial recruitment period, 172 children aged under 12 years were admitted to hospital with a diagnosis of TBI and participated in the initial data collection. At 16 years post-TBI, 66 participants could not be located, and 38 declined to participate (not interested / too busy), 2 were deceased and 11 had incomplete datasets. Thus for the current paper 55 young adults (m age = 23.82 years; range = 16.25 - 30.58 years; SD = 3.85) from the original TBI sample yielded data across the acute, 10-year and 16-year time points and were included in analyses. Participating and non-participating samples were compared on demographic and injury characteristics, including socio-economic status, gender, age at injury and the

length of period of altered consciousness, to examine potential biases in the 16-year follow-up sample, with no significant differences identified ( $p>.05$ ).

## Measures

Details of the child's medical and developmental history, and family demographic information were collected at study enrolment. Severity groups were derived from a combination of measures, including period of altered consciousness on the Glasgow Coma Scale (GCS)<sup>49</sup>, and presence of radiological and neurological abnormalities. This resulted in the following groups: (i) mild TBI ( $n = 15$ ): GCS on admission 13–15, no evidence of mass lesion on CT/MRI scans, and no neurologic deficits; (ii) moderate TBI ( $n = 29$ ): GCS on admission 9–12, and/or mass lesion or other evidence of specific injury on CT/MRI, and/or neurological impairment; and (iii) severe TBI ( $n = 11$ ): GCS on admission 3–8, and mass lesion or other evidence of specific injury on CT/MRI, and/or neurological impairment.

### *Pre-injury.*

The *Vineland Adaptive Behavior Scale* (VABS)<sup>50</sup> provides parent report of a child's level of adaptive function and was collected at time one to represent the child's pre-injury functioning. The present study utilised the Total Adaptive Behavior score ( $M = 100$ ,  $SD = 15$ ), and the Daily Living Skills and Socialization indexes as measures of pre-injury function.

Pre-injury family environment was measured using the parent-report *Family Functioning Questionnaire* (FFQ)<sup>51</sup>. Each item was rated on a 6-point scale where 1 = totally agree to 6 = totally disagree. Three factors are derived from the measure: Conflict, Intimacy and Parenting Style, with higher scores reflecting more of that characteristic. The Intimacy factor was utilised for statistical analyses because it represents a measure of family cohesion<sup>52</sup>.



*Acute post-injury.*

The IQ assessment employed at the acute time point varied depending on the child's age. Thus *Bayley Scales of Infant Development* children aged <30 months; *Wechsler Preschool and Primary Scale of Intelligence Revised*<sup>54</sup> for children aged 30 months to 6.5 years; and *Wechsler Intelligence Scale for Children - Third Edition*<sup>55</sup> children aged >6.5 years were all used. Full scale IQ scores (FSIQ), were used in analyses (M = 100, SD = 15).

*Daniel's Scale of Occupational Prestige*<sup>56</sup> was used as a rating of family socio-economic status (SES) at the acute time-point. Ratings are made on a seven-point scale where a higher score denotes lower SES.

*10 years post injury.*

Based on previous findings from the longitudinal study two measures of executive functioning were used to explore its role in prediction of problem behaviour<sup>57</sup>. The *Behavior Rating Inventory of Executive Function* (BRIEF)<sup>58</sup>, Metacognition and Behavioral Regulation index scores, and the General Executive Composite Score (M = 50, SD = 10) were calculated on the basis of parent or close other ratings at the 10 year time-point. Higher scores represent greater dysfunction, and scores >65 indicate functioning at a level of clinical concern.

The *20 Questions Task* from the DKEFS test (20Q)<sup>59</sup> was used as a direct assessment of abstract reasoning as it measures abstract thinking as well as problem solving and the utilization of feedback. The abstraction scaled score was used in analyses (M = 10, SD = 3).

*16 years post-injury.*

The *Adult Behavior Checklist* (ABCL)<sup>60</sup> consists of 126 behavior problem items that are evaluated by a significant other for the preceding six months. Statements are scored on a three-level rating scale ranging from not true to very true (M=50; SD=10; borderline/clinical range if score  $\geq 65$  for the syndrome scales;  $\geq 60$  for the domain) with a higher score indication of greater impairments. As the outcome measure in the present study, the domain Externalizing Behavior was used. It comprises three syndrome scales: Rule Breaking (13 items; e.g., gets drunk, in trouble with law), Aggressive Behavior (16 items; e.g., mean to others, threatens people) and Intrusive Behavior (6 items; e.g., brags, demands attention, shows off). The ABCL has been proven reliable in terms of test-retest correlations and internal consistency of scales<sup>60</sup>, and has good inter-rater reliability for most scales<sup>61</sup>.

The *Latrobe Communication Questionnaire* (LCQ)<sup>62</sup> is a 30-item subjective assessment that reflects the four domains Quantity, Quality, Relation and Manner of everyday communication. Each item has four levels of response ranging from (1) "Never or rarely" to (4) "Usually or always" with a higher score reflecting more frequent communication difficulty. Data analyses employed a total score, reflecting overall communication perceived by a significant other.

The *Advanced Clinical Solutions Social Perception subtest* (ACS)<sup>63</sup> measures skills associated with the comprehension of social communication. It consists of three emotion perception tests yielding four scores: Affect Naming, Prosody-Faces, Prosody-Pairs, and, collectively, the Emotion Perception Total score. Age-adjusted scaled scores ( $\underline{M}=10$ ,  $\underline{SD}=3$ ) for each of these test scores were employed in analyses.

In addition, respondents were asked whether they had 'received intervention of any kind (e.g. speech and language, motor, cognitive)'. Those who stated that they had received an intervention were asked to specify the intervention through an open-ended response.

## Procedure

The current study was approved by the Human Research Ethics Committee of RCH, Melbourne, Australia. Children were enrolled in the study during their initial hospital admission, and were evaluated at various time points: 0-3 (acute), 6, 12, and 30 months, and 5, 10 and 16-years post injury. At each wave of data collection, young people and families enrolled in the original study were sent tracing letters that included a detailed description of the study, and were asked to provide written informed consent, in keeping with hospital ethics procedures. Neuropsychological assessments and questionnaires were administered at each time point by a qualified child psychologist over a two hour period.

## Data analysis

All data were entered into SPSS statistical software (Version 21.0; SPSS, Inc., Chicago, IL) and screened for violations of normality. An alpha level of  $p < 0.05$  was used to indicate significance, and effect sizes were calculated using Cohen's  $d$ .

The calculation of individual impairment ratings was based on the ABCL Externalizing Behavior composite scale, and Chi-square tests were employed to examine the proportion young adults in each severity group demonstrating impairment at 16-years post injury. For the broadband scales, scores  $> 63$  are considered clinically significant and scores of 60-63 are in the borderline range for clinical significance.

Predictors of 16-year behavioral outcomes were examined using a series of regression analyses as follows;

(1). Preliminary univariate regressions were employed to examine relations between EB and all independent variables. Variables that were unrelated to EB at this step were excluded from subsequent analyses.

(2). Four separate multivariate adjusted regression models were employed to

investigate relationships between EB and variables related to (a) pre-injury adaptive functioning; (b) injury-related factors/acute intellectual functioning; (c) executive functioning/interventions at 10-years post-TBI; and (d) social cognition and communication at 16-years post injury.

(3). Variables that remained statistically significant in each of the adjusted models were entered into the final adjusted multivariate regression model.

## Results

### 3.1. Demographic and injury characteristics.

There were no significant differences across severity groups with respect to age at 16-year assessment, age-at-injury, pre-injury adaptive abilities, FSIQ, SES or family function (Table 1). Groups differed on gender ( $\chi^2(2, 55) = 7.51, p = .023$ ), such that there were a significantly greater proportion of males in the severe TBI group than the mild and moderate TBI groups. As expected, all severity groups differed for GCS-24 hours,  $F(2,48) = 21.92, p < .001$ .

**Table 1 about here**

### 3.2. Externalizing symptoms at 16-years post-TBI.

Table 2 displays the total number and proportion of TBI participants found to have clinical or sub-threshold levels of externalizing symptoms. Pearson Chi Square analyses revealed no significant association between externalizing symptoms and injury severity,  $\chi^2(2, 55) = .20, p = .91$ .

**Table 2 about here**

### 3.3. Predictors of ABCL Externalizing: Pre-injury adaptive functioning.

**Unadjusted model.** Preliminary univariate regression analyses revealed significant relationships between ABCL Externalizing and pre-injury Vineland Adaptive Functioning ( $p=.004$ ), including the Daily Living Skills ( $p=.040$ ), and Socialization ( $p=.007$ ). There was no significant relations between ABCL Externalizing and Vineland Communication ( $p=.106$ ), Family SES ( $p=.724$ ) or Family Intimacy ( $p=.188$ ).

**Adjusted model.** The multivariate adjusted model was moderately significant  $F(3,45) = 3.27$ ,  $p = .03$ , however due to high colinearity between the independent variables, there were no significant individual pre-injury predictors of ABCL Externalizing (Table 3).

### Table 3 about here

#### 3.4. Predictors of ABCL Externalizing: Injury-related factors and acute intellectual functioning.

Univariate regression analysis revealed a significant relation between ABCL Externalizing and FSIQ Time 1,  $F(1,42) = 11.80$ ,  $p = .001$ . (Table 4). There was no significant associations between ABCL Externalizing and Glasgow Coma Score (GCS;  $p = .491$ ), age at injury ( $p = .287$ ), neurological signs ( $p = .363$ ) or surgical intervention ( $p = .577$ ).

### Table 4 about here

#### 3.5. Predictors of ABCL Externalizing: Executive functioning/interventions at 10-years post-TBI.

**Unadjusted model.** Univariate regression analyses revealed significant relations between ABCL Externalizing and BRIEF Behavioural Regulation Index ( $p = .026$ ) and 20-questions Abstract Reasoning, ( $p = .031$ ).

ABCL Externalizing was not significantly associated with BRIEF Global Executive Composite ( $p=.066$ ), BRIEF Metacognition ( $p=.128$ ) or access to interventions by 10-years post-injury ( $p=.427$ ).

**Adjusted model.** The multivariate adjusted model was moderately significant  $F(2,27) = 4.90$ ,  $p=.02$ , with BRIEF Behaviour Regulation Index the single significant predictor (Table 5).

**Table 5 about here**

### 3.6. Predictors of ABCL Externalizing: Socio-affective functioning.

**Unadjusted model.** Univariate regression analyses revealed significant relations between ABCL Externalizing and ACS Social Perception Total Score ( $p=.02$ ) and LCQ Proxy Report ( $p<.001$ ).

**Adjusted model.** The multivariate adjusted model was highly significant  $F(2,39) = 10.11$ ,  $p<.001$ , with LCQ Proxy report the single significant predictor (Table 6).

**Table 6 about here**

### 3.7. Predictors of ABCL Externalizing: Final adjusted model.

Due to high collinearity between measures of pre-injury adaptive functioning (Table 3), the final adjusted model evaluated the respective contributions of each pre-injury variable via three separate multivariate regression models.

As shown in Table 7, Model 1 was highly significant,  $F(4,36) = 9.03$ ,  $p<.001$ , with more frequent externalizing behavior related to poorer adaptive functioning, lower FSIQ (time 1) and more frequent social communication difficulty.

Model 2 ( $F(4,36) = 9.03, p < .001$ ) and Model 3 ( $F(4,38) = 9.97, p < .001$ ) were highly significant. In addition to the significant predictors identified in Model 1, pre-injury daily living skills and pre-injury socialization emerged as significant individual predictors of externalizing outcome in Model 2 and Model 3 respectively.

## Table 7 about here

## Discussion

The aim of the present longitudinal prospective study was to 1) investigate rates of EB in young adults with pediatric TBI; and 2) evaluate the respective contributions of a variety of injury and non-injury related risk and resilience factors at various time points to variability in EB outcomes 16 years after pediatric TBI.

There was partial support for all hypotheses. Relative to population expectations, rates of EB were significantly elevated among young adult survivors of pediatric TBI. Moreover, more frequent EB at 16-years post-injury was associated with poorer pre-injury adaptive functioning, reduced acute intellectual functioning and poorer pragmatic communication skills.

## Outcomes

Our results show that, by young adulthood, one of every four young people with a history of pediatric TBI had developed clinical or sub-threshold levels of externalising behaviour. Rates of EB in our sample compare to reported prevalence rates of 5-10% in the general population<sup>64,65</sup>, indicating a heightened risk of EB in young adulthood following pediatric TBI and mirroring previously identified associations between pediatric TBI and life-course persistent offending behaviors<sup>13</sup>.

In keeping with the premises of the HMSC model, EB was linked to a range of pre-injury and post-injury risk factors, discernible in the acute, adolescent and young adult phases. The contribution of pre-injury adaptive functioning to very long term social outcome is consistent with previous reports<sup>66</sup>, and may indicate that the influence of early brain injury interacts with a pre-existing vulnerability (i.e., '*double hazard theory*'<sup>67</sup>) to heighten risk for maladaptive behavior in the very-long-term post TBI. Moreover, the relationship between EB and FSIQ converges with previous literature in non-clinical samples<sup>68,69</sup> to suggest that higher levels of intellectual functioning may represent a source of resilience that buffers the risk of behavioral dysfunction in the long-term post injury.

In keeping with previous research<sup>30</sup>, pragmatic communication was the strongest and most significant predictor EB in young adults with pediatric TBI. The finding that poorer pragmatic communication was associated with more frequent EB may be interpreted from a diathesis stress perspective<sup>47</sup>. In line with the HMSC model, it may be that difficulty using and ascribing meaning to everyday social discourse contributes to rejection or alienation by interactive partners at the level of the social interaction. In this context, failure to negotiate the complex demands of everyday discourse is likely to elicit distress, reflected in EBs (e.g., aggression, rule breaking, intrusive conduct) which further limit the individual's capacity to negotiate the normative developmental goals of young adulthood.

Contrary to expectations, clinical or sub-threshold levels of EB problems were not associated with injury severity but were instead equally apparent across young adults with TBI of all severity levels. This finding is counter to previous studies that link more severe TBI to elevated risk for violent offending<sup>6,70</sup> and custodial sentences<sup>5,9</sup>, and suggests that early clinical indicators of injury severity have limited prognostic utility for longer term behavioral outcomes at least where injuries are sustained during childhood, where the brain is rapidly developing and has potential for reorganization.



Moreover, while previous reports have linked EB and/or persistent offending to factors such as social disadvantage and poorer executive functioning<sup>32-34</sup>, these factors did not significantly contribute to EB in the final adjusted model. While these non-significant relationships may to some extent reflect small sample size, they may also indicate that, after prolonged recovery and increasing time since injury, executive function and indices of pre-injury environment become less important prognostic indicators of outcome.

### Limitations

*Sample.* Attrition and sample bias are potential risks with this prospective, longitudinal study. Due to work commitments, travel distance or current life events, some young adult participants were not able to participate. Nevertheless, comparison of the participating and non-participating families indicated no systematic differences with the exception that the non-participant group had lower SES.

*Measures.* The source of information, proxy-report completed by the significant (either parent or partner/close friend), may also affect the level of reported externalizing symptoms. As young people become more independent, parents may be less knowledgeable about the young person's psychosocial functioning. While Green et al.<sup>71</sup> reported a fair-to-excellent agreement on psychosocial functioning between the adolescent with a pediatric TBI and their parent, Rosema et al.<sup>72</sup> showed that, during the transition into adulthood, the young adult with pediatric TBI did agree with parent report on the more observable behaviors, such as drug and alcohol use, social and communication skills, however, they did not concur on levels of internalizing symptoms, aggressive behavior and thought problems. Therefore self-report as well as direct such as a structured interview is recommended to obtain a more complete representation of risk and resilience factors of EBs following pediatric TBI.

Clinical implications and future research

Relative to population expectations<sup>64,65</sup>, a greater proportion of young adults with TBI demonstrated clinical or sub-threshold levels of EBs, which may in turn place these young people at greater risk for maladaptive behavior trajectories characterized by rule breaking, anti-social behaviour and offending. These findings have implications for policy implementation in the youth justice system. For example, screening and assessment for TBI within youth justice services may increase understanding of factors that may lead young people to offend, and assist in identifying young offenders who may benefit from relevant interventions, such as psycho-education and rehabilitation programs that specifically target social communicative dysfunction that persists into the long-term post injury<sup>30</sup>. Surprisingly, and in contrast to the largely medical model used to predict outcomes post-injury, injury characteristics were less important than environmental and pre-injury factors in determining outcome. Environmental factors, in particular, may be seen as potentially modifiable risk factors, offering an opportunity for early intervention to reduce risk of long term problems in this group.

The heightened prevalence of clinical or sub-threshold levels of EB in our sample underscores the need for provision of such early preventative interventions, as well as long term follow up and psycho-education for young people with TBI. For example, initial assessments in the acute and chronic stages of injury may assist to identify children presenting with risk factors, such as poorer pre-injury adaptive functioning and reduced IQ, and direct these children to appropriate services. An awareness of such factors should be shared with primary health care providers and schools, so as to support follow-up provision and further monitoring of relevant factors. This will also offer the means to provide continued engagement with parents and young people regarding the potential medium and

long term impact of TBI on behavioural and functional difficulties that they may not readily associate with the injury.

In addition, further research is needed to identify factors that may be protective against problematic levels of externalising behaviour despite the presence of pre-injury and acute risk factors. Person-oriented, qualitative case study approaches can complement group-level analyses, and offer an opportunity to evaluate how injury and non-injury related risk and resilience factors may interact to contribute to externalising behaviour in young adulthood.

## Conclusion

In summary, results suggest that young adults with pediatric TBI are at elevated risk for externalizing trajectories characterized by aggression, rule breaking and intrusive conduct. In line with the HMSC model, more frequent externalizing behaviour was linked to a range of pre-and post injury risk and resilience factors, including poorer pre-injury adaptive functioning, reduced IQ and more frequent pragmatic communication difficulty. These findings underscore the need for screening and assessment of TBI among young offenders, and suggest that early and long-term targeted interventions may be required to address risk factors for EBs in children and young people with TBI.

## References

1. Ilie G, Boak A, Adlaf EM, Asbridge M, Cusimano, MD. Prevalence and correlates of traumatic brain injuries among adolescents. *JAMA* 2013;309:2550-2.
2. Segalowitz SJ, Lawson S. Subtle symptoms associated with self-reported mild head injury. *J Learn Disabil.* 1995;28:309–319.

- 446 3. Segalowitz SJ, Brown D. Mild head injury as a source of learning disabilities. *J Learn*  
447 *Disabil* 1991;24:551–558.
- 448 4. Frost RB, Farrer TJ, Primosch M, Hedges DW. Prevalence of traumatic brain injury in the  
449 general adult population: A meta-analysis. *Neuroepidemiology* 2013;40:154–159.
- 450 5. Davies RC, Williams WH, Hinder D, Burgess CNW, Mounce LTA. Self-reported  
451 Traumatic Brain Injury and Post Concussion Symptoms in incarcerated youth: A dose  
452 response relationship. *J Head Trauma Rehabil.* 2012;7:E21-7
- 453 6. Kenny DT, Lennings CJ. The relationship between head injury and violent offending in  
454 juvenile detainees. *Contemporary Issues Crime Just.* 2007;107:1–15.
- 455 7. Moore E, Indig D, Haysom L. Traumatic Brain Injury, Mental Health, Substance Use, and  
456 Offending Among Incarcerated Young People. *J Head Trauma Rehabil.* 2014;29:239-247
- 457 8. Williams WH, Cordan G, Mewse AJ, Tonks J, Burgess CNW. Self-reported traumatic  
458 brain injury in male young offenders: a risk factor for re-offending, poor mental health and  
459 violence? *Neuropsychol Rehabil.* 2010;20:801–812.
- 460 9. Williams WH, Mewse AJ, Tonks J, Mills S, Burgess CN, Cordan G. Traumatic brain  
461 injury in a prison population: prevalence and risk for re-offending. *Brain Inj.* 2010;24:1184-  
462 1188
- 463 10. Schofield PW, Butler G, Hollis SJ, Smith NE, Lee SJ, Kelso WM. Neuropsychiatric  
464 correlates of traumatic brain injury (TBI) among Australian prison entrants. *Brain Inj.*  
465 2006;20:1409–1418
- 466 11. Ferguson PL, Pickelsimer EE, Corrigan JD, Bogner JA, Wald M. Prevalence of traumatic  
467 brain injury among prisoners in South Carolina. *J Head Trauma Rehabil.* 2012;27:E11-20.
- 468 12. Moffitt T. Adolescent limited and life course persistent antisocial behaviour: a  
469 developmental taxonomy. *Psychol Rev.* 1993;100:674–701.

- 470 13. Raine A, Moffitt TE, Caspi A, Loeber R, Stouthamer-Loeber M, Lynam D.  
471 Neurocognitive impairments in boys on the life-course persistent antisocial path. *J Abnorm*  
472 *Psychol.* 2005;114:38-49.
- 473 14. Mulvey EP, Steinberg L, Fagan J, et al. Theory and Research on Desistance from  
474 Antisocial Activity among Serious Adolescent Offenders. *Youth Violence Juv Justice.*  
475 2004;2:213.
- 476 15. Bottoms AE, Shapland J. Steps towards desistance among male young adult recidivists .  
477 IN: Farrall S, Hough M, Maruna S, Sparks R, eds. *Escape routes: contemporary perspectives*  
478 *on life after punishment.* London: Routledge, 2011.
- 479 16. Tasker RC, Salmond CH, Westland AG, et al. Head circumference and brain and  
480 hippocampal volume after severe traumatic brain injury in childhood. *Pediatr. Res.* 2005; 58:  
481 302-308.
- 482 17. Wilde EA, Hunter JV, Newsome MR, et al. Frontal and temporal morphometric findings  
483 in children after moderate to severe traumatic brain injury. *J. Neurotrauma.* 2005; 22: 333-  
484 334.
- 485 18. Green RE, Turner GR, Thompson WF. Deficits in emotion perception in adults with  
486 recent traumatic brain injury. *Neuropsychologia.* 2004; 42: 133-141.
- 487 19. Sesma HW, Slomine BS, Ding R, McCarthy ML. Children's Health After Trauma  
488 (CHAT) Study Group. Executive functioning in the first year after pediatric traumatic brain  
489 injury. *Pediatrics.* 2008;121:E1686-95
- 490 20. Catroppa C, Anderson VA, Morse SA, Haritou F, Rosenfeld JV. Outcome and predictors  
491 of functional recovery 5 years following pediatric traumatic brain injury. *J Pediatr Psychol.*  
492 2008;33:707-718

- 493 21. Fay TB, Yeates KO, Wade SL, Drotar D, Stancin T, Taylor HG. Predicting longitudinal  
494 patterns of functional deficits in children with traumatic brain injury. *Neuropsychology*.  
495 2009;23:271–282.
- 496 22. Ryan NP, Anderson V, Godfrey C, et al. Predictors of very-long-term sociocognitive  
497 function after pediatric traumatic brain injury: evidence for the vulnerability of the immature  
498 "social brain". *J Neurotrauma*. 2014;31:649-57
- 499 23. Morgan AB, Lilienfeld SO. A meta-analytic review of the relation between antisocial  
500 behavior and neuropsychological measures of executive function. *Clin Psychol Rev*.  
501 2000;20:113–156.
- 502 24. Ogilvie JM, Stewart AL, Chan RCK, Shum DHK. Neuropsychological measures of  
503 executive function and antisocial behavior: A meta-analysis. *Criminology*. 2011;49:1063–  
504 1107
- 505 25. Brower MC, Price BH. Neuropsychiatry of frontal lobe dysfunction in violent and  
506 criminal behaviour: a critical review. *Neurol Neurosurg Psychiatry*. 2001;71:720-726
- 507 26. Tonks J, Slater A, Frampton I, Wall SE, Yates P, Williams WH. The development of  
508 emotion and empathy skills after childhood brain injury. *DevMed Child Neurol*. 2009; 51:8-  
509 16.
- 510 27. Tonks J, Yates P, Frampton I, Williams WH, Harris D, Slater A. Resilience and the  
511 mediating effects of executive dysfunction after childhood brain injury: a comparison  
512 between children aged 9-15 years with brain injury and non-injured controls. *Brain Inj*.  
513 2011;25:870–81
- 514 28. Beitchman JH, Douglas L, Wilson B, et al. Adolescent substance use disorders: Findings  
515 from a 14-year follow-up of speech/language impaired and control children. *J Clin Child*  
516 *Psychol*. 1999;28:312–321

- 517 29. Brownlie EB, Beitchman JH, Escobar M, et al. Early language impairment and young  
518 adult delinquent and aggressive behavior. *J Abnorm Child Psychol.* 2004;32:453–467
- 519 30. Ryan NP, Anderson V, Godfrey C, et al. Social communication mediates the relationship  
520 between emotion perception and externalizing behaviors in young adult survivors of pediatric  
521 traumatic brain injury. *Int J Dev Neurosci.* 2013;31:811-819
- 522 31. Snow PC, Powell MB. Oral language competence in incarcerated young offenders: Links  
523 with offending severity. *Int. J. Speech Lang. Pathol.* 2011;13:480–489
- 524 32. Farrington DP. The development of offending and antisocial behaviour from childhood:  
525 key findings from the Cambridge study in delinquent development.  
526 *J Child Psychol Psychiatry.* 1995;36:929–964
- 527 33. Office of the Surgeon General *Youth Violence: A Report of the Surgeon General.*  
528 Washington, DC: U.S. Department of Health and Human Services, Office of the Secretary,  
529 Office of Public Health and Science, Office of the Surgeon General, 2001.
- 530 34. Prior D, Paris, A. *Preventing Children's Involvement in Crime and Anti Social*  
531 *Behaviour: A Literature Review.* London: Department for Education and Skills, 2004.
- 532 35. Kurowski BG, Taylor HG, Yeates KO, Walz NC, Stancin T, Wade SL. Caregiver ratings  
533 of long-term executive dysfunction and attention problems after early childhood traumatic  
534 brain injury: family functioning is important. *PM R.* 2011;3:836-45.
- 535 36. Woods DT, Catroppa C, Barnett P, Anderson VA. Parental disciplinary practices  
536 following acquired brain injury in children. *Dev Neurorehabil.* 2011;14(5):274-82.
- 537 37. Wade SL, Cassedy A, Walz NC, Taylor HG, Stancin T, Yeates KO. The relationship of  
538 parental warm responsiveness and negativity to emerging behavior problems following  
539 traumatic brain injury in young children. *Dev Psychol;* 2011;47:119–33.

- 540 38. Yeates KO, Taylor HG, Walz NC, Stancin T, Wade SL. The family environment as a  
541 moderator of psychosocial outcomes following traumatic brain injury in young children.  
542 *Neuropsychology*. 2010;24:345-56.
- 543 39. Catroppa C, Anderson V. Recovery of educational skills following pediatric head injury.  
544 *Pediatr Rehabil*. 1999;3:167–175
- 545 40. Catroppa C, Anderson VA, Muscara F, et al Educational skills: Long-term outcome and  
546 predictors following paediatric traumatic brain injury. *Neuropsychol Rehabil*. 2009;19:716-  
547 732.
- 548 41. Ewing-Cobbs L, Barnes M, Fletcher JM, Levin HS, Swank PR, Song J. Modeling of  
549 longitudinal academic achievement scores after pediatric traumatic brain injury. *Dev*  
550 *Neuropsychol*. 2004;25:107–133
- 551 42. Ewing-Cobbs L, Prasad MR, Kramer L, et al. Late intellectual and academic outcomes  
552 following traumatic brain injury sustained during early childhood. *J Neurosurg*.  
553 2006;105:287–296
- 554 43. Dockrell J, Lindsay G, Palikara O, Cullen MA. *Raising the Achievements of Children and*  
555 *Young People with Specific Speech and Language Difficulties and other Special Educational*  
556 *Needs through School to Work and College*. London: Department for Education and Skills,  
557 2007.
- 558 44. Hines PJ, Wible B, McCartney M. Learning to read, reading to learn. *Science*  
559 2010;23:447.
- 560 45. Snow P, Powell M. Youth (in)justice: Oral language competence in early life and risk for  
561 engagement in antisocial behaviour in adolescence, *Australian Institute of Criminology*,  
562 *Trends and issues in crime and criminal justice*, 2012.
- 563 46. Li L, Liu J. The effect of pediatric traumatic brain injury on behavioral outcomes: a  
564 systematic review. *Dev Med Child Neurol*. 2013;55:37-45



- 565 47. Yeates KO, Bigler ED, Dennis M, et al. Social outcomes in childhood brain disorder: A  
566 heuristic integration of social neuroscience and developmental psychology. *Psychol Bull.*  
567 2007;133:535-556
- 568 48. Anderson V, Godfrey C, Rosenfeld JV, Catroppa C. 10 years outcome from childhood  
569 traumatic brain injury. *Int J Dev Neurosci*, 2012;30:217–224
- 570 49. Teasdale, G., & Jennett, B. (1974). Assessment of coma and impaired consciousness: A  
571 practical scale. *The Lancet*, 1974;2:81–83.
- 572 50. Sparrow S, Balla DA, & Cicchetti DV. *Vineland Adaptive Behavior Scales: Interview*  
573 *Edition. Survey Form Manual*. Circle Pines, MN: American Guidance Services Resources;  
574 1984.
- 575 51. Noller P. *ICPS Family Functioning Scales*. Unpublished manuscript. University of  
576 Queensland; 1988.
- 577 52. Catroppa, C., Anderson, V., Godfrey, C., & Rosenfeld, JV. Attentional skills 10 years  
578 post-paediatric traumatic brain injury (TBI). *Brain injury*, 2011; 25: 858-869.
- 579 53. Bayley N. *Bayley Scales of Infant Development: Birth to Two Years*. San Antonio, TX:  
580 The Psychological Corporation; 1969.
- 581 54. Wechsler D. *Manual for the Preschool and Primary Intelligence Scale—Revised*. New  
582 York, NY: Psychological Corporation; 1989.
- 583 55. Wechsler D. *Manual for the Wechsler Scale of Children’s Intelligence—III*. New York,  
584 NY: Psychological Corporation; 1991.
- 585 56. Daniel A. *Power, Privilege and Prestige: Occupations in Australia*. Melbourne,  
586 Australia: Longman-Cheshire; 1983.
- 587 57. Beauchamp, M., Catroppa, C., Godfrey, C., Morse, S., Rosenfeld, J.V. & Anderson, V.  
588 (2011) Selective Changes in Executive Functioning Ten Years After Severe Childhood  
589 Traumatic Brain Injury. *Dev Neuropsych.*, 2011;36:578–595

590

591 Catroppa

592 58. Gioia GA, Isquith PK, Guy SC, & Kenworthy L. *Behavior Rating Inventory of Executive*  
593 *Function*. Odessa, FL: Psychological Assessment; 2000.

594 59. Delis, D., Kaplan, E., & Kramer, JH. *Delis-Kaplan executive functions system*. San  
595 Antonio, TX: The Psychological Corporation; 2001.

596 60. Achenbach, T.M., Rescorla, L.A., *Manual for the ASEBA adult forms and profiles*.  
597 University of Vermont, Research Center for Children, Youth and Families, Burlington, VT;  
598 2003

599 61. Tenneij, N.H., & Koot, H.M., (2007). A preliminary investigation into the utility of the  
600 Adult Behavior checklist in the assessment of psychopathology in people with low IQ. *J*  
601 *Appl Res Intellect Disabil*;2007;2(5):391–400.

602 62. Douglas, J.M., O'Flaherty, C.A., Snow, P.C., (2000) Measuring perception of  
603 communicative ability: the development and evaluation of the La Trobe Communication  
604 Questionnaire. *Aphasiology*,2000;14(3):251-268.

605 63. Pearson, 2009. *Advanced Clinical Solutions for the WAIS-IV and WMS-IV*. San Anto-nio,  
606 TX; 2009.

607 64. Moffitt TE. Natural histories of delinquency. IN: Weitekam EGM, Kerner HJ. (Eds)  
608 *Cross-national longitudinal research on human development and criminal behaviour*.  
609 Dordrecht, Netherlands: Kluwer Academic Publishers. 3-61, 1994.

610 65. Bor W, McGee TR, Hayatbakhsh R, Najman J. *The young adult outcomes of childhood*  
611 *and adolescent antisocial behaviour: an Australian cohort study*. Australian Institute of  
612 Criminology, Criminology Research Council: Canberra, 2007

- 613 66. Catroppa C, Godfrey C, Rosenfeld JV, Hearps SJC, Anderson, V. Functional recovery 10  
614 years following pediatric traumatic brain injury: Outcomes and predictors. *J Neurotraum*,  
615 2012;29(16):2539-2547.
- 616 67. Breslau N. Does brain dysfunction increase children's vulnerability to environmental  
617 stress? *Arch Gen Psychiat* 1990;47:15-20.
- 618 68. Farrington DP. Juvenile delinquency. IN: Coleman JC. (Ed.) *The School Years*, 2nd ed.  
619 (pp. 123-163). London: Routledge, 1992.
- 620 69. Henry B, Moffitt TE. Neuropsychological and neuroimaging studies of juvenile  
621 delinquency and adult criminal behaviour. IN: Stoff D, Breiling J, Maser, JD. (Eds),  
622 *Handbook of Antisocial Behaviour*, pp. 280-288. New York: John Wiley and Sons, 1997
- 623 70. Fazel S, Lichtenstein P, Grann M, Långström N. Risk of violent crime in individuals with  
624 epilepsy and traumatic brain injury: a 35-year Swedish population study. *PLoS Med*.  
625 2011;8:e1001150.
- 626 71. Green L, Godfrey C, Soo C, Anderson V, Catroppa, C. Agreement between parent-  
627 adolescent rating on psychosocial outcome and quality-of-life following childhood traumatic  
628 brain injury. *Dev. Neurorehabil*. 2012;15:105–113.
- 629 72. Rosema S, Muscara F, Anderson V, Godfrey C, Eren S, Catroppa C. Agreement on and  
630 predictors of long-term psychosocial development 16 years post-childhood traumatic brain  
631 injury. *J Neurotraum*, 2014;31:899-905.

*Table 1. Characteristics of the TBI sample according to injury severity.*

	Mild TBI ( <i>n</i> = 15)	Moderate TBI ( <i>n</i> = 29)	Severe TBI ( <i>n</i> = 11)
<i>Demographics</i>			
No. males, <i>n</i> (%) <sup>a</sup>	8 (53)	15 (52)	7 (64)
Injury age (years), <i>M</i> ( <i>SD</i> )	7.89 (3.68)	6.37 (3.38)	6.27 (3.36)
Age at 16-year follow up, (years), <i>M</i> ( <i>SD</i> )	24.49 (4.24)	23.66 (3.63)	23.41 (4.02)
<i>Pre-injury function</i>			
VABS: PRE, <i>M</i> ( <i>SD</i> )	111.64 (20.22)	111.67 (16.15)	106.73 (16.75)
FFQ: PRE, <i>M</i> ( <i>SD</i> )	62.62 (13.99)	65.10 (5.31)	66.55 (5.09)
SES: acute, <i>M</i> ( <i>SD</i> )	3.85 (.93)	4.45 (1.06)	3.97 (1.08)
<i>Acute injury factors</i>			
GCS 24-hours, <i>M</i> ( <i>SD</i> ) <sup>a</sup>	14.83 (.39)	12.85 (2.40)	8.90 (2.51)
FSIQ: Acute, <i>M</i> ( <i>SD</i> )	100.93 (13.86)	102.85 (15.33)	97.67 (16.27)

<sup>a</sup> Denotes statistical significance,  $p < .05$ .

*Table 2. Proportion of participants with clinical or sub-threshold levels of externalizing symptoms as a function of injury severity.*

	Mild TBI ( <i>n</i> = 15)	Moderate TBI ( <i>n</i> = 29)	Severe TBI ( <i>n</i> = 11)	Total ( <i>n</i> = 55)
Impaired, <i>n</i> (%)	3 (20)	7 (24)	3 (27)	13 (24)

*Table 3. Predictors of ABCL Externalizing: Pre-injury adaptive functioning.*

	ABCL Externalizing			
	<i>B</i>	<i>SE</i>	<i>p</i> value	95% CI
Pre-injury Adaptive function	-.11	.15	.44	[-.41, .18]
Pre-injury Daily Living	.02	.13	.90	[-.24, .27]
Pre-injury Socialization	-.07	.10	.47	[-.28, .13]

*Table 4. Predictors of ABCL Externalizing: Acute Intellectual Functioning*

	ABCL Externalizing			
	<i>B</i>	<i>SE</i>	<i>p</i> value	95% CI
FSIQ Time 1	-.22	.06	.001	[-.35, -.09]

*Table 5. Predictors of ABCL Externalizing: Executive function at 10-years post-TBI.*

	ABCL Externalizing			
	<i>B</i>	<i>SE</i>	<i>p</i> value	95% CI
BRIEF BR	.16	.08	.05	[.01, .33]
20-questions	-.48	.31	.13	[-1.11, .16]

*Table 6. Predictors of ABCL Externalizing: Socio-affective function at 16-years post-TBI.*

	ABCL Externalizing			
	<i>B</i>	SE	<i>p</i> value	95% CI
ACS Social Perception	-.29	.21	.18	[-.72, .14]
LCQ Proxy Report	.35	.10	.01	[.14, .56]

*Table 7: Predictors of ABCL Externalizing: Final Adjusted Model.*

	ABCL Externalizing			
	<i>B</i>	<i>SE</i>	<i>p</i> value	95% CI
<b>Model 1</b>				
Pre-injury Adaptive Composite	-.09	.05	.05*	[-.19, .00]
FSIQ Acute	-.13	.06	.02*	[-.25, -.02]
BRIEF BR 10-years	.08	.07	.26	[-.06, .22]
LCQ Proxy 15-years	.28	.09	.003*	[.11, .46]
<b>Model 2</b>				
Pre-injury Daily Living Skills	-.11	.05	.05*	[-.23, -.01]
FSIQ Acute	-.15	.05	.01*	[-.26, -.04]
BRIEF BR 10-years	.09	.07	.22	[-.06, .24]
LCQ Proxy 15-years	.28	.09	.003*	[.10, .46]
<b>Model 3</b>				
Pre-injury Socialization	-.10	.04	.04*	[-.19, -.01]
FSIQ Acute	-.14	.06	.02*	[-.25, -.02]
BRIEF BR 10-years	.08	.06	.25	[-.06, .21]
LCQ Proxy 15-years	.29	.08	.002*	[.12, .46]

\*Denotes statistically significant relationship,  $p < .05$ .