

Bipartitions of Highly Connected Tournaments

Kim, Jaehoon; Kuhn, Daniela; Osthus, Deryk

DOI:

[10.1137/15M1006611](https://doi.org/10.1137/15M1006611)

License:

Creative Commons: Attribution (CC BY)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Kim, J, Kuhn, D & Osthus, D 2016, 'Bipartitions of Highly Connected Tournaments', *SIAM Journal on Discrete Mathematics*, vol. 30, no. 2, pp. 895–911. <https://doi.org/10.1137/15M1006611>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Eligibility for repository: Checked on 6/6/2016

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

BIPARTITIONS OF HIGHLY CONNECTED TOURNAMENTS

JAEHOON KIM, DANIELA KÜHN, DERYK OSTHUS

ABSTRACT. We show that if T is a strongly $10^9 k^6 \log(2k)$ -connected tournament, there exists a partition A, B of $V(T)$ such that each of $T[A]$, $T[B]$ and $T[A, B]$ is strongly k -connected. This provides solutions to tournament analogues of two partition conjectures of Thomassen regarding highly connected graphs. We also discuss spanning linkages as well as non-separating subdivisions in highly connected tournaments.

1. INTRODUCTION

1.1. Partitions of highly connected tournaments. The study of graph partitions where the resulting subgraphs inherit the properties of the original graph has a long history with some surprises and numerous open problems, see e.g. the survey [7]. For example, a classical result of Hajnal [1] and Thomassen [9] implies that for every k there exists an integer $f(k)$ such that every $f(k)$ -connected graph has a vertex partition into sets A and B so that both A and B induce k -connected graphs. A related conjecture of Thomassen [12] states that for every k there is an $f(k)$ such that every $f(k)$ -connected graph G has a bipartition A, B so that the spanning bipartite graph $G[A, B]$ is k -connected. It is not hard to show that one cannot achieve both the above properties simultaneously in a highly connected graph. (For example, consider the bipartite graph with vertex classes A and B where $A := [n]$, B consists of all ℓ -element subsets of $[n]$ and $a \in A$ is joined to $b \in B$ if and only if $a \in b$.) On the other hand, our main result states that for tournaments, we can find a single partition which achieves both the above properties. Below we denote by $T[A, B]$ the bipartite subdigraph of T which consists of all edges between A and B but no others.

Theorem 1.1. *Let T be a tournament and $k \in \mathbb{N}$. If T is strongly $10^9 k^6 \log(2k)$ -connected, there exists a partition V_1, V_2 of $V(T)$ such that each of $T[V_1]$, $T[V_2]$ and $T[V_1, V_2]$ is strongly k -connected.*

We have made no attempt to optimize the bound on the connectivity in Theorem 1.1. (It would be straightforward to obtain minor improvements at the expense of more careful calculations.) On the other hand, it would be interesting to obtain the correct order of magnitude for the connectivity bound.

Kühn, Osthus and Townsend [4] earlier proved the weaker result that every strongly $10^8 k^6 \log(4k)$ -connected tournament T has a vertex partition V_1, V_2 such that $T[V_1]$ and $T[V_2]$ are both strongly k -connected (with some control over the sizes of V_1 and V_2). This proved a conjecture of Thomassen. [4] raised the question whether this can be extended to digraphs.

Date: January 27, 2016.

The research leading to these results was partially supported by the European Research Council under the European Union's Seventh Framework Programme (FP/2007–2013) / ERC Grant Agreements no. 306349 (J. Kim and D. Osthus) as well as 258345 (D. Kühn).

As described later, our proof of Theorem 1.1 develops ideas in [4]. These in turn are based on the concept of robust linkage structures which were introduced in [2] to prove a conjecture of Thomassen on edge-disjoint Hamilton cycles in highly connected tournaments. Further (asymptotically optimal) results leading on from these approaches were obtained by Pokrovskiy [5, 6].

1.2. Subdivisions and linkages. The famous Lovász path removal conjecture states that for every $k \in \mathbb{N}$ there exists $g(k) \in \mathbb{N}$ such that for every pair x, y of vertices in a $g(k)$ -connected graph G we can find an induced path P joining x and y in G for which $G \setminus V(P)$ is k -connected. In [11], Thomassen proved a tournament version of this conjecture. Here, we generalize his argument to observe that highly connected tournaments contain a non-separating subdivision of any given digraph H (with prescribed branch vertices). The case when $d = 2$ and $m = 1$ corresponds to the result in [11].

Theorem 1.2. *Let $k, d, m \in \mathbb{N}$. Suppose that T is a strongly $(k + m(d + 2))$ -connected tournament, that D is a set of d vertices in T , that H is a digraph on d vertices and m edges and that ϕ is a bijection from $V(H)$ to D . Then T contains a subdivision H^* of H such that*

- (i) *for each $h \in V(H)$ the branch vertex of H^* corresponding to h is $\phi(h)$,*
- (ii) *$T \setminus V(H^*)$ is strongly k -connected,*
- (iii) *for every edge e of H , the path P_e of H^* corresponding to e is backwards-transitive.*

Here a directed path $P = x_1 \dots x_t$ in a tournament T is *backwards-transitive* if $x_i x_j$ is an edge of T whenever $i \geq j + 2$. The graph version of Theorem 1.2 is still open and would follow from the following beautiful conjecture of Thomassen [10].

Conjecture 1.3. *For every $k \in \mathbb{N}$ there exists $f(k) \in \mathbb{N}$ such that if G is a $f(k)$ -connected graph and $M \subseteq V(G)$ consists of k vertices then there exists a partition V_1, V_2 of $V(G)$ such that $M \subseteq V_1$, both $G[V_1]$ and $G[V_2]$ are k -connected, and each vertex in V_1 has at least k neighbours in V_2 .*

The case $|M| = 2$ would already imply the path removal conjecture. The case $M = \emptyset$ was proved in [3]. It implies the existence of non-separating subdivisions (without prescribed branch vertices) in highly connected graphs. Clearly, Theorem 1.1 implies a tournament version of Conjecture 1.3. (Indeed, to see this apply Theorem 1.1 to obtain a partition V_1, V_2 such that each of $T[V_1]$, $T[V_2]$ and $T[V_1, V_2]$ is strongly $2k$ -connected. So in particular, each vertex in V_2 has at least $2k$ out-neighbours and at least $2k$ in-neighbours in V_1 . By moving all vertices in $M \cap V_2$ to V_1 we obtain a partition V'_1, V'_2 such that both $T[V'_1]$ and $T[V'_2]$ are strongly k -connected, and actually $T[V'_1, V'_2]$ is too.)

The next theorem guarantees a spanning linkage in a highly connected tournament. It was proved by Thomassen [11] with a super-exponential bound on the connectivity. He asked whether a linear bound suffices. Here we reduce the bound to a polynomial one. Pokrovskiy [5] showed that a linear bound suffices to guarantee a linkage if we do not require it to be spanning.

Theorem 1.4. *Let $k \in \mathbb{N}$. Suppose that T is a strongly $(k^2 + 3k)$ -connected tournament and that $x_1, \dots, x_k, y_1, \dots, y_k$ are vertices in T such that $x_i \neq y_i$ for all $i \in [k]$ and all the pairs (x_i, y_i) are distinct. Then T contains pairwise internally disjoint paths P_i from x_i to y_i such that $\{x_1, \dots, x_k, y_1, \dots, y_k\} \cap V(P_i) = \{x_i, y_i\}$ and $V(T) = \bigcup_{i=1}^k V(P_i)$.*

Both Theorem 1.2 and 1.4 can be deduced from Theorem 1.1, but with weaker bounds. (For example, to deduce Theorem 1.2 from Theorem 1.1, we first consider a partition V_1, V_2 as in Theorem 1.1 and move all vertices of $D \cap V_2$ into V_1 to obtain a new partition V'_1, V'_2 . Then

$T[V'_1]$ is still highly connected and thus highly linked (cf. Theorem 2.2). We can use the latter property to find H^* satisfying (i) and (iii) in $T[V'_1]$. We now move the remaining vertices of V'_1 into V'_2 to obtain a set satisfying (ii). To derive Theorem 1.4 we proceed similarly except that at the end we use a result of Thomassen [8] that Theorem 1.4 holds for $k = 1$.) However, in Section 4 we adapt the argument from [11] to obtain a short direct proof of both Theorems 1.2 and 1.4.

2. NOTATION AND TOOLS

We write $\mathbb{N} := \{1, 2, \dots\}$ for the set of all natural numbers. Given $k \in \mathbb{N}$, we let $[k] := \{1, \dots, k\}$, $[k, k + \ell] := \{k, \dots, k + \ell\}$ and $\log k := \log_2 k$. We only consider digraphs which contain no loops and at most two edges between every pair of vertices, at most one in each direction. We write $V(G)$ and $E(G)$ for the set of vertices and the set of edges in a digraph G . We let $|G| := |V(G)|$. If $u, v \in V(G)$ we write uv for the directed edge from u to v . We write $N_G^-(v)$ and $N_G^+(v)$ for the in- and out-neighbourhoods of a vertex v in G . (So $v \notin N_G^-(v) \cup N_G^+(v)$.) We write $d_G^-(v) := |N_G^-(v)|$ and $d_G^+(v) := |N_G^+(v)|$ for the in-degree and the out-degree of v . We write $\delta^-(G)$ and $\delta^+(G)$ for the minimum in-degree and the minimum out-degree of G and let $\delta^0(G) := \min\{\delta^-(G), \delta^+(G)\}$. A set $A \subseteq V(G)$ *in-dominates* a set $B \subseteq V(G)$ if for every vertex $b \in B$ there exists a vertex $a \in A$ such that $ba \in E(G)$. Similarly, we say that A *out-dominates* B if for every vertex $b \in B$ there exists a vertex $a \in A$ such that $ab \in E(G)$. We say that a tournament T is *transitive* if we may enumerate its vertices v_1, \dots, v_m such that $v_i v_j \in E(T)$ if and only if $i < j$. In this case we call v_1 the *source* of T and v_m the *sink* of T . When referring to subpaths of tournaments, we always mean that these paths are directed (i.e. consistently oriented). The *length* of a path is the number of its edges. We say that a path P is *odd* if its length is odd, and *even* if its length is even. We say that two paths are disjoint if they are vertex-disjoint. A tournament T is *strongly k -connected* if $|T| > k$ and for every set $F \subseteq V(T)$ with $|F| < k$ and every ordered pair x, y of vertices in $V(T) \setminus F$ there exists a path from x to y in $T - F$. A tournament T is called *k -linked* if $|T| \geq 2k$ and whenever $x_1, \dots, x_k, y_1, \dots, y_k$ are $2k$ distinct vertices of T there exist disjoint paths P_1, \dots, P_k such that P_i is a directed path from x_i to y_i for each $i \in [k]$.

We now collect the tools which we need in our proof of Theorem 1.1. The following proposition is a straightforward consequence of the definition of linkedness.

Proposition 2.1. *Let $k \in \mathbb{N}$. Then a tournament T is k -linked if and only if $|T| \geq 2k$ and whenever $(x_1, y_1), \dots, (x_k, y_k)$ are ordered pairs of (not necessarily distinct) vertices of T , there exist distinct internally disjoint paths P_1, \dots, P_k such that for all $i \in [k]$ we have that P_i is a directed path from x_i to y_i and that $\{x_1, \dots, x_k, y_1, \dots, y_k\} \cap V(P_i) = \{x_i, y_i\}$.*

We will also use the following bound from [5] on the connectivity which forces a tournament to be highly linked.

Theorem 2.2. *For each $k \in \mathbb{N}$ every strongly $452k$ -connected tournament is k -linked.*

The following two lemmas guarantee that every tournament contains almost out-dominating and almost in-dominating sets which are not too large. (A similar observation was also used in [2], see Lemmas 8.3 and 8.4.)

Lemma 2.3. *Let T be a tournament, let $v \in V(T)$ and $c \in \mathbb{N}$ with $c \geq 2$. Suppose that $d_T^-(v) \geq 2^{c-1}$. Then there exist disjoint sets $A, E \subseteq V(T)$ and a vertex $a \in A$ such that the following properties hold:*

- (i) $2 \leq |A| \leq c$ and $T[A]$ is a transitive tournament with source a and sink v ,
- (ii) $A \setminus \{a\}$ out-dominates $V(T) \setminus (A \cup E)$,
- (iii) $|E| \leq (1/2)^{c-2} d_T^-(v)$.

Proof. Let $v_1 := v$. Roughly speaking, we will find A by choosing vertices v_1, \dots, v_i such that the size of their common in-neighbourhood (i.e. the intersection of their individual in-neighbourhoods) is minimised at each step. More precisely, suppose inductively that for some $1 \leq i < c$ we have already found a set $A_i = \{v_1, \dots, v_i\}$ and a set W_i such that the following holds:

- (a) $T[A_i]$ is a transitive tournament with sink v_1 ;
- (b) $W_i = \emptyset$ or $W_i = \{a\}$ for some vertex a . Moreover, if $W_i = \{a\}$ then $E_i \cup A_i \subseteq N^+(a)$, where $E_i := \bigcap_{j=1}^i N^-(v_j) \setminus W_i$.
- (c) $|E_i| \leq \frac{1}{2^{i-1}} d^-(v)$. Moreover, $|E_i| > 0$ if $W_i = \emptyset$.

Note that (a)–(c) hold for $i = 1$ if we let $A_1 := \{v_1\}$ and $W_1 = \emptyset$.

We first consider the case that $|E_i| \leq \frac{1}{2^{c-2}} d^-(v)$. If $W_i = \emptyset$, choose any vertex $a \in E_i$. Else let a be the vertex in W_i . In both cases let $A := A_i \cup \{a\}$ and $E := E_i \setminus \{a\}$. Then A and E satisfy (i)–(iii).

So suppose next that $|E_i| > \frac{1}{2^{c-2}} d^-(v)$. (Note that in particular, this means that $|E_i| \geq 2$.) By averaging, it follows that E_i must contain a vertex x of in-degree at most $|E_i|/2$ in $T[E_i]$. If the in-degree of x in $T[E_i]$ is nonzero or $W_i \neq \emptyset$, let $v_{i+1} := x$. Else let v_{i+1} be a vertex of in-degree at most $|E_i \setminus \{x\}|/2$ in $T[E_i \setminus \{x\}]$, and let $W_{i+1} := \{x\}$ (note that we can find such a v_{i+1} as $|E_i| \geq 2$). Now let $A_{i+1} := \{v_1, \dots, v_{i+1}\}$ and let $E_{i+1} := (E_i \cap N^-(v_{i+1})) \setminus W_{i+1}$. Then $T[A_{i+1}]$ is a transitive tournament with sink v_1 and

$$|E_{i+1}| \leq \frac{1}{2} |E_i| \leq \frac{1}{2^i} d^-(v).$$

So we have shown that (a)–(c) hold with $i+1$ playing the role of i . By repeating this construction, we will eventually find A and E satisfying (i)–(iii). (Indeed, note that we must be in the first case for some $i < c$, in particular this implies that $|A| \leq c$). \square

The next lemma follows immediately from Lemma 2.3 by reversing the orientations of all edges.

Lemma 2.4. *Let T be a tournament, let $v \in V(T)$ and $c \in \mathbb{N}$ with $c \geq 2$. Suppose that $d_T^+(v) \geq 2^{c-1}$. Then there exist disjoint sets $B, E \subseteq V(T)$ and a vertex $b \in B$ such that the following properties hold:*

- (i) $2 \leq |B| \leq c$ and $T[B]$ is a transitive tournament with sink b and source v ,
- (ii) $B \setminus \{b\}$ in-dominates $V(T) \setminus (B \cup E)$,
- (iii) $|E| \leq (1/2)^{c-2} d_T^+(v)$.

We will also need the following observation, which guarantees a small set Z of vertices in a tournament such that every vertex outside Z has many out- and in-neighbours in Z .

Proposition 2.5. *Let $k, n \in \mathbb{N}$ and let T be a tournament on $n \geq 4$ vertices. Then there is a set $Z \subseteq V(T)$ of size $|Z| \leq 3k \log n$ such that each vertex in $V(T) \setminus Z$ has at least k out-neighbours and at least k in-neighbours in Z .*

Proof. We may assume that $n \geq 3k \log n$. We will use the fact that every tournament on n vertices contains an in-dominating set of size at most $c := \lceil \log n \rceil \leq (3 \log n)/2$. (This can

be proved by choosing the vertices x_1, x_2, \dots in the in-dominating set one by one, similarly as in the proof of Lemma 2.3: at the i th step we let x_i be a vertex with the smallest out-degree in $T[\bigcap_{j < i} N^+(x_j)]$. Choose an in-dominating set V_1 in T of size at most c . Now consider the tournament $T - V_1$. Choose an in-dominating set V_2 in $T - V_1$ with size at most c . Continue in this way to obtain disjoint sets V_1, \dots, V_k . Proceed similarly to obtain disjoint sets U_1, \dots, U_k , each of size at most c , such that each U_i is an out-dominating set in $T - (U_1 \cup \dots \cup U_{i-1})$. We can take $Z := V_1 \cup \dots \cup V_k \cup U_1 \cup \dots \cup U_k$. \square

3. PROOF OF THEOREM 1.1

Let $X := \{x_1, x_2, \dots, x_{6k}\} \subseteq V(T)$ consist of $6k$ vertices whose in-degree in T is as small as possible, and let $Y := \{y_1, y_2, \dots, y_{6k}\}$ be a set of $6k$ vertices in $V(T) \setminus X$ whose out-degree in T is as small as possible. Define

$$\hat{\delta}^-(T) := \min_{v \in V(T) \setminus X} d_T^-(v) \quad \text{and} \quad \hat{\delta}^+(T) := \min_{v \in V(T) \setminus Y} d_T^+(v).$$

Let $c := \lceil \log(120k^2) \rceil + 2 \leq 9k$. Apply Lemmas 2.3 and 2.4 with parameter c repeatedly (removing the dominating sets each time) to obtain disjoint sets of vertices A_1, A_2, \dots, A_{6k} , B_1, B_2, \dots, B_{6k} and sets of vertices $E_{A_1}, \dots, E_{B_{6k}}$ satisfying the following properties for all $i \in [6k]$, where we write $D := \bigcup_{i=1}^{6k} (A_i \cup B_i)$,

- (D1) $2 \leq |A_i| \leq c$ and $T[A_i]$ is a transitive tournament with sink x_i and source a_i ,
- (D2) $2 \leq |B_i| \leq c$ and $T[B_i]$ is a transitive tournament with source y_i and sink b_i ,
- (D3) $A_i \setminus \{a_i\}$ out-dominates $V(T) \setminus (D \cup E_{A_i})$ in T ,
- (D4) $B_i \setminus \{b_i\}$ in-dominates $V(T) \setminus (D \cup E_{B_i})$ in T ,
- (D5) $|E_{A_i}| \leq (1/2)^{c-2} \hat{\delta}^-(T)$,
- (D6) $|E_{B_i}| \leq (1/2)^{c-2} \hat{\delta}^+(T)$.

Indeed, to see that we can find such sets satisfying (D1)–(D6), we first apply Lemma 2.3 with $x_1, T - (Y \cup X \setminus \{x_1\})$ playing the roles of v, T to obtain sets A_1 and E_{A_1} and a vertex $a_1 \in A_1$. Next apply Lemma 2.3 again with $x_2, T - (A_1 \cup Y \cup X \setminus \{x_2\})$ playing the roles of v, T to obtain A_2, E_{A_2} and $a_2 \in A_2$. Continue in this way. (To obtain B_i, E_{B_i} and $b_i \in B_i$ apply Lemma 2.4 instead of Lemma 2.3.)

Let

$$E_A := \bigcup_{i \in [6k]} E_{A_i}, \quad E_B := \bigcup_{i \in [6k]} E_{B_i} \quad \text{and} \quad E := E_A \cup E_B.$$

Note that

$$(3.1) \quad |E_A| \leq 6k \left(\frac{1}{2}\right)^{c-2} \hat{\delta}^-(T) \leq \frac{\hat{\delta}^-(T)}{20k} \quad \text{and} \quad |E_B| \leq \frac{\hat{\delta}^+(T)}{20k}$$

by our choice of c . Moreover, we may assume that $|E_A| \leq |E_B|$. (The case $|E_A| > |E_B|$ follows by a symmetric argument.) In particular, this implies that

$$(3.2) \quad |E| \leq |E_A| + |E_B| \leq 2|E_B| \leq \frac{\hat{\delta}^+(T)}{10k}.$$

In our proof of Theorem 1.1 we will iteratively colour the vertices of T with colours α and β , and at each step V_α will consist of all vertices of colour α and V_β is defined similarly. (If at some stage a vertex of T is coloured, we will never change its colour again.) At the end of our argument, every vertex of T will be coloured either with α or with β , i.e. V_α, V_β will form

a partition of $V(T)$. Our aim is to colour the vertices in such a way that we can take $V_1 := V_\alpha$ and $V_2 := V_\beta$.

We say a path P is *alternating* if the colour of the vertices on P alternates as we move along P . P is *monochromatic* if all vertices of P have the same colour.

At each step and for each $\gamma \in \{\alpha, \beta\}$, we call a vertex $v \in V_\gamma$ *forwards-safe* if for any set $F \not\ni v$ of at most $k-1$ vertices, there is a directed monochromatic path (possibly of length 0) in $T[V_\gamma \setminus F]$ from v to $V(T) \setminus (D \cup E_B \cup F)$. Similarly, we say that $v \in V_\gamma$ is *backwards-safe* if for any set $F \not\ni v$ of at most $k-1$ vertices, there is a directed monochromatic path (possibly of length 0) in $T[V_\gamma \setminus F]$ from $V(T) \setminus (D \cup E_A \cup F)$ to v .

We call a vertex $v \in V_\gamma$ *alternating-forwards-safe* if for any set $F \not\ni v$ of at most $k-1$ vertices, there is a directed alternating path (possibly of length 0) in $T-F$ from v to $V(T) \setminus (D \cup E_B \cup F)$. Similarly, we say that $v \in V_\gamma$ is *alternating-backwards-safe* if for any set $F \not\ni v$ of at most $k-1$ vertices, there is a directed alternating path (possibly of length 0) in $T-F$ from $V(T) \setminus (D \cup E_A \cup F)$ to v .

We say that a vertex v is *safe* if it is safe in all four respects.

In our proof of Theorem 1.1 we will often use the following properties (for each $\{\gamma, \delta\} = \{\alpha, \beta\}$), which follow immediately from the definitions:

- (S1) all coloured vertices in $V(T) \setminus (D \cup E)$ are safe,
- (S2) all coloured vertices in $V(T) \setminus (D \cup E_B)$ are forwards-safe as well as alternating-forwards-safe and all coloured vertices in $V(T) \setminus (D \cup E_A)$ are backwards-safe as well as alternating-backwards-safe,
- (S3) if $v \in V_\gamma$ has at least k forwards-safe out-neighbours of colour γ then v itself is forwards-safe, the analogue holds if v has at least k backwards-safe in-neighbours of colour γ ,
- (S4) if $v \in V_\gamma$ has at least k alternating-forwards-safe out-neighbours of colour δ with $\delta \neq \gamma$ then v itself is alternating-forwards-safe, the analogue holds if v has at least k alternating-backwards-safe in-neighbours of colour δ ,
- (S5) if $v \in V_\gamma$ is safe and in the next step we colour some more (previously uncoloured) vertices then v is still safe.

In what follows, by a (partial) colouring of the vertices of T we always mean a colouring with colours α and β in which all the vertices in

$$D_1 := \bigcup_{i \in [k]} (A_i \cup B_i) \cup \bigcup_{i \in [3k+1, 5k]} (A_i \setminus \{a_i\}) \cup \bigcup_{i \in [3k+1, 4k] \cup [5k+1, 6k]} (B_i \setminus \{b_i\}) \\ \cup \{a_i \mid i \in [2k+1, 3k] \cup [5k+1, 6k]\} \cup \{b_i \mid i \in [2k+1, 3k] \cup [4k+1, 5k]\}$$

are coloured α , and all the vertices in $D_2 := D \setminus D_1$ are coloured β .

Claim 0: *Suppose that there are paths P_1, \dots, P_{6k} of T satisfying the following properties:*

- for each $i \in [6k]$ the path P_i joins b_i to a_i ,
- the paths P_1, \dots, P_{6k} are disjoint from each other and meet D only in their endvertices.

Suppose that we have coloured all vertices of T such that

- every vertex in $D_1 \cup V(P_1) \cup \dots \cup V(P_k)$ is coloured α ,
- every vertex in $D_2 \cup V(P_{k+1}) \cup \dots \cup V(P_{2k})$ is coloured β ,
- P_{2k+1}, \dots, P_{6k} are alternating,
- every vertex is safe.

Then the sets $V_1 := V_\alpha$ and $V_2 := V_\beta$ form a partition of $V(T)$ as required in Theorem 1.1.

Note that the conditions of Claim 0 imply that P_i must be an even path for $i \in [2k + 1, 4k]$ and an odd path for $i \in [4k + 1, 6k]$ (see Figure 1).




Figure 1: Colour patterns of the paths P_1, \dots, P_{6k} in the case when $k = 1$.

To prove Claim 0, we first show that $T[V_\alpha]$ is strongly k -connected. So consider any set F of at most $k - 1$ vertices and any two vertices $x, y \in V_\alpha \setminus F$. We need to check that $T[V_\alpha \setminus F]$ contains a path from x to y . Since x is forwards-safe there exists a path Q_x in $T[V_\alpha \setminus F]$ from x to some vertex $x' \in V_\alpha \setminus (D \cup E_B \cup F)$. Similarly, since y is backwards-safe there exists a path Q_y in $T[V_\alpha \setminus F]$ from some vertex $y' \in V_\alpha \setminus (D \cup E_A \cup F)$ to y . Let $i \in [k]$ be such that F avoids $A_i \cup V(P_i) \cup B_i$. Since $x' \notin D \cup E_B$, (D4) implies that x' sends an edge to B_i . Similarly, since $y' \notin D \cup E_A$, (D3) implies that y' receives an edge from A_i . Altogether this implies that $T[V(Q_x) \cup V(Q_y) \cup A_i \cup V(P_i) \cup B_i] \subseteq T[V_\alpha \setminus F]$ contains a path from x to y , as desired.

A similar argument shows that V_β is strongly k -connected too. It remains to show that $T[V_\alpha, V_\beta]$ is strongly k -connected. Consider any set F of at most $k - 1$ vertices and any two vertices $x, y \in V(T) \setminus F$. We will show that there is an alternating path between x and y avoiding F . Since x is alternating-forwards-safe there exists an alternating path Q_x in $T[V_\alpha, V_\beta] - F$ from x to some vertex $x' \in V(T) \setminus (D \cup E_B \cup F)$. Similarly, since y is alternating-backwards-safe there exists a path Q_y in $T[V_\alpha, V_\beta] - F$ from some vertex $y' \in V(T) \setminus (D \cup E_A \cup F)$ to y . We now choose an index i as follows:

- If $x', y' \in V_\alpha$, let $i \in [2k + 1, 3k]$ be such that F avoids $A_i \cup V(P_i) \cup B_i$.
- If $x', y' \in V_\beta$, let $i \in [3k + 1, 4k]$ be such that F avoids $A_i \cup V(P_i) \cup B_i$.
- If $x' \in V_\alpha$ and $y' \in V_\beta$, let $i \in [4k + 1, 5k]$ be such that F avoids $A_i \cup V(P_i) \cup B_i$.
- If $x' \in V_\beta$ and $y' \in V_\alpha$, let $i \in [5k + 1, 6k]$ be such that F avoids $A_i \cup V(P_i) \cup B_i$.

Since $x' \notin D \cup E_B$, (D4) implies that x' sends an edge to $B_i \setminus \{b_i\}$. Similarly, since $y' \notin D \cup E_A$, (D3) implies that y' receives an edge from $A_i \setminus \{a_i\}$. Altogether this implies that $T[V(Q_x) \cup V(Q_y) \cup A_i \cup V(P_i) \cup B_i] \subseteq T - F$ contains an alternating path from x to y , as desired. This completes the proof of Claim 0.

Claim 1: Consider a partial colouring of $V(T)$ and let C denote the set of previously coloured vertices. (So $D \subseteq C$.) Let $Z \subseteq V(T) \setminus (X \cup Y)$ and $N \subseteq V(T) \setminus Z$ and suppose that $9k^2|Z| + |C \cup N| \leq 5 \cdot 10^8 k^6 \log(2k)$. Then for every colouring of the vertices in $Z \setminus C$ there is a set $Z' \subseteq V(T) \setminus (Z \cup N \cup C)$ and a colouring of the vertices in Z' such that every vertex in $Z \cup Z'$ is safe and $|Z \cup Z'| \leq 9k^2|Z|$.

To prove Claim 1, note that the strong $10^9 k^6 \log(2k)$ -connectivity of T implies that $\delta^0(T) \geq 10^9 k^6 \log(2k)$. Hence

$$(3.3) \quad \hat{\delta}^-(T) - 5k|E_A| \stackrel{(3.1)}{\geq} \frac{\hat{\delta}^-(T)}{2} \geq \frac{\delta^0(T)}{2} \geq 5 \cdot 10^8 k^6 \log(2k),$$

and similarly

$$(3.4) \quad \hat{\delta}^+(T) - 5k|E| \stackrel{(3.2)}{\geq} \frac{\hat{\delta}^+(T)}{2} \geq 5 \cdot 10^8 k^6 \log(2k).$$

Consider any colouring of $Z \setminus C$. For each vertex $z \in Z$ in turn we greedily choose $2k$ uncoloured in-neighbours outside $N \cup E_A$, and colour k of them α and the remaining k by β . (We do not modify C in this process.) To see that we can choose all these vertices to be distinct from each other, note that the total number of vertices we wish to choose is $2k|Z|$ and

$$|C \cup N \cup Z| + 2k|Z| \leq 5 \cdot 10^8 k^6 \log(2k) \stackrel{(3.3)}{\leq} \hat{\delta}^-(T) - |E_A|.$$

For each vertex in Z as well as for each of the $2k|Z|$ vertices that we coloured in the previous step in turn, we greedily choose $2k$ uncoloured out-neighbours outside $N \cup E$, and colour k of them by α and the remaining k by β . To see that we can choose all these vertices to be distinct from each other, note that the total number of vertices we wish to choose is $2k(1+2k)|Z|$ and

$$|C \cup N \cup Z| + 2k|Z| + 2k(1+2k)|Z| \leq |C \cup N| + 9k^2|Z| \leq 5 \cdot 10^8 k^6 \log(2k) \stackrel{(3.4)}{\leq} \hat{\delta}^+(T) - |E|.$$

Let Z' be the set of vertices outside $C \cup Z$ that we coloured. Then $Z' \cap N = \emptyset$. Moreover, by (S1) and (S2) all vertices in Z' are safe. Together with (S3) and (S4) this in turn implies that all vertices in Z are safe. This completes the proof of Claim 1.

Recall that we have already coloured all the vertices in D_1 by α and all the vertices in D_2 by β . Step by step, we will now colour further vertices of T . Our final aim is to arrive at a colouring of $V(T)$ which is as described in Claim 0. The first step is to colour some more vertices in order to achieve that all the coloured vertices are safe. In what follows, when saying that we colour some additional vertices we always mean that these vertices are uncoloured so far.

Claim 2: *We can colour some additional vertices of T in such a way that every coloured vertex is safe and the set C_1 consisting of all vertices coloured so far satisfies $|C_1| \leq 1500k^4$.*

To prove Claim 2, for every $v \in \{x_1, \dots, x_{6k}, y_1, \dots, y_{6k}\}$ in turn, we greedily choose $2k$ uncoloured in-neighbours and $2k$ uncoloured out-neighbours, all distinct from each other, and colour k of the in-neighbours and k of the out-neighbours by α and the remaining $2k$ in/out-neighbours by β .

Let Z^* denote the set of $4k \cdot 12k = 48k^2$ new vertices we just coloured and let $Z := Z^* \cup (D \setminus (X \cup Y))$. Then $|Z| \leq |Z^*| + |D| \leq 48k^2 + c \cdot 12k \leq 156k^2$. Apply Claim 1 with $N := \emptyset$ to find a set Z' of uncoloured vertices and a colouring of these vertices such that all the vertices in $Z \cup Z'$ are safe and $|Z \cup Z'| \leq 9k^2 \cdot |Z| \leq 1500k^4$. Our choice of Z^* and (S3), (S4) together now imply that the vertices in $X \cup Y$ are safe as well. This completes the proof of Claim 2.

Suppose that P is a path whose endvertices are already coloured, but whose internal vertices are still uncoloured. We say that we colour (the internal vertices of) P in an *alternating manner consistent with its endvertices* if the colouring results in an alternating path. (So for example, if the endvertices of P have the same colour, then P needs to be an even path.)

Claim 3: *There are paths P_1, P_2, \dots, P_{6k} of T satisfying the following properties:*

- (i) for each $i \in [6k]$, the path P_i joins b_i to a_i ,
- (ii) the paths P_1, \dots, P_{6k} are disjoint from each other and meet C_1 only in their endvertices,
- (iii) we can colour the internal vertices of P_1, \dots, P_k by α , the internal vertices of P_{k+1}, \dots, P_{2k} by β and the internal vertices of P_{2k+1}, \dots, P_{6k} in an alternating manner consistent with their endvertices and can colour some additional vertices such that the set C_4 of all coloured vertices satisfies the following properties:
 - (a) all vertices in C_4 are safe,
 - (b) there is a set $C^0 \subseteq C_4$ such that the number of coloured vertices outside C^0 is at most $3 \cdot 10^7 k^6 \log(2k)$,
 - (c) every vertex outside C_4 which has an in-neighbour in C^0 has at least k in-neighbours of each colour, and every vertex outside C_4 which has an out-neighbour in C^0 has at least k out-neighbours of each colour.

We will prove Claim 3 via a sequence of subclaims. For $i \in [6k]$ we define an i -path to be a directed path from the sink b_i of B_i to the source a_i of A_i whose internal vertices lie outside C_1 . Ideally, we would like to find disjoint i -paths P_i (one for each $i \in [6k]$) such that the following properties hold:

- (1) we can colour all the internal vertices of P_1, \dots, P_k by α , the internal vertices of P_{k+1}, \dots, P_{2k} by β and the internal vertices of P_{2k+1}, \dots, P_{6k} in an alternating manner consistent with their endvertices,
- (2) by colouring some additional vertices we can achieve that all coloured vertices are safe.

For each $i \in [6k]$ we will first try to find a short i -path P_i such that all these i -paths are disjoint and such that for each $i \in [2k+1, 6k]$ the length of the path P_i has the correct parity in order to ensure that the internal vertices of P_i can be coloured in an alternating manner consistent with the endvertices of P_i (so P_i needs to be even for $i \in [2k+1, 4k]$ and odd for $i \in [4k+1, 6k]$). We will then colour the vertices on these short i -paths as well as some additional vertices such that (1) and (2) are satisfied for the set I_{short} of all indices i for which we have been able to choose a short i -path (see Claim 3.1). This provides some of the paths required in Claim 3. To find the remaining paths, for all $i \notin I_{short}$ we will choose $10^5 k^4 \log(2k)$ i -paths $Q_{i,1}, \dots, Q_{i,10^5 k^4 \log(2k)}$ such that all these paths are internally disjoint from each other. For each $i \notin I_{short}$ with $i \in [2k]$ there will be three distinct indices $j_{i,1}, j_{i,2}, j_{i,3} \in [10^5 k^4 \log(2k)]$ such that the path P_i required in Claim 3 will consist of an initial segment of $Q_{i,j_{i,1}}$, a middle segment of $Q_{i,j_{i,2}}$, a final segment of $Q_{i,j_{i,3}}$ as well as two edges joining these three segments. Similarly, for each $i \notin I_{short}$ with $i \in [2k+1, 6k]$ the path P_i required in Claim 3 will either be one of the $Q_{i,j}$ or will consist of an initial segment of $Q_{i,j_{i,1}}$ and a final segment of $Q_{i,j_{i,2}}$ as well as an edge joining these two segments.

We will now choose the short i -paths. Let $\mathcal{P}_{short}^{correct}$ be a collection of i -paths satisfying the following properties:

- (P1) for each $i \in [6k]$, $\mathcal{P}_{short}^{correct}$ contains at most one i -path,
- (P2) all the paths in $\mathcal{P}_{short}^{correct}$ are disjoint from each other,
- (P3) each path has length at most $10k + 10$,
- (P4) for each $i \in [2k+1, 6k]$ for which $\mathcal{P}_{short}^{correct}$ contains an i -path, this path P_i has the correct parity, meaning that P_i is even if $i \in [2k+1, 4k]$ and odd if $i \in [4k+1, 6k]$,
- (P5) subject to the above conditions, $|\mathcal{P}_{short}^{correct}|$ is as large as possible.

Let $I_{short}^{correct}$ be the set of all those indices $i \in [6k]$ for which $\mathcal{P}_{short}^{correct}$ contains an i -path, and let P_i denote this i -path. Let $V_{short}^{correct}$ be the set of all internal vertices of the P_i for all $i \in I_{short}^{correct}$.

Moreover, set $I_{long} := [6k] \setminus I_{short}^{correct}$. Recall that the definition of an i -path implies that all the vertices in $V_{short}^{correct}$ are uncoloured so far (i.e. $V_{short}^{correct} \cap C_1 = \emptyset$).

Claim 3.1: *We may colour all vertices in $V_{short}^{correct}$ as well as some additional vertices of T such that the following properties hold:*

- (i) *for each $i \in I_{short}^{correct}$, all the vertices on P_i are coloured α if $i \in [k]$ and β if $i \in [k+1, 2k]$,*
- (ii) *for each $i \in I_{short}^{correct} \setminus [2k]$, P_i is coloured in an alternating manner consistent with its endvertices,*
- (iii) *the set C_2 consisting of all vertices coloured so far has size $|C_2| \leq 4000k^4$ and all vertices in C_2 are safe,*
- (iv) *for each $i \in I_{long}$, any i -path whose internal vertices lie in $V(T) \setminus C_2$ is either $b_i a_i$ or has length at least $10k + 10$.*

To prove Claim 3.1, consider any $i \in I_{short}^{correct}$ and colour all internal vertices of P_i by α if $i \in [k]$, by β if $i \in [k+1, 2k]$, and in an alternating manner consistent with the endvertices of P_i if $i \in [2k+1, 6k]$ (this is possible by (P4)). Note that $|V_{short}^{correct}| \leq 6k(10k+9) \leq 120k^2$. Together with Claim 1 (applied with $N := \emptyset$ and $Z := V_{short}^{correct}$) and the fact that $|C_1| \leq 1500k^4$ by Claim 2 this implies Claim 3.1(i)–(iii), with room to spare in (iii). Indeed, the set C'_2 of vertices coloured so far has size $|C'_2| \leq 3000k^4$.

We will now colour some additional vertices to ensure that (iv) holds too. Consider any $i \in I_{long}$. If there exists an i -path P whose internal vertices lie in $V(T) \setminus C'_2$ and whose length is at most $10k+9$, then P must have incorrect parity, i.e. P is odd if $i \in [2k+1, 4k]$ and even if $i \in [4k+1, 6k]$. Note that there cannot be two such i -paths of length at least two which are internally disjoint from each other. Indeed, if $P = v_1 \dots v_a$ and $P' = v'_1 \dots v'_{a'}$ are two such i -paths which are internally disjoint, then we may assume that $v_2 v'_2 \in E(T)$ and so $v_1 v_2 v'_2 v'_3 \dots v'_a$ is an i -path of length at most $10k+10$ with the correct parity which is disjoint from all the other paths in $\mathcal{P}_{short}^{correct}$, a contradiction to (P5).

Let $\mathcal{P}_{short}^{incorrect}$ be a collection of i -paths (with $i \in I_{long}$) whose internal vertices lie in $V(T) \setminus C'_2$ and whose length is at least two and at most $10k+9$, such that all these paths are disjoint from each other and, subject to these properties, such that $|\mathcal{P}_{short}^{incorrect}|$ is as large as possible. Let $V_{short}^{incorrect}$ be the set of all internal vertices on these paths. Thus $|V_{short}^{incorrect}| \leq 4k \cdot (10k+8) \leq 100k^2$. Colour all vertices in $V_{short}^{incorrect}$ with α and apply Claim 1 again (with $N := \emptyset$ and $Z := V_{short}^{incorrect}$). Then the set C_2 of all vertices coloured so far satisfies $|C_2| \leq 3000k^2 + 9k^2 \cdot 100k^2 \leq 4000k^4$, so (iii) still holds. Moreover, now (iv) holds too. (Indeed, recall that for each $i \in I_{long}$ any i -path P whose length is at least two and at most $10k+9$ must have the incorrect parity, and so P cannot be internally disjoint from all the paths in $\mathcal{P}_{short}^{incorrect}$ (by the maximality of $\mathcal{P}_{short}^{incorrect}$ and our argument in the previous paragraph).) This completes the proof of Claim 3.1.

Claim 3.1(iii) implies that all uncoloured vertices together with the a_i and b_i for all $i \in I_{long}$ induce a strongly $(7 \cdot 452 \cdot 10^5 k^5 \log(2k))$ -connected subtournament T' of T (with some room to spare). Theorem 2.2 implies that T' is $7 \cdot 10^5 k^5 \log(2k)$ -linked. Together with Proposition 2.1 this implies that for each $i \in I_{long}$ we can find $10^5 k^4 \log(2k)$ i -paths in T' such that all these $10^5 k^4 \log(2k) |I_{long}|$ paths have length at least two and are internally disjoint from each other and such that the internal vertices on all these paths lie outside C_2 . We choose this collection of $10^5 k^4 \log(2k) |I_{long}|$ paths such that the set V_{long} of all internal vertices on these paths is as small as possible. For all $i \in I_{long}$ and all $j \in [10^5 k^4 \log(2k)]$, let $Q_{i,j}$ denote the j th i -path we

chose. Write $Q_{i,j} = q_{i,j}^0 q_{i,j}^1 \dots q_{i,j}^{|Q_{i,j}|}$, so that $q_{i,j}^0$ is b_i and $q_{i,j}^{|Q_{i,j}|}$ is a_i . Claim 3.1(iv) implies that each $Q_{i,j}$ must have length at least $10k + 10$. Moreover, the minimality of $|V_{long}|$ implies the following:

- (Q1) the interior of each $Q_{i,j}$ induces a backwards-transitive path,
- (Q2) if $v \in V(T) \setminus (C_2 \cup V_{long})$ is an out-neighbour of $q_{i,j}^s$, then v is also an out-neighbour of $q_{i,j}^{s'}$ for all $s' \geq s + 3$,
- (Q3) if $v \in V(T) \setminus (C_2 \cup V_{long})$ is an in-neighbour of $q_{i,j}^s$, then v is also an in-neighbour of $q_{i,j}^{s'}$ for all $s' \leq s - 3$.

Let $\text{int}(Q_{i,j}) := q_{i,j}^1 \dots q_{i,j}^{|Q_{i,j}|-1}$ denote the interior of $Q_{i,j}$. Let $Q_{i,j}^1, \dots, Q_{i,j}^9$ be disjoint segments of $\text{int}(Q_{i,j})$ such that $\text{int}(Q_{i,j}) = Q_{i,j}^1 \dots Q_{i,j}^9$, $|Q_{i,j}^1| = |Q_{i,j}^2| = |Q_{i,j}^8| = |Q_{i,j}^9| = k$, $|Q_{i,j}^3| = |Q_{i,j}^7| = k + 2$ and $|Q_{i,j}^4| = |Q_{i,j}^6| = 2k + 2$. So $|Q_{i,j}^5| = |Q_{i,j}| - 10k - 8 \geq 3$. We let

$$Q_{i,j}^0 := Q_{i,j}^1 \cup Q_{i,j}^2 \cup Q_{i,j}^3 \cup Q_{i,j}^7 \cup Q_{i,j}^8 \cup Q_{i,j}^9$$

and write

$$V_{long}^0 := \bigcup_{(i,j) \in I_{long} \times [10^5 k^4 \log(2k)]} V(Q_{i,j}^0) \quad \text{and} \quad \bar{V}_{long} := \bigcup_{(i,j) \in I_{long} \times [10^5 k^4 \log(2k)]} V(Q_{i,j}^0 \cup Q_{i,j}^4 \cup Q_{i,j}^6).$$

Thus $V_{long}^0 \subseteq \bar{V}_{long} \subseteq V_{long}$ and

$$|V_{long}^0| \leq |\bar{V}_{long}| \leq (10k + 8) \cdot 6k \cdot 10^5 k^4 \log(2k) \leq 2 \cdot 10^7 k^6 \log(2k).$$

Claim 3.2: *There exist disjoint index sets $I_{R,\alpha}, I_{R,\beta} \subseteq I_{long} \times [10^5 k^4 \log(2k)]$ such that, writing*

$$R_\alpha := \bigcup_{(i,j) \in I_{R,\alpha}} V(Q_{i,j}^0) \quad \text{and} \quad R_\beta := \bigcup_{(i,j) \in I_{R,\beta}} V(Q_{i,j}^0),$$

for each $(i, j) \in I_{long} \times [10^5 k^4 \log(2k)]$ every vertex in $V(Q_{i,j}^0) \setminus (R_\alpha \cup R_\beta)$ has at least k in-neighbours and at least k out-neighbours in each of R_α and R_β . Also $|I_{R,\alpha}|, |I_{R,\beta}| \leq 100k \log(2k)$ and $|R_\alpha|, |R_\beta| \leq 1000k^2 \log(2k)$.

To prove Claim 3.2, apply Proposition 2.5 to $T[V_{long}^0]$ to find a set $Z_\alpha \subseteq V_{long}^0$ with $|Z_\alpha| \leq 3k \log |V_{long}^0| \leq 100k \log(2k)$ and such that every vertex in $V_{long}^0 \setminus Z_\alpha$ has at least k out-neighbours and k in-neighbours in Z_α . Let $I_{R,\alpha} := \{(i, j) : V(Q_{i,j}^0) \cap Z_\alpha \neq \emptyset\}$ and $I' := (I_{long} \times [10^5 k^4 \log(2k)]) \setminus I_{R,\alpha}$. We now consider $W := \bigcup_{(i,j) \in I'} V(Q_{i,j}^0)$. By Proposition 2.5 applied to $T[W]$, there exists a set $Z_\beta \subseteq W$ with $|Z_\beta| \leq 3k \log |W| \leq 100k \log(2k)$ and such that every vertex in $W \setminus Z_\beta$ has at least k out-neighbours and in-neighbours in Z_β . Let $I_{R,\beta} := \{(i, j) \in I' : V(Q_{i,j}^0) \cap Z_\beta \neq \emptyset\}$.

Let R_α and R_β be as defined in the statement of Claim 3.2. Then by definition of $I_{R,\alpha}$ and $I_{R,\beta}$, for each $(i, j) \in I_{long} \times [10^5 k^4 \log(2k)]$ every vertex in $V(Q_{i,j}^0) \setminus (R_\alpha \cup R_\beta)$ has at least k in-neighbours and at least k out-neighbours in each of R_α and R_β . Also $|R_\alpha|, |R_\beta| \leq (6k + 4) \cdot 100k \log(2k) \leq 1000k^2 \log(2k)$. This completes the proof of Claim 3.2.

Let $I_R := I_{R,\alpha} \cup I_{R,\beta}$, $R := R_\alpha \cup R_\beta$ and

$$R^{4,6} := \bigcup_{(i,j) \in I_R} V(Q_{i,j}^4 \cup Q_{i,j}^6).$$

Claim 3.3: *We may colour all vertices in $R_\alpha \cup R_\beta \cup R^{4,6}$ as well as some additional vertices lying outside \bar{V}_{long} such that*

- (i) *all vertices in R_α are coloured α , all vertices in R_β are coloured β ,*
- (ii) *for each $(i, j) \in I_R$ and each $s \in \{4, 6\}$, $Q_{i,j}^s$ is an alternating path,*
- (iii) *all coloured vertices are safe,*
- (iv) *the set C_3 consisting of all vertices coloured so far has size $|C_3| \leq 4 \cdot 10^4 k^4 \log(2k)$.*

To prove Claim 3.3, colour the vertices in $R_\alpha \cup R_\beta \cup R^{4,6}$ such that (i) and (ii) hold. Apply Claim 1 with C_2 , $R_\alpha \cup R_\beta \cup R^{4,6}$, $\bar{V}_{long} \setminus (R_\alpha \cup R_\beta \cup R^{4,6})$ playing the roles of C , Z , N to obtain a set $Z' \subseteq V(T) \setminus (\bar{V}_{long} \cup C_2)$ and a colouring of the vertices in Z' such that every vertex in $R_\alpha \cup R_\beta \cup R^{4,6} \cup Z'$ is safe and

$$\begin{aligned} |C_3| &\leq |C_2| + |R_\alpha \cup R_\beta \cup R^{4,6} \cup Z'| \leq 4000k^4 + 9k^2 \cdot (2 \cdot 1000k^2 \log(2k) + (4k + 4) \cdot 200k \log(2k)) \\ &\leq 4 \cdot 10^4 k^4 \log(2k). \end{aligned}$$

This completes the proof of Claim 3.3.

Claim 3.4: *For each $i \in I_{long}$ there is an i -path P_i such that the following properties hold:*

- (i) *P_i has no internal vertices in C_3 , and P_i and $P_{i'}$ are disjoint whenever $i \neq i'$,*
- (ii) *if $i \in I_{long} \cap [2k]$, then there exists three distinct indices $j_{i,1}, j_{i,2}, j_{i,3} \in [10^5 k^4 \log(2k)]$ such that $P_i = b_i Q_{i,j_{i,1}}^1 Q_{i,j_{i,1}}^2 q_{i,j_{i,1}}^{2k+1} Q_{i,j_{i,2}}^3 \cdots Q_{i,j_{i,2}}^7 q_{i,j_{i,2}}^{|Q_{i,j_{i,3}}| - 2k - 1} Q_{i,j_{i,3}}^8 Q_{i,j_{i,3}}^9 a_i$,*
- (iii) *if $i \in I_{long} \cap [2k + 1, 6k]$, then either $P_i = Q_{i,j_i}$ for some $j_i \in [10^5 k^4 \log(2k)]$ or there exist distinct $j_{i,1}, j_{i,2} \in [10^5 k^4 \log(2k)]$ such that $P_i = b_i Q_{i,j_{i,1}}^1 \cdots Q_{i,j_{i,1}}^4 q_{i,j_{i,1}}^{5k+5} Q_{i,j_{i,2}}^5 \cdots Q_{i,j_{i,2}}^9 a_i$,*
- (iv) *P_i is even if $i \in I_{long} \cap [2k + 1, 4k]$ and odd if $i \in I_{long} \cap [4k + 1, 6k]$.*

(Recall that $q_{i,j_{i,1}}^{2k+1}$ is the first vertex of $Q_{i,j_{i,1}}^3$, $q_{i,j_{i,3}}^{|Q_{i,j_{i,3}}| - 2k - 1}$ is the last vertex of $Q_{i,j_{i,3}}^7$ and $q_{i,j_{i,1}}^{5k+5}$ is the first vertex of $Q_{i,j_{i,1}}^5$.) To prove Claim 3.4, note that since $|C_3| \leq 4 \cdot 10^4 k^4 \log(2k) < 10^5 k^4 \log(2k) - 5$, for each $i \in I_{long}$ there are at least five paths $Q_{i,s_{i,1}}, Q_{i,s_{i,2}}, Q_{i,s_{i,3}}, Q_{i,s_{i,4}}, Q_{i,s_{i,5}}$ whose internal vertices avoid C_3 .

Suppose first that $i \in I_{long} \cap [2k]$. Consider the subtournament T_i of T spanned by $q_{i,s_{i,t}}^{|Q_{i,s_{i,t}}| - 2k - 1}$ for $t = 1, 2, 3, 4, 5$. T_i contains at least two vertices of out-degree at least two, assume they are $q_{i,s_{i,1}}^{|Q_{i,s_{i,1}}| - 2k - 1}, q_{i,s_{i,2}}^{|Q_{i,s_{i,2}}| - 2k - 1}$. We may also assume that $q_{i,s_{i,1}}^{2k+1}$ sends an edge to $q_{i,s_{i,2}}^{2k+1}$. Finally, since $q_{i,s_{i,2}}^{|Q_{i,s_{i,2}}| - 2k - 1}$ has at least two outneighbours in T_i , we may assume $q_{i,s_{i,2}}^{|Q_{i,s_{i,2}}| - 2k - 1}$ sends an edge to $q_{i,s_{i,3}}^{|Q_{i,s_{i,3}}| - 2k - 1}$. We set $j_{i,t} := s_{i,t}$ and let P_i be as described in Claim 3.4(ii), see Figure 2.

So suppose next that $i \in I_{long} \cap [2k + 1, 4k]$. If $Q_{i,s_{i,t}}$ is an even path for $t = 1$ or $t = 2$ we take it to be P_i . So suppose that these two paths are odd. We may assume that $q_{i,s_{i,1}}^{5k+5}$ sends an edge to $q_{i,s_{i,2}}^{5k+5}$. We set $j_{i,1} := s_{i,1}$ and $j_{i,2} := s_{i,2}$ and let P_i be as described in Claim 3.4(iii). If $i \in I_{long} \cap [4k + 1, 6k]$, we define P_i similarly, see Figure 3. This completes the proof of Claim 3.4.




Figure 2: Colour patterns of the paths $\text{int}(Q_{i,j'})$ with $i \in I_{\text{long}} \cap [k]$ in the case when $k = 1$ and $Q_{i,j}$ does not intersect C_3 . The thick arrows indicate $\text{int}(P_i)$.




Figure 3: Colour patterns of the paths $\text{int}(Q_{i,j'})$ with $i \in I_{\text{long}} \cap [4k + 1, 5k]$ in the case when $k = 1$, $Q_{i,j,t}$ is an even path for $t \in \{1, 2\}$ and $Q_{i,j}$ does not intersect C_3 . The thick arrows indicate $\text{int}(P_i)$.

We are now ready to prove Claim 3. For each $i \in I_{\text{long}}$, let P_i be as given by Claim 3.4. We will colour all those vertices on the paths $Q_{i,j}$ with $(i, j) \in I_{\text{long}} \times [10^5 k^4 \log(2k)]$ which are uncoloured so far as follows.

For each $i \in I_{\text{long}} \cap [2k]$, we colour all internal vertices of P_i by α if $i \leq k$ and by β if $i > k$. For each $i \in I_{\text{long}} \cap [k]$, we also colour all vertices in $(Q_{i,j}^1 \cup Q_{i,j}^9) \setminus (V(P_i) \cup R)$ by α and all vertices in $(Q_{i,j}^2 \cup Q_{i,j}^3 \cup Q_{i,j}^7 \cup Q_{i,j}^8) \setminus (V(P_i) \cup R)$ by β (for all $j \in [10^5 k^4 \log(2k)]$), see Figure 2. Similarly, for each $i \in I_{\text{long}} \cap [k + 1, 2k]$, we colour all vertices in $(Q_{i,j}^1 \cup Q_{i,j}^9) \setminus (V(P_i) \cup R)$ by β and all vertices in $(Q_{i,j}^2 \cup Q_{i,j}^3 \cup Q_{i,j}^7 \cup Q_{i,j}^8) \setminus (V(P_i) \cup R)$ by α . For each $i \in I_{\text{long}} \cap [2k]$, we colour all vertices in $(Q_{i,j}^4 \cup Q_{i,j}^6) \setminus (V(P_i) \cup R^{4,6})$ by α .

For each $i \in I_{\text{long}} \cap [2k + 1, 6k]$, we colour all internal vertices of P_i in an alternating manner consistent with the endvertices of P_i . (Claim 3.4(iv) ensures that this is possible.) For all $j \in [10^5 k^4 \log(2k)]$ we also colour all vertices in $Q_{i,j}^0 \cup Q_{i,j}^4 \cup Q_{i,j}^6 \setminus (V(P_i) \cup R \cup R^{4,6})$ in an alternating manner, see Figure 3. (That is, if $b_i = q_{i,j}^0$ is coloured α , we colour $q_{i,j}^s$ by α for all even numbers $s \leq 5k + 4$, and colour $q_{i,j}^s$ by β for all odd numbers $s \leq 5k + 4$. We colour each vertex x in $(Q_{i,j}^6 \cup \dots \cup Q_{i,j}^9) \setminus (V(P_i) \cup R \cup R^{4,6})$ in a similar way, depending on the colour of a_i and the distance of x to a_i in $Q_{i,j}$.)

Now all uncoloured vertices of V_{long} belong to $Q_{i,j}^5$ for some i, j . We let C^0 be the union of $V(Q_{i,j}^5)$ over all $(i, j) \in I_{\text{long}} \times [10^5 k^4 \log(2k)]$. We colour all uncoloured vertices in C^0 by α , and let C_4 denote the set consisting of all the vertices coloured so far. Note that $|C_4 \setminus C^0| \leq$

$|C_3| + |\overline{V}_{long}| \leq 4 \cdot 10^4 k^4 \log(2k) + 2 \cdot 10^7 k^6 \log(2k) \leq 3 \cdot 10^7 k^6 \log(2k)$. Together with Claim 3.1 this implies that parts (i), (ii) and (iii)(b) of Claim 3 hold.

We now show that all the vertices on the paths $Q_{i,j}$ are safe. Together with Claim 3.3(iii) this will imply that all vertices in C_4 are safe, i.e. Claim 3(iii)(a) will hold. Consider first any vertex $v \in V_{long}^0$. If $v \in R$, then v is safe by Claim 3.3(iii). If $v \notin R$, then by Claim 3.2 v has at least k out-neighbors and at least k in-neighbours in each of R_α and R_β , so it has k safe out-neighbours and k safe in-neighbours of each colour. Thus v is safe by (S3) and (S4). So all the the vertices in V_{long}^0 are safe.

Note that if $(i, j) \notin I_R$, then $V(Q_{i,j}^1 \cup Q_{i,j}^2 \cup Q_{i,j}^3) \setminus \{q_{i,j}^{3k+2}\}$ contains at least k vertices of each colour and so does $V(Q_{i,j}^7 \cup Q_{i,j}^8 \cup Q_{i,j}^9) \setminus \{q_{i,j}^{|Q_{i,j}|-3k-2}\}$. (Recall that $q_{i,j}^{3k+2}$ is the final vertex of $Q_{i,j}^3$ and that $q_{i,j}^{|Q_{i,j}|-3k-2}$ is the initial vertex of $Q_{i,j}^7$.)

Now consider a vertex $v \in \overline{V}_{long} \setminus V_{long}^0$, and let i, j be such that $v \in V(Q_{i,j}^4 \cup Q_{i,j}^6)$. If $v \in R^{4,6}$, then v is safe by Claim 3.3(iii). If $v \notin R^{4,6}$, then $(i, j) \notin I_R$. But by (Q1) all vertices in $Q_{i,j}^1 \cup Q_{i,j}^2 \cup Q_{i,j}^3$ (apart from possibly the final vertex of $Q_{i,j}^3$) are out-neighbours of v , so v has k safe out-neighbours coloured α and k safe out-neighbours coloured β . Similarly, all vertices in $Q_{i,j}^7 \cup Q_{i,j}^8 \cup Q_{i,j}^9$ (apart from possibly the initial vertex of $Q_{i,j}^7$) are in-neighbours of v , so v has k safe in-neighbours coloured α and k safe in-neighbours coloured β . Hence v is safe.

Now consider any vertex $v \in V(Q_{i,j}^5)$. If $(i, j) \notin I_R$, a similar argument as above shows that v is safe. If $(i, j) \in I_R$, then by (Q1) all vertices in $Q_{i,j}^4$ (apart from possibly its final vertex) are out-neighbours of v , and all vertices in $Q_{i,j}^6$ (apart from possibly its initial vertex) are in-neighbours of v . Together with Claim 3.3(ii),(iii) this shows that v has k safe out-neighbours and k safe in-neighbours of each colour. So v is safe. This completes the proof of Claim 3(iii)(a).

To check Claim 3(iii)(c), note that if a vertex v outside C_4 has an out-neighbour in C^0 , then by (Q3) all vertices in $Q_{i,j}^1 \cup Q_{i,j}^2 \cup Q_{i,j}^3 \cup Q_{i,j}^4$ (apart from possibly the last two vertices of $Q_{i,j}^4$) are out-neighbours of v . Thus v has at least k out-neighbours of each colour. In a similar way one can use (Q2) to show that v also has k in-neighbours of each colour. This completes the proof of Claim 3.

Claim 4: *We can colour all uncoloured vertices in such a way that every vertex is safe.*

To prove Claim 4, we colour all the vertices outside C_4 one by one. We first deal with all vertices in $E_A \setminus C_4$ (see STEP 1), then we move to the vertices in $E_B \setminus C_4$ (see STEP 2). Finally, we colour all the remaining vertices (see STEP 3). We let $Z_A := \emptyset$. While dealing with each vertex in $E_A \setminus C_4$ in turn (i.e. during STEP 1), we will update Z_A . At each substep, Z_A will satisfy the following properties:

- (a) Z_A consists of coloured vertices and $Z_A \cap (C_4 \cup E_A) = \emptyset$,
- (b) every coloured vertex lies in $C_4 \cup E_A \cup Z_A$,
- (c) $|Z_A| \leq 2k|E_A|$.

STEP 1. *We can colour all vertices in $E_A \setminus C_4$ as well as some set Z_A of additional vertices in such a way that all the vertices in $E_A \setminus C_4$ are backwards-safe and alternating-backwards-safe and Z_A satisfies (a)–(c).*

Consider each vertex $v \in E_A \setminus C_4$ in turn. Suppose first that v has $2k$ uncoloured in-neighbours v_1, v_2, \dots, v_{2k} outside E_A . We colour k of them by α and k of them by β and replace Z_A by $Z_A \cup \{v_1, v_2, \dots, v_{2k}\}$. We also colour v with α . Note that (S2) implies that v_1, v_2, \dots, v_{2k} are

backwards-safe and alternating-backwards-safe. Together with (S3) and (S4) this shows that v is backwards-safe and alternating-backwards-safe.

So suppose that v has less than $2k$ uncoloured in-neighbours outside E_A . Recall from Claim 3(iii)(b) that at most $3 \cdot 10^7 k^6 \log(2k)$ vertices in C_4 lie outside the set C^0 . Together with (3.3) and (c) this shows that

$$\hat{\delta}^-(T) - |E_A \cup Z_A| \geq \hat{\delta}^-(T) - 3k|E_A| \geq 5 \cdot 10^8 k^6 \log(2k) \geq 2k + |C_4 \setminus C^0|.$$

Since all coloured vertices lie in $C_4 \cup E_A \cup Z_A$, this implies that v has an in-neighbour in C^0 . But now Claim 3(iii)(c) implies that v has k in-neighbours of colour α and k in-neighbours of colour β in C_4 . Since all the vertices in C_4 are safe, this implies that v becomes backwards-safe and alternating-backwards-safe by colouring v with α .

Note that we add at most $2k$ vertices to Z_A for each vertex $v \in E_A \setminus C_4$. So at the end of STEP 1, we will still have that $|Z_A| \leq 2k|E_A|$. Since by (S2) every vertex outside E_B is forwards-safe and alternating-forwards-safe, after STEP 1, all vertices in $E_A \setminus E_B$ will be safe, while the vertices in $E_A \cap E_B$ might only be backwards-safe and alternating-backwards-safe.

Let $Z_B := \emptyset$. While dealing with each vertex in $E_B \setminus C_4$ in turn during STEP 2, we will update Z_B . At each substep, Z_B will satisfy the following properties (where $Z := Z_A \cup Z_B$):

- (a') Z_B consists of coloured vertices and $Z_B \cap (C_4 \cup E \cup Z_A) = \emptyset$,
- (b') every coloured vertex lies in $C_4 \cup E \cup Z$,
- (c') $|Z_B| \leq 2k|E_B|$ and so $|Z| \leq 4k|E|$.

STEP 2. *We can colour all uncoloured vertices in $E_B \setminus C_4$ as well as some set Z_B of additional vertices in such a way that all the vertices in $E_B \setminus C_4$ are safe and Z_B satisfies (a')–(c').*

Consider each vertex $v \in E_B \setminus C_4$ in turn. If $v \notin E_A$, then v is backwards-safe and alternating-backwards-safe by (S2). If $v \in E_A$, then by STEP 1 v is also backwards-safe and alternating-backwards-safe.

Suppose first that v has $2k$ uncoloured out-neighbours v_1, v_2, \dots, v_{2k} outside E . We colour k of them by α and k of them by β . We replace Z_B by $Z_B \cup \{v_1, v_2, \dots, v_{2k}\}$. If v is uncoloured, we colour v with α . Then (S2)–(S4) together imply that v becomes safe.

So suppose that v has less than $2k$ uncoloured out-neighbours outside E . Note that

$$\hat{\delta}^+(T) - |E \cup Z| \geq \hat{\delta}^+(T) - 5k|E| \geq 5 \cdot 10^8 k^6 \log(2k) \geq 2k + |C_4 \setminus C^0|$$

by (3.4), (c') and Claim 3(iii)(b). Since all coloured vertices lie in $C_4 \cup E \cup Z$, this implies that v has an out-neighbour in C^0 . But now Claim 3(iii)(c) implies that v has k out-neighbours of colour α and k out-neighbours of colour β in C_4 . Since all the vertices in C_4 are safe, this implies that v becomes safe by colouring v with α (in case v is still uncoloured).

Note that we add at most $2k$ vertices to Z_B for each vertex in $E_B \setminus C_4$. Thus we always have that $|Z_B| \leq 2k|E_B|$ and so $|Z| \leq 4k|E|$. After STEP 2, all vertices in $C_4 \cup E$ are safe.

STEP 3. *By colouring all the remaining uncoloured vertices with α , every vertex becomes safe.*

This follows immediately from (S2).

This completes the proof of Claim 4 and thus of Theorem 1.1. □

4. SPANNING LINKEDNESS AND NON-SEPARATING SUBDIVISIONS

The following lemma generalizes a result of Thomassen [11]. Theorems 1.2 and 1.4 then both follow easily by an inductive application of Lemma 4.1.

Lemma 4.1. *Let k, d be nonnegative integers. Let x, y, z_1, \dots, z_d be any distinct vertices in a strongly $(k + d + 4)$ -connected tournament T and let P be a shortest xy -path in $T - \{z_1, \dots, z_d\}$. Then $T - (V(P) \setminus X)$ is strongly k -connected for any (possibly empty) subset $X \subseteq \{x, y\}$.*

Proof. Write $P := x_0x_1 \dots x_m$ with $x = x_0$ and $y = x_m$. Note that P must be a backwards-transitive path. If P has length at most two, the result trivially holds. So suppose that P has length more than two. Note that in this case it suffices to show that $T - V(P)$ is strongly k -connected (otherwise we consider $x' \in \{x, x_1\}, y' \in \{y, x_{m-1}\}$ and proceed through the argument with x', y' playing the role of x, y). So suppose $T - V(P)$ is not strongly k -connected. Then there exist a partition of $V(T) \setminus V(P)$ into nonempty sets S, S_1, S_2 such that $|S| \leq k - 1$ and no vertex in S_2 sends an edge to S_1 . Since $T - (S \cup \{z_1, \dots, z_d\})$ is strongly 5-connected, there are five paths P_1, \dots, P_5 from S_2 to S_1 which are internally disjoint and do not intersect $S \cup \{z_1, \dots, z_d\}$. We may assume that the P_i are backwards-transitive. Moreover, the interior of each P_i is nonempty and is contained in $V(P)$. Altogether, this means that the intersection of P_i and $T[V(P)]$ is either a segment of P or a path of the form x_jx_ℓ with $j \geq \ell + 2$ or of the form $x_jx_{j+1}x_{j-1}$ or $x_jx_{j-2}x_{j-1}$. We let p be the largest number such that some P_i contains an edge ux_p from S_2 to x_p and we let q be the smallest number such that some P_i contains an edge x_qv from x_q to S_1 . Note that $p \geq q + 4$. Then the path obtained from P by deleting $x_{q+1}x_{q+2} \dots x_{p-1}$ and adding x_qvux_p is shorter than P , a contradiction. \square

Proof of Theorem 1.2. Write $D = \{w_1, \dots, w_d\}$. We proceed by induction on m . For $m = 1$, the assertion holds by Lemma 4.1 applied with $d - 2$ playing the role of d . Suppose that $m \geq 2$ and that the assertion holds for $m - 1$. Consider any edge $uv \in E(H)$. Without loss of generality, we may assume that $\phi(u) = w_1$ and $\phi(v) = w_2$. Then we apply Lemma 4.1 (with $d - 2$ playing the role of d) to find a w_1w_2 -path P whose interior does not intersect D and so that $T' := T - V(\text{int}(P))$ is strongly $(k + (m - 1)(d + 2))$ -connected. Now by the induction hypothesis, we can find a subdivision H_* of $H \setminus \{uv\}$ in T' which satisfies (i)–(iii) (with T' playing the role of T). Finally, let $H^* := H_* \cup \text{int}(P)$. Then H^* satisfies all requirements. \square

Proof of Theorem 1.4. We proceed by induction on k . For $k = 1$, the assertion was proven by Thomassen [8]. Assume that $k \geq 2$ and that the assertion holds for $k - 1$. Let $Z := \{x_1, \dots, x_{k-1}, y_1, \dots, y_{k-1}\}$ and let $X := \{x_k, y_k\} \cap Z$. We can now apply Lemma 4.1 with $d = |Z \setminus X|$ to find a x_ky_k -path P avoiding $Z \setminus X$ so that $T[W]$ is strongly $((k - 1)^2 + 3(k - 1))$ -connected, where $W := V(T) \setminus (V(P) \setminus X)$. Now by the induction hypothesis, we can find P_1, \dots, P_{k-1} in $T[W]$ so that P_i is a path from x_i to y_i and $W = \bigcup_{i=1}^{k-1} V(P_i)$. Let $P_k := P$. Then P_1, \dots, P_k are as desired. \square

5. ACKNOWLEDGEMENTS

We are grateful to Jørgen Bang-Jensen for helpful comments on an earlier version of this paper. Moreover, we thank the referees for a careful reading of the paper.

REFERENCES

- [1] P. Hajnal, Partition of graphs with condition on the connectivity and minimum degree, *Combinatorica* **3** (1983), 95–99.
- [2] D. Kühn, J. Lapinskas, D. Osthus and V. Patel, Proof of a conjecture of Thomassen on Hamilton cycles in highly connected tournaments, *Proc. London Math. Soc.* **109** (2014), 733–762.
- [3] D. Kühn and D. Osthus, Partitions of graphs with high minimum degree or connectivity, *J. Combinatorial Theory B* **88** (2003), 29–43.
- [4] D. Kühn, D. Osthus and T. Townsend, Proof of a tournament partition conjecture and an application to 1-factors with prescribed cycle lengths, *Combinatorica*, to appear.
- [5] A. Pokrovskiy, Highly linked tournaments, *J. Combinatorial Theory B* **115** (2015), 339–347.
- [6] A. Pokrovskiy, Edge disjoint Hamiltonian cycles in highly connected tournaments, preprint, arxiv:1406.7556.
- [7] A. Scott, Judicious partitions and related problems, *Surveys in Combinatorics 2005, London Math. Soc. Lecture Note Ser.* **327**, Cambridge Univ. Press, Cambridge, (2005) 95–117.
- [8] C. Thomassen, Hamiltonian-connected tournaments, *J. Combinatorial Theory B* **28** (1980), 142–163.
- [9] C. Thomassen, Graph decomposition with constraints on the connectivity and minimum degree, *J. Graph Theory* **7** (1983), 165–167.
- [10] C. Thomassen, Graph decomposition with applications to subdivisions and path systems modulo k , *J. Graph Theory* **7** (1983), 261–271.
- [11] C. Thomassen, Connectivity in tournaments in, *Graph Theory and Combinatorics, A volume in honour of Paul Erdős (B. Bollobás, ed.)*, Academic Press, London (1984), 305–313.
- [12] C. Thomassen, Configurations in graphs of large minimum degree, connectivity, or chromatic number, *Combinatorial Mathematics: Proceedings of the Third International Conference (New York, 1985)*, vol. 555 of *Ann. New York Acad. Sci.*, New York, 1989, 402–412.

Jaehoon Kim, Daniela Kühn, Deryk Osthus
School of Mathematics
University of Birmingham
Edgbaston
Birmingham
B15 2TT
UK

E-mail addresses: {kimJS, d.kuhn, d.osthus}@bham.ac.uk