

## Biosynthesis of mycobacterial arabinogalactan: identification of a novel alpha(1-3) arabinofuranosyltransferase

Birch, Helen; Alderwick, Luke; Bhatt, Apoorva; Rittmann, D; Krumbach, K; Singh, Albel; Bai, Y; Lowary, TL; Eggeling, L; Besra, Gurdyal

DOI:

[10.1111/j.1365-2958.2008.06354.x](https://doi.org/10.1111/j.1365-2958.2008.06354.x)

*Citation for published version (Harvard):*

Birch, H, Alderwick, L, Bhatt, A, Rittmann, D, Krumbach, K, Singh, A, Bai, Y, Lowary, TL, Eggeling, L & Besra, G 2008, 'Biosynthesis of mycobacterial arabinogalactan: identification of a novel alpha(1-3) arabinofuranosyltransferase', *Molecular Microbiology*, vol. 69, no. 5, pp. 1191-206.  
<https://doi.org/10.1111/j.1365-2958.2008.06354.x>

[Link to publication on Research at Birmingham portal](#)

### General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

**Biosynthesis of mycobacterial arabinogalactan: identification of a novel  
?(1?3) arabinofuranosyltransferase**

Journal:	<i>Molecular Microbiology</i>
Manuscript ID:	MMI-2008-07576.R1
Manuscript Type:	Research Article
Date Submitted by the Author:	12-May-2008
Complete List of Authors:	<p>Birch, Helen; University of Birmingham, School of Biosciences            Alderwick, Luke; University of Birmingham, School of Biosciences            Bhatt, Apoorva; University of Birmingham, School of Biosciences            Rittmann, Doris; Institute for Biotechnology, Research Centre            Juelich            Krumbach, Karin; Institute for Biotechnology, Research Centre            Juelich            Singh, Albel; University of Birmingham, School of Biosciences            Bai, Yu; University of Alberta, Alberta Ingenuity Centre for            Carbohydrate Science and Department of Chemistry            Lowary, Todd; University of Alberta, Alberta Ingenuity Centre for            Carbohydrate Science and Department of Chemistry            Eggeling, Lothar; Research Centre, Biotechnology            Besra, Gurdya; University of Birmingham, School of Biosciences</p>
Key Words:	Mycobacterium tuberculosis, Corynebacterium glutamicum, arabinogalactan, arabinofuranosyltransferase, cell wall

**Biosynthesis of mycobacterial arabinogalactan: identification of a novel  $\alpha(1\rightarrow3)$   
arabinofuranosyltransferase**

Helen L. Birch<sup>1‡</sup>, Luke J. Alderwick<sup>1‡</sup>, Apoorva Bhatt<sup>1</sup>, Doris Rittmann<sup>2</sup>, Karin Krumbach<sup>2</sup>,  
Albel Singh<sup>1</sup>, Yu Bai<sup>3</sup>, Todd L. Lowary<sup>3</sup>, Lothar Eggeling<sup>2</sup>, and Gurdyal S. Besra<sup>1\*</sup>

<sup>1</sup>School of Biosciences, University of Birmingham, Edgbaston, Birmingham, B15 2TT, UK

<sup>2</sup>Institute for Biotechnology 1, Research Centre Juelich, D-52425 Juelich, Germany

<sup>3</sup>Alberta Ingenuity Centre for Carbohydrate Science and Department of Chemistry, University of  
Alberta, Canada AB T6G 2G2

\*To whom correspondence should be addressed; Tel: (0)121 415 8125 ; Fax: (0)121 414 5925;

Email: [g.besra@bham.ac.uk](mailto:g.besra@bham.ac.uk)

‡These authors contributed equally to the work

**Key words:** *Corynebacterium glutamicum*, *Mycobacterium tuberculosis*, arabinogalactan,  
arabinofuranosyltransferase, cell wall

**Running Title:** Identification of a novel arabinofuranosyltransferase

## Summary

The cell wall mycolyl-arabinogalactan-peptidoglycan complex is essential in mycobacterial species, such as *Mycobacterium tuberculosis* and is the target of several anti-tubercular drugs. For instance, ethambutol targets arabinogalactan biosynthesis through inhibition of the arabinofuranosyltransferases Mt-EmbA and Mt-EmbB. A bioinformatics approach identified putative integral membrane proteins, MSMEG2785 in *Mycobacterium smegmatis*, Rv2673 in *Mycobacterium tuberculosis* and NCgl1822 in *Corynebacterium glutamicum*, with 10 predicted transmembrane domains and a glycosyltransferase motif (DDX), features that are common to the GT-C superfamily of glycosyltransferases. Deletion of *M. smegmatis* MSMEG2785 resulted in altered growth and glycosyl linkage analysis revealed the absence of AG  $\alpha(1\rightarrow3)$ -linked arabinofuranosyl (Araf) residues. Complementation of the *M. smegmatis* deletion mutant was fully restored to a wild type phenotype by MSMEG2785 and Rv2673, and as a result, we have now termed this previously uncharacterized open reading frame, arabinofuranosyltransferase C (*aftC*). Enzyme assays using the sugar donor  $\beta$ -D-arabinofuranosyl-1-monophosphoryl-decaprenol (DPA) and a newly synthesized linear  $\alpha(1\rightarrow5)$ -linked Ara<sub>5</sub> neoglycolipid acceptor together with chemical identification of products formed, clearly identified AftC as a branching  $\alpha(1\rightarrow3)$  arabinofuranosyltransferase. This newly discovered glycosyltransferase sheds further light on the complexities of *Mycobacterium* cell wall biosynthesis, such as in *M. tuberculosis* and related species and represents a potential new drug target.

Introduction

Tuberculosis (TB) affects a third of the world population and causes 1.8 million fatalities annually (Dye, 2006). The spread of TB has been facilitated in recent years due to the susceptibility of HIV infected individuals to *Mycobacterium tuberculosis*, the etiological agent of TB (Paolo and Nosanchuk, 2004). The problem has also been compounded by the emergence of multi-drug resistant TB (MDR-TB) (Kaye and Frieden, 1996) and extensively drug-resistant (XDR)-TB strains (Shah *et al.*, 2007). *M. tuberculosis* and other mycobacteria have a distinct cell wall which has a lipid-rich outer layer that is highly impermeable (Minnikin, 1982). One of the major components of this outer envelope are mycolic acids, long chain  $\alpha$ -alkyl,  $\beta$ -hydroxy fatty acids that are essential for bacterial survival (Vilcheze *et al.*, 2000; Portevin *et al.*, 2004; Bhatt *et al.*, 2005; Parish *et al.*, 2007). These are found either esterified to the non-reducing termini of arabinogalactan (AG), or are present as trehalose esters, such as trehalose dimycolate (TDM) (Brennan and Nikaido, 1995; Dover *et al.*, 2004).

A common feature of members of the *Corynebacteriaceae* is that they all possess this unusual cell wall architecture (McNeil *et al.*, 1990, 1991; Besra *et al.*, 1995). Apart from mycolic acids, the cell wall is dominated by a second macromolecule, an essential heteropolysaccharide termed arabinogalactan (AG), which is linked to both mycolic acids and peptidoglycan, forming the mycolyl-arabinogalactan-peptidoglycan (mAGP) complex (Daffé *et al.*, 1990; McNeil *et al.*, 1990; McNeil *et al.*, 1991; Besra *et al.*, 1995). The formation of the arabinan domain ( $\alpha$ 1 $\rightarrow$ 5,  $\alpha$ 1 $\rightarrow$ 3 and  $\beta$ 1 $\rightarrow$ 2 glycosyl linkages) of AG results from the subsequent addition of arabinofuranose (Araf) residues by a set of unique arabinofuranosyltransferases including, the Emb proteins of which 3 paralogues exist in *Mycobacterium avium* (Belanger *et al.*, 1996) and

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

*M. tuberculosis* (Telenti *et al.*, 1997), AftA (Alderwick *et al.*, 2006a) and AftB (Seidel *et al.*, 2007a). The lipid linked sugar donor  $\beta$ -D-arabinofuranosyl-1-monophosphoryldecaprenol (DPA) (Wolucka *et al.*, 1994; Lee *et al.*, 1995; Lee *et al.*, 1997), serves as the substrate molecule for these complex membrane bound glycosyltransferases.

The anti-tuberculosis drug ethambutol (EMB) was shown to specifically inhibit AG biosynthesis (Takayama and Kilburn, 1989). The precise molecular target of EMB occupies the *embCAB* locus in *M. tuberculosis* (Telenti *et al.*, 1997). To further define the role of EmbCAB proteins in cell wall arabinan biosynthesis, *embA*, *embB* and *embC* were individually inactivated in *Mycobacterium smegmatis* (Escuyer *et al.*, 2001; Zhang *et al.*, 2003). All three mutants were viable, however, the non-reducing terminal Ara<sub>6</sub> motif which is the template for mycolylation in AG (McNeil *et al.*, 1991) was altered in both the Ms-*embA* and Ms-*embB* mutants (Escuyer *et al.*, 2001), whilst Ms-*embC* was shown to be involved in the formation of the arabinan domains of lipoarabinomannan (LAM) (Zhang *et al.*, 2003). Attempts to obtain deletion mutants of *embA* (Amin *et al.*, 2008) and *embB* in *M. tuberculosis* and *embAB* in *M. smegmatis* have proved unsuccessful (G.S. Besra, unpublished results). In contrast, deletion of the single Cg-*emb* orthologue and chemical analysis of the cell wall revealed a novel truncated AG structure possessing only terminal (t)-Araf residues with a corresponding loss of cell wall bound mycolic acids (Alderwick *et al.*, 2005). The presence of a novel enzyme responsible for “priming” the galactan domain for further elaboration by Emb proteins led to the identification of AftA (Alderwick *et al.*, 2006a). Recently, a retaining GT-C enzyme was identified, now termed AftB, which is responsible for the attachment of terminal  $\beta(1\rightarrow2)$  Araf residues, and marks the “end point” for AG arabinan biosynthesis (Figure 1) before decoration with mycolic acids (Seidel *et al.*, 2007a).

It is clear that additional arabinofuranosyltransferases involved in AG and LAM biosynthesis still remain to be identified. Indeed, Liu and Musheginan (2003) identified fifteen members of the GT-C superfamily residing in *M. tuberculosis*, representing candidates involved in the biosynthesis of cell wall related glycans and lipoglycans (Liu and Musheginan, 2003). We have continued our earlier studies (Alderwick *et al.*, 2006a; Alderwick *et al.*, 2006b; Seidel *et al.*, 2007a) to identify genes required for the biosynthesis of the core structural elements of the mAGP complex by studying mutants of *M. smegmatis* and the orthologous genes and enzymes of *M. tuberculosis* and *C. glutamicum*. Herein, we present MSMEG2785, Rv2673c and NCgl1822 as a new distinct arabinofuranosyltransferase of the GT-C superfamily, which is responsible for the transfer of Ara<sub>f</sub> residues from DPA to the arabinan domain to form  $\alpha(1\rightarrow3)$ -linked Ara<sub>f</sub> residues, which result in the branched arabinan domain distal to the non-reducing terminal Ara<sub>6</sub> motif characteristic of mycobacterial AG.

## Results

### Genome comparison of the Rv2673 locus

The arabinofuranosyltransferases EmbA, EmbB, and EmbC are vital for *M. tuberculosis* and represent a target for the established drug EMB (Mikusova *et al.*, 1995; Belanger *et al.*, 1996; Telenti *et al.*, 1997). Structural considerations of these proteins and a search for new drug targets resolved that more than 16 related proteins are present in *M. tuberculosis*, possibly also acting as glycosyltransferases (Liu and Mushegian, 2003). In our systematic analysis of GT-C glycosyltransferases, focusing on those present in *M. tuberculosis* and *C. glutamicum*, we have previously identified the arabinofuranosyltransferases AftA (Alderwick *et al.*, 2006a) and AftB (Seidel *et al.*, 2007a), as well as several  $\alpha$ -mannosyltransferases (Mishra *et al.*, 2007; Mishra *et al.*, 2008). Each of these glycosyltransferases plays a specific yet decisive role in cell wall biosynthesis and assembly. *In silico* analysis of one of the putative glycosyltransferases of *M. tuberculosis*, Rv2673, highlighted that orthologues are present in a range of species belonging to the sub-order *Corynebacteriineae*, including the families *Mycobacteriaceae*, *Corynebacteriaceae* and *Nocardiaceae* (Figure 2A). Furthermore, the organization of the gene locus is largely retained. The adjacent genes are largely of unknown function. *RibD* encodes a bifunctional deaminase-reductase domain, followed by a gene product containing a hydrolase domain, which is however absent in *Corynebacterium*, and downstream of Rv2673 a gene of unknown function is present. The wide distribution of Rv2673, its syntenic organization, and the fact that it is retained even in *M. leprae*, strongly indicates a fundamental function of its product. According to our experimental analysis (see below) we annotated this gene as arabinofuranosyltransferase C (*aftC*).



AftC of *M. tuberculosis* is 433 amino acid residues long. It is a hydrophobic protein and is predicted to possess 10 transmembrane-spanning segments (Figure 2B). However, in contrast to AftA, AftB or EmbC, it is characterized by the absence of a periplasmic carboxyterminal extension. The amino acid sequence among the *Corynebacteriaceae* is very well conserved, and there are 43% identical residues shared by the *M. tuberculosis* and *C. glutamicum* proteins. The degree of conservation is particularly high in the loop regions, for instance between helix 1 and 2, 3 and 4, or 6 and 7 (Figure 2B). The fully conserved aspartyl (D) and glutamyl (E) residues, which we propose to be involved in catalysis or substrate binding, are located in the first extended loop region (Liu and Mushegian, 2003), as we have demonstrated for similarly located aspartyl (D) residues of Cg-Emb and AftB (Seidel *et al.*, 2007a; Seidel *et al.*, 2007b). Interestingly, the long transmembrane helix 8 is well conserved and it is within this region that there is a strong identity to a membrane protein of *Vibrio parahaemolyticus* (CpsG). Furthermore, this gene is located in a gene cluster involved in the biosynthesis of a capsular polysaccharide within this pathogen (Guvener and McCarter, 2003).

**Construction and growth of mutants**

In order to delete *aftC* and study for possible consequences we generated a null mutant of *M. smegmatis* mc<sup>2</sup>155 MSMEG2785 (ortholog of Rv2673) using specialized transduction (Figure 3A). In contrast to our *C. glutamicum* studies (see below) growth of *M. smegmatis*Δ*aftC* in comparison to *M. smegmatis* was poor in liquid medium (Figure 3B) and sensitive to the addition of Tween-80 on agar plates (>0.005%). Complementation of *M. smegmatis*Δ*aftC* with either pMV261-Ms-*aftC* or pMV261-Mt-*aftC* restored the mutant to a wild type phenotype (Figure 3B). On solid media *M. smegmatis*Δ*aftC* had a smooth and glossy appearance in comparison to

the typical crenulated colony morphology found for wild type *M. smegmatis* (Figure 3C) and failed to stain as 'acid-fast' positive (data not shown). In addition, susceptibility of *M. smegmatis* $\Delta$ *aftC* to EMB and the hydrophobic antibiotics rifampicin and chloramphenicol was enhanced (minimal inhibitory concentration of 2, 100 and 10  $\mu$ g/ml for wild type *M. smegmatis* in comparison to 0.4, 4 and 5  $\mu$ g/ml for *M. smegmatis* $\Delta$ *aftC*, respectively) indicating increased permeability and that *M. smegmatis* $\Delta$ *aftC* had an altered cell wall. To study the function of the corynebacterial AftC the non-replicative plasmid pK19mobsacB $\Delta$ *aftC* was constructed. This was used to transform *C. glutamicum* to kanamycin resistance, indicating integration in its chromosome (Supplementary Figure S1A). Loss of vector was obtained by selection for sucrose-resistance yielding clones with *aftC* deleted. A PCR analysis with primer pairs P5 and P6 resulted in the expected fragment of 2160 bp for the wild type and of 1065 bp for the deletion mutant, which was termed *C. glutamicum* $\Delta$ *aftC*. Colonies of this mutant were more erose compared to the usual glossy appearance of the wild type colony (data not shown). In contrast to *M. smegmatis* $\Delta$ *aftC* the growth of the *C. glutamicum* $\Delta$ *aftC* mutant on the salt medium CGXII possessed only a slightly reduced growth rate of 0.32 h<sup>-1</sup>, whereas, that of the wild type *C. glutamicum* was 0.39<sup>-1</sup>h (Supplementary Figure S1B).

**mAGP analyses from *M. smegmatis*, *M. smegmatis* $\Delta$ *aftC*, *M. smegmatis* $\Delta$ *aftC* pMV261-Ms-*aftC*, *M. smegmatis* $\Delta$ *aftC* pMV261-Mt-*aftC*, *C. glutamicum* and *C. glutamicum* $\Delta$ *aftC***

To study the function of mycobacterial *aftC* deletion, defatted cells were analyzed qualitatively for AG esterified mycolic acids and cell wall associated lipids from an equivalent starting amount of biomass for each strain due to differences in growth rate (Figure 3B). As expected, *M. smegmatis* exhibited a typical profile of cell wall bound  $\alpha$ ,  $\alpha'$  and epoxy-mycolic acid methyl

esters (MAMEs), whereas, these products were drastically reduced in *M. smegmatis* $\Delta$ *aftC* (Figure 4A). In addition, complementation of *M. smegmatis* $\Delta$ *aftC* with either pMV261-Ms-*aftC* or pMV261-Mt-*aftC* (Figure 4A), led to the restoration of normal 'levels' of cell wall bound mycolic acids. Analysis of cell wall associated lipids in several independent experiments highlighted an apparent increase in TDM for the *aftC* deletion mutant. This was confirmed quantitatively through [<sup>14</sup>C]acetate labeling of cultures and equal loading of radioactivity of extractable free lipids from *M. smegmatis*, *M. smegmatis* $\Delta$ *aftC* and the complemented *M. smegmatis* $\Delta$ *aftC* strain using plasmids pMV261-Ms-*aftC* and pMV261-Mt-*aftC* (Figure 4B). Typically, wild type *M. smegmatis* synthesized 5250 cpm, whereas *M. smegmatis* $\Delta$ *aftC* afforded 14676 cpm of TDM after equivalent loading of radioactivity and quantitative analysis by phosphorimaging. Complementation of *M. smegmatis* $\Delta$ *aftC* with either pMV261-Ms-*aftC* or pMV261-Mt-*aftC* restored the phenotype of the deletion mutant back to that of wild type *M. smegmatis* (Figure 4B). These results demonstrated that Ms-*aftC* and Mt-*aftC* are involved in a key aspect of arabinan biosynthesis, whereby deletion substantially perturbs tethering of mycolic acids to AG, which results in an increase in TDM production.

The cell wall core (mAGP) was prepared from *M. smegmatis* and *M. smegmatis* $\Delta$ *aftC* as described (Daffé *et al.*, 1990; Besra *et al.*, 1995; Alderwick *et al.*, 2005) and the ratio of Ara to Gal in mAGP determined by gas chromatography (GC) analysis of alditol acetates (Daffé *et al.*, 1990; Besra *et al.*, 1995; Escuyer *et al.*, 2001; Alderwick *et al.*, 2005) (Figure 5). The glycosyl composition was calculated based on a single rhamnosyl (Rha) residue per AG chain (McNeil *et al.*, 1990). The glycosyl compositional analysis revealed a relative molar ratio of Rha:Ara:Gal of 1:71:31 and an Ara:Gal ratio of 2.3:1 which is in accord with previous data (Escuyer *et al.*, 2001). The *M. smegmatis* $\Delta$ *aftC* mutant yielded AG with a significant reduction in Ara content

concomitant with a relative increase in the amount of Gal. The *M. smegmatis* $\Delta$ *aftC* yielded an AG with a Rha:Ara:Gal ratio of 1:22:56 and an Ara:Gal ratio of 0.4:1. Complementation of *M. smegmatis* $\Delta$ *aftC* with either pMV261-Ms-*aftC* or pMV261-Mt-*aftC*, restored the Rha:Ara:Gal ratio to that of wild type *M. smegmatis*. Gas chromatography mass spectrometry (GC/MS) analysis of per-*O*-methylated alditol acetate derivatives prepared from *M. smegmatis* and *M. smegmatis* $\Delta$ *aftC* indicated the complete absence of 3,5-Araf branching residues and a significant reduction in *t*-Araf, 2-Araf and 5-Araf-linkages (Figure 6). Complementation of *M. smegmatis* $\Delta$ *aftC* with either plasmid encoding Ms-*aftC* or Mt-*aftC* restored the glycosyl linkage profile to that of wild type *M. smegmatis* (Figure 6). These results demonstrate that MSMEG2785 and Rv2673, are functionally equivalent and are involved in the synthesis of 3,5-Araf branching residues. Interestingly, LAM preparations from *M. smegmatis* $\Delta$ *aftC* were truncated in size on SDS-PAGE analysis to 'full-size' LAM from wild type *M. smegmatis*. Further purification and detailed chemical analyses of LAM from the *aftC* mutant strain are currently being undertaken and will be reported separately (H.L. Birch, unpublished results).

In contrast to the mycolic acid studies performed with the mycobacterial *aftC* deletion mutant, *C. glutamicum* $\Delta$ *aftC* cells were analyzed quantitatively for AG esterified corynomycolic acids due to similar growth rates between strains (Supplementary Figure S1B). Wild type *C. glutamicum* exhibited the known profile of corynomycolic acid methyl esters (CMAMEs, 35345 cpm) (Supplementary Figure S2), whereas, cell wall bound CMAMEs were significantly reduced in *C. glutamicum* $\Delta$ *aftC* (8023 cpm). The above data was reassuring as the qualitative (*M. smegmatis* $\Delta$ *aftC*) and quantitative (*C. glutamicum* $\Delta$ *aftC*) analyses were comparable in terms of a reduction in cell wall bound mycolic acids (Figure 4A and Supplementary Figure S2). Importantly, these results have also shown that Cg-*aftC* is involved in a key aspect of arabinan

1  
2 biosynthesis, whereby deletion perturbs tethering of corynomycolic acids to AG. The GC/MS  
3 profiles of per-*O*-methylated alditol acetate derivatives of *C. glutamicum* and *C.*  
4 *glutamicum* $\Delta$ *aftC* are shown in Supplementary Figure S3 with *C. glutamicum* $\Delta$ *aftC* also clearly  
5 devoid of 3,5-Araf branching residues.  
6  
7  
8  
9  
10

11  
12  
13  
14 ***In vitro* arabinofuranosyltransferase activity with extracts of *M. smegmatis*, *M.***  
15 ***smegmatis* $\Delta$ *aftC* and complemented strains**  
16  
17

18  
19  
20  
21 Initial attempts to develop an *in vitro* assay using either purified recombinant expressed AftC or  
22 *E. coli* membranes expressing *aftC*, have thus far proved unsuccessful, probably due to the  
23 hydrophobic nature of the protein. In an alternative approach, we assessed the capacity of  
24 membrane preparations from *M. smegmatis*, *M. smegmatis* $\Delta$ *aftC* and *M. smegmatis* $\Delta$ *aftC*  
25 complemented with pMV261-Mt-*aftC* to catalyze arabinofuranosyltransferase activity in the  
26 presence of exogenous synthetic acceptors (Lee *et al.*, 1997; Seidel *et al.*, 2007a).  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37

38 We first assessed whether *M. smegmatis* $\Delta$ *aftC* was deficient in  $\alpha(1\rightarrow5)$  and  $\beta(1\rightarrow2)$   
39 arabinofuranosyltransferase activity using an  $\alpha$ -D-Araf-(1 $\rightarrow$ 5)- $\alpha$ -D-Araf-*O*-(CH<sub>2</sub>)<sub>7</sub>CH<sub>3</sub> (Ara<sub>2</sub>)  
40 synthetic acceptor (Lee *et al.*, 1997) and DP[<sup>14</sup>C]A as a sugar donor based on an established  
41 assay format for determining  $\alpha(1\rightarrow5)$  and  $\beta(1\rightarrow2)$  arabinofuranosyltransferase activities (Lee *et*  
42 *al.*, 1998). TLC/autoradiographic analysis of the products which were only synthesized in the  
43 presence of Ara<sub>2</sub>, when assayed with *M. smegmatis* membranes resulted in the formation of two  
44 products (A and B) (Figure 7A and B). The enzymatic synthesis of products A and B are  
45 consistent with our previous studies using mycobacterial (Lee *et al.*, 1997) and corynebacterial  
46 (Seidel *et al.*, 2007a) membrane preparations resulting in trisaccharide products as a result of  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

$\alpha(1\rightarrow5)$  and  $\beta(1\rightarrow2)$  Ara<sub>f</sub> linkages to the Ara<sub>2</sub> acceptor (Figure 7A). Addition of EMB in several experiments, even at high concentrations of up to 1 mg/ml to the reaction mixture, resulted in a decrease in only the *in vitro* synthesized  $\alpha$ -D-[<sup>14</sup>C]Ara<sub>f</sub>-(1 $\rightarrow$ 5)- $\alpha$ -D-Ara<sub>f</sub>-(1 $\rightarrow$ 5)- $\alpha$ -D-Ara<sub>f</sub>-O-(CH<sub>2</sub>)<sub>7</sub>CH<sub>3</sub> product A (Figure 7A and B). Assays performed with membranes from *M. smegmatis* $\Delta$ *aftC* and the pMV261-Mt-*aftC* complemented strain using the Ara<sub>2</sub> synthetic acceptor gave a similar profile to that of wild type *M. smegmatis* (Figure 7B). The data clearly show that the *M. smegmatis* $\Delta$ *aftC* strain possesses comparable levels of EMB-sensitive  $\alpha(1\rightarrow5)$  and EMB-resistant  $\beta(1\rightarrow2)$  arabinofuranosyltransferase activity.

The lack of  $\alpha(1\rightarrow3)$  arabinofuranosyltransferase activity in the previously reported Ara<sub>2</sub> and  $\alpha$ -D-Ara<sub>f</sub>-(1 $\rightarrow$ 5)- $\alpha$ -D-Ara<sub>f</sub>-(1 $\rightarrow$ 5)- $\alpha$ -D-Ara<sub>f</sub>-O-(CH<sub>2</sub>)<sub>7</sub>CH<sub>3</sub> (Ara<sub>3</sub>) acceptor based assays (Lee *et al.*, 1997) required the development of an arabinofuranosyltransferase assay using the Ara-extended synthetic acceptor  $\alpha$ -D-Ara<sub>f</sub>-(1 $\rightarrow$ 5)- $\alpha$ -D-Ara<sub>f</sub>-(1 $\rightarrow$ 5)- $\alpha$ -D-Ara<sub>f</sub>-(1 $\rightarrow$ 5)- $\alpha$ -D-Ara<sub>f</sub>-(1 $\rightarrow$ 5)- $\alpha$ -D-Ara<sub>f</sub>-O-(CH<sub>2</sub>)<sub>8</sub>NH<sub>2</sub> (Ara<sub>5</sub>) (Supplementary Experimental and Supplementary Figure S4) and DP[<sup>14</sup>C]A as a sugar donor (Lee *et al.*, 1998). TLC/autoradiographic analysis of the products which are only synthesized in the presence of Ara<sub>5</sub>, when assayed with *M. smegmatis* membranes resulted in the formation of a single product X (Figure 8A) through the transfer of a single [<sup>14</sup>C]Ara<sub>f</sub> residue, with a retardation factor (*R<sub>f</sub>*) consistent with a synthetic Ara<sub>6</sub> acceptor (Appelmelk *et al.*, 2008) standard (Figure 8B). In addition, the synthesis of product X in over-expression studies was enhanced. Consistently from two independent membrane preparations and assays performed in triplicate from *M. smegmatis* pMV261-Mt-*aftC* produced product X (6453 cpm) in comparison to membranes from wild type *M. smegmatis* (4289 cpm) in the above assays demonstrating that AftC was functionally involved in the synthesis of product X. The inclusion of EMB in several experiments, even at high concentrations of up to 1 mg/ml to the

reaction mixture did not inhibit the synthesis of this *in vitro* synthesized [ $^{14}\text{C}$ ]Ara $\beta$ -Ara $_5$  (Figure 8A, Product X) illustrating that the Ara $_5$  acceptor was not extended *via* an EMB-sensitive  $\alpha(1\rightarrow5)$  arabinofuranosyltransferase. Interestingly, membranes prepared from the *M. smegmatis* $\Delta\text{aftC}$  strain were unable to synthesize the *in vitro* product to the same level of activity that was observed with wild type membranes prepared from *M. smegmatis* (Figure 8A). This was to be expected, since our earlier *in vivo* and *in vitro* studies would have anticipated residual Ara $_6$  product formation, considering that *M. smegmatis* $\Delta\text{aftC}$  possesses  $\beta(1\rightarrow2)$  arabinofuranosyltransferase activity. Assays performed with membranes from the *M. smegmatis* $\Delta\text{aftC}$  pMV261-Mt-*aftC* complemented strain, gave a similar profile to that of wild type *M. smegmatis* (Figure 8A).

To establish that the Ara $_5$  acceptor is being utilized by two different arabinofuranosyltransferases, presumably establishing  $\beta(1\rightarrow2)$  and  $\alpha(1\rightarrow3)$  linkages, assays similar to that used before were scaled up (see Experimental Procedures) and product X extracted and purified through preparative TLC for each membrane preparation. GC (Sasaki *et al.*, 2005) and GC/MS (Alderwick *et al.*, 2005) analyses of the partially per-*O*-methylated, per-*O*-acetylated alditol acetate derivatives of product X in assays performed with *M. smegmatis* membranes revealed the addition of  $\beta(1\rightarrow2)$  [ $R_t$  11.75 min;  $m/z$  129, 130, 161, 190] and  $\alpha(1\rightarrow3)$  [ $R_t$  12.39 min;  $m/z$  118, 129, 130, 190, 202, 233] linked Ara $\beta$  residues (Figure 9A and B). Therefore, the product migrating below Ara $_5$  and co-incident with the Ara $_6$  acceptor standard on TLC (Figure 8A and B) is in fact a mixture of two products (Figure 9B). The addition of  $\beta(1\rightarrow2)$ -linked Ara $\beta$  residues can be attributed to the function of AftB. The presence of  $\alpha(1\rightarrow3)$ -linked Ara $\beta$  residues in this assay using an Ara $_5$  acceptor clearly highlights the role of a novel arabinofuranosyltransferase(s) capable of functioning in an  $\alpha(1\rightarrow3)$  capacity. Importantly, the



1  
2  
3 level of  $\alpha(1\rightarrow3)$  activity when the Ara<sub>5</sub> acceptor is incubated with membranes prepared from *M.*  
4  
5 *smegmatis* $\Delta$ *aftC* is completely abolished (Figure 9A). However,  $\beta(1\rightarrow2)$  activity is clearly  
6  
7 present in *M. smegmatis* $\Delta$ *aftC* (Figure 9A). In addition, *M. smegmatis* $\Delta$ *aftC* complemented with  
8  
9 pMV261-Mt-*aftC* restores  $\alpha(1\rightarrow3)$  arabinofuranosyltransferase activity to wild type *M.*  
10  
11 *smegmatis* (Figure 9A). The results clearly establish both from *in vivo* and *in vitro* experiments  
12  
13 that AftC catalyzes the addition of an  $\alpha(1\rightarrow3)$ -Araf unit *via* an  $\alpha(1\rightarrow3)$   
14  
15 arabinofuranosyltransferase and that this enzyme is also resistant to EMB (Figure 8A).  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



## Discussion

The mAGP complex represents one of the most important cell wall components of the *Corynebacteriaceae* and is essential for the viability of *M. tuberculosis* (Vilcheze *et al.*, 2000; Pan *et al.*, 2001; Gande *et al.*, 2004; Mills *et al.*, 2004). It is therefore not surprising that one of the most effective anti-mycobacterial drugs, EMB, targets its synthesis through inhibition of AG biosynthesis. However, the emergence of MDR-TB and XDR-TB has accelerated the need to discover new drug targets (Brennan and Nikaido, 1995). One of the strategies is to identify genes involved in AG biosynthesis. Based on this strategy we previously identified the presence of a new “priming” enzyme, now termed AftA, which would link the initial Ara<sub>f</sub> unit with the C-5 OH of a  $\beta(1\rightarrow6)$  linked Gal<sub>f</sub> of a pre-synthesized galactan core (Alderwick *et al.*, 2005), and more recently identified the AftB enzyme responsible for  $\beta(1\rightarrow2)$  Ara<sub>f</sub> residues.

The previously described Emb (Alderwick *et al.*, 2005), AftA (Alderwick *et al.*, 2006a) and AftB proteins (Seidel *et al.*, 2007a) are distinct arabinofuranosyltransferases. Thus, despite some functional relationship, these glycosyltransferases have inherent specific features as evident from the insensitivity of AftA and AftB towards EMB, whereas the single Cg-Emb (Alderwick *et al.*, 2005; Radmacher *et al.*, 2005), and Mt-Emb proteins are sensitive towards EMB (Telenti *et al.*, 1997; Belanger *et al.*, 1996). The number of arabinofuranosyltransferases that are required for mycobacterial arabinan biosynthesis has been a matter of speculation to date depending on how the arabinan chains are assembled. The primary structure of AG (Besra *et al.*, 1995; Daffé *et al.*, 1990) would suggest at least five distinct arabinofuranosyltransferases are required for the complete formation of AG. Interestingly, *M. smegmatis* *embA* and *embB* mutants were found to possess reduced amounts of the non-reducing terminal disaccharide  $\beta$ -D-Ara<sub>f</sub>-(1 $\rightarrow$ 2)- $\alpha$ -D-Ara<sub>f</sub>

and result in the removal of the dominant terminal non-reducing Ara<sub>6</sub> branched motif in the mutant being replaced by a linear Ara<sub>4</sub> motif (Escuyer *et al.*, 2001). The authors of this study concluded that the *M. smegmatis* *embA* and *embB* mutants result in a lack of 3-arm branching off the main  $\alpha(1\rightarrow5)$ -arabinan chain proximal to the non-reducing and attachment site of mycolic acids in AG (Escuyer *et al.*, 2001). Initially, it was proposed that the  $\beta$ -D-Araf-(1 $\rightarrow$ 2)- $\alpha$ -D-Araf disaccharide was assembled using EmbA and EmbB. However, the recent identification of AftB, the development of specific *in vitro* assays in combination with mutant strains, and recent structural studies have fuelled speculation that EmbA/B act as  $\alpha(1\rightarrow5)$  arabinofuranosyltransferases (Bhamidi *et al.*, 2008; Seidel *et al.*, 2007a; Alderwick *et al.*, 2005).

In this study, we have identified MSMEG2785 (also Rv2673 and NCgl1822), which we have termed AftC, as a novel branching arabinofuranosyltransferase. More precisely, AftC catalyzes the addition of  $\alpha(1\rightarrow3)$  Araf residues as shown through both *in vivo* and *in vitro* experiments, ultimately resulting in 3,5-Araf residues after further  $\alpha(1\rightarrow5)$  extension, characteristic of AG. For instance, incubation of membranes prepared from *M. smegmatis* with DP[<sup>14</sup>C]A and a linear  $\alpha(1\rightarrow5)$ -Ara<sub>5</sub> neoglycolipid acceptor resulted in the synthesis of an Ara<sub>6</sub> product. Further chemical characterization of the product by glycosyl linkage analysis established that the  $\alpha(1\rightarrow5)$ -Ara<sub>5</sub> acceptor was extended *via* an EMB resistant  $\alpha(1\rightarrow3)$  arabinofuranosyltransferase giving rise to 3-linked Araf residues and corroborated our earlier cell wall analysis of the *M. smegmatis* $\Delta$ *aftC* mutant. Since, it is now established that only  $\alpha(1\rightarrow5)$  arabinofuranosyltransferase(s) are EMB-sensitive it can be further speculated that EmbA and EmbB function in the assembly of the linear  $\alpha(1\rightarrow5)$  arabinan segments as presented in Figure 10, which is in accordance with previous data and the phenotype of a Cg-Emb mutant

(Alderwick *et al.*, 2005). It is clear that further studies are required to establish the precise role of EmbA and EmbB in mycobacteria.

The analysis of the *M. smegmatis*Δ*aftC* mutant to date and based on the Ara:Gal ratio would suggest that the residual arabinan segment in the mutant consists of approximately five Ara<sub>f</sub> residues:  $\beta$ -D-Ara<sub>f</sub>-(1→2)-α-D-Ara<sub>f</sub>-(1→5)-α-D-Ara<sub>f</sub>-(1→5)-α-D-Ara<sub>f</sub>-(1→5)-α-D-Ara<sub>f</sub> located at three branches on the galactan chain (Alderwick *et al.*, 2005; Besra *et al.*, 1995). This is consistent with the recent primary structure of AG (Bhamidi *et al.*, 2008), with a 'non-variable' terminal non-reducing Ara<sub>17</sub> motif, introduction of a 3,5-Ara<sub>f</sub> residue distal to this non-reducing end by AftC and further extension by a linear α(1→5)Ara<sub>f</sub> domain (Figure 10). The latter appears to be variable (up to 12/13 residues). However, based on *M. smegmatis*Δ*aftC* and the subsequent Ara:Gal compositional analysis a dominant Ara<sub>22</sub>/Ara<sub>23</sub> motif would be consistent with recent (Bhamidi *et al.*, 2008) and previous (Besra *et al.*, 1995) structural data on AG and this is represented in terms of biosynthetic considerations in Figure 10. It is also possible that AftC or a second distinct α(1→3) arabinofuranosyltransferase (shown as AftD in Figure 10) may be involved in late stages of AG synthesis i.e. the non-reducing Ara<sub>6</sub> motif and is consistent with our data and the model presented in Figure 10.

The discovery of AftC has now shed new light on the key arabinofuranosyltransferases to build an arabinan domain typical for *Corynebacterianae*. In this context, the genomic organization in the genomes of the *Corynebacterianae* sequenced is intriguing, revealing high synteny of the *M. tuberculosis* *aftC* locus to the maps of all other *Mycobacterium* and *Corynebacterium* species. The identification of new cell wall biosynthetic drug targets is of great importance, especially with the emergence of MDR-TB. This newly discovered DPA dependent arabinofuranosyl

transferase represents, along with a straightforward *in vitro* enzyme assay, a promising candidate for further exploitation as a potential drug target.

For Peer Review

Experimental procedures

**Bacterial strains and growth conditions.** *C. glutamicum* ATCC 13032 (referred to the remainder of the text as *C. glutamicum*) and *Escherichia coli* DH5 $\alpha$ mc $\alpha$ r were grown in Luria-Bertani broth (LB, Difco) at 30°C and 37°C, respectively. The recombinant strains generated in this study were grown on complex Brain Heart Infusion medium (BHI, Difco), and the salt medium CGXII used for *C. glutamicum* as described (Eggeling and Bott, 2005). Kanamycin and ampicillin were used at a concentration of 50  $\mu$ g/ml. *M. smegmatis* strains were grown in Tryptic Soy Broth (TSB; Difco) containing 0.005 % Tween80 (TSBT). Solid media were made by adding 1.5 % agar to the above-mentioned broths. The concentrations of antibiotics used for *M. smegmatis* were 100  $\mu$ g/ml for hygromycin and 20  $\mu$ g/ml for kanamycin. Minimal inhibitory concentrations were determined by plating cells on solid media supplemented with various concentrations of EMB, rifampicin and chloramphenicol. The minimal inhibitory concentration was defined as the first concentration of drug that would inhibit 100% of growth after 5 days of incubation (Belanger *et al.*, 1996). *M. tuberculosis* H37Rv DNA was obtained from the NIH Tuberculosis Research Materials and Vaccine Testing Contract at Colorado State University. All other chemicals were of reagent grade and obtained from Sigma-Aldrich.

**Construction of plasmids and strains.** Approximately 1 kb of upstream and downstream flanking sequences of *MSMEG2785* were PCR amplified from *M. smegmatis* mc<sup>2</sup>155 genomic DNA using the primer pairs MS2785LL (TTTTTTTTCCATAAATTGGATCCGCTGACCGACCTCATC) and MS2785LR (TTTTTTTTCCATTTCTTGGCGAGCCCGAGCTTGAAGTTG), and MS2785RL (TTTTTTTTCCATAGATTGGTTCCTGCTGCTGTCCCTTGG) and MS2785RR

(TTTTTTTTCCATCTTTTGGCGAACTCAGCGGCGATTAC), respectively (all primers are given in 5' to 3' direction). Following restriction digestion of the primer incorporated *Van91I* sites, the PCR fragments were cloned into *Van91I*-digested p0004S to yield the knockout plasmid p $\Delta$ MSMEG2785 which was then packaged into the temperature sensitive mycobacteriophage phAE159 as described previously (Bardarov *et al.*, 2002) to yield phasmid DNA of the knockout phage ph $\Delta$ MSMEG2785. Generation of high titre phage particles and specialized transduction were performed as described earlier (Bardarov *et al.*, 2002; Stover *et al.*, 1991). Deletion of MSMEG2785 in one hygromycin resistant transductant was confirmed by Southern blot. To enable expression of MSMEG2785 and Rv2673, in the deletion mutant, these were amplified using primer pairs designed for subsequent cloning into the mycobacterial-shuttle vector pMV261 (Stover *et al.*, 1991). All cloned fragments were verified by sequencing.

To construct the deletion vector pK19mobsacB $\Delta$ *afnC* (NCgl1822), crossover PCR was applied with primer pairs AB (A, CGTTAAGCTTCGATCTTGTTGATGTGTGGCATCACACG; B, CCCATCCACTAACTTAAACAGCGCCATCAACAACATGG) and CD (C, TGTTTAAGTTTAGTGGATGGGTGATCCAACGCACGACCATC; D, GCATGGATCCACGCATACCGAGGGAAAGATCTTC) and *C. glutamicum* genomic DNA as template. Both amplified products were used in a second PCR with primer pairs AD to generate a 656 bp fragment consisting of sequences adjacent to Cg-*afnC*, which was ligated with BamHI-HindIII-cleaved pK19mobsacB. All plasmids were confirmed by sequencing. The chromosomal deletion of Cg-*afnC* was performed as described previously using two rounds of positive selection (Schafer *et al.*, 1994), and its successful deletion was verified by use of two different primer pairs.

**Isolation of the mAGP complex, glycosyl composition and linkage analysis of alditol acetates by GC and GC/MS.** The thawed cells were resuspended in phosphate buffered saline containing 2% Triton X-100 (pH 7.2), disrupted by sonication and centrifuged at 27000 x g (Besra *et al.*, 1995; Alderwick *et al.*, 2005). The pelleted material was extracted three times with 2% SDS in phosphate buffered saline at 95°C for 1 h, washed with water, 80% (v/v) acetone in water, and acetone, and finally lyophilized to yield a highly purified cell wall preparation (Besra *et al.*, 1995; Alderwick *et al.*, 2005). Cell wall or per-*O*-methylated cell wall preparations (Alderwick *et al.*, 2005) were hydrolyzed in 2 M TFA, reduced with NaB<sup>2</sup>H<sub>4</sub> and the resultant alditols per-*O*-acetylated and examined by GC and GC/MS as described previously (Besra *et al.*, 1995; Alderwick *et al.*, 2005).

**Extraction and analysis of cell wall bound mycolic acids.** In terms of *M. smegmatis* strains equivalent amounts of freeze-dried bacilli (100 mg) were processed as described previously (Seidel *et al.*, 2007a) following two consecutive CHCl<sub>3</sub>/CH<sub>3</sub>OH/H<sub>2</sub>O (10:10:3, v/v/v) extractions for 4 h at 50°C in the analysis of cell wall associated lipids, and cell wall bound MAMEs. Alternatively, *M. smegmatis* and *C. glutamicum* cultures (5 ml) were grown and metabolically labelled at mid-logarithmic phase of growth using 1 µCi/ml [1,2-<sup>14</sup>C]acetate (50-62 mCi/mmol, GE Healthcare, Amersham Bioscience) for 4 h at either 30°C or 37°C with gentle shaking, harvested, washed and freeze-dried. Cells were then extracted by two consecutive extractions with 2 ml of CHCl<sub>3</sub>/CH<sub>3</sub>OH/H<sub>2</sub>O (10:10:3, v/v/v) for 4 h at 50°C to provide cell wall associated lipids and analyzed as described previously (Seidel *et al.*, 2007a). The crude lipid extracts were resuspended in CHCl<sub>3</sub>:CH<sub>3</sub>OH (2:1) and equivalent aliquots (50,000 cpm) analyzed by TLC using silica gel plates (5735 silica gel 60F<sub>254</sub>, Merck) developed in CHCl<sub>3</sub>:CH<sub>3</sub>OH:NH<sub>4</sub>OH (80:20:2, v/v/v) to separate [<sup>14</sup>C]-labeled TDM and phospholipids (Mikusova *et al.*, 1995).

Lipids were visualized by autoradiography by overnight exposure of Kodak X-Omat AR film to the TLC plates to reveal labelled lipids, quantified by phosphorimaging and compared to know standards (Mikusova *et al.*, 1995). The bound MAMEs/CMAMEs from the above de-lipidated extracts were released by the addition of 2 ml of 5% aqueous solution of tetra-butyl ammonium hydroxide followed by overnight incubation at 100°C. After cooling, water (2 ml), CH<sub>2</sub>Cl<sub>2</sub> (4 ml) and CH<sub>3</sub>I (500 µl) were added and mixed thoroughly for 30 min. The lower organic phase was recovered following centrifugation and washed three times with water (4 ml), dried and resuspended in diethyl ether (4 ml). After centrifugation the clear supernatant was again dried and resuspended in CH<sub>2</sub>Cl<sub>2</sub> (100 µl). An aliquot (5 µl) from each strain was subjected to scintillation counting and an equivalent (5 µl) aliquot analyzed by TLC using silica gel plates (5735 silica gel 60F<sub>254</sub>, Merck), developed in petroleum ether/acetone (95:5, v/v) and either visualized by autoradiography by exposure of Kodak X-Omat AR film to the TLC plates to reveal [14C]-labeled MAMEs/CMAMEs, or charred following spraying with 5% molybdophosphoric acid in ethanol at 100°C and compared to know standards.

**Arabinofuranosyltransferase activity with membrane preparations of *M. smegmatis*, *M. smegmatis* pMV261-Mt-*aftC*, *M. smegmatis*Δ*aftC* and *M. smegmatis*Δ*aftC* pMV261-Mt-*aftC*.**

Membranes were prepared as described previously (Lee *et al.*, 1997; Alderwick *et al.*, 2006a) and resuspended in 50 mM MOPS (pH 7.9), containing 5 mM β-mercaptoethanol and 10 mM MgCl<sub>2</sub> (buffer A) to a final concentration of 15-10 mg/ml. The neoglycolipid acceptors used in this study were α-D-Araf-(1→5)-α-D-Araf-(1→5)-α-D-Araf-(1→5)-α-D-Araf-(1→5)-α-D-Araf-O-(CH<sub>2</sub>)<sub>8</sub>NH<sub>2</sub> (Ara<sub>5</sub>, see Supplementary Material) and α-D-Araf-(1→5)-α-D-Araf-O-(CH<sub>2</sub>)<sub>7</sub>CH<sub>3</sub> (Ara<sub>2</sub>) (Lee *et al.*, 1995; Lee *et al.*, 1998). The acceptors (either Ara<sub>2</sub> or Ara<sub>5</sub>) and DP[<sup>14</sup>C]A (Lee *et al.*, 1995; Lee *et al.*, 1998) (stored in CHCl<sub>3</sub>/CH<sub>3</sub>OH, 2:1, v/v) were aliquoted into 1.5 ml



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

epENDORF tubes to a final concentration of 2 mM and 200,000 cpm (90  $\mu$ M), respectively, and dried under nitrogen. The arabinofuranosyltransferase assay was carried out as described previously (Lee *et al.*, 1997) with modifications. IgePal<sup>TM</sup> (Sigma-Aldrich) was added (0.1%, v/v) with the appropriate amount of buffer A (final volume 80  $\mu$ l). Tubes were sonicated for 15 min to resuspend lipid linked substrates and then mixed with the remaining assay components, which included membrane protein from either *M. smegmatis*, *M. smegmatis* pMV261-Mt-*aftC* *M. smegmatis* $\Delta$ *aftC* or *M. smegmatis* $\Delta$ *aftC* pMV261-Mt-*aftC* (1 mg), 1 mM ATP, 1 mM NADP and in some cases EMB (0-1 mg/ml). Assays were incubated for 1 h at 37°C and quenched by the addition of 533  $\mu$ l CHCl<sub>3</sub>/CH<sub>3</sub>OH (1:1, v/v). After mixing and centrifugation at 27000 x g for 15 min at 4°C, the supernatant was removed and dried under nitrogen. The residue was then resuspended in 700  $\mu$ l of CH<sub>3</sub>CH<sub>2</sub>OH/H<sub>2</sub>O (1:1, v/v) and loaded onto a 1 ml SepPak strong anion exchange cartridge (Supelco), pre-equilibrated with CH<sub>3</sub>CH<sub>2</sub>OH/H<sub>2</sub>O (1:1, v/v). The column was washed with 2 ml CH<sub>3</sub>CH<sub>2</sub>OH and the eluate collected, dried and partitioned between the two phases arising from a mixture of *n*-butanol (3 ml) and water (3 ml). The resulting organic phase was recovered following centrifugation at 3,500 x g and the aqueous phase again extracted twice with 3 ml of water-saturated *n*-butanol. The pooled extracts were back-washed twice with *n*-butanol-saturated water (3 ml). The *n*-butanol fraction was dried and resuspended in 200  $\mu$ l butanol. The extracted radiolabeled material was quantified by liquid scintillation counting using 10 % of the labeled material and 5 ml of EcoScintA (National Diagnostics, Atlanta). The incorporation of [<sup>14</sup>C]Ara<sub>f</sub> was determined by subtracting counts present in control assays (incubations in the absence of acceptor). The remaining labeled material was subjected to thin-layer chromatography (TLC) using either isopropanol:acetic acid:water (8:1:1, v/v/v) for the assays utilizing the Ara<sub>5</sub> acceptor or CHCl<sub>3</sub>:CH<sub>2</sub>OH:H<sub>2</sub>O:NH<sub>4</sub>OH (65:25:3.6:0.5, v/v/v/v) in the case of the Ara<sub>2</sub> acceptor on aluminum-backed Silica Gel 60 F<sub>254</sub> plates (Merck, Darmstadt,

Germany). Autoradiograms were obtained by exposing TLCs to X-ray film (Kodak X-Omat) for 3 days.

**Characterization of  $\alpha(1\rightarrow3)$ -arabinofuranosyltransferase activity with membranes prepared from *M. smegmatis*, *M. smegmatis* $\Delta$ *aftC* and *M. smegmatis* $\Delta$ *aftC* pMV261-Mt-*aftC*.** Large-scale reaction mixtures containing cold DPA (200  $\mu$ g, 0.75 mM) (Lee *et al.*, 1997) and 50 mM of the acceptor Ara<sub>5</sub> were mixed and given an initial incubation at 37°C with membranes prepared from either *M. smegmatis*, *M. smegmatis* $\Delta$ *aftC* or *M. smegmatis* $\Delta$ *aftC* pMV261-Mt-*aftC* for 1 h. The assays were replenished with fresh membranes (1 mg) and re-incubated for 1 h at 37 °C with the entire process repeated thrice. Products were extracted from reaction mixtures by *n*-butanol/water phase separation as described earlier to extract products. Products were applied to preparative TLC plates, developed in isopropanol:acetic acid:water (8:1:1, v/v/v) and sprayed with 0.01% 1,6-diphenylhexatriene in petroleum-ether:acetone (9:1, v/v), and the products localized under long-wave (366 nm) UV light (Lee *et al.*, 1997). The plate was then re-developed in toluene to remove the reagent and the bands recovered from the plates by extraction with *n*-butanol. The butanol phases were washed with water saturated with *n*-butanol and the dried products subjected to GC (Sasaki *et al.*, 2005) and GC/MS as described (Lee *et al.*, 1997; Alderwick *et al.*, 2006a).

**Acknowledgments**

HLB is a Medical Research Council Quota Student. GSB acknowledges support in the form of a Personal Research Chair from Mr. James Bardrick, Royal Society Wolfson Research Merit Award, as a former Lister Institute-Jenner Research Fellow, the Medical Research Council and The Wellcome Trust (081569/Z/06/Z). AB is supported by a Career Development Award from the Medical Research Council. The contributions of TLL and YB to this work were supported by the Alberta Ingenuity Centre for Carbohydrate Science and the Natural Sciences and Engineering Research Council of Canada. The authors would also like to thank Tsungda Hsu (Albert Einstein College of Medicine, New York) for the gift of plasmid p0004S.

**Abbreviations**

The abbreviations used are: AG, Arabinogalactan; Ara, arabinose; Cg, *C. glutamicum*; 2D-TLC, two-dimensional thin-layer chromatography; DPA, decaprenol phosphoarabinose; EMB, ethambutol; *f*, furanose; Gal, galactose; GC, gas chromatography; GC/MS, gas chromatography mass spectrometry; LB, Luria-Bertani; MAME, mycolic acid methyl ester; mAGP, mycolyl-arabinogalactan-peptidoglycan; Man-LAM, mannose capped LAM; MS, *M. smegmatis*; Mt, *M. tuberculosis*; OD, Optical Density; TLC, thin-layer chromatography.

## Figure Legends

**Fig. 1. Biosynthetic pathway leading to arabinan formation in *M. tuberculosis* AG.**

**Fig. 2. Comparison of the *aftC* locus within the *Corynebacteriaceae*.** (A) The locus in the bacteria analyzed consists of *aftC* which in *M. tuberculosis* has the locus tag Rv2673 and in *C. glutamicum* NCgl1822. The genomic region displayed encompasses 7 kb, and orthologous genes are highlighted accordingly. Abbreviations: *M. marinum*, *Mycobacterium marinum*; *M. av subsp. par.*, *Mycobacterium avium subsp. paratuberculosis*; *C. efficiens*, *Corynebacterium efficiens*; *C. jeikeium*, *Corynebacterium jeikeium*; *Nocardia farcina*, *Nocardia farcina* IFM 10152; *Rhodococcus*, *Rhodococcus* sp. strain RHA1. (B) AftC is a hydrophobic protein predicted to span the membrane 10 times and the transmembrane helices are numbered accordingly. The lower part of the figure shows the degree of conservation of the orthologues given in A as analysed by the DIALIGN method (Brudno *et al.*, 2003). Also shown is the approximate position of the fully conserved aspartyl (D) and glutamyl (E) residues.

**Fig. 3. Generation of a *MSMEG2785* null mutant.** (A) A map of the *MSMEG2785* region in the parental *M. smegmatis* strain and its corresponding region in the  $\Delta$ *MSMEG2785* mutant. *res*,  $\gamma\delta$  resolvase site; *hyg*, hygromycin resistance gene from *Streptomyces hygroscopicus*; *sacB*, sucrose counter-selectable gene from *Bacillus subtilis*. Digoxigenin-labelled probes were derived from ~1kb upstream and downstream flanking sequences that were used to construct the knockout plasmid, and are indicated by thick lines with square ends. *SacI* digested bands expected in a Southern blot are indicated in roman numerals with sizes in brackets. The inset shows the Southern blot of *SacI* digested genomic DNA from the two strains with expected

bands indicated by arrows. **(B)** Growth of wild type of *M. smegmatis* ( $\diamond$ ), *M. smegmatis* $\Delta$ *aftC* ( $\square$ ), *M. smegmatis* $\Delta$ *aftC* pMV261-Ms-*aftC* ( $\triangle$ ), and *M. smegmatis* $\Delta$ *aftC* pMV261-Mt-*aftC* ( $\circ$ ) on TSB medium. **(C)** Colony morphology of wild type *M. smegmatis* and *M. smegmatis* $\Delta$ *aftC* on TSB/agar plates. Black bar represents 1 mm.

**Fig. 4. Analysis of cell wall associated lipids and bound MAMEs from *M. smegmatis*, *M. smegmatis* $\Delta$ *aftC*, *M. smegmatis* $\Delta$ *aftC* pMV261-Ms-*aftC* and *M. smegmatis* $\Delta$ *aftC* pMV261-Mt-*aftC*.** **(A)** Analysis of cell wall bound MAMEs from *M. smegmatis*, *M. smegmatis* $\Delta$ *aftC*, *M. smegmatis* $\Delta$ *aftC* pMV261-Ms-*aftC* and *M. smegmatis* $\Delta$ *aftC* pMV261-Mt-*aftC*. The bound mycolic acids from an equivalent amount of freeze-dried cells (100 mg), which were initially de-lipidated using two consecutive extractions of  $\text{CHCl}_3$ : $\text{CH}_3\text{OH}$ : $\text{H}_2\text{O}$  (10/10/3; v/v/v) at 50°C for 4 h, were released by the addition of tetra-butylammonium hydroxide at 100°C overnight, and methylated as described in the “Experimental Procedures”. An equivalent aliquot from each strain was subjected to TLC using silica gel plates (5735 silica gel 60F<sub>254</sub>, Merck), and developed in petroleum ether/acetone (95:5, v/v) and charred to reveal MAMEs and compared to known standards (Gande *et al.*, 2004). **(B)** Quantitative analysis of extractable [<sup>14</sup>C]-lipids from *M. smegmatis*, *M. smegmatis* $\Delta$ *aftC*, *M. smegmatis* $\Delta$ *aftC* pMV261-Ms-*aftC* and *M. smegmatis* $\Delta$ *aftC* pMV261-Mt-*aftC*. Lipids were extracted from cells by a series of organic washes as described in ‘Experimental Procedures’ (Seidel *et al.*, 2007a). An equivalent aliquot (50, 000 cpm) from each strain was subjected to TLC using silica gel plates (5725 silica gel 60F<sub>254</sub>, Merck) developed in  $\text{CHCl}_3$ : $\text{CH}_3\text{OH}$ : $\text{NH}_4\text{OH}$  (80:20:2, v/v/v) and quantified using phosphorimaging and compared to known standards (Mikusova *et al.*, 1995) after exposure to Kodak X-Omat film for 24 hours.

**Fig. 5. GC analysis of cell walls of *M. smegmatis*, *M. smegmatis* $\Delta$ *aftC*, *M. smegmatis* $\Delta$ *aftC* pMV261-*Ms-aftC* and *M. smegmatis* $\Delta$ *aftC* pMV261-*Mt-aftC*.** Samples of purified cell walls were hydrolyzed with 2M TFA, reduced, per-*O*-acetylated and analyzed as described under “Experimental Procedures” (Besra *et al.*, 1995; Alderwick *et al.*, 2005).

**Fig. 6. GC/MS analysis of cell walls of *M. smegmatis*, *M. smegmatis* $\Delta$ *aftC*, *M. smegmatis* $\Delta$ *aftC* pMV261-*Ms-aftC* and *M. smegmatis* $\Delta$ *aftC* pMV261-*Mt-aftC*.** Samples of per-*O*-methylated cell walls were hydrolyzed with 2M TFA, reduced, per-*O*-acetylated and analyzed as described under “Experimental Procedures” (Besra *et al.*, 1995; Alderwick *et al.*, 2005).

**Fig. 7. Arabinofuranosyltransferase activity utilizing an Ara<sub>2</sub> acceptor and membranes prepared from *M. smegmatis*, *M. smegmatis* $\Delta$ *aftC* and *M. smegmatis* $\Delta$ *aftC* pMV261-*Mt-aftC*.** (A) Biosynthetic reaction scheme of products A and B formed in arabinofuranosyltransferase assays using the neoglycolipid Ara<sub>2</sub> acceptor. (B) Arabinofuranosyltransferase activity was determined using the synthetic Ara<sub>2</sub> acceptor in a cell-free assay with and without EMB (1 mg/ml) as previously described (Lee *et al.*, 1997). The products of the assay were resuspended prior to scintillation counting (10 %) and the remaining subjected to TLC using silica gel plates (5735 silica gel 60F<sub>254</sub>, Merck) in CHCl<sub>3</sub>:CH<sub>3</sub>OH:H<sub>2</sub>O:NH<sub>4</sub>OH (65/25/3.6/0.5, v/v/v/v) with the reaction products visualized by autoradiography. The TLC autoradiogram is representative of several independent experiments.

**Fig. 8. Arabinofuranosyltransferase activity utilizing an Ara<sub>5</sub> acceptor and membranes prepared from *M. smegmatis*, *M. smegmatis* $\Delta$ *aftC* and *M. smegmatis* $\Delta$ *aftC* pMV261-*Mt-aftC*.** (A) Arabinofuranosyltransferase activity was determined using the synthetic Ara<sub>5</sub> acceptor

in a cell-free assay with and without EMB (1 mg/ml). The products reflective of three independent enzyme preparations and assays were resuspended prior to scintillation counting (10 %) and the remaining subjected to TLC using silica gel plates (5735 silica gel 60F<sub>254</sub>, Merck) in isopropanol:acetic acid:water (8/1/1/, v/v/v) with the reaction product X visualized by autoradiography. The TLC autoradiogram is representative of three independent experiments. (B) Ara<sub>5</sub> and Ara<sub>6</sub> (Appelmeik *et al.*, 2008) acceptor standards were subjected to TLC using silica gel plates (5735 silica gel 60F<sub>254</sub>, Merck) in isopropanol:acetic acid:water (8/1/1/, v/v/v) with the reaction products visualized by staining with  $\alpha$ -naphthol followed by charring.

**Fig. 9. GC characterization of *in vitro* synthesized product X from the arabinofuranosyltransferase assays utilizing the Ara<sub>5</sub> acceptor.** (A) GC analysis of the partially per-*O*-methylated, per-*O*-acetylated alditol acetate derivative of product X obtained from assays containing membranes prepared from either *M. smegmatis*, *M. smegmatis* $\Delta$ *aftC* or *M. smegmatis* $\Delta$ *aftC* pMV261-Mt-*aftC* (Sasaki *et al.*, 2005). (B) Panel illustrates the structure(s) of product X.

**Fig. 10. Mycobacterial arabinan biosynthesis and the role of AftC.**



## References


- Alderwick, L.J., Radmacher, E., Seidel, M., Gande, R., Hitchen, P.G., Morris, H.R., Dell, A., Sahm, H., Eggeling, L., and Besra, G.S. (2005) Deletion of *Cg-emb* in corynebacterianeae leads to a novel truncated cell wall arabinogalactan, whereas inactivation of *Cg-ubiA* results in an arabinan-deficient mutant with a cell wall galactan core. *J Biol Chem* **280**: 32362-32371.
- Alderwick, L.J., Seidel, M., Sahm, H., Besra, G.S., and Eggeling, L. (2006a) Identification of a novel arabinofuranosyltransferase (AftA) Involved in cell wall arabinan biosynthesis in *Mycobacterium tuberculosis*. *J Biol Chem* **281**: 15653-15661.
- Alderwick, L.J., Dover, L.G., Seidel, M., Gande, R., Sahm, H., Eggeling, L., and Besra, G.S. (2006b) Arabinan-deficient mutants of *Corynebacterium glutamicum* and the consequent flux in decaprenylmonophosphoryl-D-arabinose metabolism. *Glycobiology* **16**: 1073-1081.
- Amin, A.G., Goude, R., Shi, L., Zhang, J., Chatterjee, D., and Parish, T. (2008) EmbA is an essential arabinosyltransferase in *Mycobacterium tuberculosis*. *Microbiology* **154**: 240-248.
- Appelmek, B.J., den Dunnen, J., Driessen, N.N., Ummels, R., Pak, M., Nigou, J., Larrouy-Maumus, G., Gurcha, S.S., Movahedzadeh, F., Geurtsen, J., Brown, E.J., Eysink Smeets, M.M., Besra, G.S., Willemsen, P.T.J., Lowary, T.L., van Kooyk, Y., Maaskant, J.J., Stoker, N.G., van der Ley, P., Puzo, G., Vandenbroucke-Grauls, J.E., Wieland, C.W., van der Poll, T., Geijtenbeek, T., van der Sar, A.M., and Bitter, W. (2008) The mannose cap of mycobacterial lipoarabinomannan does not dominate the *Mycobacterium*-host interaction. *Cellular Microbiology* **10**: 930-944.
- Bardarov, S., Bardarov Jr, S., Jr., Pavelka Jr, M.S., Jr., Sambandamurthy, V., Larsen, M., Tufariello, J., Chan, J., Hatfull, G., and Jacobs Jr, W.R., Jr. (2002) Specialized transduction: an efficient method for generating marked and unmarked targeted gene disruptions in *Mycobacterium tuberculosis*, *M. bovis* BCG and *M. smegmatis*. *Microbiology* **148**: 3007-3017.
- Belanger, A.E., Besra, G.S., Ford, M.E., Mikusova, K., Belisle, J.T., Brennan, P.J., and Inamine, J.M. (1996) The *embAB* genes of *Mycobacterium avium* encode an arabinosyl transferase involved in cell wall arabinan biosynthesis that is the target for the antimycobacterial drug ethambutol. *Proc Natl Acad Sci U S A* **93**: 11919-11924.
- Besra, G.S., Khoo, K.H., McNeil, M.R., Dell, A., Morris, H.R., and Brennan, P.J. (1995) A new interpretation of the structure of the mycolyl-arabinogalactan complex of *Mycobacterium tuberculosis* as revealed through characterization of oligoglycosylalditol fragments by fast-atom bombardment mass spectrometry and <sup>1</sup>H nuclear magnetic resonance spectroscopy. *Biochemistry* **34**: 4257-4266.
- Bhamidi, S., Scherman, M.S., Rithner, C.D., Prenni, J.E., Chatterjee, D., Khoo, K.H., and McNeil, M.R. (2008) The identification and location of succinyl residues and the characterization of the interior arabinan region allows for a model of the complete primary structure of *Mycobacterium tuberculosis* mycolyl arabinogalactan. *J Biol Chem.* **283**: 12992-13000.
- Bhatt, A., Kremer, L., Dai, A.Z., Sacchettini, J.C., and Jacobs, W.R., Jr. (2005) Conditional depletion of KasA, a key enzyme of mycolic acid biosynthesis, leads to mycobacterial cell lysis. *J Bacteriol* **187**: 7596-7606.





- Brennan, P.J., and Nikaido, H. (1995) The envelope of mycobacteria. *Annu Rev Biochem* **64**: 29-63.
- Brudno, M., Chapman, M., Gottgens, B., Batzoglou, S., and Morgenstern, B. (2003) Fast and sensitive multiple alignment of large genomic sequences. *BMC Bioinformatics* **4**: 66.
- Daffé, M., Brennan, P.J., and McNeil, M. (1990) Predominant structural features of the cell wall arabinogalactan of *Mycobacterium tuberculosis* as revealed through characterization of oligoglycosyl alditol fragments by gas chromatography/mass spectrometry and by  $^1\text{H}$  and  $^{13}\text{C}$  NMR analyses. *J Biol Chem* **265**: 6734-6743.
- Dover, L.G., Cerdano-Tarraga, A.M., Pallen, M.J., Parkhill, J., and Besra, G.S. (2004) Comparative cell wall core biosynthesis in the mycolated pathogens, *Mycobacterium tuberculosis* and *Corynebacterium diphtheriae*. *FEMS Microbiol Rev* **28**: 225-250.
- Dye, C. (2006) Global epidemiology of tuberculosis. *Lancet* **367**: 938-940.
- Eggeling, L., and Bott, M. (2005) *Handbook of Corynebacterium glutamicum*.: Boca Raton, FL 33487-2742, CRC Press, Taylor Francis Group.
- Escuyer, V.E., Lety, M.A., Torrelles, J.B., Khoo, K.H., Tang, J.B., Rithner, C.D., Frehel, C., McNeil, M.R., Brennan, P.J., and Chatterjee, D. (2001) The role of the *embA* and *embB* gene products in the biosynthesis of the terminal hexaarabinofuranosyl motif of *Mycobacterium smegmatis* arabinogalactan. *J Biol Chem* **276**: 48854-48862.
- Gande, R., Gibson, K.J., Brown, A.K., Krumbach, K., Dover, L.G., Sahm, H., Shioyama, S., Oikawa, T., Besra, G.S., and Eggeling, L. (2004) Acyl-CoA carboxylases (*accD2* and *accD3*), together with a unique polyketide synthase (*Cg-pks*), are key to mycolic acid biosynthesis in *Corynebacteriaceae* such as *Corynebacterium glutamicum* and *Mycobacterium tuberculosis*. *J Biol Chem* **279**: 44847-44857.
- Guvener, Z.T., and McCarter, L.L. (2003) Multiple regulators control capsular polysaccharide production in *Vibrio parahaemolyticus*. *J Bacteriol* **185**: 5431-5441.
- Kaye, K., and Frieden, T.R. (1996) Tuberculosis control: the relevance of classic principles in an era of acquired immunodeficiency syndrome and multidrug resistance. *Epidemiol Rev* **18**: 52-63.
- Lee, R.E., Mikusova, K., Brennan, P.J., and Besra, G.S. (1995) Synthesis of the arabinose anomer  $\beta$ -D-arabinofuranosyl-1-monophosphoryldecaprenol, development of a basic arabinosyltransferase assay, and identification of ethambutol as an arabinosyltransferase inhibitor. *J. Am. Chem. Soc* **117**: 11829-11832.
- Lee, R.E., Brennan, P.J., and Besra, G.S. (1997) Mycobacterial arabinan biosynthesis: the use of synthetic arabinoside acceptors in the development of an arabinosyl transfer assay. *Glycobiology* **7**: 1121-1128.
- Lee, R.E., Brennan, P.J., and Besra, G.S. (1998) Synthesis of  $\beta$ -D-arabinofuranosyl-1-monophosphoryl polyprenols: examination of their function as mycobacterial arabinosyl transferase donors. *Bioorg Med Chem Lett* **8**: 951-954.
- Liu, J., and Mushegian, A. (2003) Three monophyletic superfamilies account for the majority of the known glycosyltransferases. *Protein Sci* **12**: 1418-1431.
- McNeil, M., Daffe, M., and Brennan, P.J. (1990) Evidence for the nature of the link between the arabinogalactan and peptidoglycan of mycobacterial cell walls. *J Biol Chem* **265**: 18200-18206.
- McNeil, M., Daffe, M., and Brennan, P.J. (1991) Location of the mycolyl ester substituents in the cell walls of mycobacteria. *J Biol Chem* **266**: 13217-13223.
- Mikusova, K., Slayden, R.A., Besra, G.S., and Brennan, P.J. (1995) Biogenesis of the mycobacterial cell wall and the site of action of ethambutol. *Antimicrob Agents Chemother* **39**: 2484-2489.

- Mills, J.A., Motichka, K., Jucker, M., Wu, H.P., Uhlik, B.C., Stern, R.J., Scherman, M.S., Vissa, V.D., Pan, F., Kundu, M., Ma, Y.F., and McNeil, M. (2004) Inactivation of the mycobacterial rhamnosyltransferase, which is needed for the formation of the arabinogalactan-peptidoglycan linker, leads to irreversible loss of viability. *J Biol Chem* **279**: 43540-43546.
- Minnikin, D.E. (1982) Lipids: Complex lipids, Their Chemistry, Biosynthesis and Roles. In *The Biology of the Mycobacteria*. Ratledge, C. and Stanford, J. (eds). London: Academic Press, pp. 95-184.
- Mishra, A.K., Alderwick, L.J., Rittmann, D., Tatituri, R.V., Nigou, J., Gilleron, M., Eggeling, L., and Besra, G.S. (2007) Identification of an  $\alpha(1\rightarrow6)$  mannopyranosyltransferase (MptA), involved in *Corynebacterium glutamicum* lipomanann biosynthesis, and identification of its orthologue in *Mycobacterium tuberculosis*. *Mol Microbiol* **65**: 1503-1517.
- Mishra, A.K., Alderwick, L.J., Rittmann, D., Wang, C., Bhatt, A., Jacobs, W.R. Jr., Takayama, K., Eggeling, L., and Besra, G.S. (2008) Identification of a novel  $\alpha(1\rightarrow6)$  mannopyranosyltransferase (MptB) from *Corynebacterium glutamicum* by deletion of a conserved gene, NCgl1505, affords a lipomanann- and lipoarabinomann-deficient mutant. *Mol Microbiol*. **Paper in press**.
- Pan, F., Jackson, M., Ma, Y., and McNeil, M. (2001) Cell wall core galactofuran synthesis is essential for growth of mycobacteria. *J Bacteriol* **183**: 3991-3998.
- Paolo, W.F., Jr., and Nosanchuk, J.D. (2004) Tuberculosis in New York city: recent lessons and a look ahead. *Lancet Infect Dis* **4**: 287-293.
- Parish, T., Roberts, G., Laval, F., Schaeffer, M., Daffe, M., and Duncan, K. (2007) Functional complementation of the essential gene *fabG1* of *Mycobacterium tuberculosis* by *Mycobacterium smegmatis* *fabG* but not *Escherichia coli* *fabG*. *J Bacteriol* **189**: 3721-3728.
- Portevin, D., De Sousa-D'Auria, C., Houssin, C., Grimaldi, C., Chami, M., Daffe, M., and Guilhot, C. (2004) A polyketide synthase catalyzes the last condensation step of mycolic acid biosynthesis in mycobacteria and related organisms. *Proc Natl Acad Sci U S A* **101**: 314-319.
- Radmacher, E., Stansen, K.C., Besra, G.S., Alderwick, L.J., Maughan, W.N., Hollweg, G., Sahm, H., Wendisch, V.F., and Eggeling, L. (2005) Ethambutol, a cell wall inhibitor of *Mycobacterium tuberculosis*, elicits L-glutamate efflux of *Corynebacterium glutamicum*. *Microbiology* **151**: 1359-1368.
- Sasaki, G.L., Iacaomini, M., and Gorin, P.A.J. (2005) Methylation-GC-MS analysis of arabinofuranose-and galactofuranose-containing structures; rapid synthesis of partially O-methylated alditol acetates. *An. Acad. Bras. Ciênc.*, **77**: 223-234
- Schafer, A., Tauch, A., Jager, W., Kalinowski, J., Thierbach, G., and Puhler, A. (1994) Small mobilizable multi-purpose cloning vectors derived from the *Escherichia coli* plasmids pK18 and pK19: selection of defined deletions in the chromosome of *Corynebacterium glutamicum*. *Gene* **145**: 69-73.
- Seidel, M., Alderwick, L.J., Birch, H.L., Sahm, H., Eggeling, L., and Besra, G.S. (2007a) Identification of a novel arabinofuranosyltransferase AftB involved in a terminal step of cell wall arabinan biosynthesis in *Corynebacterianae*, such as *Corynebacterium glutamicum* and *Mycobacterium tuberculosis*. *J Biol Chem* **282**: 14729-14740.
- Seidel, M., Alderwick, L.J., Sahm, H., Besra, G.S., and Eggeling, L. (2007b) Topology and mutational analysis of the single Emb arabinofuranosyltransferase of *Corynebacterium glutamicum* as a model of Emb proteins of *Mycobacterium tuberculosis*. *Glycobiology* **17**: 210-219.


- 1
- 2
- 3 Shah, N.S., Wright, A., Bai, G.H., Barrera, L., Boulahbal, F., Martin-Casabona, N.,
- 4 Drobniewski, F., Gilpin, C., Havelkova, M., Lepe, R., Lumb, R., Metchock, B., Portaels,
- 5 F., Rodrigues, M.F., Rusch-Gerdes, S., Van Deun, A., Vincent, V., Laserson, K., Wells,
- 6 C., and Cegielski, J.P. (2007) Worldwide emergence of extensively drug-resistant
- 7 tuberculosis. *Emerg Infect Dis* **13**: 380-387.
- 8
- 9 Stover, C.K., de la Cruz, V.F., Fuerst, T.R., Burlein, J.E., Benson, L.A., Bennett, L.T., Bansal,
- 10 G.P., Young, J.F., Lee, M.H., Hatfull, G.F., and et al. (1991) New use of BCG for
- 11 recombinant vaccines. *Nature* **351**: 456-460.
- 12
- 13 Takayama, K., and Kilburn, J.O. (1989) Inhibition of synthesis of arabinogalactan by ethambutol
- 14 in *Mycobacterium smegmatis*. *Antimicrob Agents Chemother* **33**: 1493-1499.
- 15
- 16 Telenti, A., Philipp, W.J., Sreevatsan, S., Bernasconi, C., Stockbauer, K.E., Wiele, B., Musser,
- 17 J.M., and Jacobs, W.R., Jr. (1997) The *emb* operon, a gene cluster of *Mycobacterium*
- 18 *tuberculosis* involved in resistance to ethambutol. *Nat Med* **3**: 567-570.
- 19
- 20 Vilcheze, C., Morbidoni, H.R., Weisbrod, T.R., Iwamoto, H., Kuo, M., Sacchettini, J.C., and
- 21 Jacobs, W.R., Jr. (2000) Inactivation of the *inhA*-encoded fatty acid synthase II (FASII)
- 22 enoyl-acyl carrier protein reductase induces accumulation of the FASII end products and
- 23 cell lysis of *Mycobacterium smegmatis*. *J Bacteriol* **182**: 4059-4067.
- 24
- 25 Wolucka, B.A., McNeil, M.R., de Hoffmann, E., Chojnacki, T., and Brennan, P.J. (1994)
- 26 Recognition of the lipid intermediate for arabinogalactan/arabinomannan biosynthesis
- 27 and its relation to the mode of action of ethambutol on mycobacteria. *J Biol Chem* **269**:
- 28 23328-23335.
- 29
- 30 Zhang, N., Torrelles, J.B., McNeil, M.R., Escuyer, V.E., Khoo, K.H., Brennan, P.J., and
- 31 Chatterjee, D. (2003) The Emb proteins of mycobacteria direct arabinosylation of
- 32 lipoarabinomannan and arabinogalactan via an N-terminal recognition region and a C-
- 33 terminal synthetic region. *Mol Microbiol* **50**: 69-76.
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60




117x252mm (300 x 300 DPI)




52x96mm (600 x 600 DPI)



212x323mm (300 x 300 DPI)




173x298mm (400 x 400 DPI)




160x336mm (400 x 400 DPI)






348x258mm (300 x 300 DPI)


review




220x241mm (400 x 400 DPI)



238x148mm (400 x 400 DPI)



262x340mm (300 x 300 DPI)



254x190mm (300 x 300 DPI)

review