

Narrative-based computational modelling of the Gp130/JAK/STAT signalling pathway

Guerriero, ML; Dudka, A; Underhill-Day, Nicholas; Heath, John; Priami, C

DOI:

[10.1186/1752-0509-3-40](https://doi.org/10.1186/1752-0509-3-40)

Citation for published version (Harvard):

Guerriero, ML, Dudka, A, Underhill-Day, N, Heath, J & Priami, C 2009, 'Narrative-based computational modelling of the Gp130/JAK/STAT signalling pathway', *BMC systems biology*, vol. 3, pp. 40. <https://doi.org/10.1186/1752-0509-3-40>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

id	description	react	alt
LIF pathway → type I receptor gp130:LIFR			
11	if gp130.LIF is bound and LIFR.LIF is not bound and gp130.OSM is not bound and LIFR.OSM is not bound and LIFR is not dimer and gp130.typeI is not dimer and gp130.typeII is not dimer then LIFR dimerizes with gp130 on typeI	11	2
12	if gp130.LIF is bound and LIFR.LIF is not bound and gp130.OSM is not bound and LIFR.OSM is not bound and LIFR is dimer and gp130.typeI is dimer and gp130.typeII is not dimer and LIFR.Y981 is not phospho and gp130.Y767 is not phospho then LIFR dedimerizes from gp130 on typeI	12	
13	if LIFR.LIF is bound and gp130.LIF is not bound and LIFR.OSM is not bound and gp130.OSM is not bound and LIFR is not dimer and gp130.typeI is not dimer and gp130.typeII is not dimer then LIFR dimerizes with gp130 on typeI	11	4
14	if LIFR.LIF is bound and gp130.LIF is not bound and LIFR.OSM is not bound and gp130.OSM is not bound and LIFR is dimer and gp130.typeI is dimer and gp130.typeII is not dimer and LIFR.Y981 is not phospho and gp130.Y767 is not phospho then LIFR dedimerizes from gp130 on typeI	12	
OSM pathway → type I receptor gp130:LIFR			
15	if gp130.OSM is bound and LIFR.OSM is not bound and gp130.LIF is not bound and LIFR.LIF is not bound and LIFR is not dimer and gp130.typeI is not dimer and gp130.typeII is not dimer then LIFR dimerizes with gp130 on typeI	11	6
16	if gp130.OSM is bound and LIFR.OSM is not bound and gp130.LIF is not bound and LIFR.LIF is not bound and LIFR is dimer and gp130.typeI is dimer and gp130.typeII is not dimer and LIFR.Y981 is not phospho and gp130.Y767 is not phospho then LIFR dedimerizes from gp130 on typeI	12	
17	if LIFR.OSM is bound and gp130.OSM is not bound and LIFR.LIF is not bound and gp130.LIF is not bound and LIFR is not dimer and gp130.typeI is not dimer and gp130.typeII is not dimer then LIFR dimerizes with gp130 on typeI	11	8
18	if LIFR.OSM is bound and gp130.OSM is not bound and LIFR.LIF is not bound and gp130.LIF is not bound and LIFR is dimer and gp130.typeI is dimer and gp130.typeII is not dimer and LIFR.Y981 is not phospho and gp130.Y767 is not phospho then LIFR dedimerizes from gp130 on typeI	12	
OSM pathway → type II receptor gp130:OSMR			
19	if gp130.OSM is bound and OSMR.OSM is not bound and gp130.LIF is not bound and OSMR.LIF is not bound and OSMR is not dimer and gp130.typeI is not dimer and gp130.typeII is not dimer then OSMR dimerizes with gp130 on typeII	11	6
20	if gp130.OSM is bound and OSMR.OSM is not bound and gp130.LIF is not bound and OSMR.LIF is not bound and OSMR is dimer and gp130.typeI is not dimer and gp130.typeII is dimer and OSMR.Y917 is not phospho and gp130.Y767 is not phospho then OSMR dedimerizes from gp130 on typeII	12	
21	if OSMR.OSM is bound and gp130.OSM is not bound and OSMR.LIF is not bound and gp130.LIF is not bound and OSMR is not dimer and gp130.typeI is not dimer and gp130.typeII is not dimer then OSMR dimerizes with gp130 on typeII	11	10
22	if OSMR.OSM is bound and gp130.OSM is not bound and OSMR.LIF is not bound and gp130.LIF is not bound and OSMR is dimer and gp130.typeI is not dimer and gp130.typeII is dimer and OSMR.Y917 is not phospho and gp130.Y767 is not phospho then OSMR dedimerizes from gp130 on typeII	12	