UNIVERSITY^{OF} BIRMINGHAM University of Birmingham Research at Birmingham

Wnt4 and LAP2alpha as Pacemakers of Thymic Epithelial Senescence

Kvell, K; Varecza, Z; Bartis, D; Hesse, S; Parnell, Sonia; Anderson, Graham; Jenkinson, Eric; Pongracz, JE

DOI: 10.1371/journal.pone.0010701

Document Version Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Kvell, K, Varecza, Z, Bartis, D, Hesse, S, Parnell, S, Anderson, G, Jenkinson, E & Pongracz, JE 2010, 'Wnt4 and LAP2alpha as Pacemakers of Thymic Epithelial Senescence', *PLoS ONE*, vol. 5, no. 5, pp. e10701. https://doi.org/10.1371/journal.pone.0010701

Link to publication on Research at Birmingham portal

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

•Users may freely distribute the URL that is used to identify this publication.

•Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.

•User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?) •Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Wnt4 and LAP2alpha as Pacemakers of Thymic Epithelial Senescence

Krisztian Kvell¹, Zoltan Varecza¹, Domokos Bartis¹, Sebastian Hesse¹, Sonia Parnell², Graham Anderson², Eric J. Jenkinson², Judit E. Pongracz^{1,2}*

1 Department of Medical Biotechnology, Institute for Immunology and Biotechnology, University of Pecs, Pecs, Hungary, 2 Division of Immunity and Infection, Department of Anatomy, Institute for Biomedical Research, University of Birmingham, Birmingham, United Kingdom

Abstract

Age-associated thymic involution has considerable physiological impact by inhibiting *de novo* T-cell selection. This impaired T-cell production leads to weakened immune responses. Yet the molecular mechanisms of thymic stromal adipose involution are not clear. Age-related alterations also occur in the murine thymus providing an excellent model system. In the present work structural and molecular changes of the murine thymic stroma were investigated during aging. We show that thymic epithelial senescence correlates with significant destruction of epithelial network followed by adipose involution. We also show in purified thymic epithelial cells the age-related down-regulation of Wnt4 (and subsequently FoxN1), and the prominent increase in LAP2 α expression. These senescence-related changes of gene expression are strikingly similar to those observed during mesenchymal to pre-adipocyte differentiation of fibroblast cells suggesting similar molecular background in epithelial cells. For molecular level proof-of-principle stable LAP2 α and Wnt4-over-expressing thymic epithelial cell lines were established. LAP2 α over-expression provoked a surge of PPAR γ expression, a transcription factor expressed in pre-adipocytes. In contrast, additional Wnt4 decreased the mRNA level of ADRP, a target gene of PPARy. Murine embryonic thymic lobes have also been transfected with LAP2a- or Wnt4-encoding lentiviral vectors. As expected LAP2 α over-expression increased, while additional Wnt4 secretion suppressed PPAR γ expression. Based on these pioneer experiments we propose that decreased Wnt activity and increased LAP2 α expression provide the molecular basis during thymic senescence. We suggest that these molecular changes trigger thymic epithelial senescence accompanied by adipose involution. This process may either occur directly where epithelium can trans-differentiate into pre-adipocytes; or indirectly where first epithelial to mesenchymal transition (EMT) occurs followed by subsequent pre-adipocyte differentiation. The latter version fits better with literature data and is supported by the observed histological and molecular level changes.

Citation: Kvell K, Varecza Z, Bartis D, Hesse S, Parnell S, et al. (2010) Wnt4 and LAP2alpha as Pacemakers of Thymic Epithelial Senescence. PLoS ONE 5(5): e10701. doi:10.1371/journal.pone.0010701

Editor: Immo A. Hansen, New Mexico State University, United States of America

Received February 4, 2010; Accepted April 27, 2010; Published May 18, 2010

Copyright: © 2010 Kvell et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: Research was supported by the following grants: The Wellcome Trust grant No.: 079415 (grant-holders: J.E.P., G.A. and E.J.J.), 'Science Please' Research Team on Innovation grant No.: SROP-4.2.2/08/1/2008-0011 (grant holder: J.E.P.) and OTKA (Hungarian Scientific Research Fund) type: PD (post-doctoral) grant No.: 78310 (grant-holder: K.K.). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: judit.e.pongracz@aok.pte.hu

Introduction

Thymic senescence

Thymic senescence begins early, around late puberty. This process is called adipose involution, as the thymus is invaded by adipose tissue [1]. Due to decrease in thymic epithelial tissue mass, the thymus can no longer support the same output of T-cell production [2]. Therefore peripheral blood T lymphocyte composition exhibits the dominance of memory T lymphocytes resulting in impaired responses towards novel, particularly viral infections [3,4,5]. Since the thymic epithelium has a key role in deleting auto-reactive T-cell clones, functional impairment increases the chances of developing auto-immune disease [6]. If we were able to slow down or even stop the loss of thymic epithelium the elderly would have a better chance to address lateonset autoimmune diseases and viral infections. However, despite studies of thymic senescence, the molecular mechanism of thymic aging remains elusive.

Signaling pathways of thymic epithelial cell development and maintenance

Understanding signaling mechanisms that regulate tissue development and maintenance of thymic epithelial cells might reveal the process of adipose involution. Certainly, maintenance and functional integrity of the thymic stroma requires stimuli through Notch, BMP, and Wnt signaling pathways [7,8,9,10,11]. Undoubtedly, the Wnt family of secreted glycoproteins is one of the best analyzed among the required ligands [12]. Most members of the nineteen known Wnt glycoproteins have been implicated in both the development of embryonic thymus and the maintenance of adult thymic epithelium [13]. In the thymus, Wnt ligands originate primarily from thymic epithelial cells and activate a highly complex signaling network via ten G-protein dependent receptors called Frizzleds (Fz), and their co-receptors of lowdensity lipoprotein receptor-related proteins 5/6 called LRP5/6 [14,15]. The actual constellation of ligands, receptors, co-receptors and further regulatory molecules define Wnt-mediated effects.

Recent studies have highlighted Wnt4 as responsible for the direct up-regulation of FoxN1, a key transcription factor responsible for the differentiation of thymic epithelial cells and the subsequent maintenance of thymic epithelial identity [13]. Interestingly, the Wnt/ β -catenin pathway is known to efficiently block the adipocyte differentiation program in mesenchymal elements like fibroblasts [16,17,18,19].

Trans-differentiation of fibroblasts into adipocytes

Studies with fibroblast cells have also revealed that fibroblast to pre-adipocyte transformation is strongly connected to LAP2a, the member of the LAP2 protein family [17]. To date there are 7 classified intranuclear LAP2 polypeptides marked by the Greek alphabet. They are all splice variants of the same LAP2 gene previously called thymopoietin. While most splice variants associate with the nuclear envelope, LAP 2α is involved in several nucleoplasmic activities including cell-cycle control and differentiation [20,21]. LAP2 α is synthesized in the cytoplasm and is then transported into the nucleus by a PKC-dependent mechanism [22]. The mere over-expression of LAP2 α in fibroblasts is known to directly up-regulate PPARy expression, an acknowledged marker and key transcription factor of pre-adipocyte differentiation [17]. In pre-adipocytes PPAR γ expression is followed by an increase of ADRP expression (adipose differentiation-related protein) a known direct target gene of PPAR γ . Although LAP2 α over-expression alone initiates pre-adipocyte differentiation in fibroblasts, it is not sufficient to complete the adipocyte differentiation program in the absence of additional stimuli [17].

Results and Discussion

Disintegration of epithelial network

Senescence exhibits characteristic histological changes in both the human and mouse thymus [1,23]. In order to demonstrate this process the thymic lobes of 1 month and 1 year old BALB/c mice were analyzed (see Figures 1A and 1B). In young adult mice, histology revealed strict segregation of epithelial cell compartments by staining for medullary (EpCAM1⁺⁺, Ly51⁻) and cortical (EpCAM1⁺, Ly51⁺⁺) epithelial cellular subsets (Figure 1A). This shows high level of morphological integrity just preceding puberty/early adulthood. However, the highly organized structure disintegrates and becomes chaotic by the age of 1 year (Figure 1B). By this age the previously shown strict cortico-medullary delineation becomes disintegrated, degenerative vacuoles appear surrounded by areas showing strong co-staining with both epithelial markers. There are also other large cellular areas that lack staining with either epithelial marker, a pattern completely absent at the young adult age.

Staining of extracellular matrix components of fibroblast origin (ER-TR7^{++}) was also performed on cryostate thymic sections of 2



Figure 1. Disintegration of epithelial network. Figure 1A demonstrates cryostat section of 1 month, whereas figure 1B presents cryostat section of 1 year old BALB/c mouse thymus. Staining pattern: anti-EpCAM1-FITC (green), anti-Ly51-PE (red), DAPI (blue). 'M' marks medullary (EpCAM1⁺⁺, Ly51⁻), while 'C' marks cortical (EpCAM1⁺, Ly51⁺⁺) epithelial compartments on Figure 1A. Single asterisk (*) marks degenerative vacuoles, while double asterisk (**) mark the loss of epithelial staining on Figure 1B. Figure 1C (lower left) shows cryostate section of 2 month, whereas figure 1D (lower right) demonstrates cryostate section of 9 month old BALB/c mouse thymus. Staining pattern: anti-EpCAM1-FITC, ER-TR7-PE, DAPI (blue). The EpCAM1⁺⁺ thymic medulla is outlined by continuous line on Figures 1C and 1D for easier visualization. doi:10.1371/journal.pone.0010701.g001

month and 9 month old BALB/c mice to identify epithelial and mesenchymal elements in young adult and aging thymic lobes. The above ages were selected to check additional time points and more precisely map the timeframe of thymic physiological senescence (see Figures 1C and 1D). The staining patterns are strikingly different at the two ages examined. In the 2 month old thymic tissue section a-EpCAM1 and ER-TR7-staining show little tendency for co-localization. In stark contrast, by the age of 9 months a-EpCAM1 and ER-TR7-staining show significant overlap within the thymic medulla, a phenomenon completely absent at earlier ages.

Adipose involution

To demonstrate how the disorganization of thymic epithelial network is followed by the emergence of adipocytes, thymic sections of 1.5 year old GFP-transgenic BALB/c mice were analyzed. This mouse strain develops and reproduces exactly like control BALB/c mice, and the thymic epithelial function and thymocyte maturation is indistinguishable from wild type controls [24]. However, due to the ubiquitous and strong EF1 promoter-driven transgene transcription, bright GFP expression offers a native, green-colored, cytoplasmic staining for all the cells in these mice. Thymic sections of senescent GFP-transgenic mice were co-stained with LipidTox Red to identify adipocytes. Histology shows the presence of relatively large, inflated cells in which the green-colored (GFP-containing) cytoplasm is pushed to the periphery by red-staining neutral lipid deposits, a pattern characteristic of adipose cells (see Figure 2).

Molecular changes of thymic epithelium

Having presented structural changes of thymic epithelial senescence, we set out to investigate the underlying molecular events. In order to detect gene expression changes, thymic epithelial cells were purified from 1 month and 1 year old BALB/c mice based on EpCAM1 expression (MACS separation). Following cDNA synthesis, quantitative RT-PCR analysis was performed. Several genes including Wnt4, FoxN1, PPAR γ , ADRP, lamin1 and LAP2 α were tested (Table 1 lists primer sequences and characteristics, see Figures 3A–D for changes in gene expression). Figure 3A shows that the expression of both Wnt4 and FoxN1 decreases in thymic epithelial cells. Highly decreased level (or total absence in some cases) of FoxN1 could be



Figure 2. Adipose involution. Figure 2 shows adipose involution over cryostat section of 1.5 year old GFP-transgenic BALB/c mouse thymus. Staining pattern: GFP (green), LipidTox Red (red). doi:10.1371/journal.pone.0010701.g002

the consequence of strong Wnt4 down-regulation by the age of 1 year, indicating that thymic epithelial cells can down-regulate FoxN1 expression while maintaining that of epithelial cell surface markers like EpCAM1 [13]. At the same time, mRNA levels of pre-adipocyte differentiation markers PPARy and ADRP rise with age in the same, EpCAM1-positive cell population (Figure 3C). This finding is in harmony with histological data demonstrating the emergence of adipocytes in the thymic lobes of senescent mice (Figure 2). The expression of lamin1, a key component of the nuclear lamina remains unaffected during senescence in thymic epithelial cells; whereas, the expression of LAP2 α increases significantly (see Figure 3B). This degree of dissociation between lamin1 and LAP2 α expression is of note and suggests functional differences despite conventionally anticipated association of lamin1 and LAP2 molecular family members. The measured LAP2 α up-regulation associated with age-related adipose involution is, however, in perfect agreement with other literature data suggesting the pre-adipocyte differentiation-promoting effect of LAP 2α in fibroblasts [17]. This is the first report to show that such, normally fibroblast associated molecular changes occur in purified thymic epithelial cells. In the literature, epithelial-mesenchymal transition is associated with differential expression of E- and Ncadherin [25]. While E-cadherin decreases, N-cadherin normally compensates for the loss of E-cadherin expression. To investigate whether the first step towards pre-adipocyte differentiation is the epithelial-mesenchymal transition of epithelial cells, gene expression changes of E-cadherin and N-cadherin were measured (Figure 3D). While E-cadherin mRNA levels significantly decreased, N-cadherin gene expression showed a slight increase, indicating that EMT might be the initial step in epithelial cell transition to become pre-adipocytes.

Transgenic cell lines

Stable LAP2a over-expressing or Wnt4-secreting transgenic TEP1 cell lines were established using lentiviral transgenesis. The use of a primary-derived model cell line provides the advantage of absolute purity, the complete lack of other cell types that could potentially affect the gene expression profile of epithelial cells [26]. The established transgenic cell lines proliferated normally and did not show obvious signs of phenotypic changes (data not shown). In contrast to morphology, quantitative RT-PCR analysis revealed that LAP2 α over-expression triggers an immense surge of PPAR γ expression (Figure 4). Such an increase in mRNA level suggests that this is not a plain quantitative, but rather a qualitative change. ADRP a direct target gene of PPARy was also up-regulated albeit to a lesser extent (Figure 4). On the other hand in Wnt4-secreting TEP1 cells the mRNA level of both PPAR γ and ADRP was decreased (Figure 4). In the TEP1 cell line the expression of FoxN1 could not be addressed as it is very low/undetectable and remains as such with all the tested treatments (data not shown).

Transfected embryonic thymic organ cultures

To confirm the involvement of LAP2 α and Wnt4 during adipogenesis through their direct effect on PPAR γ expression in primary cells, murine thymic lobes were isolated from timed pregnancies at E12. Thymic lobes at the age of E12 provide an excellent experimental setting where the thymus has just been formed and there is no sign of aging. Furthermore, thymic lobes at this stage are also small enough to be both cultured and transfected as a whole, nutrients and virions have free access to most of the cells in the lobe without the need of disrupting any intercellular connection or tissue matrix [8,27]. The isolated lobes were therefore transfected with lentiviral vectors encoding GFP (mock), Wnt4 or LAP2 α and were cultured for 4 days *in vitro*. Q-PCR was

Table 1. List of gene	specific PCR primers.
-----------------------	-----------------------

Gene	Forward primer	Reverse primer
β-actin	5'-TGG CGC TTT TGA CTC AGG A -3'	5'-GGG AGG GTG AGG GAC TTC C - 3'
Wnt4	5'-CTC AAA GGC CTG ATC CAG AG - 3'	5'-TCA CAG CCA CAC TTC TCC AG - 3'
LAP2α	5'-TGA ACT GCA GGC AGC TAA GA-3'	5'-TCA TAG CTA GAC TCT GAG G-3'
Lamin1	5' - TGA GTA CAA CCT GCG CTC AC -3'	5' - TGA CTA GGT TGT CCC CGA AG -3'
ΡΡΑRγ	5' - CCC AAT GGT TGC TGA TTA CAA A -3'	5' - AAT AAT AAG GTG GAG ATG CAG GTT CT -3'
ADRP	5' - CGC CAT CGG ACA CTT CCT TA -3'	5' - GTG ATG GCA GGC GAC ATC T -3'
E-cadherin	5'- AAG TGA CCG ATG ATG ATG CC -3'	5'- CTT CAT TCA CGT CTA CCA CGT -3'
N-cadherin	5' - GTG GAG GCT TCT GGT GAA AT - 3'	5' - CTG CTG GCT CGC TGC TT - 3'
FoxN1	Applied Biosystems TaqMan probe PN4351272 (Mm00477457_m1)	

doi:10.1371/journal.pone.0010701.t001

performed to confirm over-expression of LAP2 α and Wnt4 in the embryonic thymic lobes as a result of lentiviral transgenesis (Figure 5A) and their effect on PPAR γ expression was also analyzed (Figure 5B). The level of over-expression was confirmed following transfection with both LAP2 α - and Wnt4-encoding viral vectors. Q-PCR analysis revealed that LAP2 α over-expression triggers an increase of PPAR γ expression, whereas additional Wnt4 secretion suppresses PPAR γ level (Figure 5B). The latter Wnt4-mediated suppression of PPAR γ expression in cultures of E12 thymic embryonic lobes was also confirmed by treatment with Wnt4-containing supernatants of Wnt4 over-expressing TEP1 cell line (data not shown). Interestingly, the expression of FoxN1 did not decrease in LAP2 α over-expressing thymic lobes (data not shown), possibly due to high levels of Wnt4 in the embryonic thymic tissue preserving FoxN1 status. Our molecular studies using E12 thymic lobes confirmed our data obtained with the TEP1 cell lines, that even in embryonic thymic tissue preadipocyte differentiation markers can be up-regulated in the presence of LAP2 α , indicating that the process can be dissected and controlled at a molecular level.

Conclusion

Here we show that with senescence, thymic epithelial Wnt4 secretion decreases, possibly below a threshold level that is required to maintain the identity of established thymic epithelial cells. This is measured by the loss of FoxN1 expression, a key transcription factor defining thymic epithelial cell identity. However, these epithelial cells still express cell surface markers



Figure 3. Molecular changes in thymic epithelium. Figures 3A–D demonstrate gene expression changes of MACS purified thymic epithelial cells measured by Q-PCR. Please note that the Y-axis scale is logarithmic. Error bars show ±1 SD. doi:10.1371/journal.pone.0010701.g003



Figure 4. Confirmation in transgenic thymic cell lines. Figure 4 shows gene expression changes of LAP2 α and Wnt4 over-expressing transgenic TEP1 cells measured by Q-PCR. Please note that Y-axis scale is logarithmic. Error bars show ± 1 SD. doi:10.1371/journal.pone.0010701.g004

characteristic for thymic epithelial cells - i.e. EpCAM1. Wnt4 deprivation opens up an opportunity for trans-differentiation into pre-adipocytes. The simultaneous increase in $LAP2\alpha$ expression provides the necessary signal that pushes dedifferentiated thymic epithelial cells to differentiate into preadipocytes, as detected by increased mRNA levels of PPAR γ and ADRP.

We propose two different mechanisms for the process of adipose involution (see Figure 6). The first allows for the direct initiation of pre-adipocyte differentiation from de-differentiated thymic epithelial cells due to the down-regulation of Wnt4 and up-regulation of LAP2a. Although we cannot rule out this first model, we favor the second model where the process occurs indirectly: dedifferentiation of thymic epithelial cells triggers EMT first, and then the resulting fibroblasts undergo the conventional route of differentiation program towards adipocyte-lineage commitment. The latter model certainly fits better with current literature of EMT [28] and is also supported by our histological and molecular results. Co-localization of a-EpCAM1 and ER-TR7-staining in the aging thymic medulla (Figure 1D) confirms that in the 9 month old thymus there are cells expressing the EpCAM1 marker as a legacy of their primary origin, and also secreting ER-TR7-positive extracellular matrix components, a function conventionally attributed to fibroblast cells. Moreover, Q-PCR data obtained with cDNA samples of MACS-purified thymic epithelial cells also



Figure 6. Model for thymic epithelial senescence. Figure 6 demonstrates our molecular level model of thymic adipose involution. Decreasing Wnt4 and increasing LAP2 levels promote epithelial cells to differentiate into pre-adipocytes either directly or indirectly via EMT. doi:10.1371/journal.pone.0010701.g006

demonstrate an age-related shift in cadherin expression levels characteristic for EMT (Figure 3D) providing additional evidence for the active process of EMT during thymic epithelial senescence.

Our model of thymic epithelial senescence is based on data obtained with mice undergoing physiological senescence. This is the first model for the molecular basis of the thymic epithelium to undergo adipose involution. This model withstands molecular level proof-of-principle using both a model cell line and primary embryonic thymic organ cultures rendered transgenic by lentiviral transgenesis.

Perspectives

Further experiments, however, are required. We plan using inducible, LAP2a-transgenic mice to allow us precise temporalspatial over-expression of LAP2a in adult thymic epithelium to model and decisively verify the role of LAP2 α in pre-adipocyte trans-differentiation in vivo exploiting our experience in establishing transgenic animals [24,29]. If LAP2 α proves to be a master regulator of thymic adipose involution in vivo too, this knowledge appoints LAP 2α as target molecule for directed rejuvenation of the thymic epithelial structure and function. This rejuvenation process could theoretically reinforce naïve T-cell output to reach young adult levels that could ameliorate senescence-related immunological disorders like impaired antiviral defense and late-onset autoimmune diseases.



Level of PPARg expression

Figure 5. Confirmation in transfected thymic lobes. Figures 5A-B present gene expression changes measured by Q-PCR from cDNA of murine thymic lobes transfected at E12 and cultured for 4 days in vitro. Please note that Y-axis scale is linear. Error bars show ±1 SD. doi:10.1371/journal.pone.0010701.g005

Methods

Cell lines and mice

The 293T (ATCC: CRL-11268) and TEP1 [26] cell lines were cultured in DMEM supplemented with 10% FCS, penicillin, streptomycin and β -mercapto-ethanol (Lonza Walkersville). For the experiments we used thymic lobes from timed pregnancies at E12, and also from adult BALB/c mice at 4 week and 1 year of age, and from 1.5 year old GFP-transgenic BALB/c-mice. Mice were bred in our animal facility; all animal work has been conducted according to relevant national and international guidelines following approval of ethics committee of the University of Pecs. Senescent animals developed and aged normally, without any treatment.

Transgenic cell, organ and animal models

The GFP-transgenic BALB/c model was created using lentiviral transgenesis as published by our group [24]. The Wnt4 sequence was purchased and subcloned from an Origene (Origene) vector containing human full-length Wnt4 cDNA. The full-length murine LAP2 α cDNA containing plasmid was a kind gift of Dr. Simon Amos. The GFP (mock), LAP2 α or Wnt4 over-expressing TEP1 cell lines or E12 thymic lobes were generated using lentiviral vectors that were prepared as described previously [30]. Following overnight lentiviral transfection the thymic lobes were transferred over Nucleopore Track-Etch Membranes (Whatman) and were cultured in DMEM supplemented with 20% FCS, penicillin, streptomycin, ciprofloxacin, amphotericin-B and β -mercapto-ethanol (Lonza Walkersville).

Histology using fluorescent antibodies, proteins and dyes

Sections (9 μ m) of frozen thymic lobes of BALB/c mice were fixed in cold acetone, then dried and blocked using 5% BSA in PBS for 20 min before staining with a-Ly51-PE (clone 6C3), a-EpCAM-FITC (clone G8.8), ER-TR7-PE antibodies and DAPI. Thymic sections of GFP-transgenic mice were fixed in 4% paraformaldehyde before staining with LipidTOX Red following the manufacturer's instructions (Invitrogen). The sections were analyzed using an Olympus BX61 microscope equipped with a CCD camera and AnalySIS software.

References

- Marinova TT (2005) Epithelial framework reorganization during human thymus involution. Gerontology 51: 14–18.
- Ribeiro RM, Perelson AS (2007) Determining thymic output quantitatively: using models to interpret experimental T-cell receptor excision circle (TREC) data. Immunol Rev 216: 21–34.
- Grubeck-Loebenstein B (2009) Fading Immune Protection in Old Age: Vaccination in the Elderly. J Comp Pathol.
- Chidgey A, Dudakov J, Seach N, Boyd R (2007) Impact of niche aging on thymic regeneration and immune reconstitution. Semin Immunol 19: 331– 340.
- Gui J, Zhu X, Dohkan J, Cheng L, Barnes PF, et al. (2007) The aged thymus shows normal recruitment of lymphohematopoietic progenitors but has defects in thymic epithelial cells. Int Immunol 19: 1201–1211.
- Hsu HC, Mountz JD (2003) Origin of late-onset autoimmune disease. Immunol Allergy Clin North Am 23: 65–82, vi.
- Bleul C, Boehm T (2005) BMP signaling is required for normal thymus development. J Immunol 175: 5213–5221.
- Pongracz J, Hare K, Harman B, Anderson G, Jenkinson E (2003) Thymic epithelial cells provide Wnt signals. Eur J Immunol 33: 1949–1956.
- Osada M, Ito E, Fermin HA, Vazquez-Cintron E, Venkatesh T, et al. (2006) The Wnt signaling antagonist Kremen1 is required for development of thymic architecture. Clin Dev Immunol 13: 299–319.
- Kuraguchi M, Wang XP, Bronson RT, Rothenberg R, Ohene-Baah NY, et al. (2006) Adenomatous polyposis coli (APC) is required for normal development of skin and thymus. PLoS Genet 2: e146.
- Anderson G, Pongracz J, Parnell S, Jenkinson EJ (2001) Notch ligand-bearing thymic epithelial cells initiate and sustain Notch signaling in thymocytes independently of T cell receptor signaling. Eur J Immunol 31: 3349–3354.

Separation and enrichment of thymic epithelial cells

Thymic lobes were digested with type F collagenase from *C. hystolyticum* (Sigma) for 30 min, then washed with DMEM 10% FCS. Cell suspensions were then labeled with anti-EpCAM1-FITC (clone G8.8) and washed with MACS-buffer followed by incubation with anti-FITC micro-beads (Miltenyi Biotec), the EpCAM⁺-cells were used for total RNA isolation and subsequent quantitative PCR analysis. The cells were purified using MACS LS separation columns (Miltenyi Biotec).

RNA isolation, preparation of cDNA, Q-PCR analysis

Total RNA was isolated the RNAII kit (Macherey-Nagel), including an on column DNA digestion step. cDNA was constructed using the high capacity RNA to cDNA kit (Applied Biosystems). For Q- PCR analysis, we used an AB7500 platform and either SYBR green or TaqMan PCR master mix (Applied Biosystems). Gene expression was normalized to β -actin. The sequences and data of primers are listed in Table 1.

Statistical analysis

All experiments were performed on three occasions, representative experiments are shown. Measures were obtained in triplicates; data are presented as mean ± 1 SD by error bars.

Acknowledgments

The supernatant of ER-TR7 hybridoma clone was originally donated by Dr. Willem van Ewijk to Dr. Peter Balogh, who provided it for the authors. The authors are grateful to Prof. S. Amos (Institute of Hematology, Chaim Sheba Medical Center, Tel-Hashomer, Israel) for providing the murine LAP2 α construct and Prof. E. L. Cooper (Laboratory of Comparative Neuroimmunology, Department of Neurobiology, David Geffen School of Medicine at UCLA, University of California, Los Angeles, USA) for critically and carefully reading the manuscript.

Author Contributions

Conceived and designed the experiments: KK GA EJJ JEP. Performed the experiments: KK ZV DB SH SP. Analyzed the data: KK ZV DB SH SP GA EJJ JEP. Contributed reagents/materials/analysis tools: KK GA EJJ JEP. Wrote the paper: KK JEP.

- Mikels AJ, Nusse R (2006) Whits as ligands: processing, secretion and reception. Oncogene 25: 7461–7468.
- Balciunaite G, Keller M, Balciunaite E, Piali L, Zuklys S, et al. (2002) Wnt glycoproteins regulate the expression of FoxN1, the gene defective in nude mice. Nat Immunol 3: 1102–1108.
- Gordon MD, Nusse R (2006) Wnt signaling: multiple pathways, multiple receptors, and multiple transcription factors. J Biol Chem 281: 22429–22433.
- Schweizer L, Varmus H (2003) Wnt/Wingless signaling through beta-catenin requires the function of both LRP/Arrow and frizzled classes of receptors. BMC Cell Biol 4: 4.
- Torday JS, Rehan VK (2006) Up-regulation of fetal rat lung parathyroid hormone-related protein gene regulatory network down-regulates the Sonic Hedgehog/Wnt/betacatenin gene regulatory network. Pediatr Res 60: 382–388.
- Dorner D, Vlcek S, Foeger N, Gajewski A, Makolm C, et al. (2006) Laminaassociated polypeptide 2alpha regulates cell cycle progression and differentiation via the retinoblastoma-E2F pathway. J Cell Biol 173: 83–93.
- Christodoulides C, Lagathu C, Sethi JK, Vidal-Puig A (2009) Adipogenesis and WNT signalling. Trends Endocrinol Metab 20: 16–24.
- Moldes M, Zuo Y, Morrison RF, Silva D, Park BH, et al. (2003) Peroxisomeproliferator-activated receptor gamma suppresses Wnt/beta-catenin signalling during adipogenesis. Biochem J 376: 607–613.
- Berger R, Theodor L, Shoham J, Gokkel E, Brok-Simoni F, et al. (1996) The characterization and localization of the mouse thymopoletin/lamina-associated polypeptide 2 gene and its alternatively spliced products. Genome Res 6: 361–370.
- Hutchison CJ, Alvarez-Reyes M, Vaughan OA (2001) Lamins in disease: why do ubiquitously expressed nuclear envelope proteins give rise to tissue-specific disease phenotypes? J Cell Sci 114: 9–19.

- Dreger M, Otto H, Neubauer G, Mann M, Hucho F (1999) Identification of phosphorylation sites in native lamina-associated polypeptide 2 beta. Biochemistry 38: 9426–9434.
- Oksanen A (1971) Multilocular fat in thymuses of rats and mice associated with thymus involution: a light- and electron-microscope and histochemical study. J Pathol 105: 223–226.
- Kvell K, Czompoly T, Hiripi L, Balogh P, Kobor J, et al. Characterisation of eGFP-transgenic BALB/c mouse strain established by lentiviral transgenesis. Transgenic Res 19: 105–112.
- Seike M, Mizutani H, Sudoh J, Gemma A (2009) Epithelial to mesenchymal transition of lung cancer cells. J Nippon Med Sch 76: 181.
- Beardsley TR, Pierschbacher M, Wetzel GD, Hays EF (1983) Induction of T-Cell Maturation by a Cloned Line of Thymic Epithelium (TEPI) 10.1073/

pnas.80.19.6005. Proceedings of the National Academy of Sciences 80: $6005{-}6009.$

- Pongracz JE, Parnell SM, Jones T, Anderson G, EJ. J (2006) Overexpression of ICAT highlights a role for catenin-mediated canonical Wnt signalling in early T cell development. Eur J Immunol 36: 2376–2383.
- Friedl P, Gilmour D (2009) Collective cell migration in morphogenesis, regeneration and cancer. Nat Rev Mol Cell Biol 10: 445–457.
- Hiripi L, Negre D, Cosset FL, Kvell K, Czompoly T, et al. Transgenic rabbit production with simian immunodeficiency virus-derived lentiviral vector. Transgenic Res.
- Kvell K, Nguyen TH, Salmon P, Glauser F, Werner-Favre C, et al. (2005) Transduction of CpG DNA-stimulated primary human B cells with bicistronic lentivectors. Mol Ther 12: 892–899.