

Home-based Reach-to-Grasp training for people after stroke is feasible: a pilot randomised controlled trial

Turton, A; Cunningham, Paul; van Wijck, Frederike; Smartt, H; Rogers, CA; Sackley, Catherine; Jowett, Sue; Wolf, Steven L; Wheatley, Keith; van Vliet, P

DOI:

[10.1177/0269215516661751](https://doi.org/10.1177/0269215516661751)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Turton, A, Cunningham, P, van Wijck, F, Smartt, H, Rogers, CA, Sackley, C, Jowett, S, Wolf, SL, Wheatley, K & van Vliet, P 2016, 'Home-based Reach-to-Grasp training for people after stroke is feasible: a pilot randomised controlled trial', *Clinical Rehabilitation*. <https://doi.org/10.1177/0269215516661751>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Home-based Reach-to-Grasp training for people after stroke is feasible: a pilot randomised controlled trial

Supplementary Tables

Table i): RTG treatment received and adherence to self-practice

	Randomised to RtG (n=24)		Randomised to Usual care (n=23)		Total (n=47)		Number of missing values	
	Median	IQR	Median	IQR	Median	IQR	RTG	Usual care
TREATMENT INTENSITY (THERAPIST'S VISITS)								
Completed all 14 RtG intervention visits	16	66.7%			16	66.7%	0	
Number of visits out of 14 completed per participant	14	(13.0, 14.0)	10.5	(5.0, 14.0)	13.5	(10.0, 14.0)	0	5
Total treatment duration (hours)	8.4	(7.7, 9.3)	5.3	(4.2, 8.0)	8.1	(5.5, 9.1)	0	6
Average upper limb treatment per visit (mins) *	38.5	(35.4, 48.8)	38.6	(30.0, 45.0)	38.6	(35.0, 47.7)	0	6
Average visit duration (mins) *	60.7	(59.8, 62.1)	38.6	(30.0, 45.0)	56.1	(42.9, 60.8)	0	6
Total repetitions (all participants)	2198.5	(1249.0, 2943.0)	110	(54.0, 283.0)	1260	(283.0, 2576.0)	0	12
Repetitions per visit	157	(96.1, 210.6)	16.3	(6.3, 24.3)	102.2	(24.3, 197.9)	0	12
ADHERENCE (SELF PRACTICE RtG)								
Number of days practice out of 42	14	(6.0, 26.0)			14	(6.0, 26.0)	3	
Total number of reps	642	(170.0, 2080.0)			642	(170.0, 2080.0)	7	
Number of reps per day*	70	(47.3, 141.2)			70	(47.3, 141.2)	7	
Total time (hours)	6.5	(1.5, 12.8)			6.5	(1.5, 12.8)	3	
Time per day (mins) *	30	(22.2, 44.5)			30	(22.2, 44.5)	3	

Notes:

* Averages generated using only days or visits where relevant data had been recorded (no records available for 3 RTG participants, partial records for the remaining 21)

Table ii): Functional Assessment: Wolf Motor Function Test (WMFT)

		Randomised to RtG (n=24)		Randomised to Usual care (n=23)		Total (n=47)		Missing			
		Median	IQR	Median	IQR	Median	IQR	Not attempted		Failed (too slow)	
								RTG n	Usual care n	RTG n	Usual care n
Number of tasks completed in under 120s	Baseline	6	(3.0, 11.5)	4	(3.0, 10.0)	6	(3.0, 10.0)	0	0		
	7 weeks	8	(5.0, 14.0)	5	(3.0, 11.0)	7	(3.0, 12.0)	1	4		
	3 months	10	(5.0, 13.0)	6.5	(3.0, 11.0)	8	(3.0, 12.0)	1	5		
	6 months	8.5	(4.5, 13.5)	6	(3.0, 14.0)	7	(3.0, 14.0)	8	10		
Median time for tasks 1 to 6	Baseline	2.5	(2.0, 5.0)	2	(2.0, 2.5)	2.5	(2.0, 5.0)	0	0	15	18
	7 weeks	2	(1.3, 2.5)	2	(2.0, 4.0)	2	(1.5, 3.0)	1	2	11	14
	3 months	2	(2.0, 2.5)	2.5	(1.0, 2.5)	2	(2.0, 2.5)	1	4	9	12
	6 months	3	(2.0, 3.0)	2	(1.0, 3.0)	2	(2.0, 3.0)	8	10	7	7
Median time for tasks 8 to 13	Baseline	8.5	(3.8, 16.3)	13	(3.8, 21.3)	9.3	(3.8, 20.0)	0	0	20	19
	7 weeks	5	(3.0, 8.0)	8	(2.0, 16.0)	5.5	(3.0, 8.0)	1	2	16	18
	3 months	6	(3.0, 7.0)	3	(2.8, 4.5)	4	(3.0, 7.0)	1	4	17	15
	6 months	6.5	(4.5, 9.3)	4	(2.5, 11.5)	5	(4.0, 10.5)	8	8	12	10
Median time for tasks 15 to 17	Baseline	17	(9.5, 30.0)	18.5	(12.0, 25.0)	17	(12.0, 25.0)	0	0	20	21
	7 weeks	8	(7.5, 10.0)	13	(12.0, 14.0)	10	(8.0, 12.0)	1	4	19	17
	3 months	8	(6.5, 10.5)	8	(5.0, 11.0)	8	(6.0, 11.0)	1	5	19	16
	6 months	14	(12.0, 16.0)	6	(5.0, 7.0)	7	(6.0, 12.0)	8	8	14	12
Median quality for all tasks	Baseline	1	(1.0, 3.0)	1	(1.0, 3.0)	1	(1.0, 3.0)	0	0		
	7 weeks	3	(1.0, 5.0)	1	(0.0, 3.0)	2	(0.0, 4.0)	3	5		
	3 months	3	(0.0, 5.0)	1	(0.0, 5.0)	2	(0.0, 5.0)	2	5		
	6 months	2	(0.0, 4.5)	0.5	(0.0, 5.0)	1	(0.0, 5.0)	8	9		
Median box weight (lb)	Baseline	1	(0.0, 3.0)	0.5	(0.0, 2.0)	1	(0.0, 2.0)	1	1		
	7 weeks	2	(0.0, 6.0)	0	(0.0, 1.5)	1	(0.0, 4.0)	3	3		
	3 months	2	(0.0, 6.0)	1	(0.0, 4.0)	1	(0.0, 6.0)	1	4		
	6 months	2	(0.0, 6.0)	0.5	(0.0, 6.0)	2	(0.0, 6.0)	9	9		
Median grip strength (kg)	Baseline	0	(0.0, 5.0)	0	(0.0, 3.0)	0	(0.0, 4.0)	1	0		
	7 weeks	2	(0.0, 7.0)	0	(0.0, 2.0)	0.5	(0.0, 5.5)	1	2		
	3 months	3	(0.0, 10.0)	0	(0.0, 2.0)	0	(0.0, 9.0)	2	5		

	Randomised to RtG (n=24)		Randomised to Usual care (n=23)		Total (n=47)		Missing				
	Median	IQR	Median	IQR	Median	IQR	Not attempted		Failed (too slow)		
							RTG	Usual care	RTG	Usual care	
							n	n	n	n	
	6 months	1	(0.0, 8.0)	0	(0.0, 8.0)	0	(0.0, 8.0)	8	8		
Median time for all tasks	Baseline	8	(2.0, 10.0)	7.5	(4.0, 11.0)	8	(4.0, 10.0)	0	0	21	21
	7 weeks	3	(2.5, 4.0)	5	(3.0, 7.0)	3	(3.0, 5.0)	1	4	19	17
	3 months	2.5	(2.0, 3.0)	2.5	(2.0, 3.0)	2.5	(2.0, 3.0)	1	5	19	16
	6 months	4	(4.0, 4.0)	2	(1.0, 3.0)	2.5	(1.5, 3.5)	8	10	15	10

Table iii): Functional Assessment: Stroke Impact Scale (SIS)

	Randomised to RtG (n=24)		Randomised to Usual care (n=23)		Total (n=47)		Number of missing values	
	Median	IQR	Median	IQR	Median	IQR	RTG	Usual care
Physical								
Baseline	41	(31.0, 56.0)	50	(31.0, 63.0)	44	(31.0, 63.0)	0	0
7 weeks	50	(38.0, 63.0)	53	(44.0, 56.0)	50	(44.0, 63.0)	1	1
3 months	50	(38.0, 63.0)	50	(38.0, 56.0)	50	(38.0, 63.0)	2	2
6 months	44	(38.0, 50.0)	38	(31.0, 56.0)	44	(31.0, 56.0)	9	7
Memory and thinking								
Baseline	77	(51.5, 94.5)	86	(54.0, 96.0)	79	(54.0, 96.0)	0	0
7 weeks	86	(71.0, 96.0)	75	(61.0, 100.0)	82	(68.0, 96.0)	1	1
3 months	82	(64.0, 96.0)	89	(79.0, 98.0)	89	(79.0, 96.0)	3	3
6 months	75	(54.0, 93.0)	80.5	(50.0, 100.0)	75	(54.0, 100.0)	9	7
Emotion								
Baseline	62.5	(57.0, 67.0)	58	(47.0, 67.0)	61	(53.0, 67.0)	0	0
7 weeks	61	(61.0, 69.0)	61	(53.0, 69.0)	61	(56.0, 69.0)	2	1
3 months	64	(53.0, 69.0)	61	(56.0, 65.5)	62.5	(56.0, 67.0)	2	3
6 months	67	(61.0, 69.0)	56	(50.0, 62.5)	61	(53.0, 67.0)	11	7
ADL								
Baseline	48	(35.0, 65.0)	50	(40.0, 68.0)	50	(37.5, 68.0)	1	2
7 weeks	51.5	(35.0, 70.0)	60	(45.0, 75.0)	57	(40.0, 72.5)	2	1
3 months	53	(33.0, 73.0)	61.5	(54.0, 72.5)	58.5	(45.0, 73.0)	2	3
6 months	57	(40.0, 75.0)	60	(50.0, 75.0)	57	(50.0, 75.0)	9	8
Mobility								
Baseline	61	(32.0, 78.0)	75	(50.0, 86.0)	69	(42.0, 81.0)	0	0
7 weeks	63.5	(44.0, 89.0)	75	(67.0, 89.0)	72	(58.0, 89.0)	2	2
3 months	81	(44.0, 89.0)	83	(72.0, 89.0)	83	(54.5, 89.0)	3	4
6 months	75	(53.0, 92.0)	78	(67.0, 89.0)	78	(64.0, 89.0)	9	8
Hand function								
Baseline	0	(0.0, 15.0)	0	(0.0, 5.0)	0	(0.0, 10.0)	0	0
7 weeks	5	(0.0, 25.0)	0	(0.0, 20.0)	0	(0.0, 22.5)	1	2
3 months	0	(0.0, 35.0)	0	(0.0, 22.5)	0	(0.0, 30.0)	3	3
6 months	0	(0.0, 30.0)	7.5	(0.0, 20.0)	0	(0.0, 25.0)	11	7

	Randomised to RtG (n=24)		Randomised to Usual care (n=23)		Total (n=47)		Number of missing values	
	Median	IQR	Median	IQR	Median	IQR	RTG	Usual care
Participation								
Baseline	41	(34.0, 53.0)	34	(25.0, 41.0)	38	(28.0, 47.0)	1	0
7 weeks	50	(38.0, 66.0)	47	(36.0, 59.0)	50	(38.0, 59.0)	1	3
3 months	50	(34.0, 66.0)	50	(38.0, 56.0)	50	(36.0, 61.0)	3	4
6 months	47	(34.0, 63.0)	48.5	(31.0, 66.0)	47	(31.0, 66.0)	11	9
Stroke recovery								
Baseline	40	(30.0, 50.0)	50	(40.0, 60.0)	45	(33.0, 60.0)	0	0
7 weeks	50	(40.0, 60.0)	50	(40.0, 60.0)	50	(40.0, 60.0)	1	2
3 months	50	(40.0, 60.0)	50	(40.0, 65.0)	50	(40.0, 60.0)	2	3
6 months	55	(30.0, 60.0)	50	(40.0, 65.0)	50	(35.0, 65.0)	9	8