

## GPCR-styrene maleic acid lipid particles (GPCR-SMALPs)

Wheatley, Mark; Charlton, Jack; Jamshad, Mohammed; Routledge, Sarah J; Bailey, Sian; La-Borde, Penelope J; Azam, Maria T; Logan, Richard T; Bill, Roslyn M; Dafforn, Timothy; Poyner, David R

DOI:

[10.1042/BST20150284](https://doi.org/10.1042/BST20150284)

License:

None: All rights reserved

*Document Version*

Peer reviewed version

*Citation for published version (Harvard):*

Wheatley, M, Charlton, J, Jamshad, M, Routledge, SJ, Bailey, S, La-Borde, PJ, Azam, MT, Logan, RT, Bill, RM, Dafforn, T & Poyner, DR 2016, 'GPCR-styrene maleic acid lipid particles (GPCR-SMALPs): their nature and potential', *Biochemical Society Transactions*, vol. 44, no. 2, pp. 619-623. <https://doi.org/10.1042/BST20150284>

[Link to publication on Research at Birmingham portal](#)

### **Publisher Rights Statement:**

The final version of record is available at DOI: [10.1042/BST20150284](https://doi.org/10.1042/BST20150284)  
Eligibility for repository: Checked on 18/7/2016

### **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

### **Take down policy**

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

# **GPCR-Styrene Maleic Acid Lipid Particles (GPCR-SMALPs): their nature and potential.**

Mark Wheatley\*<sup>1</sup>, Jack Charlton\*, Mohammed Jamshad\*, Sarah J. Routledge<sup>†‡</sup>, Sian Bailey\*, Penelope J. La-Borde\*, Maria T. Azam\*, Richard T. Logan\*, Roslyn M. Bill<sup>†</sup>, Tim R. Dafforn\*, David R. Poyner<sup>†</sup>.

\*School of Biosciences, University of Birmingham, Edgbaston, Birmingham, B15 2TT, UK.

<sup>†</sup>School of Life and Health Sciences, Aston University, Birmingham B4 7ET, UK.

<sup>‡</sup> Current address: Department of Pharmacology, University of Cambridge, Tennis Court Road, Cambridge CB2 1PD

<sup>1</sup>Address for correspondence: Professor Mark Wheatley  
School of Biosciences  
University of Birmingham  
Birmingham  
B15 2TT  
UK

tel: 00-44-121-4143981

e-mail: [m.wheatley@bham.ac.uk](mailto:m.wheatley@bham.ac.uk)

## ABSTRACT

G-protein-coupled receptors (GPCRs) form the largest class of membrane proteins and are an important target for therapeutic drugs. These receptors are highly dynamic proteins sampling a range of conformational states in order to fulfil their complex signalling roles. In order to fully understand GPCR signalling mechanisms it is necessary to extract the receptor protein out of the plasma membrane. Historically this has universally required detergents which inadvertently strip away the annulus of lipid in close association with the receptor and disrupt lateral pressure exerted by the bilayer. Detergent-solubilised GPCRs are very unstable which presents a serious hurdle to characterisation by biophysical methods. A range of strategies have been developed to ameliorate the detrimental effect of removing the receptor from the membrane including amphipols and reconstitution into nanodiscs stabilised by membrane scaffolding proteins but they all require exposure to detergent. Poly(styrene-co-maleic acid) (SMA) incorporates into membranes and spontaneously forms nano-scale SMA lipid particles (SMALPs), effectively acting like a 'molecular pastry cutter' to 'solubilise' GPCRs in the complete absence of detergent at any stage and with preservation of the native annular lipid throughout the process. GPCR-SMALPs have similar pharmacological properties to membrane-bound receptor, exhibit enhanced stability compared to detergent-solubilised receptors and being non-proteinaceous in nature, are fully compatible with downstream biophysical analysis of the encapsulated GPCR.

**Keywords:** SMALP, G-protein-coupled receptor (GPCR), detergent-free, adenosine receptor, membrane protein solubilisation, protein thermostability.

### Abbreviations:

A<sub>2a</sub>R, adenosine A<sub>2a</sub> receptor; A<sub>2a</sub>R-SMALP, SMALP-solubilised A<sub>2a</sub>R; A<sub>2a</sub>R-DDM, DDM-solubilised A<sub>2a</sub>R; β<sub>1</sub>-AR, β<sub>1</sub>-adrenergic receptor; β<sub>2</sub>-AR, β<sub>2</sub>-adrenergic receptor; CD, circular dichroism; DDM, *n*-dodecyl-β-D-maltopyranoside; DMPC, 1,2-dimyristoyl-*sn*-glycero-3-phosphocholine; GPCR, G-protein-coupled receptor; MAPK, mitogen-activated protein kinase; MSP, membrane scaffolding protein; SMA, poly(styrene-co-maleic acid); SMALP, SMA lipid particle; V<sub>1a</sub>R, V<sub>1a</sub> vasopressin receptor.

## **Introduction.**

G-protein-coupled receptors (GPCRs) form the largest class of membrane proteins in the human genome with >800 unique GPCRs. These receptors regulate a plethora of physiological responses which has made them the primary focus for therapeutic intervention. Such is the importance of GPCRs to cell signalling and health that ~50% of all prescribed drugs and ~25% of the top-selling drugs generate their effects by modulating the function of members of this receptor family [1]. The natural activators of GPCRs are very diverse in their physico-chemical characteristics ranging from photons and small biogenic amines to peptides and large glycoproteins. Despite this pronounced diversity in the nature of the agonists, these receptors share a conserved protein architecture comprising a bundle of seven transmembrane (TM) helices linked by extracellular loops (ECLs) and intracellular loops (ICLs) [2]. GPCRs have been subdivided into families on the basis of sequence conservation [3], with three of these families being of particular importance: the rhodopsin/ $\beta$ -adrenergic receptor family (Family A), the secretin receptor family (Family B) and the metabotropic glutamate receptor family (Family C). The largest of these families by far is Family A.

## **GPCRs are highly dynamic in their conformation.**

It was long-thought that GPCRs were simply on/off switches which existed in two conformations, with the 'on' conformation being induced by the binding of an agonist. The activated receptor then activated a specific G-protein to initiate an intracellular signal. In contrast, it was thought that antagonists did not induce this conformational change but merely occupied the binding site thereby preventing activation by agonist. Subsequently, it was proposed that rather than acting as simple on/off switches, GPCRs existed in an equilibrium between the inactive (R) conformation and the active (R\*) conformation and that agonists and inverse agonists stabilised the conformations R\* and R, respectively. It is now recognised that GPCR signalling is extremely complex and that the receptors can exist in a spectrum of conformational states. Individual GPCRs can signal through multiple intracellular cascades by activating more than one class of G-protein and can also initiate G-protein-independent signalling such as  $\beta$ -arrestin-dependent GPCR activation of mitogen-activated protein kinase (MAPK) [4]. Some agonists activate multiple signalling cascades equally whereas other agonists are 'biased' in that they selectively activate one signalling pathway over another. Furthermore, the pharmacological classification of an individual ligand binding to a GPCR can be dictated by the signalling system being assayed. For example, the peptide SPG binds to the V<sub>1a</sub> vasopressin receptor (V<sub>1a</sub>R) which stimulates MAPK but blocks V<sub>1a</sub>R-mediated inositol phosphate signalling [5]. Full agonists and partial agonists possessing different sub-sets of functional groups stabilise different receptor conformations [6, 7] and establish different networks of hydrogen bonds in the binding site [8]. Consequently, instead of possessing just one R\* conformation, activated GPCRs can sample a wide spectrum of distinct active receptor conformations (R\*, R\*', R\*\* etc.), with different efficacies for different signalling systems. The precise receptor conformation stabilised by a particular ligand will dictate that ligand's pharmacological profile. A completely new level of conformational complexity is presented by allosteric ligands, which bind to sites which are discrete to the classical (orthosteric) binding site. Allosteric ligands alter the receptor conformation upon binding thereby tuning the GPCR signalling up or down [9]. From what has been said above, it can be appreciated that the normal functioning of a GPCR requires the receptor protein to populate a wide range of conformational states. Defining these multifarious conformational states and characterising the transition between states is

a pre-requisite to fully understanding GPCR signalling. Such experiments usually require purified GPCR. However, the flexible and highly dynamic nature inherent in the receptor protein presents a serious obstacle to such studies.

### **Studying isolated GPCRs – some problems.**

In common with integral membrane proteins in general, GPCRs have evolved to exist embedded in a lipid bilayer. This requires the protein to stably interact with two discrete physico-chemical environments simultaneously; i) the aqueous phase plus charged lipid headgroups of the membrane and ii) the hydrophobic membrane interior. Both of these environments contribute to the overall functional protein fold of the receptor. When GPCRs are studied *in situ* this is not a problem, but it becomes a problem when experiments require the receptor to be extracted from the complex environment of the plasma membrane. This has universally required the use of a class of surfactants commonly referred to as detergents. Although detergents are very good at molecular dispersal and in gross terms the detergent micelles approximate to the physico-chemical properties of a membrane, micelles are actually poor mimics of the plasma membrane bilayer. The solubilisation process strips away the annular lipid in close association with the protein and it is thought that this removal of lipid from membrane proteins may be the most common cause of solubilisation-induced loss of function [10]. The lipid components of native cell membranes are very heterogeneous with respect to structure, head group and acyl chains and membrane proteins can have a specific requirement for a particular lipid. It is well-established that GPCRs can be affected by the nature of juxtaposed lipid. Cholesterol can modulate receptor conformation [11] and function [12, 13]. Indeed a specific cholesterol binding site incorporating a 'cholesterol consensus motif' has been proposed for some GPCRs following the identification of cholesterol in GPCR crystal structures [14]. Very recently, it has been shown that phospholipids can act as allosteric regulators of GPCRs with phosphatidylglycerol and phosphatidylethanolamine favouring active and inactive conformational states of the  $\beta_2$ -AR respectively [15]. Disruption of the lipid bilayer by detergent not only removes these structural nuances, it also removes the lateral pressure exerted by the bilayer structure on the embedded membrane proteins. This lateral pressure has been shown to be important for maintaining the native fold and activity of membrane proteins [16, 17]. It would be expected that removal of lateral pressure would be particularly disruptive to integral membrane proteins that are conformationally dynamic, such as GPCRs. Consequently, detergent solubilisation of membrane proteins typically results in protein instability and progressive loss of function. Continued presence of detergent is required to prevent aggregation of detergent-solubilised protein despite the detrimental stability implications cited above. These problems are reflected in the relatively very small number of high-resolution crystal structures for membrane proteins compared to soluble proteins.

### **Strategies for studying isolated GPCRs.**

A significant advance was made by the development of a new detergent which was less perturbing to GPCR conformation [18] but receptor stability remained an issue. One approach to increasing the stability of GPCRs in detergent is to alter the receptor sequence to generate a modified protein with significantly greater tolerance to detergent exposure. This has been achieved by two different methodologies: i) a programme of extensive mutagenesis was used to develop a series of specific point mutations that generated a stabilised receptor (referred to as a StaR) in a specific conformation [19] and ii) a directed evolution method was developed using *E.coli* that

allowed the direct selection of GPCRs stable in a pre-selected detergent from libraries containing >100 million variants [20]. As an alternative strategy, the solubilising detergent used initially to extract the receptor has been replaced with amphipathic polymers (amphipols) [21] or by a stabilising engineered  $\beta$ -sheet peptide [22]. Sligar and collaborators have developed protocols for encapsulating detergent-solubilised GPCRs in a lipid disc stabilised by an annulus of membrane scaffolding proteins (MSPs) derived from apolipoproteins, thereby replacing the detergent micelle with a phospholipid bilayer [23].

### **Detergent-free extraction of GPCRs within a nano-scale native lipid bilayer.**

Historically there has been an absolute requirement for detergent to solubilise GPCRs despite the problems this causes. In an ideal situation, the 'solubilisation' would remove the GPCR from the membrane still embedded in a nanoscale section of the native bilayer, thereby preserving the closely-associated annular lipid and lateral pressure. This is now possible using poly(styrene-co-maleic acid) (SMA) (Figure 1) which incorporates into membranes and spontaneously forms nano-scale SMA lipid particles (SMALPs), effectively acting like a 'molecular pastry cutter'. In 2009, SMA polymer with a 2:1 ratio of styrene to maleic acid was shown to solubilise membrane proteins [24]. Biophysical analysis of SMALPs revealed that the SMA polymer forms an annulus surrounding and stabilising a disc of lipid bilayer (~10 nm diameter), possessing the expected thickness of a cell membrane, with the styrene rings of the polymer intercalated between the lipid acyl chains and the maleic acid likely to interact with the lipid headgroups [25]. The behaviour of lipids in a 2:1 SMA polymer SMALP was bilayer-like, as the transition temperature for gel to liquid phase transition of 1,2-dimyristoyl-*sn*-glycero-3-phosphocholine (DMPC) was only 1 °C lower in the SMALP. A 3:1 SMA polymer has also been used to solubilise proteins including the *Halobacteria* proton pump bacteriorhodopsin [26]. However, the DMPC transition temperature in a 3:1 SMA polymer SMALP was considerably reduced (~10 °C) indicating a perturbation of the lipid bilayer. This could be important as membrane fluidity could impact on the conformational flexibility of dynamic proteins like GPCRs. We used 2:1 SMA polymer to solubilise the adenosine A<sub>2a</sub> receptor (A<sub>2a</sub>R) and purify it to homogeneity [27]. A<sub>2a</sub>R-expressing membranes of the yeast *Pichia pastoris*, at a final concentration of 40 mg/ml (wet weight) were incubated with SMA (2.5 % w/v final concentration) for 2 h at 25 °C with gentle stirring. Non-solubilized material was sedimented at 100,000 x g for 1 h at 4 °C, to yield a supernatant containing A<sub>2a</sub>R-SMALP [27]. This was the first time that a GPCR had been solubilised and purified in the total absence of detergent at any stage. The efficiency of extraction of active A<sub>2a</sub>R expressed in HEK 293T cells using our SMALP approach was directly compared to that of n-dodecyl- $\beta$ -D-maltoside (DDM), a detergent commonly employed to solubilise GPCRs. The total binding of the A<sub>2a</sub>R-SMALP was  $2.0 \pm 0.24$  pmol/mg of protein (n=3) equivalent to a yield of  $23.3 \pm 2.7$  % compared to the original HEK 293T cell membrane preparation. This was similar to the recovery seen from detergent (DDM) solubilisation of A<sub>2a</sub>R ( $27.6 \pm 11.4$  %, n=3). However, it was noted that the recovery with A<sub>2a</sub>R-SMALP was less variable than the A<sub>2a</sub>R-DDM [27]. It has been our experience that the extraction of GPCR-SMALPs is equally effective from a range of commonly employed GPCR expression systems, including yeast (*P. pastoris*, mammalian cells (HEK293T and COS-7 cells) and insect cells (*Sf9* cells). Given that the A<sub>2a</sub>R encapsulated in the SMALP had been embedded in native membrane (Figure 2) throughout the solubilisation process, we hypothesised that it would display increased stability compared to detergent-solubilised A<sub>2a</sub>R. We made a direct comparison between the

thermostability of A<sub>2a</sub>R solubilised by DDM and A<sub>2a</sub>R-SMALP, using radioligand binding assays with [<sup>3</sup>H]ZM241385 to monitor preservation of receptor conformational integrity. We established that the A<sub>2a</sub>R-SMALP displayed increased thermostability over A<sub>2a</sub>R-DDM to a range of challenges including storage at 37 °C and storage at 4 °C. Repeated freeze-thaw cycles are particularly damaging to detergent-solubilised GPCRs. A<sub>2a</sub>R-DDM lost all binding capability after just one freeze-thaw whereas the A<sub>2a</sub>R-SMALP binding capability was undiminished after 5 freeze-thaw cycles [27]. Such is the stability of the A<sub>2a</sub>R-SMALP that it could be freeze-dried onto a tube and when re-hydrated with buffer retained ~70% of its binding capability. The improved stability of the GPCR was independent of the source of receptor as this phenomenon was observed with A<sub>2a</sub>R-SMALP generated from either the yeast *P. pastoris* or mammalian HEK 293T cells. The lipid composition of yeast and mammalian cells differ particularly with respect to cholesterol which is replaced by ergosterol in yeast [28] but the thermostability endowed by the SMALP on A<sub>2a</sub>R was comparable between the two expression systems.

### **GPCR-SMALP facilitates biophysical analysis.**

A great strength of the GPCR-SMALP is that the co-polymer stabilising the nanoparticle is non-proteinaceous in nature. Consequently it does not interfere with analysis of the encapsulated GPCR protein using biophysical methods such as CD [27]. The GPCR-SMALP would also facilitate the use of spectroscopic analysis of GPCR conformational changes using endogenous tryptophan fluorescence or introduced fluorescent moieties such as IAEDANS or bimane. The SMALP particles may however be too large for NMR currently. In contrast, a range of biophysical methods cannot be easily applied to GPCRs embedded in MSP-nanodiscs, as the discs effectively have an inherent contaminant in the form of the stabilising scaffolding protein which can interfere with studies on the reconstituted GPCR protein.

### **Conclusion and future perspectives.**

A range of self-assembly systems has been developed which can be applied to studying and purifying GPCRs, including bicelles, amphipols, MSP-stabilised nanodiscs and SMALPs [29] but only the approach exploiting SMALPs is totally detergent-free, retains lateral pressure and preserves the native annular lipid environment of the receptor throughout. It is anticipated that GPCR-SMALPs will facilitate our understanding of fundamental GPCR molecular mechanisms using biophysical techniques. GPCR-SMALPs may also have general utility in a range of receptor-based assays linked to drug discovery.

### **Funding.**

We are grateful for financial support from the Biotechnology and Biological Sciences Research Council [joint grants BB/I020349/1 and BB/I019960/1 (to MW, TRD and DRP, RMB respectively)]

### **References.**

- 1) Lagerström, M.C., and Schiöth, H.B. (2008) Structural diversity of G-protein-coupled receptors and significance for drug discovery. *Nat. Rev. Drug Discov.* **7**, 339-357

- 2) Wheatley, M., Wootten, D., Conner, M.T., Simms, J., Kendrick, R., Logan, R.T., Poyner, D.R. and Barwell, J. (2012) Lifting the lid on G-protein-coupled receptors: The role of extracellular loops. *Brit. J. Pharmacol.* **165**, 1688-1703
- 3) Schiöth, H.B. and Fredriksson, R. (2005) The GRAFS classification system of G-protein-coupled receptors in comparative perspective. *Gen. Comp. Endocrinol.* **142**, 94-101
- 4) Azzi, M., Charest, P.G., Angers, S., Rousseau, G., Kohout, T., Bouvier, M., Piñeyro, G. (2003) Beta-arrestin-mediated activation of MAPK by inverse agonists reveals distinct active conformations for G protein-coupled receptors. *Proc. Natl. Acad. Sci. USA* **100**, 11406-11411
- 5) MacKinnon, A.C., Tufail-Hanif, U., Wheatley, M., Rossi, A.G., Haslett, C., Seckl, M. and Sethi, T. (2009). Targeting V1a vasopressin receptors with [Arg6, D-Trp7,9, NmePhe8]Substance P (6-11) identifies a strategy to develop novel anti-cancer therapies. *Br. J. Pharmacol.* **156**, 36–47
- 6) Swaminath, G., Xiang, Y., Lee, T.W., Steenhuis, J., Parnot, C. and Kobilka, B.K. (2004) Sequential binding of agonists to the beta2 adrenoceptor. Kinetic evidence for intermediate conformational states. *J. Biol. Chem.* **279**, 686-691
- 7) Swaminath, G., Deupi, X., Lee, T.W., Zhu, W., Thian, F.S., Kobilka, T.S. and Kobilka, B. (2005) Probing the beta2 adrenoceptor binding site with catechol reveals differences in binding and activation by agonists and partial agonists. *J. Biol. Chem.* **280**, 22165-22171
- 8) Warne, T., Moukhametzianov, R., Baker, J.G., Nehme, R., Edwards, P.C., Leslie, A.G.W., Schertler, G.F. and Tate, C.G. (2011). The structural basis for agonist and partial agonist action on a b1-adrenergic receptor. *Nature* **469**, 241–245
- 9) Wootten, D., Christopoulos, A. and Sexton, P. (2013) Emerging paradigms in GPCR allostery: Implications for drug discovery. *Nat. Rev. Drug Discov.* **12**, 630-644
- 10) Popot, J.-L. (2010) Amphipols, nanodiscs and fluorinated surfactants: Three nonconventional approaches to studying membrane proteins in aqueous solutions. *Annu. Rev. Biochem.* **79**, 737–775
- 11) Muth, S., Fries, A., and Gimpl, G. (2011) Cholesterol-induced conformational changes in the oxytocin receptor. *Biochem. J.* **437**, 541-553
- 12) Gimpl, G., Burger, K., and Fahrenholz, F. (1997) Cholesterol as modulator of receptor function. *Biochemistry* **36**, 10959-10974
- 13) Pang, L., Graziano, M. and Wang, S. (1999) Membrane cholesterol modulates galanin-GalR2 interaction. *Biochemistry* **38**, 12003-12011
- 14) Hanson, M. A., Cherezov, V., Griffith, M. T., Roth, C. B., Jaakola, V. P., Chien, E. Y., Velasquez, J., Kuhn, P., and Stevens, R. C. (2008) A specific cholesterol binding site is established by the 2.8Å structure of the human  $\beta$ 2-adrenergic receptor. *Structure* **16**, 897-905
- 15) Dawaliby R, Trubbia C, Delporte C, Masureel M, Van Antwerpen P, Kobilka BK, Govaerts C. (2015) Allosteric regulation of G protein-coupled receptor activity by phospholipids. *Nat. Chem. Biol.* [doi: 10.1038/nchembio.1960. Epub ahead of print 16 Nov].
- 16) Charalambous, K., Miller, D., Curnow, P. and Booth, P. J. (2008) Lipid bilayer composition influences small multidrug transporters. *BMC Biochem.* **9**, 31.
- 17) Miller, D., Charalambous, K., Rotem, D., Schuldiner, S., Curnow, P. and Booth, P. J. (2009) *In vitro* unfolding and refolding of the small multidrug transporter EmrE. *J. Mol. Biol.* **393**, 815–832



- 18) Chae, P.S., Rasmussen, S.G., Rana, R.R., Gotfryd, K., Chandra, R., Goren, M.A., Kruse, A.C., Nurva, S., Loland, C.J., Pierre, Y., Drew, D., Popot, J.L., Picot, D., Fox, B.G., Guan, L., Gether, U., Byrne, B., Kobilka, B. and Gellman, S.H. (2010) Maltose-neopentyl glycol (MNG) amphiphiles for solubilization, stabilization and crystallization of membrane proteins. *Nat. Methods* **7**, 1003-1008
- 19) Serrano-Vega, M. J., Magnani, F., Shibata, Y. and Tate, C. G. (2008) Conformational thermostabilization of the beta1-adrenergic receptor in a detergent-resistant form. *Proc. Natl Acad. Sci. USA* **105**, 877-882
- 20) Scott, D.J. and Plückthun, A. (2013) Direct molecular evolution of detergent-stable G-protein-coupled receptors using polymer encapsulated cells. *J. Mol. Biol.* **425**, 662-677
- 21) Banères, J.L., Popot, J.L. and Mouillac, B. (2011) New advances in production and functional folding of G-protein-coupled receptors. *Trends Biotechnol.* **29**, 314-322
- 22) Tao, H., Lee, S.C., Moeller, A., Roy, R.S., Siu, F.Y., Zimmermann, J., Stevens, R.C., Potter, C.S., Carragher, B. and Zhang, Q. (2013) Engineered nanostructured  $\beta$ -sheet peptides protect membrane proteins. *Nat. Methods* **10**, 759-761
- 23) Bayburt, T.H. and Sligar, S.G. (2010) Membrane protein assembly into Nanodiscs. *FEBS Lett.* **584**, 1721-1727
- 24) Knowles, T.J., Finka, R., Smith, C., Lin, Y.-P., Dafforn, T., and Overduin, M. (2009) Membrane proteins solubilized intact in lipid containing nanoparticles bounded by styrene maleic acid copolymer. *J. Am. Chem. Soc.* **131**, 7484-7485
- 25) Jamshad, M., Grimard, V., Idini, I., Knowles, T.J., Dowle, M.R., Schofield, M., Sridhar, P., Lin, Y.-P., Finka, R., Wheatley, M., Thomas, O.R.T., Palmer, R.E., Overduin, M., Govaerts, C., Ruyschaert, J.-M., Edler, K.J. and Dafforn, T. (2014) Structural analysis of a nanoparticle containing a lipid bilayer used for detergent-free extraction of membrane proteins. *Nano Res.* **8**, 774-789
- 26) Orwick-Rydmark, M., Lovett, J.E., Graziadei, A., Linholm, L., Hicks, M.R. and Watts, A. (2012) Detergent-free incorporation of a seven-transmembrane receptor protein into nanosized bilayer lipodisc particles for functional and biophysical studies. *Nano Lett.* **12**, 4687-4692
- 27) Jamshad, M., Charlton, J., Lin, Y.-P., Routledge, S.J., Bawa, Z., Knowles, T.J., Overduin, M., Dekker, N., Dafforn, T.R., Bill, R.M., Poyner, D.R. and Wheatley, M. (2015) *Biosci. Rep.* **35**, e00188
- 28) Finean, J.B., Coleman, R., and Michell, R.H. (1978) *Membranes and their cellular functions*, 2nd ed., Blackwell Scientific Publications, Oxford, UK.
- 29) Jamshad, M., Lin, Y.-P., Knowles, T.J., Parslow, R.M., Harris, C., Wheatley, M., Poyner, D.R., Bill, R.M., Thomas, O.R.T., Overduin, M. and Dafforn, T.R. (2011) *Biochem. Soc. Trans.* **39**, 813-818

## FIGURE LEGENDS

**Figure 1**     **Poly(styrene-co-maleic acid).**

**Figure 2**     **An A<sub>2a</sub>R-SMALP.**

A schematic representation of an A<sub>2a</sub>R-SMALP viewed from within the plane of the membrane. Only the receptor (coloured) and lipid molecules (grey) are shown.




Figure 1




Figure 2.

