

On special cases of the generalized max-plus eigenproblem

Butkovic, Peter; Jones, Daniel

DOI:
[10.1137/15M1041031](https://doi.org/10.1137/15M1041031)

License:
Creative Commons: Attribution (CC BY)

Document Version
Publisher's PDF, also known as Version of record

Citation for published version (Harvard):
Butkovic, P & Jones, D 2016, 'On special cases of the generalized max-plus eigenproblem', *S I A M Journal on Matrix Analysis and Applications*, vol. 37, no. 3, pp. 1002–1021. <https://doi.org/10.1137/15M1041031>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

ON SPECIAL CASES OF THE GENERALIZED MAX-PLUS EIGENPROBLEM*

PETER BUTKOVIČ† AND DANIEL JONES‡

Abstract. We study the generalized eigenproblem $A \otimes x = \lambda \otimes B \otimes x$, where $A, B \in \mathbb{R}^{m \times n}$ in the max-plus algebra. It is known that if A and B are symmetric, then there is at most one generalized eigenvalue, but no description of this unique candidate is known in general. We prove that if $C = A - B$ is symmetric, then the common value of all saddle points of C (if any) is the unique candidate for λ . We also explicitly describe the whole spectrum in the case when B is an outer product. It follows that when A is symmetric and B is constant, the smallest column maximum of A is the unique candidate for λ . Finally, we provide a complete description of the spectrum when $n = 2$.

Key words. matrix, max-plus algebra, generalized eigenproblem, spectrum

AMS subject classifications. 15A18, 15A80, 93C55

DOI. 10.1137/15M1041031

1. Introduction. We start with four motivational examples. They are all variants of a model called the *multiprocessor interactive system*.

Example 1. Products P_1, \dots, P_m are prepared using n processors, with every processor potentially contributing to the completion of each product. It is assumed that every processor can work for all products simultaneously and that all these actions on a processor start as soon as the processor starts to work. Let a_{ij} be the duration of the work of the j th processor needed to complete the partial product for P_i ($i = 1, \dots, m; j = 1, \dots, n$). Let us denote by x_j the starting time of the j th processor ($j = 1, \dots, n$). Then all partial products for P_i ($i = 1, \dots, m$) will be ready at time

$$\max(x_1 + a_{i1}, \dots, x_n + a_{in}).$$

Hence if b_1, \dots, b_m are given completion times of the products that have to be met exactly, then the starting times have to satisfy the system of equations

$$\max(x_1 + a_{i1}, \dots, x_n + a_{in}) = b_i, \quad \forall i = 1, \dots, m.$$

If we denote $a \oplus b = \max(a, b)$ and $a \otimes b = a + b$, and the pair of operations (\oplus, \otimes) is extended to matrices and vectors in the same way as in linear algebra, then this can be written as a compact equation:

$$(1) \quad A \otimes x = b.$$

The matrix $A = (a_{ij})$ is called the *production matrix*.

*Received by the editors September 24, 2015; accepted for publication (in revised form) by W.-W. Lin May 10, 2016; published electronically August 4, 2016. This research was supported by EPSRC grant EP/J00829X/1.

<http://www.siam.org/journals/simax/37-3/M104103.html>

†Corresponding author. School of Mathematics, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK (p.butkovic@bham.ac.uk).

‡School of Mathematics, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK (danjones1729@hotmail.co.uk).

Example 2. Consider the system in which processors P_1, \dots, P_n work interactively and in stages. In each stage all processors simultaneously produce components necessary for the work of some or all of the other processors in the next stage. Let $x_i(k)$ denote the starting time of the k th stage on P_i ($i = 1, \dots, n$), and let a_{ij} denote the duration of the operation at which processor P_j prepares the component necessary for processor P_i in the $(k+1)$ st stage ($i, j = 1, \dots, n$). Then, avoiding any delay, we have

$$(2) \quad x_i(k+1) = \max(x_1(k) + a_{i1}, \dots, x_n(k) + a_{in}) \quad (i = 1, \dots, n; k = 0, 1, \dots).$$

In the max-algebraic notation, (2) gets the form

$$x_i(k+1) = a_{i1} \otimes x_1(k) \oplus \dots \oplus a_{in} \otimes x_n(k) \quad (i = 1, \dots, n; k = 0, 1, \dots).$$

or, in the matrix-vector form,

$$x(k+1) = A \otimes x(k) \quad (k = 0, 1, \dots).$$

We say that the system reaches a *steady regime* [15], [16] if it eventually moves forward in regular steps; that is, for some λ and k_0 we have $x(k+1) = \lambda \otimes x(k)$ for all $k \geq k_0$. Equivalently, the time between the starts of consecutive stages will eventually stabilize and be the same constant for every processor. If this happens, then we have

$$A \otimes x(k) = \lambda \otimes x(k) \quad \forall k \geq k_0,$$

and so $x(k)$ is a max-plus eigenvector of A with associated eigenvalue λ .

Example 3. Now suppose that in addition to the assumption of Example 1, k other machines independently prepare partial products for products Q_1, \dots, Q_m , and the duration and starting times are b_{ij} and y_j , respectively. Then the *synchronization problem* is to find starting times of all $n+k$ machines so that each pair (P_i, Q_i) ($i = 1, \dots, m$) is completed at the same time. This task is equivalent to solving the system of equations

$$(3) \quad \max(x_1 + a_{i1}, \dots, x_n + a_{in}) = \max(y_1 + b_{i1}, \dots, y_k + b_{ik}) \quad (i = 1, \dots, m).$$

Again, using max-algebra and denoting $K = \{1, \dots, k\}$ we can write this system as a system of max-linear equations:

$$(4) \quad \sum_{j \in N}^{\oplus} a_{ij} \otimes x_j = \sum_{j \in K}^{\oplus} b_{ij} \otimes y_j \quad (i = 1, \dots, m).$$

In the matrix-vector notation it has the form

$$A \otimes x = B \otimes y$$

or, specifically,

$$A \otimes x = B \otimes x.$$

Example 4. A variant of (3) is the task when $n = k$ and the starting times are linked; for instance, it is required that they be the same or, more generally, that there be a fixed interval between the starting times of the first and second systems; that is, the starting times x_j, y_j of each pair of machines differ by the same value. If we denote this (unknown) value by λ , then $y = \lambda \otimes x$ and the equations read

$$(5) \quad \max(x_1 + a_{i1}, \dots, x_n + a_{in}) = \max(\lambda + x_1 + b_{i1}, \dots, \lambda + x_n + b_{in})$$

for $i = 1, \dots, m$. In max-algebraic notation, this system gets the form of a “generalized eigenproblem”:

$$A \otimes x = \lambda \otimes B \otimes x.$$

The examples above give rise to the following systems:

$$(6) \quad A \otimes x = b,$$

$$(7) \quad A \otimes x = \lambda \otimes x,$$

$$(8) \quad A \otimes x = B \otimes x,$$

and

$$(9) \quad A \otimes x = \lambda \otimes B \otimes x.$$

Systems (6) are historically the first problem studied in max-algebra [14], and their solution set can easily be described [12] both algebraically (using residuation) and combinatorially (in terms of set coverings; see below). For the origins of max-algebra see also [26], [28], and [22].

Systems (7) have also been intensively studied since the 1960s [15] (see also [22], [19]). It is known [19], [13], [12] that every $n \times n$ matrix over $\mathbb{R} \cup \{-\infty\}$ has up to n eigenvalues, with the maximum cycle mean of A always being a (biggest) eigenvalue. All eigenvalues and bases of all eigenspaces can be found in $O(n^3)$ time.

Systems (8) have been studied since 1978 [7], [8], [9], [10], [11]. It has been proved that the solution set is finitely generated [10]. These systems have been shown to be equivalent to mean payoff games [2]. A number of solution methods exist [17], [20], [3], [27]. Although none of them is polynomial, this problem is known to be in $NP \cap co-NP$ [5], and it is therefore expected that eventually a polynomial solution method will be found.

Systems (9) for fixed λ reduce to (8), so the effort is usually concentrated on finding the spectrum. In contrast to (7), no polynomial method seems to be known in general for finding even a single eigenvalue or deciding whether such a value exists. The task of finding the spectrum is made more complicated by the fact that on one hand, there may be no eigenvalue at all, and on the other hand, there is no upper limit for the number of eigenvalues—there may even be a continuum of them, and any union of closed intervals is the spectrum for some generalized eigenproblem [25]. Nevertheless, there is a pseudopolynomial method for finding the whole spectrum based on Newton-type iterations [21]. This problem was studied for the first time independently in [6] and [18] and then by many other authors. A fast method for narrowing the search for a generalized eigenvalue is presented in [12, section 9.3].

Paper [6] presents a number of conditions that are either necessary or sufficient for the existence of either a solution or a unique solution to (9). In particular, it follows that there is at most one generalized eigenvalue if A and B are finite and symmetric. The paper also includes specific conditions and a graphical method for 2×2 matrices.

Note that systems (6)–(9) may be considered over $\mathbb{R} \cup \{-\infty\}$. The aim of the present paper is to study the spectrum of (9) for finite matrices (that is, matrices over \mathbb{R}), more precisely, we do the following:

(a) We prove that if $C = A - B$ is symmetric, then the common value of all saddle points of C (if any) is the unique candidate for λ (section 3).

(b) We explicitly describe the whole spectrum provided that B is an outer product (section 4).

(c) We show that it follows that when A is symmetric and B is constant, the smallest column maximum of A is the unique candidate for λ (section 4).

(d) Finally, we provide an alternative, algebraic description of the whole spectrum and eigenspaces when x is two-dimensional (section 5).

2. Prerequisites. In this section we give the definitions and some basic results which will be used in the formulations and proofs of our results. For the proofs and more information about max-algebra, the reader is referred to [1], [4], [12], and [23].

We assume everywhere that $m, n \geq 1$ are natural numbers and denote $M = \{1, \dots, m\}$ and $N = \{1, \dots, n\}$; the symbol \mathbb{R} stands for $\mathbb{R} \cup \{-\infty\}$. We use the conventions $\max \emptyset = -\infty$ and $\min \emptyset = +\infty$.

It will be useful to first recall a simple property of matrices, well known in the theory of matrix games. Let $A = (a_{ij}) \in \mathbb{R}^{m \times n}$, and denote $v_1(A) = \max_{i \in M} \min_{j \in N} a_{ij}$ (usually called “gain-floor”) and $v_2(A) = \min_{j \in N} \max_{i \in M} a_{ij}$ (“loss-ceiling”). A pair $(r, s) \in M \times N$ is called a *saddle point* in A if

$$\max_{i \in M} a_{is} = a_{rs} = \min_{j \in N} a_{rj}.$$

THEOREM 5 (see [24]). *The inequality $v_1(A) \leq v_2(A)$ holds for every $A \in \mathbb{R}^{m \times n}$. The equality holds if and only if there is a saddle point in A . If (r, s) is any saddle point, then $v_1(A) = a_{rs} = v_2(A)$ (and this value is called the value of the game A).*

We will use the following notation:

$$M_j(A) = \left\{ r \in M; a_{rj} = \max_{i \in M} a_{ij} \right\}, \quad j \in N,$$

$$N_i(A) = \{ j \in N; i \in M_j(A) \}, \quad i \in M.$$

We will also write M_j, N_i instead of $M_j(A), N_i(A)$ if no confusion can arise. The following will be useful.

PROPOSITION 6. $\bigcup_{j \in N} M_j = M$ if and only if $N_i \neq \emptyset$ for every $i \in M$.

Proof. The proof is straightforward from the definitions. \square

If $a, b \in \mathbb{R}$, then we set

$$a \oplus b = \max(a, b)$$

and

$$a \otimes b = a + b.$$

Throughout the paper we denote $-\infty$ by ε (the neutral element with respect to \oplus), and for convenience we also denote by the same symbol any vector with all components $-\infty$, and a matrix with all entries $-\infty$. If $a \in \mathbb{R}$, then the symbol a^{-1} stands for $-a$. The symbol a^k ($k \geq 1$ integer) stands for the iterated product $a \otimes a \otimes \dots$ in which the symbol a appears k times (that is, ka in conventional notation). By *max-algebra* (recently also called “tropical linear algebra”) we understand the analogue of linear algebra developed for the pair of operations (\oplus, \otimes) , extended to matrices and vectors as in conventional linear algebra. That is, if $A = (a_{ij})$, $B = (b_{ij})$, and $C = (c_{ij})$ are matrices of compatible sizes with entries from \mathbb{R} , we write $C = A \oplus B$ if $c_{ij} = a_{ij} \oplus b_{ij}$ for all $i, j \in N$ and write $C = A \otimes B$ if

$$c_{ij} = \bigoplus_{k \in N} a_{ik} \otimes b_{kj} = \max_k (a_{ik} + b_{kj})$$

for all i and $j \in N$. If $\alpha \in \overline{\mathbb{R}}$, then $\alpha \otimes A = (\alpha \otimes a_{ij})$. Although the use of the symbols \otimes and \oplus is common in max-algebra, we will apply the usual convention of not writing the symbol \otimes . Thus, in what follows the symbol \otimes will not be used and, unless explicitly stated otherwise, all multiplications indicated are in max-algebra.

A vector or matrix is called *finite* if all its entries are real numbers. A square matrix is called *diagonal* if all its diagonal entries are real numbers and off-diagonal entries are ε . More precisely, if $x = (x_1, \dots, x_n)^T \in \mathbb{R}^n$, then $\text{diag}(x_1, \dots, x_n)$ or just $\text{diag}(x)$ is the $n \times n$ diagonal matrix

$$\begin{pmatrix} x_1 & \varepsilon & \dots & \varepsilon \\ \varepsilon & x_2 & \dots & \varepsilon \\ \vdots & \vdots & \ddots & \vdots \\ \varepsilon & \varepsilon & \dots & x_n \end{pmatrix}.$$

The matrix $\text{diag}(0)$ is called the *unit matrix* and denoted by I . Obviously, $AI = IA = A$ whenever A and I are of compatible sizes. A matrix obtained from a diagonal matrix (unit matrix) by permuting the rows and/or columns is called a *generalized permutation matrix* (*permutation matrix*). It is known that in max-algebra, generalized permutation matrices are the only type of invertible matrices [16], [12]. Clearly,

$$(\text{diag}(x_1, \dots, x_n))^{-1} = \text{diag}(x_1^{-1}, \dots, x_n^{-1}).$$

The following theorem is probably the historically first result in max-algebra; here we denote the following for $A \in \overline{\mathbb{R}}^{m \times n}$ and $b \in \overline{\mathbb{R}}^n$:

$$S(A, b) = \{x \in \overline{\mathbb{R}}^n; Ax = b\}.$$

THEOREM 7 (see [14], [12]). *If $A \in \overline{\mathbb{R}}^{m \times n}$ is a matrix with no ε columns, $b \in \mathbb{R}^m$, and $B = (\text{diag}(b))^{-1}A$, then $S(A, b) \neq \emptyset$ if and only if $\bigcup_{j \in N} M_j(B) = M$.*

COROLLARY 8. *If $A \in \overline{\mathbb{R}}^{m \times n}$ is a matrix with no ε columns, $b \in \mathbb{R}^m$, and $B = (\text{diag}(b))^{-1}A$, then $S(A, b) \neq \emptyset$ if and only if $N_i(B) \neq \emptyset$ for every $i \in M$.*

PROPOSITION 9. *If $b \in \mathbb{R}^m$ is a constant vector, then $M_j(A) = M_j(B)$ for all $j \in N$ and, consequently, also $N_i(A) = N_i(B)$ for all $i \in M$.*

Proof. The proof is straightforward from the definitions. \square

Suppose that $A = (a_{ij}) \in \overline{\mathbb{R}}^{m \times n}$ and $B = (b_{ij}) \in \overline{\mathbb{R}}^{m \times n}$ are given. The *generalized eigenproblem* for (A, B) is the task of finding $x \in \overline{\mathbb{R}}^n, x \neq \varepsilon$ (*generalized eigenvector* or just *eigenvector*) and $\lambda \in \mathbb{R}$ (*generalized eigenvalue* or just *eigenvalue*) such that

$$(10) \quad Ax = \lambda Bx.$$

Note that the case $\lambda = \varepsilon$ is trivial and is not discussed here. We denote

$$V(A, B, \lambda) = \{x \in \overline{\mathbb{R}}^n; Ax = \lambda Bx, x \neq \varepsilon\},$$

$$\Lambda(A, B) = \{\lambda \in \mathbb{R}; V(A, B, \lambda) \neq \emptyset\}.$$

The set $\Lambda(A, B)$ will be called the *spectrum* of the pair (A, B) . In the rest of the paper we will assume that $A, B \in \mathbb{R}^{m \times n}$. It is easy to see that then a generalized eigenvector exists if and only if a finite generalized eigenvector exists. Note that some

statements remain valid if the finiteness requirement is removed or replaced by the condition that there are no ε columns.

The next two statements have previously been proved and provide useful information about the spectrum. Here and in the rest of the paper we denote

$$C = A - B = (c_{ij}),$$

$$\underline{\lambda}(C) = \max_{i \in M} \min_{j \in N} c_{ij},$$

and

$$\bar{\lambda}(C) = \min_{i \in M} \max_{j \in N} c_{ij}.$$

We will use shorthand $\underline{\lambda}$ or $\bar{\lambda}$ if no confusion can arise.

PROPOSITION 10 (see [12]). $\Lambda(A, B) \subseteq [\underline{\lambda}, \bar{\lambda}]$ holds for any $A, B \in \mathbb{R}^{m \times n}$.

The interval $[\underline{\lambda}, \bar{\lambda}]$ will be called the *feasibility interval* for the generalized eigenproblem.

The following statement follows from the results in [6].

PROPOSITION 11. If $A, B \in \mathbb{R}^{n \times n}$ are symmetric, then $|\Lambda(A, B)| \leq 1$.

This paragraph presents a sketch of a numerical method of [21] for solving the generalized eigenproblem for arbitrary matrices A and B . However, it will not be used in this paper and may be skipped. The method is based on the following idea. Let us define (see [16]) min-algebra over $\mathbb{R} \cup \{+\infty\}$ by

$$a \oplus' b = \min(a, b)$$

and

$$a \otimes' b = a \otimes b$$

for all a and b except when one of a and b is $+\infty$ and the other $-\infty$. In this case,

$$+\infty \otimes -\infty = -\infty \otimes +\infty = -\infty$$

but

$$+\infty \otimes' -\infty = -\infty \otimes' +\infty = +\infty.$$

We also define $A^\# = -A^T$. As in max-algebra, we extend the pair of operations (\oplus', \otimes') to matrices and vectors. We will not write the operator \otimes' , and for matrices the convention applies that the product is in min-algebra whenever it follows the symbol $\#$; otherwise it is in max-algebra. In this way a residuated pair of operations (a special case of Galois connection) has been defined, namely,

$$Ax \leq y \iff x \leq A^\#y,$$

for all x, y . Due to isotonicity, $Ax \leq y$ implies $A(A^\#y) \leq y$, and it follows immediately that a one-sided system $A \otimes x = b$ has a solution if and only if $A(A^\#b) = b$. It will be convenient to denote $P_A(z) = A(A^\#z)$ for any A and z . Thus, finding a solution to a two-sided system with separated variables $A \otimes x = B \otimes y$ then means finding a z such that $P_AP_B(z) = z = P_BP_A(z)$ or, equivalently, $P_AP_B(z) \oplus' P_BP_A(z) = z$. The system $A \otimes x = \lambda \otimes B \otimes x$ is equivalent to

$$C(\lambda)x = Dy,$$

where

$$C(\lambda) = \begin{pmatrix} A \\ \lambda + B \end{pmatrix}, \quad D = \begin{pmatrix} I \\ I \end{pmatrix},$$

and I is a unit matrix of an appropriate size. The function

$$h_\lambda(x) = (\lambda \oplus A^\# Bx) \oplus' (-\lambda + B^\# Ax)$$

is an order-preserving, additively homogeneous and continuous map. As such, it has a largest “eigenvalue”

$$r(h_\lambda) = \max \{ \rho; h_\lambda(x) = \rho + x \}.$$

It is proved that $\Lambda(A, B)$ coincides with

$$\{ \lambda; r(h_\lambda) = 0 \}.$$

It is also proved that the function $s(\lambda) = r(h_\lambda)$ is piecewise-linear and Lipschitz continuous, and therefore in the case of integer matrices A and B its zero-level set can be found by a pseudopolynomial number of calls to an oracle that computes the value of a mean payoff game.

3. Generalized eigenproblem for symmetric matrices with saddle point.

Note that in general even if the premise of Proposition 11 is satisfied, it is not clear what the unique candidate for the generalized eigenvalue is, and even if such a candidate is known, it is not clear how to check in polynomial time whether it is such an eigenvalue.

Let $C \in \mathbb{R}^{m \times n}$ be any matrix. Clearly, $\underline{\lambda}(C)$ coincides with $v_1(C)$. Although in general $\bar{\lambda}(C)$ is different from $v_2(C)$ we observe that for $C^T = (c_{ji}) = (c'_{ij})$ we have

$$\begin{aligned} \bar{\lambda} &= \min_{i \in M} \max_{j \in N} c_{ij} \\ &= \min_{j \in M} \max_{i \in N} c_{ji} \\ &= \min_{j \in M} \max_{i \in N} c'_{ij} \\ &= v_2(C^T). \end{aligned}$$

Hence $\underline{\lambda}(C) = v_1(C)$ and $\bar{\lambda}(C) = v_2(C)$ if C is symmetric, in particular, when $C = A - B$ and both A and B are symmetric. Thus, if C is symmetric and has a saddle point, say (r, s) , then $\underline{\lambda} = a_{rs} = \bar{\lambda}$ and so either a_{rs} is the unique generalized eigenvalue or there is no such eigenvalue. The next two examples confirm that both cases are possible.

Example 12. If

$$A = \begin{pmatrix} 2 & 1 \\ 1 & 0 \end{pmatrix}, \quad B = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix},$$

then $C = A$ is symmetric and has saddle point $(1, 2)$ of value 1, which is therefore the unique candidate for a generalized eigenvalue. If there was an associated eigenvector $x = (x_1, x_2)^T$, we may assume without loss of generality $x_1 = 0$, and the individual terms in Ax and λBx are as described in the following matrices:

$$\begin{pmatrix} 2 & 1 + x_2 \\ 1 & x_2 \end{pmatrix}, \quad \begin{pmatrix} 1 & 1 + x_2 \\ 1 & 1 + x_2 \end{pmatrix}.$$

The first equation implies $1 + x_2 \geq 2$ and the second $1 + x_2 \leq 1$; thus $\Lambda(A, B) = \emptyset$.

Example 13. If

$$A = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \quad B = \begin{pmatrix} -2 & -1 \\ 0 & 0 \end{pmatrix},$$

then C is the same as in the previous example and so $\lambda = 1$ is the unique candidate for a generalized eigenvalue. Now the zero vector $x = (0, 0)^T$ is an associated eigenvector, and thus $\Lambda(A, B) = \{1\}$.

The following example shows that $\Lambda(A, B)$ may be nonempty even if C is symmetric and has no saddle point.

Example 14. If

$$A = \begin{pmatrix} 1 & 2 \\ 3 & 1 \end{pmatrix}, \quad B = \begin{pmatrix} 1 & 0 \\ 1 & 0 \end{pmatrix},$$

then

$$C = \begin{pmatrix} 0 & 2 \\ 2 & 1 \end{pmatrix}$$

is a symmetric matrix without a saddle point, but $\lambda = 2$ is a generalized eigenvalue with associated eigenvector $x = (0, 1)^T$.

We summarize as follows.

THEOREM 15. *If $A, B \in \mathbb{R}^{n \times n}$, $C = A - B$ is a symmetric matrix with a saddle point (r, s) , and $\Lambda(A, B) \neq \emptyset$, then c_{rs} is the unique generalized eigenvalue for (A, B) .*

The authors are not aware of any polynomial method for checking that the saddle point value of $A - B$ is a generalized eigenvalue of (A, B) .

4. Generalized eigenproblem when B is an outer product. Let us consider the generalized eigenproblem (10), where B is a (max-algebraic) outer product of two vectors, say $v = (v_1, \dots, v_m)^T \in \mathbb{R}^m$ and $w = (w_1, \dots, w_n)^T \in \mathbb{R}^n$. Thus, we can write

$$B = (b_{ij}) = vw^T = (v_i + w_j).$$

Let $V = \text{diag}(v)$ and $W = \text{diag}(w)$. Then (10) reads

$$Ax = \lambda vw^T x$$

and is equivalent to

$$V^{-1}Ax = \lambda V^{-1}Bx$$

or

$$V^{-1}Ax = \lambda \begin{pmatrix} 0 \\ \vdots \\ 0 \end{pmatrix} w^T x.$$

Set $x = W^{-1}y$, where $y = (y_1, \dots, y_n)^T$; then (10) is equivalent to

$$V^{-1}AW^{-1}y = \lambda \begin{pmatrix} 0 \\ \vdots \\ 0 \end{pmatrix} (y_1 \oplus y_2 \oplus \dots \oplus y_n).$$

Here the right-hand side is actually equal to

$$\lambda \begin{pmatrix} 0 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & 0 \end{pmatrix} y,$$

and thus in system (10) with B an outer product it can be assumed without loss of generality that B is a zero matrix (a matrix with all entries equal to 0). In such a system the right-hand side of each equation is $\lambda(x_1 \oplus x_2 \oplus \cdots \oplus x_n)$, and thus an $x \neq \varepsilon$ satisfying $Ax = \lambda 0x$ exists if and only if there is a $z \in \mathbb{R}$ satisfying

$$(11) \quad A'x = \begin{pmatrix} z \\ \vdots \\ z \end{pmatrix},$$

where A' is obtained from A by adding an extra row whose every entry is λ , that is,

$$A' = \begin{pmatrix} A \\ \lambda \ \dots \ \lambda \end{pmatrix} = (a'_{ij}).$$

Clearly, A' is an $(m+1) \times n$ matrix. Theorem 7 and Proposition 9 enable us to solve such systems, and we use them in the next two propositions. It will be useful to denote

$$M' = M \cup \{m+1\},$$

$$M'_j = \left\{ r \in M'; a'_{rj} = \max_{i \in M'} a'_{ij} \right\}, \quad j \in N,$$

and

$$N'_i = \{j \in N; i \in M'_j\}, \quad i \in M'.$$

PROPOSITION 16. *Let $A \in \mathbb{R}^{m \times n}$. Then $\lambda \in \Lambda(A, 0)$ if and only if $\bigcup_{j \in N} M'_j = M'$.*

Proof. The proof follows straightforwardly from the previous discussion, Theorem 7, and Proposition 9. \square

PROPOSITION 17. $\Lambda(A, 0) \subseteq [\lambda_0, \lambda_1]$ holds for every $A = (a_{ij}) \in \mathbb{R}^{m \times n}$, where

$$\lambda_0 = \min_{j \in N} \max_{i \in M} a_{ij} \text{ and}$$

$$\lambda_1 = \max_{j \in N} \max_{i \in M} a_{ij}.$$

Proof. A real number λ is in $\Lambda(A, 0)$ if and only if the system (11) has a solution for some $z \in \mathbb{R}$. This is a one-sided system whose solvability does not depend on z because the right-hand side is a constant vector. The solvability criterion is given in Theorem 7. We apply this condition to A' using Proposition 9, first to each of the first m rows and then to the last row:

$$(12) \quad (\forall i \in M) (\exists j \in N) (\forall r \in M) a_{ij} \geq a_{rj} \oplus \lambda$$

and

$$(13) \quad (\exists j \in N) (\forall i \in M) \lambda \geq a_{ij}.$$

The latter is equivalent to $\lambda \geq \min_{j \in N} \max_{i \in M} a_{ij}$, which proves the lower bound. The first is equivalent to the requirement that for every $i \in M$ there exists a $j \in N$ satisfying

$$a_{ij} \geq \max_{r \in M} a_{rj} \oplus \lambda.$$

Since

$$\max_{r \in M} a_{rj} \oplus \lambda \geq \max_{r \in M} a_{rj} \geq a_{ij},$$

it follows that

$$a_{ij} = \max_{r \in M} a_{rj} \oplus \lambda = \max_{r \in M} a_{rj}$$

and $\lambda \leq \max_{r \in M} a_{rj} \leq \max_{j \in N} \max_{i \in M} a_{ij}$, which proves the upper bound. \square

Proposition 17 does not provide any tool for checking whether $\Lambda(A, 0)$ is nonempty. We give this answer next.

PROPOSITION 18. *The following statements are equivalent for every $A = (a_{ij}) \in \mathbb{R}^{m \times n}$:*

- (a) $\Lambda(A, 0) \neq \emptyset$,
- (b) $\bigcup_{j \in N} M_j = M$,
- (c) $N_i(A) \neq \emptyset$ for every $i \in M$, and
- (d) $\lambda_0 \in \Lambda(A, 0)$.

Proof. The equivalence of (b) and (c) has been shown in Proposition 6. We prove (b) \Rightarrow (d) \Rightarrow (a) \Rightarrow (b). Suppose (b) is true. There is an index $k \in N$ such that

$$\lambda_0 = \max_{i \in M} a_{ik} \leq \max_{i \in M} a_{ij}$$

holds for every $j \in N$. Hence, for $\lambda = \lambda_0$ and for every $j \in N$ we have $M_j \subseteq M'_j$ and $m+1 \in M'_k$. It follows that

$$\bigcup_{j \in N} M'_j \supseteq \bigcup_{j \in N} M_j \cup \{m+1\} = M \cup \{m+1\} = M',$$

and thus

$$\bigcup_{j \in N} M'_j = M',$$

and so (11) has a solution for $\lambda = \lambda_0$, which proves (d) by Proposition 16.

The second implication is trivial, so suppose now that $\lambda \in \Lambda(A, 0)$. Hence, (11) has a solution with this value of λ , and thus $\bigcup_{j \in N} M'_j = M'$. Let $i \in M$; then $i \in M'_j$ for some $j \in N$ and therefore also $i \in M_j$ because for any $j \in N$ the set M'_j either coincides with M_j or is $M_j \cup \{m+1\}$ or is just $\{m+1\}$. Statement (b) now follows. \square

COROLLARY 19. *If $A \in \mathbb{R}^{n \times n}$ is symmetric, then either $\Lambda(A, 0)$ is empty or $\{\lambda_0\}$.*

Proposition 18 shows that the lower bound for the spectrum specified in Proposition 17 is tight for every matrix with nonempty spectrum. In general this is not true about the upper bound. For instance, if

$$A = \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix}, \quad B = 0,$$

then $C = A$ is symmetric, and so $\Lambda(A, B) = \{1\}$ by Proposition 18 and Corollary 19. However, $\lambda_1 = 2 > 1$.

In order to give exact bounds we now introduce the following:

$$L = \min_{i \in M} \min_{j \in N_i} a_{ij}$$

and

$$U = \min_{i \in M} \max_{j \in N_i} a_{ij}.$$

PROPOSITION 20. $L = \lambda_0$ for any $A \in \mathbb{R}^{m \times n}$.

Proof. There exist $r \in M$ and $s \in N$ such that

$$\lambda_0 = \min_{j \in N} \max_{i \in M} a_{ij} = \max_{i \in M} a_{is} = a_{rs}.$$

So a_{rs} is the smallest column maximum in A . Let $k \in M$. The quantity $\min_{l \in N_k} a_{kl}$ is the smallest of all column maxima appearing in row k of A (recall that this value is $+\infty$ if $N_k = \emptyset$). Hence $\min_{l \in N_k} a_{kl} \geq a_{rs}$, and therefore also $L = \min_{k \in M} \min_{l \in N_k} a_{kl} \geq a_{rs} = \lambda_0$.

On the other hand, $L = \min_{k \in M} \min_{l \in N_k} a_{kl} \leq \min_{l \in N_r} a_{rl} = a_{rs} = \lambda_0$. \square

PROPOSITION 21. $\Lambda(A, 0) \subseteq [L, U]$ holds for every $A = (a_{ij}) \in \mathbb{R}^{m \times n}$ and $\{L, U\} \subseteq \Lambda(A, 0)$ whenever $\Lambda(A, 0) \neq \emptyset$.

Proof. The lower bounds in Propositions 17 and 21 coincide by Proposition 20, so we need only prove the upper bound and its tightness. Suppose without loss of generality that $\Lambda(A, 0) \neq \emptyset$ and so $N_i \neq \emptyset$ for every $i \in M$. Let $\lambda = U = \min_{i \in M} \max_{j \in N_i} a_{ij}$. Then $\lambda = \max_{i \in M} a_{is}$ for some $s \in N$, and thus $m+1 \in M'_s$. At the same time if $r \in M$, then $\max_{j \in N_r} a_{rj} \geq \min_{i \in M} \max_{j \in N_i} a_{ij} = \lambda$. Therefore, $r \in M'_j$ for some $j \in N_r$. This shows that $U \in \Lambda(A, 0)$ by Proposition 16.

Suppose now that $\lambda > U$. Then $\lambda > \max_{j \in N_i} a_{ij}$ for some $i \in M$. Hence, $\lambda > a_{ij}$ for all $j \in N_i$ and so $i \notin M'_j$ for all $j \in N_i$. Since also $i \notin M_j$ for all $j \notin N_i$ and $M_j \subseteq M'_j$ for every $j \in N_i$, we have that $i \notin M_j$ for all $j \in N$, and thus $\lambda \notin \Lambda(A, 0)$ by Proposition 18. \square

THEOREM 22. $\Lambda(A, 0) = [L, U]$ holds for every $A = (a_{ij}) \in \mathbb{R}^{m \times n}$.

Proof. If $\Lambda(A, 0) = \emptyset$, then by Proposition 18 $N_i = \emptyset$ for some $i \in M$ and so $U = -\infty$, $L = +\infty$, and $[L, U] = \emptyset$.

Suppose now $\Lambda(A, 0) \neq \emptyset$. Due to Proposition 21 we may also assume that $L < U$, and we need only prove that $(L, U) \subseteq \Lambda(A, 0)$. Let $\lambda \in (L, U)$. If $i \in M$, then $\lambda < U \leq \max_{j \in N_i} a_{ij} = a_{it}$ for some $t \in N_i$. Hence, $i \in M_t = M'_t$ and so $i \in \bigcup_{j \in N} M'_j$. On the other hand, $\lambda > L \geq a_{rs}$, where $r \in M, s \in N_r$, hence $\lambda = \max_{i \in M'} a'_{is}$, and thus $m+1 \in M'_s$. We conclude that $\bigcup_{j \in N} M'_j = M'$ and so $\lambda \in \Lambda(A, 0)$ by Proposition 16. \square

5. Two-dimensional generalized eigenproblem. In this section we give a complete description of the spectrum of (A, B) and all eigenspaces, where $A, B \in \mathbb{R}^{m \times 2}$.

The case when $x_1 = \varepsilon$ reduces the generalized eigenproblem to the question of whether the second columns of A and B are proportional (and the coefficient of proportionality is then the unique generalized eigenvalue). This holds similarly for $x_2 = \varepsilon$, so we will restrict our attention to the task of finding finite x . By homogeneity of $V(A, B, \lambda)$ we can assume that $x_1 = 0$. We will therefore study the one-variable problem

$$(14) \quad a_{i1} \oplus a_{i2}x_2 = \lambda b_{i1} \oplus \lambda b_{i2}x_2, \quad i \in M.$$

Before we answer the main question, we will show in subsection 5.1 how to find all solutions of two-sided systems

$$(15) \quad Ax = Bx, x \in \mathbb{R}^2$$

for $A, B \in \mathbb{R}^{m \times 2}$, and then in subsection 5.2 we show how to solve (14) for $m = 2$. The following technical lemma will be useful.

LEMMA 23 (cancellation rule). *Let $v, w, a, b \in \mathbb{R}, a < b$. Then for any real x we have*

$$v \oplus ax = w \oplus bx$$

if and only if

$$v = w \oplus bx.$$

Proof. The proof is straightforward; see also [12, Lemma 7.4.1]. \square

This section can be seen as an independent generalization of the results for 2×2 matrices in [6] to $m \times 2$ matrices.

5.1. Two-dimensional two-sided systems. Suppose that a system

$$(16) \quad a_{i1} \oplus a_{i2}x_2 = b_{i1} \oplus b_{i2}x_2; i \in M$$

is given (as explained before, $x_1 = 0$ is assumed without loss of generality). Let us denote

$$V_i = \{x_2 \in \mathbb{R}; a_{i1} \oplus a_{i2}x_2 = b_{i1} \oplus b_{i2}x_2\}, \quad i \in M,$$

and $V = \bigcap_{i \in M} V_i$.

We will explicitly describe each V_i . Let us apply the cancellation rule of Lemma 23 to (16). For every $i \in M$ there are either two, or one, or no cancellations.

(i) If there is no cancellation, then

$$a_{i1} = b_{i1} \quad \text{and} \quad a_{i2} = b_{i2}$$

and so $V_i = \mathbb{R}$.

(ii) If there is exactly one cancellation, then we can assume without loss of generality that it takes place on the left-hand side, and we consider two cases.

Either $a_{i1} < b_{i1}$ and $a_{i2} = b_{i2}$ so that (16) reduces to

$$a_{i2}x_2 = b_{i1} \oplus a_{i2}x_2$$

yielding $V_i = [a_{i2}^{-1}b_{i1}, +\infty)$, or $a_{i1} = b_{i1}$ and $a_{i2} < b_{i2}$ so that (16) reduces to

$$a_{i1} = a_{i1} \oplus b_{i2}x_2$$

yielding $V_i = (-\infty, a_{i1}b_{i2}^{-1}]$.

(iii) If there are two cancellations and they take place on the same side, then this side becomes ε yielding $V_i = \emptyset$. If the two cancellations take place on different sides, then either $a_{i1} > b_{i1}$ and $a_{i2} < b_{i2}$ so that (16) reduces to

$$a_{i1} = b_{i2}x_2$$

yielding $V_i = \{a_{i1}b_{i2}^{-1}\}$, or $a_{i1} < b_{i1}$ and $a_{i2} > b_{i2}$ yielding similarly $V_i = \{a_{i2}^{-1}b_{i1}\}$.

Since each V_i obtained above is a closed interval (including possibly a singleton or empty set) and can be found in a constant number of operations, the intersection $V = \bigcap_{i \in M} V_i$ is also a closed interval (including possibly a singleton or empty set) and can be found in $O(m)$ time.

We conclude with the following.

PROPOSITION 24. *The solution set to (15) is of the form*

$$\left\{ \alpha(0, x_2)^T; \alpha \in \mathbb{R}, x_2 \in V \right\},$$

where V is a closed interval (including possibly a singleton or empty set) and can be found in $O(m)$ time as described above.

5.2. Generalized eigenproblem for 2×2 matrices. Our aim in this subsection is to describe the whole spectrum for the 2×2 generalized eigenproblem

$$(17) \quad \begin{aligned} a_{11} \oplus a_{12}x_2 &= \lambda b_{11} \oplus \lambda b_{12}x_2, \\ a_{21} \oplus a_{22}x_2 &= \lambda b_{21} \oplus \lambda b_{22}x_2, \end{aligned}$$

where all a_{ij} and b_{ij} are real numbers.

This is a very special case, already solved in [6], but it will be of key importance for solving the general two-dimensional case in the next subsection. We use a different methodology from that in [6] for solving the general case in subsection 5.3.

Recall that by Proposition 10, $\Lambda(A, B) \subseteq [\underline{\lambda}, \bar{\lambda}]$ (see section 2). Both $\underline{\lambda}$ and $\bar{\lambda}$ can easily be found (in $O(mn)$ time). We will therefore assume that $\underline{\lambda} < \bar{\lambda}$, since otherwise we have $\Lambda(A, B) = \emptyset$ (if $\underline{\lambda} > \bar{\lambda}$), or $\underline{\lambda} = \bar{\lambda}$ is the unique candidate for a value in $\Lambda(A, B)$, and this can be verified easily, for instance, using the tools of subsection 5.1. We will distinguish four cases and describe the spectrum in each of them. The feasibility interval $[\underline{\lambda}, \bar{\lambda}]$ has exactly one of the forms

$$\begin{aligned} &[\max(c_{11}, c_{21}), \min(c_{12}, c_{22})], \\ &[\max(c_{12}, c_{22}), \min(c_{11}, c_{21})], \\ &[\max(c_{11}, c_{22}), \min(c_{12}, c_{21})], \\ &[\max(c_{12}, c_{21}), \min(c_{11}, c_{22})]. \end{aligned}$$

Note that the first case can be equivalently described by inequalities $c_{11} < c_{12}, c_{21} < c_{22}$; this holds similarly for the other cases. The first two cases can be transformed into each other by swapping the variables x_1 and x_2 and similarly for the last two cases. So we essentially have only two cases. In fact we will deal only with the third (and thus also with the fourth) case as the first (two) will be covered by the discussion in subsection 5.3.

In what follows we denote

$$\gamma^2 = a_{12}a_{21}b_{11}^{-1}b_{22}^{-1}.$$

PROPOSITION 25. *If $c_{11} < c_{12}, c_{22} < c_{21}$, and $\underline{\lambda} < \bar{\lambda}$, then $\Lambda(A, B) = \{\hat{\gamma}\}$, where $\hat{\gamma}$ is the unique projection of γ onto $[\underline{\lambda}, \bar{\lambda}]$, that is,*

$$\hat{\gamma} = \begin{cases} \underline{\lambda} & \text{if } \gamma \leq \underline{\lambda}, \\ \gamma & \text{if } \gamma \in (\underline{\lambda}, \bar{\lambda}), \\ \bar{\lambda} & \text{if } \gamma \geq \bar{\lambda}. \end{cases}$$

Proof. Note first that by the assumptions, we have $\underline{\lambda} = \max(c_{11}, c_{22})$ and $\bar{\lambda} = \min(c_{12}, c_{21})$. Let us denote

$$S = (\underline{\lambda}, \bar{\lambda}) \cap \Lambda(A, B).$$

It is sufficient to prove the following statements:

- (i) $S \neq \emptyset \implies S = \{\gamma\}$,
- (ii) $\gamma \in (\underline{\lambda}, \bar{\lambda}) \implies \gamma \in S$,
- (iii) $\gamma \in (\underline{\lambda}, \bar{\lambda}) \implies \underline{\lambda}, \bar{\lambda} \notin \Lambda(A, B)$,
- (iv) $\gamma \leq \underline{\lambda} \implies \Lambda(A, B) = \{\underline{\lambda}\}$, and
- (v) $\gamma \geq \bar{\lambda} \implies \Lambda(A, B) = \{\bar{\lambda}\}$.

In order to prove (i) suppose $\lambda \in S$. Hence, we have

$$c_{11}, c_{22} < \lambda < c_{12}, c_{21},$$

and thus (using $c_{ij} = a_{ij}b_{ij}^{-1}$)

$$\begin{aligned} a_{11} &< \lambda b_{11}, & a_{12} &> \lambda b_{12}, \\ a_{22} &< \lambda b_{22}, & a_{21} &> \lambda b_{21}. \end{aligned}$$

Using the cancellation rule of Lemma 23 in this case, system (17) reduces to

$$(18) \quad \left. \begin{aligned} a_{12}x_2 &= \lambda b_{11} \\ a_{21} &= \lambda b_{22}x_2 \end{aligned} \right\}.$$

So $x_2 = \lambda b_{11}a_{12}^{-1}$ and $x_2 = \lambda^{-1}a_{21}b_{22}^{-1}$, from which $\lambda = \gamma$ follows.

(ii) Suppose $\gamma \in (\underline{\lambda}, \bar{\lambda})$ and put $\lambda = \gamma$. By taking $x_2 = \lambda b_{11}a_{12}^{-1} = \lambda^{-1}a_{21}b_{22}^{-1}$, we see that $\lambda \in \Lambda(A, B)$.

(iii) Suppose that $\gamma \in (\underline{\lambda}, \bar{\lambda})$ and $\lambda = \underline{\lambda} \in \Lambda(A, B)$. If $c_{11} < c_{22}$, then

$$c_{11} < c_{22} = \lambda < c_{12}, c_{21}$$

and thus,

$$\begin{aligned} a_{11} &< \lambda b_{11}, & a_{12} &> \lambda b_{12}, \\ a_{22} &= \lambda b_{22}, & a_{21} &> \lambda b_{21}. \end{aligned}$$

By cancellations and substituting λb_{22} for a_{22} , system (17) reduces to

$$(19) \quad \left. \begin{aligned} a_{12}x_2 &= \lambda b_{11} \\ a_{21} \oplus \lambda b_{22}x_2 &= \lambda b_{22}x_2 \end{aligned} \right\}.$$

So $x_2 = \lambda b_{11}a_{12}^{-1}$ and $x_2 \geq \lambda^{-1}a_{21}b_{22}^{-1}$, from which $\lambda^2 \geq \gamma^2$, a contradiction.

A contradiction is obtained in a similar way when $c_{11} > c_{22}$ or $c_{11} = c_{22}$.

The case of $\lambda = \bar{\lambda} \in \Lambda(A, B)$ is dealt with in a similar way.

(iv) Suppose $\gamma \leq \underline{\lambda}$. Due to (i) it is sufficient to prove that $\underline{\lambda} \in \Lambda(A, B)$ and $\bar{\lambda} \notin \Lambda(A, B)$. Let $\lambda = \underline{\lambda}$. It is easily verified that x_2 is a solution to (17), where

$$x_2 = \begin{cases} \lambda b_{11}a_{12}^{-1} & \text{if } c_{11} < c_{22}, \\ \lambda^{-1}a_{21}b_{22}^{-1} & \text{if } c_{11} > c_{22}, \text{ and} \\ \text{any value in } [a_{11}a_{22}^{-1}, a_{11}a_{12}^{-1}] & \text{if } c_{11} = c_{22}. \end{cases}$$

Let $\lambda = \bar{\lambda}$ and suppose $c_{12} < c_{21}$. Then $\lambda > \gamma$ and

$$c_{11}, c_{22} < \lambda = c_{12} < c_{21}$$

and thus,

$$\begin{aligned} a_{11} &< \lambda b_{11}, & a_{12} &= \lambda b_{12}, \\ a_{22} &< \lambda b_{22}, & a_{21} &> \lambda b_{21}. \end{aligned}$$

By cancellations and substituting λb_{12} for a_{12} , system (17) reduces to

$$(20) \quad \left. \begin{aligned} a_{12}x_2 &= \lambda b_{11} \oplus a_{12}x_2 \\ a_{21} &= \lambda b_{22}x_2 \end{aligned} \right\}.$$

So $x_2 \geq \lambda a_{12}^{-1} b_{11}$ and $x_2 = \lambda^{-1} b_{22}^{-1} a_{21}$, from which $\lambda^2 \leq \gamma^2$, a contradiction.

A contradiction can similarly be obtained when $c_{12} > c_{21}$ or $c_{12} = c_{21}$.

(v) The proof of this part is similar to that of (iv) and is omitted here. \square

5.3. Generalized eigenproblem: The two-dimensional case. As before, due to the finiteness of x and homogeneity, we assume that $x_1 = 0$ and therefore study system (17).

We will distinguish three cases.

Case 1: If $c_{i1} = c_{i2}$ for some $i \in M$, then by Proposition 10 this value is the unique candidate for the generalized eigenvalue. Using the method of subsection 5.1 it can be readily checked whether this is indeed the case.

Case 2: If $c_{i_1 1} < c_{i_1 2}$ and $c_{i_2 1} > c_{i_2 2}$ for some $i_1, i_2 \in M$, then the 2×2 system

$$\begin{pmatrix} a_{i_1 1} & a_{i_1 2} \\ a_{i_2 1} & a_{i_2 2} \end{pmatrix} x = \lambda \begin{pmatrix} b_{i_1 1} & b_{i_1 2} \\ b_{i_2 1} & b_{i_2 2} \end{pmatrix} x$$

has a unique eigenvalue by Proposition 25 and is therefore a unique candidate for an eigenvalue of the whole system. This can easily be checked by the method of subsection 5.1.

Case 3: If $c_{i1} < c_{i2}$ for all $i \in M$ (the case when $c_{i1} > c_{i2}$ for all $i \in M$ can be discussed similarly), then for any $i \in M$ the feasibility interval for the i th equation alone is $[c_{i1}, c_{i2}]$. Suppose $\lambda \in \Lambda(A, B) \cap (c_{i1}, c_{i2})$. Then $a_{i1} < \lambda b_{i1}$ and $a_{i2} > \lambda b_{i2}$, and the equation

$$(21) \quad a_{i1} \oplus a_{i2}x_2 = \lambda b_{i1} \oplus \lambda b_{i2}x_2$$

using cancellations reduces to

$$a_{i2}x_2 = \lambda b_{i1}.$$

Hence, $x_2 = \lambda b_{i1} a_{i2}^{-1}$, and thus the dependence of x_2 on λ over (c_{i1}, c_{i2}) is expressed by a linear function (with slope 1). This concludes the case when λ is strictly between c_{i1} and c_{i2} . To finish Case 3 suppose now that $\lambda = c_{i1} \in \Lambda(A, B)$. Then $a_{i1} = \lambda b_{i1}$ and $a_{i2} > \lambda b_{i2}$, and (21) using cancellations reduces to

$$a_{i1} \oplus a_{i2}x_2 = a_{i1}.$$

Hence $x_2 \leq a_{i1} a_{i2}^{-1}$. Similarly if $\lambda = c_{i2} \in \Lambda(A, B)$, then $x_2 \geq b_{i1} b_{i2}^{-1}$. Note that

$$\lim_{\lambda \rightarrow c_{i1}} \lambda b_{i1} a_{i2}^{-1} = a_{i1} a_{i2}^{-1} \quad \text{and} \quad \lim_{\lambda \rightarrow c_{i2}} \lambda b_{i1} a_{i2}^{-1} = b_{i1} b_{i2}^{-1},$$

and so the graph of dependence of x_2 on λ over $[c_{i1}, c_{i2}]$ is a continuous, piecewise linear map; see Figure 1. This result is consistent with the fact that $a_{i1} a_{i2}^{-1} < b_{i1} b_{i2}^{-1}$ since this is equivalent to $c_{i1} < c_{i2}$.




FIG. 1. A single equation.

Finally, we note that for the whole system we have

$$(\underline{\lambda}, \bar{\lambda}) = \bigcap_{i \in M} (c_{i1}, c_{i2}).$$

Thus if $\lambda \in \Lambda(A, B) \cap (\underline{\lambda}, \bar{\lambda})$, then $\lambda \in \Lambda(A, B) \cap (c_{i1}, c_{i2})$ for every $i \in M$, and so x_2 is the *common* value of all $\lambda b_{i1} a_{i2}^{-1}$, $i \in M$. This implies $(\underline{\lambda}, \bar{\lambda}) \subseteq \Lambda(A, B)$. We have proved the following.

PROPOSITION 26. *If $A, B \in \mathbb{R}^{m \times 2}$ and $c_{i1} < c_{i2}$ for every $i \in M$, then a generalized eigenvalue in $(\underline{\lambda}, \bar{\lambda})$ exists if and only if all values in $(\underline{\lambda}, \bar{\lambda})$ are generalized eigenvalues. This is equivalent to the requirement that all values $b_{i1} a_{i2}^{-1}$ for $i \in M$ coincide.*

If the condition in Proposition 26 is satisfied, then by continuity also $\underline{\lambda}, \bar{\lambda} \in \Lambda(A, B)$, and in this case $\Lambda(A, B) = [\underline{\lambda}, \bar{\lambda}]$. If not, then $\underline{\lambda}, \bar{\lambda}$ have to be examined separately to see if they are generalized eigenvalues. Figures 2–6 indicate that all possibilities may occur (both $\underline{\lambda}$ and $\bar{\lambda}$, exactly one, or none in $\Lambda(A, B)$).

Summarizing all cases we have that if $A, B \in \mathbb{R}^{m \times 2}$, then $\Lambda(A, B)$ can be found in $O(m)$ time and has one of the following forms (the illustrating figures are drawn for $m = 2$):

- $[\underline{\lambda}, \bar{\lambda}]$ (see Figure 2);
- $\{\underline{\lambda}, \bar{\lambda}\}$ (see Figure 3);
- $\{\lambda\}$, where $\lambda \in [\underline{\lambda}, \bar{\lambda}]$ (see Case 2 and Figures 4 and 5);
- \emptyset (see Figure 6 and Case 1).

In all cases the eigenspace associated with a fixed generalized eigenvalue is described in Proposition 24.




FIG. 2. A continuum of solutions: $\lambda \in [\underline{\lambda} = a_{21}b_{21}^{-1}, \bar{\lambda} = a_{12}b_{12}^{-1}]$, $x_2 = \lambda b_{11}a_{12}^{-1} = \lambda b_{21}a_{22}^{-1}$.




FIG. 3. Two solutions: $(\lambda, x_2) = (\underline{\lambda} = a_{21}b_{21}^{-1}, a_{21}b_{21}^{-1}b_{11}a_{12}^{-1})$ and $(\lambda, x_2) = (\bar{\lambda} = a_{12}b_{12}^{-1}, a_{12}b_{12}^{-1}b_{21}a_{22}^{-1})$.




FIG. 4. One solution: $(\lambda, x_2) = (\underline{\lambda} = a_{21}b_{21}^{-1}, a_{21}b_{21}^{-1}b_{11}a_{12}^{-1})$.




FIG. 5. One solution: $(\lambda, x_2) = (\bar{\lambda} = a_{22}b_{22}^{-1}, a_{22}b_{22}^{-1}b_{11}a_{12}^{-1})$.




FIG. 6. No solutions.

REFERENCES

- [1] M. AKIAN, R. BAPAT, AND S. GAUBERT, *Max-plus algebras*, in Handbook of Linear Algebra, Discrete Math. Appl. 39, L. Hogben, ed., Chapman & Hall/CRC, Boca Raton, FL, 2006, pp. 25/1–25/17.
- [2] M. AKIAN, S. GAUBERT, AND A. GUTERMAN, *Tropical polyhedra are equivalent to mean payoff games*, Int. J. Algebra Comput., 22 (2012), 125001.
- [3] X. ALLAMIGEON, S. GAUBERT, AND E. GOUBAULT, *Computing the vertices of tropical polyhedra using directed hypergraphs*, Discrete Comput. Geom., 49 (2013), pp. 247–279.
- [4] F. L. BACCELLI, G. COHEN, G.-J. OLSDER, AND J.-P. QUADRAT, *Synchronization and Linearity*, John Wiley, Chichester, New York, 1992.
- [5] M. BEZEM, R. NIEUWENHUIS, AND E. RODRÍGUEZ-CARBONELL, *Hard problems in max-algebra, control theory, hypergraphs and other areas*, Inform. Process. Lett., 110 (2010), pp. 133–138.
- [6] P. A. BINDING AND H. VOLKMER, *A generalized eigenvalue problem in the max algebra*, Linear Algebra Appl., 422 (2007), pp. 360–371.
- [7] P. BUTKOVIČ, *On certain properties of the systems of linear extremal equations*, Ekonom.-Mat. Obzor, 14 (1978), pp. 72–78.
- [8] P. BUTKOVIČ, *Solution of systems of linear extremal equations*, Ekonom.-Mat. Obzor, 17 (1981), pp. 402–416.
- [9] P. BUTKOVIČ, *On properties of solution sets of extremal linear programs*, Ann. Discrete Math., 19 (1984), pp. 41–54.
- [10] P. BUTKOVIČ AND G. HEGEDÜS, *An elimination method for finding all solutions of the system of linear equations over an extremal algebra*, Ekonom.-Mat. Obzor, 20 (1984), pp. 203–214.
- [11] P. BUTKOVIČ, *Necessary solvability conditions of systems of linear extremal equations*, Discrete Appl. Math., 10 (1985), pp. 19–26.
- [12] P. BUTKOVIČ, *Max-linear Systems: Theory and Algorithms*, Springer Monogr. Math., Springer-Verlag, London, 2010.
- [13] P. BUTKOVIČ, R. A. CUNINGHAME-GREEN, AND S. GAUBERT, *Reducible spectral theory with applications to the robustness of matrices in max-algebra*, SIAM J. Matrix Anal. Appl., 31 (2009), pp. 1412–1431, doi:10.1137/080731232.
- [14] R. A. CUNINGHAME-GREEN, *Process synchronisation in a steelworks—a problem of feasibility*, in Proceedings of the 2nd International Conference on Operational Research, J. Banbury and J. Maitland, eds., English University Press, London, 1960, pp. 323–328.
- [15] R. A. CUNINGHAME-GREEN, *Describing industrial processes with interference and approximating their steady-state behaviour*, Oper. Res. Quart., 13 (1962), pp. 95–100.
- [16] R. A. CUNINGHAME-GREEN, *Minimax Algebra*, Lecture Notes in Econom. and Math. Systems 166, Springer-Verlag, Berlin, 1979.

- [17] R. A. CUNINGHAME-GREEN AND P. BUTKOVIČ, *The equation $A \otimes x = B \otimes y$ over $(\max, +)$* , Theoret. Comput. Sci., 293 (2003), pp. 3–12.
- [18] R. A. CUNINGHAME-GREEN AND P. BUTKOVIČ, *Generalised eigenproblem in max algebra*, in Proceedings of the 9th IEEE International Workshop on Discrete Event Systems (WODES 2008), Göteborg, Sweden, 2008, pp. 236–241.
- [19] S. GAUBERT, *Théorie des systèmes linéaires dans les dioïdes*, Thèse, Ecole des Mines de Paris, Paris, 1992.
- [20] S. GAUBERT, R. D. KATZ, AND S. SERGEEV, *Tropical linear-fractional programming and parametric mean-payoff games*, J. Symbolic Comput., 47 (2012), pp. 1447–1478.
- [21] S. GAUBERT AND S. SERGEEV, *The level set method for the two-sided max-plus eigenproblem*, Discrete Event Dyn. Syst., 23 (2013), pp. 105–134.
- [22] M. GONDTRAN AND M. MINOUX, *Valeurs propres et vecteur propres dans les dioïdes et leur interprétation en théorie des graphes*, Bull. Direction Études Recherches Sér. C Math. Informat., 1977, no. 2, pp. 25–41.
- [23] B. HEIDERGOTT, G.-J. OLSDER, AND J. VAN DER WOUDE, *Max Plus at Work: Modeling and Analysis of Synchronized Systems: A Course on Max-Plus Algebra and Its Applications*, Princeton University Press, Princeton, NJ, 2005.
- [24] G. OWEN, *Game Theory*, 3rd ed., Academic Press, New York, 1995.
- [25] S. SERGEEV, *On the system $Ax = \lambda Bx$ in max algebra: Every system of intervals is a spectrum*, Kybernetika, 47 (2011), pp. 715–721.
- [26] N. N. VOROB'EV, *Extremal algebra of positive matrices*, Elektron. Informationsverarbeit. Kybernetik, 3 (1967), pp. 39–71 (in Russian).
- [27] E. A. WALKUP AND G. BORIELLO, *A general linear max-plus solution technique*, in Idempotency (Bristol, 1994), Publ. Newton Inst. 11, J. Gunawardena, ed., Cambridge University Press, Cambridge, UK, 1998, pp. 406–415.
- [28] K. ZIMMERMANN, *Extremální Algebra*, Research Publications 46, Department of Scientific Information, Economic Institute of the Czech Academy of Sciences, Prague, 1976 (in Czech).