

Magnesium for Atrial Fibrillation: Myth or Magic?

Kotecha, Dipak

DOI:

[10.1161/CIRCEP.116.004521](https://doi.org/10.1161/CIRCEP.116.004521)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Kotecha, D 2016, 'Magnesium for Atrial Fibrillation: Myth or Magic?', *Circulation Arrhythmia and Electrophysiology*, vol. 9, no. 9. <https://doi.org/10.1161/CIRCEP.116.004521>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Checked for eligibility: 05/10/2016

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Title: Magnesium for Atrial Fibrillation: Myth or Magic?

Running title: Magnesium for Atrial Fibrillation: Myth or Magic?

(49 characters including spaces)

Author: Dipak Kotecha, MD PhD PCAP MRCP FESC FHEA

Author affiliations: Institute of Cardiovascular Sciences, University of Birmingham, Birmingham, UK; and Monash Centre of Cardiovascular Research and Education in Therapeutics, Monash University, Melbourne, Australia.

Correspondence: d.kotecha@bham.ac.uk

Journal subject terms

Arrhythmia and Electrophysiology: Atrial Fibrillation

Quality and Outcomes: Health Services

Word count: 927

Key words: Atrial fibrillation; Magnesium, Cardioversion; Prevention

Low magnesium levels have been implicated as a risk factor for the development of atrial fibrillation.¹ However, like many interventions before it, supplementing magnesium levels does not necessarily lead to a successful cardioversion or better prevention of AF in high-risk groups. This issue is clinically important in view of the high burden that AF poses, the increasing incidence of AF, and the association of AF with adverse long-term clinical outcomes.² Not only is AF prevalent in the general population (approximately 3%, with much higher rates in older patients), but also a frequent comorbidity and risk factor in patients with a range of cardiovascular conditions, including heart failure and after cardiac surgery.^{3,4} Identifying practical and effective methods for managing AF is a clinical imperative.

The attention on magnesium as a potential anti-arrhythmic agent is founded on a small number of physiological assessments in human and animal models.⁵ Intravenous magnesium directly affects myocardial potassium channels, has voltage-dependent and indirect effects on calcium and sodium channels, prolongs the PR interval and increases the refractory period of antegrade atrioventricular node conduction.^{6,7} However, although low serum magnesium levels were associated with incident AF in the Atherosclerosis Risk in Communities (ARIC) study, dietary levels were not⁸, suggesting a causal disconnect. This was demonstrated in another large community cohort where the association of hypomagnesemia with incident AF was identified in long-term follow-up, but not present within the first 90 days.⁹

In this issue of *Circulation: Arrhythmia and Electrophysiology*, Rajagopalan and colleagues add to the growing evidence-base on this issue by performing a randomized, double-blind, placebo-controlled trial of intravenous magnesium prior to electrical cardioversion of AF.¹⁰ 261 patients were enrolled with normal magnesium levels at baseline (2.1 ± 0.2 mg/dL).

Their key finding was that one-hour conversion to sinus rhythm was similar in both the magnesium-treated patients and the placebo group (86.4% magnesium versus 86.0% placebo). They also found no difference in biphasic energy requirement, or the number of shocks needed in a ramping energy protocol.

Where does this fit in with other studies? Figure 1 displays trials and meta-analyses that have randomized patients to magnesium in a range of situations, including treatment of acute AF, prevention of AF during cardiac surgery and facilitation of electrical cardioversion.¹⁰⁻¹⁴ Although varying with respect to population and magnesium dosage, they share common features of a small sample size, short follow-up and disappointing treatment effect.

It is likely that patient selection is a major issue – AF is not just a single condition, but the endpoint of numerous pathologies.¹⁵ Future trials, in this and other areas, should target patients better in order to optimize the likelihood of demonstrating treatment effects. Another key problem is the length of follow-up. The possibility of a longer-term benefit of magnesium supplementation is unknown from the current literature.

Sample size is also important – demonstrating the effect of a treatment is challenging when immediate success rates are so high. This study, and others, may actually be underpowered when considering the heterogeneity of patients included, particularly with respect to anti-arrhythmic drug use. Nonetheless this data, in addition to other studies discussed, would suggest that magnesium is not a useful clinical therapy to improve the success rates of electrical cardioversion. Although there are data to support the use of magnesium for facilitating pharmacological cardioversion, this is largely based on retrospective analyses in patients receiving ibutilide or dofetilide as antiarrhythmic drug therapy.¹⁶⁻¹⁸

Magnesium has also been noted as a potential drug for complementing a rate control


approach. In meta-analysis of five randomized trials (n=380), patients receiving magnesium were three times as likely to reach a heart rate <100 beats/minute compared to placebo, mostly with digoxin as background therapy.¹³ In 199 patients receiving rate control for rapid AF (again mostly with digoxin), those randomized to intravenous supplemental magnesium were more likely to reach a rate of <100 beats/minute at two hours (relative risk 1.89; 95% CI 1.38-2.59).¹⁹ While these results are encouraging, magnesium has not been widely adopted as a rate control strategy, as more effective agents are available for acute ventricular rate control, such as beta-blockers, verapamil and diltiazem.²

The authors of this study are congratulated on recognizing the importance of study design and choosing to employ a randomized and blinded approach. Magnesium has shown potential benefits in observational studies, and has an inherent physiological association due to its effects on membrane potentials and ion transport. This exemplifies the problems in assigning causality and disentangling the issue of confounding in observational studies, which we recently highlighted for digoxin use in AF.²⁰ Although observational data are useful for determining epidemiological patterns, decision on treatment effects should be restricted to randomized controlled trials, where selection and performance biases can be addressed. More trials of magnesium are in-process. Completed but yet to report is a 300-patient study in Tunisia randomizing to high or low-dose magnesium infusion, to assess effects on rate and rhythm control in emergency department patients with rapid AF (NCT00965874). Currently recruiting is a double-blind placebo-controlled trial in Thailand of 128 patients to assess sinus rhythm conversion and rate control over 6 hours (NCT01049464), and a Norwegian single-blind trial of 218 patients with paroxysmal AF or flutter, with outcomes of cardioversion success at 24 hours and also 3-month AF follow-up (NCT01818583).

Further trial data may shed light on whether there is any role for magnesium in improving the management of AF patients. However at present, the available data would suggest that magnesium, as an adjunct to electrical cardioversion or for prevention, is more myth than a practical, easy (or magical) solution to the growing problem of AF.

Figure 1

Effect of magnesium on success of cardioversion or prevention of AF



Legend: Selected studies and meta-analyses of randomized controlled trials of magnesium for facilitating cardioversion or preventing AF.¹⁰⁻¹⁴ An ‘event’ is a successful cardioversion to sinus rhythm, or the absence of AF on follow-up. AF = atrial fibrillation; CI = confidence interval; Mg = magnesium; N = total number of patients; RCT = randomized controlled trial.

Conflict of Interest Disclosures

DK has received research grants from Menarini, speaker fees from Atricure and professional development support from Daiichi Sankyo.

References

1. Khan AM, Lubitz SA, Sullivan LM, Sun JX, Levy D, Vasan RS, Magnani JW, Ellinor PT, Benjamin EJ, Wang TJ. Low serum magnesium and the development of atrial fibrillation in the community: the Framingham Heart Study. *Circulation*. 2013;127:33-38
2. Kirchhof P, Benussi S, Kotecha D, Ahlsson A, Atar D, Casadei B, Castella Pericas M, Diener H, Heidbuchel H, Hendriks J, Hindricks G, Manolis AS, Oldgren J, Popescu BA, Schotten U, Van Putte B, Vardas P. 2016 ESC guidelines for the management of atrial fibrillation developed in collaboration with EACTS: The Task Force for the management of atrial fibrillation of the European Society of Cardiology (ESC). Developed with the special contribution of the European Heart Rhythm Association (EHRA) of the ESC. Endorsed by the European Stroke Organisation (ESO). *Eur Heart J*. 2016; in press: DOI: 10.1093/eurheartj/ehw1210
3. Kotecha D, Piccini JP. Atrial fibrillation in heart failure: what should we do? *Eur Heart J*. 2015;36:3250-3257
4. Steinberg BA, Zhao Y, He X, Hernandez AF, Fullerton DA, Thomas KL, Mills R, Klaskala W, Peterson ED, Piccini JP. Management of postoperative atrial fibrillation and subsequent outcomes in contemporary patients undergoing cardiac surgery: insights from the Society of Thoracic Surgeons CAPS-Care Atrial Fibrillation Registry. *Clin Cardiol*. 2014;37:7-13
5. Delva P. Magnesium and cardiac arrhythmias. *Mol Aspects Med*. 2003;24:53-62
6. Bara M, Guiet-Bara A, Durlach J. Regulation of sodium and potassium pathways by magnesium in cell membranes. *Magnes Res*. 1993;6:167-177
7. Kulick DL, Hong R, Ryzen E, Rude RK, Rubin JN, Elkayam U, Rahimtoola SH, Bhandari AK. Electrophysiologic Effects of Intravenous Magnesium in Patients with Normal Conduction Systems and No Clinical-Evidence of Significant Cardiac Disease. *Am Heart J*. 1988;115:367-373
8. Misialek JR, Lopez FL, Lutsey PL, Huxley RR, Peacock JM, Chen LY, Soliman EZ, Agarwal SK, Alonso A. Serum and dietary magnesium and incidence of atrial fibrillation in whites and in African Americans--Atherosclerosis Risk in Communities (ARIC) study. *Circ J*. 2013;77:323-329
9. Markovits N, Kurnik D, Halkin H, Margalit R, Bialik M, Lomnicki Y, Loebstein R. Database evaluation of the association between serum magnesium levels and the risk of atrial fibrillation in the community. *Int J Cardiol*. 2016;205:142-146
10. Rajagopalan B, Shah Z, Narasimha D, Bhatia A, Kim CH, Switzer DF, Gudleski GH, Curtis AB. Efficacy of intravenous magnesium in facilitating cardioversion of atrial fibrillation. *Circ Arrhythm Electrophysiol*. 2016;9:e003968
11. Chu K, Evans R, Emerson G, Greenslade J, Brown A. Magnesium sulfate versus placebo for paroxysmal atrial fibrillation: a randomized clinical trial. *Acad Emerg*

Med. 2009;16:295-300

12. Cook RC, Yamashita MH, Kearns M, Ramanathan K, Gin K, Humphries KH. Prophylactic magnesium does not prevent atrial fibrillation after cardiac surgery: a meta-analysis. *Ann Thorac Surg.* 2013;95:533-541
13. Ho KM, Sheridan DJ, Paterson T. Use of intravenous magnesium to treat acute onset atrial fibrillation: a meta-analysis. *Heart.* 2007;93:1433-1440
14. Sultan A, Steven D, Rostock T, Hoffmann B, Mullerleile K, Servatius H, Drewitz I, Luker J, Meyer P, Salukhe T, Willems S. Intravenous administration of magnesium and potassium solution lowers energy levels and increases success rates electrically cardioverting atrial fibrillation. *J Cardiovasc Electrophysiol.* 2012;23:54-59
15. Fabritz L, Guasch E, Antoniadou C, Bardinet I, Benninger G, Betts TR, Brand E, Breithardt G, Bucklar-Suchankova G, Camm AJ, Cartlidge D, Casadei B, Chua WW, Crijns HJ, Deeks J, Hatem S, Hidden-Lucet F, Kaab S, Maniadakis N, Martin S, Mont L, Reinecke H, Sinner MF, Schotten U, Southwood T, Stoll M, Vardas P, Wakili R, West A, Ziegler A, Kirchhof P. Expert consensus document: Defining the major health modifiers causing atrial fibrillation: a roadmap to underpin personalized prevention and treatment. *Nat Rev Cardiol.* 2016;13:230-237
16. Coleman CI, Sood N, Chawla D, Talati R, Ghatak A, Kluger J. Intravenous magnesium sulfate enhances the ability of dofetilide to successfully cardiovert atrial fibrillation or flutter: results of the Dofetilide and Intravenous Magnesium Evaluation. *Europace.* 2009;11:892-895
17. Kalus JS, Spencer AP, Tsikouris JP, Chung JO, Kenyon KW, Ziska M, Kluger J, White CM. Impact of prophylactic i.v. magnesium on the efficacy of ibutilide for conversion of atrial fibrillation or flutter. *Am J Health Syst Pharm.* 2003;60:2308-2312
18. Tercius AJ, Kluger J, Coleman CI, White CM. Intravenous magnesium sulfate enhances the ability of intravenous ibutilide to successfully convert atrial fibrillation or flutter. *Pacing Clin Electrophysiol.* 2007;30:1331-1335
19. Davey MJ, Teubner D. A randomized controlled trial of magnesium sulfate, in addition to usual care, for rate control in atrial fibrillation. *Ann Emerg Med.* 2005;45:347-353
20. Ziff OJ, Lane DA, Samra M, Griffith M, Kirchhof P, Lip GY, Steeds RP, Townend J, Kotecha D. Safety and efficacy of digoxin: systematic review and meta-analysis of observational and controlled trial data. *BMJ.* 2015;351:h4451