

# Bleeding risk assessment in atrial fibrillation: observations on the use and misuse of bleeding risk scores.

Lip, Gregory; Lane, Deirdre

DOI:

[10.1111/jth.13386](https://doi.org/10.1111/jth.13386)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Lip, G & Lane, D 2016, 'Bleeding risk assessment in atrial fibrillation: observations on the use and misuse of bleeding risk scores.', *Journal of Thrombosis and Haemostasis*, vol. 14, no. 9, pp. 1711-1714.

<https://doi.org/10.1111/jth.13386>

[Link to publication on Research at Birmingham portal](#)

## Publisher Rights Statement:

This is the peer reviewed version of the following article: Lip GYH, Lane DA. Bleeding risk assessment in atrial fibrillation: observations on the use and misuse of bleeding risk scores. *J Thromb Haemost* 2016; 14: 1711–4, which has been published in final form at 10.1111/jth.13386. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving

## General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

# Patient with Atrial Fibrillation; Eligible for Oral Anticoagulation

## Bleeding risk assessment

Identifies 'at risk' patients for more regular review and follow-up

### EHR and 'electronic alerts'

Low risk=No action

High risk=Patient 'flagged up' for review

'High risk' is not a reason or excuse to withhold OAC

### Review and address potentially reversible bleeding risk factors

- Uncontrolled hypertension
- Labile INRs (if on VKA)
- Concomitant use of aspirin and NSAIDs in anticoagulated patient
- Alcohol excess

For people with an increased risk of bleeding the benefit of OAC usually, but not always, outweighs the bleeding risk; thus careful monitoring of bleeding risk is important

Do not withhold OAC solely because the person is at risk of having a fall