

The Widom-Rowlinson model, the hard-core model and the extremality of the complete graph

Cohen, Emma; Csikvari, Peter; Perkins, Will; Tetali, Prasad

DOI:

[10.1016/j.ejc.2016.11.003](https://doi.org/10.1016/j.ejc.2016.11.003)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Cohen, E, Csikvari, P, Perkins, W & Tetali, P 2017, 'The Widom-Rowlinson model, the hard-core model and the extremality of the complete graph', *European Journal of Combinatorics*, vol. 62, pp. 70-76.
<https://doi.org/10.1016/j.ejc.2016.11.003>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Checked 17/11/2016

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

THE WIDOM-ROWLINSON MODEL, THE HARD-CORE MODEL AND THE EXTREMALITY OF THE COMPLETE GRAPH

EMMA COHEN, PÉTER CSIKVÁRI, WILL PERKINS, AND PRASAD TETALI

ABSTRACT. Let H_{WR} be the path on 3 vertices with a loop at each vertex. D. Galvin [4, 5] conjectured, and E. Cohen, W. Perkins and P. Tetali [2] proved that for any d -regular simple graph G on n vertices we have

$$\text{hom}(G, H_{\text{WR}}) \leq \text{hom}(K_{d+1}, H_{\text{WR}})^{n/(d+1)}.$$

In this paper we give a short proof of this theorem together with the proof of a conjecture of Cohen, Perkins and Tetali [2]. Our main tool is a simple bijection between the Widom-Rowlinson model and the hard-core model on another graph. We also give a large class of graphs H for which we have

$$\text{hom}(G, H) \leq \text{hom}(K_{d+1}, H)^{n/(d+1)}.$$

In particular, we show that the above inequality holds if H is a path or a cycle of even length at least 6 with loops at every vertex.

1. INTRODUCTION

For graphs G and H , with vertex and edge sets V_G, E_G, V_H , and E_H respectively, a map $\varphi : V_G \rightarrow V_H$ is a homomorphism if $(\varphi(u), \varphi(v)) \in E_H$ whenever $(u, v) \in E_G$. The number of homomorphisms from G to H is denoted by $\text{hom}(G, H)$. When $H = H_{\text{ind}}$, an edge with a loop at one end, homomorphisms from G to H_{ind} correspond to independent sets in the graph G , and so $\text{hom}(G, H_{\text{ind}})$ counts the number of independent sets in G .

For a given H , the set of homomorphisms from G to H correspond to valid configurations in a corresponding statistical physics model with *hard constraints* (forbidden local configurations). The independent sets of G are the valid configurations of the *hard-core model* on G , a model of a random independent set from a graph. Another notable case is when $H = H_{\text{WR}}$, a path on 3 vertices with a loop at each vertex. In this case, we can imagine a homomorphism from G to H_{WR} as a 3-coloring of the vertex set of G subject to the requirement that a blue and a red vertex cannot be adjacent (with white vertices considered unoccupied); such a coloring is called a *Widom-Rowlinson configuration* of G , from the Widom-Rowlinson model of two particle types which repulse each other [12, 1]. See Figure 1.

For a fixed graph H , it is natural to study the normalized graph parameter

$$p_H(G) := \text{hom}(G, H)^{1/|V_G|},$$

where V_G denotes the number of vertices of the graph G .

2010 *Mathematics Subject Classification*. Primary: 05C35. Secondary: 05C31, 05C70, 05C80.

Key words and phrases. graph homomorphisms, Widom-Rowlinson model, hard-core model.

The second author is partially supported by the National Science Foundation under grant no. DMS-1500219, by the MTA Rényi "Lendület" Groups and Graphs Research Group, by the ERC Consolidator Grant 648017, and by the Hungarian National Research, Development and Innovation Office, NKFIH grant K109684. Research of the last author is supported in part by the NSF grant DMS-1407657.



FIGURE 1. The target graphs for the Widom-Rowlinson model and the hard-core model.

For $H = H_{\text{ind}}$, J. Kahn [7] proved that for any d -regular bipartite graph G ,

$$p_{H_{\text{ind}}}(G) \leq p_{H_{\text{ind}}}(K_{d,d}),$$

where $K_{d,d}$ is the complete bipartite graph with classes of size d . Y. Zhao [10] showed that one could drop the condition of bipartiteness in Kahn's theorem. That is, he showed that $p_{H_{\text{ind}}}(G) \leq p_{H_{\text{ind}}}(K_{d,d})$, for *any* d -regular graph G . Y. Zhao proved his result by reducing the general case to the bipartite case with a clever trick. He proved that

$$p_{H_{\text{ind}}}(G) \leq p_{H_{\text{ind}}}(G \times K_2),$$

where $G \times K_2$ is the bipartite graph obtained by replacing every vertex u of V_G by a pair of vertices $(u, 0)$ and $(u, 1)$ and replacing every edge $(u, v) \in E_G$ by the pair of edges $((u, 0), (v, 1))$ and $((u, 1), (v, 0))$. This is clearly a bipartite graph, and if G is d -regular then $G \times K_2$ is still d -regular.

D. Galvin [4, 5] conjectured a different behavior for $H = H_{\text{WR}}$: that instead of $K_{d,d}$, the complete graph K_{d+1} maximizes $p_{H_{\text{ind}}}(G)$ among d -regular graphs G . E. Cohen, W. Perkins and P. Tetali [2] proved that this was indeed the case:

Theorem 1.1. [2] *For any d -regular simple graph G on n vertices we have*

$$p_{H_{\text{WR}}}(G) \leq p_{H_{\text{WR}}}(K_{d+1});$$

in other words,

$$\text{hom}(G, H_{\text{WR}}) \leq \text{hom}(K_{d+1}, H_{\text{WR}})^{n/(d+1)}.$$

One of the goals of this paper is to give a very simple proof of this fact¹, along with a slight generalization. We use a trick similar to that used by Y. Zhao [10, 11]. We will need the following definition:

Definition 1.2. The *extended line graph* \tilde{H} of a (bipartite) graph H has $V_{\tilde{H}} = E_H$; two edges e and f of H are adjacent in \tilde{H} if

- (a) $e = f$,
- (b) e and f share a common vertex, or
- (c) e and f are opposite edges of a 4-cycle in G .

Throughout, V_H and E_H refer to the vertex-set and edge-set, respectively, of the graph H . If H is bipartite, we use A_H and B_H to refer to the parts of a fixed bipartition. Now we can give a generalization of Theorem 1.1:

Theorem 1.3. *If \tilde{H} is the extended line graph of a bipartite graph H , then for any d -regular simple graph G on n vertices we have*

$$p_{\tilde{H}}(G) \leq p_{\tilde{H}}(K_{d+1}),$$

¹In fact, Theorem 1.1 follows from a stronger result in [2] that the Widom-Rowlinson *occupancy fraction* is maximized by K_{d+1} . We note that this stronger result also follows from the transformation below and Theorem 1 of [3].

or in other words,

$$\text{hom}(G, \tilde{H}) \leq \text{hom}(K_{d+1}, \tilde{H})^{n/(d+1)}.$$

To see that Theorem 1.3 is a generalization of Theorem 1.1 it suffices to check that H_{WR} is precisely the extended line graph of the path on 4 vertices. In Section 3 we will prove a slight generalization of Theorem 1.3 which allows for weights on the vertices of H .

2. SHORT PROOF OF THEOREM 1.1

We are not the first to notice the following connection between the Widom-Rowlinson model and the hardcore model (see, e.g., Section 5 of [1]): Given a graph G , let G' be the bipartite graph with vertex set $V_{G'} = V_G \times \{0, 1\}$, where $(u, 0)$ and $(v, 1)$ are adjacent in G' whenever either $(u, v) \in E_G$ or $u = v$. That is, G' is $G \times K_2$ with the extra edges $((u, 0), (u, 1))$ for all $u \in V_G$. We will show that

$$\text{hom}(G, H_{\text{WR}}) = \text{hom}(G', H_{\text{ind}}).$$

Indeed, consider an independent set I in G' . Color $u \in V_G$ blue if $(u, 1) \in I$, red if $(u, 0) \in I$, and white if it is neither red or blue. Note that since I was an independent set and $((u, 0), (u, 1)) \in E_{G'}$, the color of vertex u is well-defined and this coloring is in fact a Widom-Rowlinson coloring of G . This same construction also works in the other direction, so

$$\text{hom}(G, H_{\text{WR}}) = \text{hom}(G', H_{\text{ind}}).$$

If G is d -regular then G' is $(d+1)$ -regular, and $K'_{d+1} = K_{d+1, d+1}$. Applying J. Kahn's result [7] for $(d+1)$ -regular bipartite graphs, we see that if G has n vertices then

$$\begin{aligned} \text{hom}(G, H_{\text{WR}}) &= \text{hom}(G', H_{\text{ind}}) \\ &\leq \text{hom}(K_{d+1, d+1}, H_{\text{ind}})^{2n/(2(d+1))} = \text{hom}(K_{d+1}, H_{\text{WR}})^{n/(d+1)}. \end{aligned}$$

We remark that the transformation $G \rightarrow G'$ is also mentioned in [8].

3. EXTENSION

In this section we would like to point out that for every graph H there is an \tilde{H} such that

$$\text{hom}(G, \tilde{H}) = \text{hom}(G', H),$$

where G' is the bipartite graph defined in the previous section. Exactly the same argument we used for H_{WR} will work for any graph \tilde{H} constructed in this manner. Actually, the situation is even better. To give the most general version we need a definition.

Definition 3.1. Let G be a bipartite graph. Let H be another bipartite graph equipped with a weight function $\nu : V_H \rightarrow \mathbb{R}_+$. Let $\mathbb{I}_{E_H} : A_H \times B_H \rightarrow \{0, 1\}$ denote the characteristic function of E_H . Define

$$Z_b(G, H) = \sum_{\substack{\varphi: V_G \rightarrow V_H \\ \varphi(A_G) \subseteq A_H \\ \varphi(B_G) \subseteq B_H}} \prod_{(a, b) \in E_G} \mathbb{I}_{E_H}(\varphi(a), \varphi(b)) \prod_{w \in V_G} \nu(\varphi(w)),$$

(The subscript b stands for bipartite.) If G and H are not necessarily bipartite graphs, but H is a weighted graph we can still define

$$Z(G, H) = \sum_{\varphi: V_G \rightarrow V_H} \prod_{(u,v) \in E_G} \mathbb{I}_{E_H}(\varphi(u), \varphi(v)) \prod_{w \in V_G} \nu(\varphi(w)).$$

In the language of statistical physics, $Z_b(G, H)$ and $Z(G, H)$ are *partition functions*.

Somewhat surprisingly, J. Kahn's result holds even in this general case, as shown by D. Galvin and P. Tetali [6].

Theorem 3.2. [6] *For any bipartite graph H equipped with the weight function $\nu : V_H \rightarrow \mathbb{R}_+$ and $\mathbb{I}_{E_H} : A_H \times B_H \rightarrow \{0, 1\}$, and for any d -regular simple graph G on n vertices,*

$$Z_b(G, H) \leq Z_b(K_{d,d}, H)^{n/(2d)}.$$

The key observation is that for a bipartite graph H equipped with the weight function $\nu : V_H \rightarrow \mathbb{R}_+$ and characteristic function $\mathbb{I}_{E_H} : A_H \times B_H \rightarrow \{0, 1\}$, we can define a weighted graph \tilde{H} with weight function $\tilde{\nu}$ and characteristic function $\mathbb{I}_{E_{\tilde{H}}}$ such that

$$(3.1) \quad Z(G, \tilde{H}) = Z_b(G', H),$$

for any graph G (where G' is the modification of G defined in the previous section). Indeed, construct \tilde{H} with vertex set $A_H \times B_H$, edges

$$\mathbb{I}_{E_{\tilde{H}}}((a_1, b_1), (a_2, b_2)) = \mathbb{I}_{E_H}(a_1, b_2) \mathbb{I}_{E_H}(a_2, b_1),$$

and weight function

$$\tilde{\nu}(a, b) = \nu(a) \nu(b) \mathbb{I}_{E_H}(a, b).$$

In effect, the vertex set of \tilde{H} is only the edges of H (since non-edge pairs are given weight 0). Now, for a map $\varphi : G' \rightarrow H$, we can consider the map $\tilde{\varphi} : G \rightarrow \tilde{H}$ given by

$$\tilde{\varphi}(u) = (\varphi((u, 0)), \varphi((u, 1))).$$

By the construction of the graphs G' and \tilde{H} , the contribution of φ to $Z_b(G, H)$ is the same as the contribution of $\tilde{\varphi}$ to $Z(G, \tilde{H})$, and the result (3.1) follows.

Finally, applying Theorem 3.2 to the $(d+1)$ -regular graph G' yields

$$Z(G, \tilde{H}) = Z_b(G', H) \leq Z_b(K_{d,d}, H)^{2n/(2(d+1))} = Z(K_{d+1}, \tilde{H})^{n/(d+1)}.$$

Hence we have proved the following theorem.

Theorem 3.3. *For a bipartite graph $H = (A, B, E)$ with vertex weight function $\nu : V_H \rightarrow \mathbb{R}_+$ let \tilde{H} be the following weighted graph: its vertex set is $E(H)$, its edge set is defined by $((a_1, b_1), (a_2, b_2)) \in E(\tilde{H})$ if and only if $(a_1, b_2) \in E(H)$ and $(a_2, b_1) \in E(H)$, and the weight function on the vertex set is $\tilde{\nu}(a, b) = \nu(a) \nu(b)$ for $(a, b) \in E(H)$. Then for any d -regular simple graph G on n vertices we have*

$$Z(G, \tilde{H}) \leq Z(K_{d+1}, \tilde{H})^{n/(d+1)}.$$

We can obtain Conjecture 3 of [2] as a corollary by applying this theorem in the case where H is the path on 4 vertices, $a_1 b_1 a_2 b_2$, with appropriate vertex weights. Indeed, if $\nu(a_1) = 1$, $\nu(b_1) = \lambda_b$, $\nu(a_2) = \frac{\lambda_w}{\lambda_b}$, $\nu(b_2) = \frac{\lambda_r \lambda_b}{\lambda_w}$ then \tilde{H} is precisely the

Widom-Rowlinson graph with vertex weights $\lambda_b, \lambda_r, \lambda_w$. This proves that even for the vertex-weighted Widom-Rowlinson graph we have

$$Z(G, H_{\text{WR}}) \leq Z(K_{d+1}, H_{\text{WR}})^{n/(d+1)}.$$

Hence we have proved the following theorem.

Theorem 3.4. *Let H_{WR} be the Widom-Rowlinson graph with vertex weights $\lambda_b, \lambda_w, \lambda_r$. Then for any d -regular simple graph G on n vertices we have*

$$Z(G, H_{\text{WR}}) \leq Z(K_{d+1}, H_{\text{WR}})^{n/(d+1)}.$$

Now let us consider the special case when H is unweighted ($\nu \equiv 1$). In this case $\tilde{\nu}$ is just \mathbb{I}_{E_H} , so we can think of \tilde{H} as an unweighted graph with vertex set $V_{\tilde{H}} = E_H$. There is an edge in \tilde{H} between edges $e = (a_1, b_1)$ and $f = (a_2, b_2)$ of H whenever (a_1, b_2) and (a_2, b_1) are both also edges of H . This is always the case when either $a_1 = a_2$ or $b_1 = b_2$, so in particular every edge $e \in E_H = V_{\tilde{H}}$ has a self-loop in \tilde{H} , and every pair of incident edges in H are adjacent in \tilde{H} . We also get an edge $(e, f) \in E_{\tilde{H}}$ if four vertices $a_1 b_1 a_2 b_2$ are all distinct and form a 4-cycle with e and f as opposite edges. In other words, \tilde{H} is precisely the extended line graph of H . Hence as a corollary of Theorem 3.3 we have proved Theorem 1.3.

If H does not contain any 4-cycle, then \tilde{H} is simply the line graph of H with loops at every vertex. In particular, if H is a path (or even cycle of length at least 6) then \tilde{H} is again a path (or even cycle of length at least 6), but now with a loop at every vertex. Letting H° denote the graph obtained by adding a loop at every vertex of the graph H , we can write the corollary

Corollary 3.5. *If $H = C_k^\circ$ (for $k \geq 6$ even) or if $H = P_k^\circ$ (for any k), then for any d -regular graph G*

$$p_H(G) \leq p_H(K_{d+1}).$$

It is a good question how to characterize all of the graphs \tilde{H} which can be obtained this way. Note that since \tilde{H} is always fully-looped, this class has no intersection with the class of graphs found by Galvin [4]: the set of graphs H_q^ℓ obtained from a complete looped graph on q vertices with $\ell \geq 1$ loops deleted.

Remark 3.6. Let S_k be the star on k vertices. One can show (for details see [4]) that, for large enough d ,

$$p_{S_k^\circ}(K_{d+1}) < p_{S_k^\circ}(K_{d,d})$$

for $k \geq 6$. From this example we can see that in order to have $p_H(G) \leq p_H(K_{d+1})$ it is not sufficient merely for H to have a loop at every vertex.

L. Sernau [9] introduced many ideas for extending certain inequalities to a larger class of graphs. For instance, recall that the $H_1 \times H_2$ has $V_{H_1 \times H_2} = V_{H_1} \times V_{H_2}$ and $((a_1, b_1), (a_2, b_2)) \in E_{H_1 \times H_2}$ if and only if $(a_1, a_2) \in E_{H_1}$ and $(b_1, b_2) \in E_{H_2}$. Sernau noted that if H_1 and H_2 are graphs such that

$$p_{H_i}(G) \leq p_{H_i}(K_{d+1}),$$

for $i = 1, 2$, then it is also true that

$$p_{H_1 \times H_2}(G) \leq p_{H_1 \times H_2}(K_{d+1}).$$

This inequality simply follows from the identity

$$\text{hom}(G, H_1 \times H_2) = \text{hom}(G, H_1) \text{hom}(G, H_2),$$

which is explained in [9]. Surprisingly, this observation does not allow us to extend our result to any new graphs, because the product of two extended line graphs is again an extended line graph:

$$\tilde{H}_1 \times \tilde{H}_2 = \tilde{H}_{12},$$

where $H_{12} = (A_{H_1} \times A_{H_2}, B_{H_1} \times B_{H_2}, E_{H_1} \times E_{H_2})$.

4. ON A THEOREM OF L. SERNAU

Theorem 3 of [9] also provides a class of graphs for which K_{d+1} is the maximizing graph. Below we explain the relationships between our results and his theorem.

Definition 4.1. Let H and A be graphs. Then the graph H^A is defined as follows: its vertices are the maps $f : V(A) \rightarrow V(H)$ and the $(f_1, f_2) \in E(H^A)$ if $(f_1(u), f_2(v)) \in E(H)$ whenever $(u, v) \in E(A)$.

Then Sernau proved the following theorem.

Theorem 4.2. [9] *Let G be a d -regular graph, and let $F = l(H^B)$, where H is an arbitrary graph, B is a bipartite graph, and $l(H^B)$ is the graph induced by the vertices of H^B which have a loop. Then*

$$p_F(G) \leq p_F(K_{d+1}).$$

When $H = H_{ind}$, $B = K_2$ then $l(H^B) = H_{WR}$ so this also proves the conjecture of D. Galvin. Note that when $B = K_2$ then $l(H^B)$ is the extended line graph of $H \times K_2$. It is not a great surprise that these results are similar, even the proofs behind these results are strongly related to each other.

5. CONJECTURES

Let H be a simple graph, i.e., with no multiple edges or loops. Let H^o denote the graph obtained by adding a loop at each vertex of H (so for instance C_n^o denotes the n -cycle with a loop at each vertex).

Conjecture 5.1. Let G be a d -regular simple graph. Then for any $n \geq 4$

$$p_{C_n^o}(G) \leq p_{C_n^o}(K_{d+1}).$$

Conjecture 5.2. Let G be a d -regular simple graph. Then for any $d \geq 4$

$$p_{S_4^o}(G) \leq p_{S_4^o}(K_{d+1}).$$

Furthermore, for $k \geq 6$

$$p_{S_k^o}(G) \leq p_{S_k^o}(K_{d,d}).$$

Finally, for an arbitrary graph H it is not clear how to characterize the maximizers over all d -regular graphs G of $p_H(G)$. If we restrict to bipartite G , however, D. Galvin and P. Tetali proved that $p_H(G) \leq p_H(K_{d,d})$ [6]. We conjecture that this can be extended to the class of triangle-free graphs.

Conjecture 5.3. Let G be a d -regular triangle-free graph. Then for any graph H we have

$$p_H(G) \leq p_H(K_{d,d}).$$

Acknowledgments. We thank David Galvin and Luke Sernau for helpful conversations. We are also grateful to the anonymous referees for their careful reading and useful suggestions on the paper.

REFERENCES

- [1] G. R. Brightwell, O. Häggström and P. Winkler: *Nonmonotonic behavior in hard-core and Widom-Rowlinson models*, Journal of Statistical Physics, **94** (1999), pp. 415–435
- [2] E. Cohen, W. Perkins and P. Tetali: *On the Widom-Rowlinson occupancy fraction in regular graphs*, to appear in Combinatorics, Probability and Computing
- [3] E. Davies, M. Jenssen, W. Perkins, and B. Roberts: *Independent sets, matchings, and occupancy fractions*, arXiv preprint arXiv:1508.04675 (2015)
- [4] D. Galvin: *Maximizing h -colorings of a regular graph*, Journal of Graph Theory, **73**(1) (2013), pp. 66–84
- [5] D. Galvin: *Three tutorial lectures on entropy and counting*, arXiv preprint arXiv:1406.7872, 2014
- [6] D. Galvin and P. Tetali: *On weighted graph homomorphisms*, DIMACS Series in Discrete Mathematics and Theoretical Computer Science, **63** (2004), pp. 97–104
- [7] J. Kahn: *An entropy approach to the hard-core model on bipartite graphs*, Combinatorics, Probability and Computing, **10** (3) (2001), pp. 219–237
- [8] J. H. Kim, C. Lee and J. Lee: *Two approaches to Sidorenko’s conjecture*, to appear in Trans. Amer. Math. Soc.
- [9] L. Sernau: *Graph operations and upper bounds on graph homomorphism counts*, Arxiv preprint 1510.01833
- [10] Y. Zhao: *The number of independent sets in a regular graph*, Combinatorics, Probability and Computing, **19** (2) (2010), pp. 315–320
- [11] Y. Zhao: *The bipartite swapping trick on graph homomorphisms*, SIAM Journal on Discrete Mathematics, **25** (2) (2011), pp. 660–680
- [12] B. Widom and J. S. Rowlinson: *New model for the study of liquid-vapor phase transition*, J. Chem. Phys. **52** (1970), 1670–1684

SCHOOL OF MATHEMATICS, GEORGIA INSTITUTE OF TECHNOLOGY, ATLANTA, GA 30332-0160

E-mail address: `ecohen32@math.gatech.edu`

MASSACHUSETTS INSTITUTE OF TECHNOLOGY, DEPARTMENT OF MATHEMATICS, CAMBRIDGE MA 02139 & MTA-ELTE GEOMETRIC AND ALGEBRAIC COMBINATORICS RESEARCH GROUP, H-1117 BUDAPEST, PÁZMÁNY PÉTER SÉTÁNY 1/C, HUNGARY

E-mail address: `peter.csikvari@gmail.com`

SCHOOL OF MATHEMATICS, UNIVERSITY OF BIRMINGHAM, UK

E-mail address: `william.perkins@gmail.com`

SCHOOL OF MATHEMATICS AND SCHOOL OF COMPUTER SCIENCE, GEORGIA INSTITUTE OF TECHNOLOGY, ATLANTA, GA 30332-0160

E-mail address: `tetali@math.gatech.edu`