

# IncP plasmid carrying the colistin resistance gene mcr-1 in *Klebsiella pneumoniae* from hospital sewage

Zhao, Feifei; Feng, Yu; Lü, Xiaoju; McNally, Alan; Zong, Zhiyong

DOI:

[10.1128/AAC.02229-16](https://doi.org/10.1128/AAC.02229-16)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Zhao, F, Feng, Y, Lü, X, McNally, A & Zong, Z 2017, 'IncP plasmid carrying the colistin resistance gene mcr-1 in *Klebsiella pneumoniae* from hospital sewage', *Antimicrobial Agents and Chemotherapy*, vol. 61, e02229. <https://doi.org/10.1128/AAC.02229-16>

[Link to publication on Research at Birmingham portal](#)

## General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

**An IncP plasmid carrying the colistin resistance gene *mcr-1*  
in *Klebsiella pneumoniae* from hospital sewage**

**Feifei Zhao<sup>1,2</sup>, Yu Feng<sup>1,2</sup>, Xiaoju Lü<sup>1,2</sup>, Alan McNally<sup>4</sup>, Zhiyong Zong<sup>1, 2, 3§</sup>**

<sup>1</sup>*Center of Infectious Diseases, West China Hospital, Sichuan University, Chengdu, China.*

<sup>2</sup>*Division of Infectious Diseases, State Key Laboratory of Biotherapy, Chengdu, China.*

<sup>3</sup>*Department of Infection Control, West China Hospital, Sichuan University, Chengdu, China.*

<sup>4</sup>*Pathogen Research Group, Nottingham Trent University, Nottingham, UK.*

**Running title: *mcr-1* in sewage**

**Keywords:** colistin resistance; *mcr-1*; sewage; *Klebsiella pneumoniae*

§Corresponding author. Mailing address: Center of Infectious Diseases, West China Hospital (Huaxi), Guoxuexiang 37, Chengdu 610041, China. Phone: 86-28-8542-2637. Fax: 86-28-8542-3212. E-mail: [zongzhiy@scu.edu.cn](mailto:zongzhiy@scu.edu.cn)

20   **Abstract**

21    A *Klebsiella pneumoniae* strain of ST313 recovered from hospital sewage was found  
22    to carry the plasmid-borne colistin resistance gene *mcr-1*, which was bracketed by two  
23    copies of the insertion sequence IS*Apl1* on a 57-kb self-transmissible IncP type  
24    plasmid of a new IncP-1 clade. The carriage of *mcr-1* on a self-transmissible  
25    broad-host-range plasmid highlights that *mcr-1* has the potential to be spread beyond  
26    the *Enterobacteriaceae*.

Colistin is the last resort antimicrobial agents to treat the infections caused by many Gram-negative bacteria. Recently, a plasmid-borne colistin resistance gene, *mcr-I*, has been found in *Escherichia coli* and *Klebsiella pneumonia* from human and animals in China (1). A few follow-up studies have found *mcr-I*-carrying *E. coli* in many countries in Africa (2), Europe (3-8), Asia (1-3, 9, 10) and North America (9, 11). In addition, *mcr-I* has also been found in *Kluyvera ascorbata* (12) and several species of *Salmonella* (13-15). All of above findings suggest that *mcr-I* has been widely distributed and imposes an emerging threat for clinical management and public and animal health. During a screening study for the presence of colistin-resistant *Enterobacteriaceae* in hospital sewage, we found a *mcr-I*-carrying *K. pneumonia*, which is reported here.

*K. pneumoniae* strain WCHKP1511 was recovered from the influx mainstream of hospital sewage at West China Hospital, Chengdu, western China, in November 2015. Strain WCHKP1511 grew on CHROMAgar Orientation agar plates (CHROMAgar, Paris, France) containing 4 µg/ml colistin and 64 µg/ml linezolid. Species identification was established by partially sequencing the *gyrB* gene (16). Strain WCHKP1511 was resistant to colistin (MIC, 8 µg/ml), polymyxin B (MIC, 8 µg/ml), chloramphenicol (MIC, 128 µg/ml) and tetracycline (MIC, 64 µg/ml) but was susceptible to amikacin (MIC, 0.5 µg/ml), ceftazidime (MIC, ≤0.5 µg/ml), cefotaxime (MIC, 0.03 µg/ml), ciprofloxacin (MIC, 0.03 µg/ml), imipenem (MIC, 0.125 µg/ml)

and tigecycline (MIC, 1 µg/ml) as determined using the microdilution broth method following recommendations of the Clinical Laboratory Standards Institute (CLSI) (17). WCHKP1511 was susceptible to fosfomycin (MIC, 32 µg/ml) as determined using the agar dilution method following recommendations of CLSI (17). In addition, strain WCHKP1511 was resistant to ampicillin and trimethoprim-sulphamethoxazole, intermediate to amoxicillin-clavuanate, gentamicin, tobramycin and nitrofurantoin, and susceptible to aztreonam, cefazolin, cefepime, cefoxitin, ceftriaxone, ertapenem, levofloxacin and piperacillin-tazobactam as determined using the Vitek II automated system (bioMerieux, Lyon, France). Breakpoints defined by FDA and by EUCAST were used for tigecycline and colistin, respectively; otherwise, those defined by CLSI were applied. Strain WCHKP1511 was found to carry *mcr-1* by PCR and sequencing (1, 12). Although *mcr-1* has been widely found in *E. coli*, *mcr-1*-carrying *K. pneumoniae* remains uncommon and had only been found in eastern China (Jiangsu and Zhejiang provinces) before (1, 18).

Strain WCHKP1511 was subjected to the 150-bp paired-end whole genome sequencing with a ca. 200× coverage using the HiSeq 2500 Sequencer (Illumina, San Diego, CA, USA). A total of 3,784,972 reads and 567,745,800 clean bases were generated, which were assembled into 275 contigs (230 contigs ≥ 1,000 bp) in length (N50, 67,217 bp) with a 57.46% GC content using the Spades program (19).

69 WCHKP1511 belonged to ST313, which was determined by using the genomic  
70 sequence to query the multi-locus sequence typing database of *K. pneumoniae*  
71 (<http://bigsdh.web.pasteur.fr/klebsiella/klebsiella.html>). In the *K. pneumoniae* MLST  
72 database, only one ST313 strain, KML 2185, which was recovered from human blood  
73 in the Netherlands in 2007, has been deposited. *K. pneumoniae* strains carrying *mcr-1*  
74 in Jiangsu province belonged to ST25 (20), while the STs of those in Zhejiang  
75 province were unknown. ST313 was not closely related to ST25, as only 2 out of 7  
76 alleles were identical between the two STs.

77

78 Antimicrobial resistance genes were predicted using ResFinder from the Center for  
79 Genomic Epidemiology (<http://genomicepidemiology.org/>). In addition to *mcr-1*,  
80 WCHKP1511 had multiple genes mediating resistance to aminoglycosides  
81 (*aac(3)-Iva*, *aadA2*, *aph(3')-Ia* and *aph(4)-I*),  $\beta$ -lactams (*bla*<sub>TEM-135</sub> and a new *bla*<sub>SHV</sub>  
82 variant), chloramphenicol (*floR*), fosfomycin (*fosA*), quinolones (*oqxA* and *oqxB*),  
83 sulphonamides (*sul2*), trimethoprim (*dfrA12*) and tetracycline (*tet(A)*) as predicted by  
84 ResFinder. *bla*<sub>TEM-135</sub> encodes a broad-spectrum  $\beta$ -lactamase (21). Of note, although  
85 strain WCHKP1511 carried *fosA*, it was susceptible to fosfomycin, which warrants  
86 further investigation. *bla*<sub>SHV</sub> in strain WCHKP1511 is a new variant, which encodes a  
87 SHV enzyme with an amino acid difference (Thr14Asn, the position is based on the  
88 ATG start condon) from SHV-111, the closest match. The new SHV has been assigned  
89 SHV-195 by the NCBI  $\beta$ -lactamase classification system

([www.ncbi.nlm.nih.gov/pathogens/submit\\_beta\\_lactamase/](http://www.ncbi.nlm.nih.gov/pathogens/submit_beta_lactamase/)). As strain WCHKP1511 was susceptible to third generation cephalosporins, *bla*<sub>SHV-195</sub> is unlikely to encode an extended-spectrum  $\beta$ -lactamase (ESBL). The complete coding sequence of *bla*<sub>SHV-195</sub> was cloned onto the pBC SK vector (Agilent, Santa Clara, CA, USA), which was electroporated into *E. coli* DH5 $\alpha$ . *E. coli* transformants containing *bla*<sub>SHV-195</sub> were resistant to ampicillin (MIC, >256  $\mu$ g/mL) and cephalothin (32  $\mu$ g/mL) but were susceptible to aztreonam, ceftazidime, cefotaxime, cefoxitin and imipenem determined using the broth microdilution method (17). This confirmed that *bla*<sub>SHV-195</sub> encodes a broad-spectrum rather than an ESBL.

In addition, there was a predicted bleomycin-resistance gene, designated ORFble here, which was not identified by Resfinder but was identified by the Prokka annotation tool (22) and was confirmed by Protein-BLAST in strain WCHKP1511. The complete coding sequence of ORFble was amplified with primers 1511\_ble\_BamHI\_F (CGCGGATCCTTGGTTCACCATGAAGATG)/1511\_ble\_EcoRI\_R (CCGGAATTCCGGCCGATTGCTGAACAGATTA), was cloned onto pBC SK and was electroporated into *E. coli* DH5 $\alpha$ . ORFble-containing transformants were selected on LB agar plates containing 25  $\mu$ g/mL chloramphenicol and the presence of ORFble in transformants was confirmed by PCR and sequencing. However, MIC (0.25  $\mu$ g/mL) of zeocin (Thermo Fisher Scientific, Waltham, MA, USA), a bleomycin, against *E. coli* DH5 $\alpha$  transformant containing ORFble was the same as that against DH5 $\alpha$  as

determined using the broth microdilution method (17). This suggests that ORFble did not mediate resistance to bleomycin and its function remains undetermined.

Conjugation experiments were carried out in broth using azide-resistant *E. coli* strain J53 as the recipient and 2 µg/ml colistin plus 150 µg/ml sodium azide was used for selecting transconjugants. The presence of *mcr-I* in transconjugants was confirmed using PCR. In strain WCHKP1511, *mcr-I* could be transferred to *E. coli* J53 at a frequency of  $10^{-2}$  cells per donor cell by mating, suggesting that *mcr-I* was carried by a self-transmissible plasmid, which was assigned pMCR\_1511. The sequence of pMCR\_1511 was completely circularised with gaps between contigs closed by Sanger sequencing of amplicons from PCRs using primers designed based on available contig sequences. pMCR\_KP1511 was 57,278 bp in size and had no known antimicrobial resistance genes other than *mcr-I*. Unlikely the previously-described *mcr-I*-carrying IncI2 plasmid pHNSHP45 (GenBank accession number KP347127) (1), pMCR\_KP1511 belonged to the IncP type, a broad-host-range incompatibility group. Plasmid pKH-457-3-BE carrying *mcr-I* in *E. coli* from Belgium was found to have an IncP backbone (3). However, the sequence of pKH-457-3-BE was not available for further analysis. Nonetheless, it has been suggested that pKH-457-3-BE was 99% similarity and 73% coverage with the IncHI2 plasmid pHXY0908 (GenBank accession number KM877269) in *Salmonella enterica* serotype Typhimurium (8). pMCR\_KP1511 had only a 6% coverage with pHXY0908, indicating that



pMCR\_1511 was very different from pKH-457-3-BE and pKH-457-3-BE may not be a true IncP plasmid but is likely of IncHI2.

pMCR\_1511 has the typical IncP-1 plasmid backbone (23) containing the *trfA* encoding the replication initiation protein, two *par* modules for plasmid partitioning, two conjugative regions *tra* (17.6 kb) and *trb* (12.7 kb), the host-lethal protein-encoding *kil* genes and their regulator *kor* (stands for *kil*-*override*) genes, and a toxin-antitoxin *higA-B* system (Figure 1). The backbone of pMCR\_1511 was highly similar (99% identity) with that on plasmid pHNFP671 (GenBank accession number KP324830), which was an IncP plasmid in *E. coli* isolate FP671 from Guangzhou, China but did not carry *mcr-I*. IncP-1 plasmid has six assigned clades, i.e.,  $\alpha$ ,  $\beta$ ,  $\gamma$ ,  $\delta$ ,  $\epsilon$  and  $\zeta$ , among which  $\beta$  clade has  $\beta 1$  and  $\beta 2$  two subclades (24). To determine of which clade pMCR\_1511 was, the sequence of 30 genes belonging to the IncP-1 backbone was retrieved from pMCR\_1511, concatenated and then aligned to the counterparts of one representative plasmid of each clade including  $\beta 1$  and  $\beta 2$  subclades as described previously (25). Phylogenetic analysis of the IncP-1 plasmid backbone revealed that pMCR\_1511 and pHNFP671 belonged to a new IncP-1 clade (Figure S1 in the supplementary file).

When compared the sequence of pHNFP671, pMCR\_1511 has two unique regions (Figure 1). One of the unique region harbored *mcr-I* and the other contained ORFble.

153 The sequence comparison of pMCR\_1511 and pHNFP671 allowed us to analyze the  
154 genetic context of *mcr-1* in detail. Like most genetic contexts available in the  
155 GenBank, *mcr-1* was located downstream of the insertion sequence IS*Apl1* on  
156 pMCR\_1511. However, there was another IS*Apl1*, which was interrupted (see below),  
157 downstream and therefore *mcr-1* was bracketed by two copies of IS*Apl1* on  
158 pMCR\_1511. It has been known that IS*Apl1* is able to generate 2-bp direct target  
159 repeats (DR) upon insertion (26)  
160 ([https://www-is.biotoul.fr/index.html?is\\_special\\_name=ISAp11](https://www-is.biotoul.fr/index.html?is_special_name=ISAp11)). The 2-bp flanking  
161 sequences of the region bracketed by the two copies of IS*Apl1* were identical (AC,  
162 Figure 2). When the region formed by two copies of IS*Apl1* and one of the 2-bp  
163 flanking sequences were subtracted artificially, the joined sequence perfectly matched  
164 that of an open reading frame (orf) with unknown function on plasmid pHNFP671. It  
165 therefore proved that the 2-bp sequence was truly DR generated by IS*Apl1* rather than  
166 coincidence and the two IS*Apl1* formed a composite transposon to mobilize *mcr-1*  
167 gene. A very recent analysis revealed that the IS*Apl1*-formed composite transposon  
168 carrying *mcr-1* has also been seen on either the chromosome or a plasmid (IncH or  
169 unknown Inc groups) of seven *E. coli* strains (27). The IS*Apl1*-formed composite  
170 transposon carrying *mcr-1* is all located at different locations, which are also different  
171 from the location on pMCR\_1511, in the seven *E. coli* strains and is flanked by 2-bp  
172 DR in five strains (27). The previous analysis (27) and the findings in the present

study suggest that the IS*AplI*-formed composite transposon is a common vehicle for mediating the spread of *mcr-1*.

The IS*AplI* downstream of *mcr-1* on pMCR\_1511 was interrupted by the insertion of Tn3 with the characteristic 5-bp DR (Figure 2). The Tn3 was also disrupted by IS26 and most of the Tn3 was absent, which may be due to the action of IS26. It is well known that the insertion of IS26 can lead to the deletion of the adjacent sequence of the insertion site (28). Alternative explanation for the absence of most part of Tn3 is that the insertion of the second IS26 and the recombination between the two IS26 could lead to the loss of the intervening region. Although IS*AplI* downstream of *mcr-1* was interrupted by Tn3, the right-end inverted repeat (IRR) of the IS*AplI* remained intact (Figure 2). The transposase encoded by the IS*AplI* upstream of *mcr-1* had the potential to recognize the IRR of the IS*AplI* downstream of *mcr-1* and then could realize the mobilization of the region bracketed by the two copies of the IS*AplI*.

In conclusion, the plasmid-borne colistin resistance gene *mcr-1* was found in a *K. pneumoniae* of an infrequently encountered ST from hospital sewage. *mcr-1* was carried by a self-transmissible IncP plasmid, which is a broad-host-range type of plasmids and has the potential to mediate the dissemination of *mcr-1* from the *Enterobacteriaceae* to other Gram-negative bacteria such as *Pseudomonas aeruginosa*. *mcr-1* was bracketed by two copies of IS*AplI*, which were able to form a composite

transposon and represented a common mechanism for mediating the mobilization of *mcr-1*.

**Nucleotide sequence accession number.** Reads and the Whole Genome Shotgun Sequencing project of *K. pneumoniae* strain WCHKP1511 have been deposited into DDBJ/EMBL/GenBank under accession number SRR3170679 and LSMF000000000, respectively. The sequence of pMCR\_1511 has been deposited into DDBJ/EMBL/GenBank under accession number KX377410.

## Acknowledgements

The work was supported by a grant from the National Natural Science Foundation of China (project no. 81572030) and a joint grant from the National Natural Science Foundation of China (project no. 8151101182) and the Newton Advanced Fellowship, Royal Society, UK.

## References

1. Liu YY, Wang Y, Walsh TR, Yi LX, Zhang R, Spencer J, Doi Y, Tian G, Dong B, Huang X, Yu LF, Gu D, Ren H, Chen X, Lv L, He D, Zhou H, Liang Z, Liu JH, Shen J. 2015. Emergence of plasmid-mediated colistin resistance mechanism MCR-1 in animals and human beings in China: a microbiological and molecular biological study. *Lancet Infect Dis* **16**:161-168.
2. Olaitan AO, Chabou S, Okdah L, Morand S, Rolain J-M. 2016. Dissemination of the *mcr-1* colistin resistance gene. *Lancet Infect Dis* **16**:147.
3. Malhotra-Kumar S, Xavier BB, Das AJ, Lammens C, Butaye P, Goossens H. 2016. Colistin resistance gene *mcr-1* harboured on a multidrug resistant plasmid. *Lancet Infect Dis* **16**:283-284.

- 220 4. **Hasman H, Hammerum AM, Hansen F, Hendriksen RS, Olesen B, Agerso**  
221 **Y, Zankari E, Leekitcharoenphon P, Stegger M, Kaas RS, Cavaco LM,**  
222 **Hansen DS, Aarestrup FM, Skov RL.** 2015. Detection of *mcr-1* encoding  
223 plasmid-mediated colistin-resistant *Escherichia coli* isolates from human  
224 bloodstream infection and imported chicken meat, Denmark 2015. Euro  
225 Surveill **20**:doi: 10.2807/1560-7917.ES.2015.2820.2849.30085.
- 226 5. **Haenni M, Poirel L, Kieffer N, Châtre P, Saras E, Métayer V, Dumoulin R,**  
227 **Nordmann P, Madec J-Y.** 2016. Co-occurrence of extended spectrum  $\beta$   
228 lactamase and MCR-1 encoding genes on plasmids. Lancet Infect Dis  
229 **16**:281-282.
- 230 6. **Falgenhauer L, Waezsada S-E, Yao Y, Imirzalioglu C, Käsbohrer A,**  
231 **Roesler U, Michael GB, Schwarz S, Werner G, Kreienbrock L,**  
232 **Chakraborty T.** 2016. Colistin resistance gene *mcr-1* in extended-spectrum  
233  $\beta$ -lactamase-producing and carbapenemase-producing Gram-negative bacteria  
234 in Germany. Lancet Infect Dis **16**:282-283.
- 235 7. **Arcilla MS, van Hattem JM, Matamoros S, Melles DC, Penders J, de Jong**  
236 **MD, Schultz C.** 2016. Dissemination of the *mcr-1* colistin resistance gene.  
237 Lancet Infect Dis **16**:147-149.
- 238 8. **Poirel L, Kieffer N, Liassine N, Thanh D, Nordmann P.** 2016.  
239 Plasmid-mediated carbapenem and colistin resistance in a clinical isolate of  
240 *Escherichia coli*. Lancet Infect Dis **16**:281.
- 241 9. **Stoesser N, Mathers AJ, Moore CE, Day NPJ, Crook DW.** 2016. Colistin  
242 resistance gene *mcr-1* and pHNSHP45 plasmid in human isolates of  
243 *Escherichia coli* and *Klebsiella pneumoniae*. Lancet Infect Dis **16**:285-286.
- 244 10. **Suzuki S, Ohnishi M, Kawanishi M, Akiba M, Kuroda M.** 2016.  
245 Investigation of a plasmid genome database for colistin-resistance gene *mcr-1*.  
246 Lancet Infect Dis **16**:284-285.
- 247 11. **Mulvey MR, Mataseje LF, Robertson J, Nash JHE, Boerlin P, Toye B,**  
248 **Irwin R, Melano RG.** 2016. Dissemination of the *mcr-1* colistin resistance  
249 gene. Lancet Infect Dis **16**:289-290.
- 250 12. **Zhao F, Zong Z.** 26 September 2016. *Kluyvera ascorbata* carrying the *mcr-1*  
251 colistin resistance gene from hospital sewage. Antimicrob Agents Chemother  
252 10.1128/AAC.01165-16.
- 253 13. **Webb HE, Granier SA, Marault M, Millemann Y, den Bakker HC,**  
254 **Nightingale KK, Bugarel M, Ison SA, Scott HM, Loneragan GH.** 2016.  
255 Dissemination of the *mcr-1* colistin resistance gene. Lancet Infect Dis  
256 **16**:144-145.
- 257 14. **Hu Y, Liu F, Lin IYC, Gao GF, Zhu B.** 2016. Dissemination of the *mcr-1*  
258 colistin resistance gene. Lancet Infect Dis **16**:146-147.
- 259 15. **Tse H, Yuen K-Y.** 2016. Dissemination of the *mcr-1* colistin resistance gene.  
260 Lancet Infect Dis **16**:145-146.
- 261 16. **Yamamoto S, Harayama S.** 1995. PCR amplification and direct sequencing

- of *gyrB* genes with universal primers and their application to the detection and taxonomic analysis of *Pseudomonas putida* strains. *Appl Environ Microbiol* **61**:1104-1109.
17. **CLSI**. 2013. Performance Standards for Antimicrobial Susceptibility Testing; Twenty-Third Informational Supplement. M100-S23. Clinical and Laboratory Standards Institute, Wayne, PA, USA.
  18. **Du H, Chen L, Tang Y-W, Kreiswirth BN**. 2016. Emergence of the *mcr-1* colistin resistance gene in carbapenem-resistant *Enterobacteriaceae*. *Lancet Infect Dis* **16**:287-288.
  19. **Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA**. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* **19**:455-477.
  20. **Li A, Yang Y, Miao M, Chavda KD, Mediavilla JR, Xie X, Feng P, Tang YW, Kreiswirth BN, Chen L, Du H**. 2016. Complete sequences of *mcr-1*-harboring plasmids from extended spectrum  $\beta$ -lactamase (ESBL)- and carbapenemase-producing *Enterobacteriaceae* (CPE). *Antimicrob Agents Chemother* **60**:4351-4354.
  21. **Pasquali F, Kehrenberg C, Manfreda G, Schwarz S**. 2005. Physical linkage of Tn3 and part of Tn1721 in a tetracycline and ampicillin resistance plasmid from *Salmonella Typhimurium*. *J Antimicrob Chemother* **55**:562-565.
  22. **Seemann T**. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* **30**:2068-2069.
  23. **Pansegrau W, Lanka E, Barth PT, Figurski DH, Guiney DG, Haas D, Helinski DR, Schwab H, Stanisich VA, Thomas CM**. 1994. Complete nucleotide sequence of Birmingham IncP alpha plasmids. Compilation and comparative analysis. *J Mol Biol* **239**:623-663.
  24. **Sen D, Brown CJ, Top EM, Sullivan J**. 2013. Inferring the evolutionary history of IncP-1 plasmids despite incongruence among backbone gene trees. *Mol Biol Evol* **30**:154-166.
  25. **Norberg P, Bergstrom M, Jethava V, Dubhashi D, Hermansson M**. 2011. The IncP-1 plasmid backbone adapts to different host bacterial species and evolves through homologous recombination. *Nat Commun* **2**:268.
  26. **Tegetmeyer HE, Jones SC, Langford PR, Baltes N**. 2008. IS*AplI*, a novel insertion element of *Actinobacillus pleuropneumoniae*, prevents ApxIV-based serological detection of serotype 7 strain AP76. *Vet Microbiol* **128**:342-353.
  27. **Snesrud E, He S, Chandler M, Dekker JP, Hickman AB, McGann P, Dyda F**. 12 September 2016. A Model for Transposition of the Colistin Resistance Gene *mcr-1* by IS*AplI*. *Antimicrob Agents Chemother* doi: 10.1128/AAC.01457-16.
  28. **Mollet B, Iida S, Arber W**. 1985. Gene organization and target specificity of

304 the prokaryotic mobile genetic element IS26. Mol Gen Genet **201**:198-203.

305

306

## Figure legends

**Figure 1.** Genetic structure of IncP plasmid pMCR\_1511 carrying *mcr-I* and the comparison with IncP plasmid pHNFP671. Regions and genes that are indicated are ORFble (a predicted [but actually not] bleomycin resistance gene), *higA-B* (encoding a toxin/antitoxin system), *mcr-I*, *kor-par-kli* (for plasmid maintenance), *tra* and *trb* (the two conjugation-encoding regions), and *trfA* (encoding the plasmid replication initiation protein). Backbones of the two IncP plasmids are almost identical. Compared to pHNFP671, pMCR\_1511 carried two additional regions that contained either *mcr-I* or ORFble.

**Figure 2.** Genetic context of *mcr-I* on pMCR\_1511. Genetic context of *mcr-I* on the IncI2 plasmid pHNSHP45 (GenBank accession number KP347127) and the corresponding region on the IncP plasmid pHNFP671 (GenBank accession number KP324830) are shown for comparison. orfs that encode hypothetical proteins with unknown function are indicated in white except that the one disrupted by the IS*AplI*-formed composite transposon on pMCR\_1511 is shown in black ( $\Delta 1$  and  $\Delta 2$ ). Other genes shown are *nikB* (encoding relaxase of the plasmid), *ydgA* (DNA topoisomerase III), *ydfA* (transcriptional regulator), *parA* (resolvase), *bla*<sub>TEM</sub> (shown as a white arrow in Tn3) and *traB* (conjugative protein). The 2-bp direct repeat (GA) abutting the IS*AplI*-*mcr-I*-*pho* region on pHNSHP45 and the 2-bp direct repeat (AC) abutting the IS*AplI*-formed composite transposon on pMCR\_1511 are shown. On



328 pMCR\_1511, the IS*AplI* downstream of *mcr-1* was interrupted by Tn3, which was  
329 interrupted by IS26.