

Catching ghosts with a coarse net

Petrovskaya, Natalia; Petrovskii, Sergei

DOI:

[10.1098/rsif.2016.0855](https://doi.org/10.1098/rsif.2016.0855)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Petrovskaya, N & Petrovskii, S 2017, 'Catching ghosts with a coarse net: use and abuse of spatial sampling data in detecting synchronization', *Journal of The Royal Society Interface*, vol. 14, no. 127, 20160855.
<https://doi.org/10.1098/rsif.2016.0855>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Final Version of Record available at: <http://dx.doi.org/10.1098/rsif.2016.0855>

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Catching Ghosts with a Coarse Net: Use and Abuse of Spatial Sampling Data in Detecting Synchronization

Natalia Petrovskaya^a and Sergei Petrovskii^{b1}

^a School of Mathematics, University of Birmingham,
Birmingham, B15 2TT, U.K.
Email: n.b.petrovskaya@bham.ac.uk

^b Department of Mathematics, University of Leicester,
Leicester, LE1 7RH, U.K.
Email: sp237@le.ac.uk

Abstract

Synchronization of population dynamics in different habitats is a frequently observed phenomenon. A common mathematical tool to reveal synchronization is the (cross)correlation coefficient between time courses of values of the population size of a given species where the population size is evaluated from spatial sampling data. The corresponding sampling net or grid is often coarse, i.e. it does not resolve all details of the spatial configuration, and the evaluation error – i.e. the difference between the true value of the population size and its estimated value – can be considerable. We show that this estimation error can make the value of the correlation coefficient very inaccurate or even irrelevant. We consider several population models to show that the value of the correlation coefficient calculated on a coarse sampling grid rarely exceeds 0.5, even if the true value is close to 1, so that the synchronization is effectively lost. We also observe ‘ghost synchronization’ when the correlation coefficient calculated on a coarse sampling grid is close to 1 but in reality the dynamics are not correlated. Finally, we suggest a simple test to check the sampling grid coarseness and hence to distinguish between the true and artifactual values of the correlation coefficient.

Keywords: sparse data; sampling; coarse grid; data analysis; correlation coefficient; ghost synchronization

¹Corresponding author.

1 Introduction

Evaluation of properties of a spatially extended system from sparse spatial data is an inherent problem in many applications across science and engineering [9, 13, 54]. The term “sparse data” usually refers to a situation where the information about a spatial distribution of a certain quantity (e.g. the concentration of a chemical substance in the environment) is available only at nodes of a certain discrete grid or net, and the number of grid nodes is not large enough to resolve the details of the heterogeneous distribution of the substance and/or the spatial configuration of the system. Often sparse spatial data are used to evaluate an integral property of the system – say, a ‘total mass’, the exact meaning of which depends on particular application. Here we mention just a few examples: in water quality engineering, it is the total amount of a harmful chemical substance occasionally released into the environment [9]; in geology, it is the total stock of a valuable mineral [13, 62]; in medicine, it is the total mass of a tumor [21]; in ecology, it is the population size of a dangerous pest [45, 54, 56]. A generic problem for the above examples, as well as for other similar situations where researchers or engineers have to deal with sparse data, is that, due to the apparent loss of information, the evaluation of the total mass can produce a result of unacceptably low accuracy. In its turn, that can lead to wrong decisions and dramatic consequences, e.g. a large reserve of a valuable mineral is overlooked, a dangerous tumor is not treated timely, etc.

In this paper, we consider the problem of sparse data in the context of ecological applications. Information about population abundance is routinely used in ecology for many different purposes such as to assess the ecosystem state and biodiversity [4, 35], to identify and monitor endangered species [20, 55], to monitor pests [10], to account for some important properties of population dynamics (e.g. spatial patterning [1, 11] or synchronization [7, 24, 37, 57, 61]), etc. While theoretical studies usually assume that this information – e.g. as quantified by the population size or the population density – is readily available with any required precision, estimation of population abundance in empirical studies is rarely straightforward. For instance, evaluation of the population size by direct counting is only possible for some species and only in relatively small habitats. Much more often, collection of relevant field data involve a certain sampling procedure [54]. Samples are taken across space with the intention to provide an estimate of the population density of a given species at some particular locations inside a given area or habitat [36, 56]; see Fig. 1. The accuracy of the population abundance evaluation depends on the accuracy of the local population density estimate at the location of a given sample and on the sample size (i.e. the total number of samples in a given census). It can also depend on the way how the local data are pulled together or ‘integrated’ to produce an estimate of the population size [41, 43].

We mention here that the accuracy of the local data depends on the nature of the samples which, in its turn, depends on the species traits. For instance, for soil-dwelling insects, sampling is often done by taking soil cores [10, 31, 56]. The insect count in a given soil core gives an almost perfect estimate of the local population density. For flying insects or insects walking/crawling on the surface, their census is usually done by installing traps and subsequently analyzing trap counts, a procedure that inevitably introduces a certain error [45, 46]. The precision of the local population density estimation and the number of samples collected in a population census, albeit not being




Figure 1: (left) A sketch of sampling on a coarse grid along a 1D transect. The dashed curve shows a hypothetical population distribution which is only known at the location r_1, \dots, r_N of samples. (right) Population density $u(x, y)$ of a carabid beetle species in a farm field obtained from data collected on a fine grid of 16×16 traps (data from [1]). The large black dots show the location of the nodes of a hypothetical coarse sampling grid of 3×3 nodes. Apparently, in both cases shown in the left and right panels, the sampling data obtained on a coarse grid would miss some important details of the population distributions.

entirely independent², are affected by largely different factors. Since in this paper our main goal is to understand how the reliability of conclusions about population dynamics (in particular, synchronization) can be affected by the samples size, we assume that local data are precise. The only source of the evaluation error is then the coarseness of the sampling grid.

In sampling procedures, both the location of the samples and their total number (to which altogether we refer as a sampling grid) used in any one given census or population survey are chosen based on a variety of reasons. A closer look at the corresponding arguments reveals that, while in some cases they are theory-based (e.g. based on the analysis of variances [47, 59]) in many other cases the properties of the sampling grid are decided based on a rule of thumb [12]. The question hence arises as to when – i.e. under what conditions – the population size estimated from data collected over a given sampling grid is accurate enough to provide a reliable information for any conclusions about the system properties and/or dynamics. In our previous work [39, 41, 43, 45], we showed that the above question “when” has a relative rather than absolute meaning as the evaluation accuracy depends on the spatial pattern of the population distribution. The same sampling grid may provide a very accurate estimate of the population size for one spatial population distribution but can lead to a completely wrong result for another distribution.

The main goal of this study is to consider how the quality of sampling data affects the conclusions on the presence or absence of synchronization between population fluctuations in different habitats. Synchronization is frequently reported in the literature [7, 24, 28, 37, 57, 61]; however,

²e.g. budget constraints can make it difficult to combine high-quality (and hence more expensive) local sampling with large number of samples.

technical details of the data collection, such as the properties of the sampling grid are often omitted. Also, the effect of the spatial pattern in the distribution of the corresponding populations usually remains obscure. Therefore, at least in some cases, the question may arise as to whether the observed synchronization is really as strong as it is reported. We will focus on the rather common case where the sampling grid is coarse, i.e. where the total number of samples taken in a given area is not large. We will show that sampling over a coarse sampling grid can lead to wrong conclusions, because synchronization remains undetected. We will also show that sampling over a coarse grid can result in a ‘ghost synchronization’ where synchronization is seen in the data whilst in reality the monitored populations are not synchronized.

2 Mathematical framework

Consider a generic case where a certain population described by its population density $u(\mathbf{r}, t)$ is sampled in a given domain A . The domain may be the species habitat or it may be defined by some external factors or tasks³. Although the results of our analysis are going to be rather general and not restricted to any particular species or taxa, for the convenience of interpretation we will mostly talk about invertebrates, e.g. insects, worms or slugs. In case of an invertebrate population, its sampling is usually done by installing traps and then analyzing the trap counts [8, 15, 45] or by taking soil cores and counting the number of individuals in each core [10, 56]. For simplicity, we assume that all samples in the given census are taken over a sufficiently small period of time, so that the population density would not undergo any significant change during the census duration. In mathematical terms, this standard sampling procedure provides the information about the population density as a set of values $u(\mathbf{r}_1, t), u(\mathbf{r}_2, t), \dots, u(\mathbf{r}_N, t)$ where \mathbf{r}_i is the location of the i th sample, N is the number of samples in the given census and t is the timing of the census. This information is then processed to produce a certain index that quantifies the population abundance in the given area. Quite often, the sampling data are used to obtain an estimate of the population size ω_A or the average population density $\bar{u} = \omega_A/Z_A$ where Z_A is the area of the domain. There are several ways to calculate the average density [43]; in this study, we consider the method commonly used in empirical ecology [54, 56] to estimate the average by calculating the arithmetic mean of the sampled values:

$$\bar{u} \approx \frac{1}{N} \sum_{i=1}^N u(\mathbf{r}_i, t). \quad (1)$$

In case the census is done regularly, say at times t_1, t_2, \dots, t_k , application of (1) to the samples collected in each census results in a time series of the average population densities, i.e. $S = \{\bar{u}^{(1)}, \bar{u}^{(2)}, \dots, \bar{u}^{(k)}\}$. This time series can then be further analyzed depending on the purpose of the study. When the study focuses on synchronization, sampling procedure is applied to a number of domains/habitats to produce the corresponding number of different time series. For example, sampling in domains A and B would produce two time series, $S_A = \{\bar{u}_A^{(1)}, \bar{u}_A^{(2)}, \dots, \bar{u}_A^{(k)}\}$ and

³For instance, in the context of integrated pest management such a domain can be a farm field.

$S_B = \{\bar{u}_B^{(1)}, \bar{u}_B^{(2)}, \dots, \bar{u}_B^{(k)}\}$, respectively. The standard statistical tool to reveal synchronization is the (cross)correlation coefficient [28, 48, 49, 57]:

$$\rho_{AB}(k, N) = \frac{\sum_{i=1}^k (\bar{u}_A^{(i)} - \mu_A)(\bar{u}_B^{(i)} - \mu_B)}{\sqrt{\left(\sum_{i=1}^k (\bar{u}_A^{(i)} - \mu_A)^2\right) \left(\sum_{i=1}^k (\bar{u}_B^{(i)} - \mu_B)^2\right)}}, \quad (2)$$

where μ_A and μ_B are the sample means of the time series S_A and S_B :

$$\mu_A = \frac{1}{k} \sum_{i=1}^k \bar{u}_A^{(i)}, \quad \mu_B = \frac{1}{k} \sum_{i=1}^k \bar{u}_B^{(i)}. \quad (3)$$

It follows from (2) that $-1 \leq \rho \leq 1$ where $0 < \rho \leq 1$ corresponds to correlation and $-1 \leq \rho < 0$ to anti-correlation.

Depending on the calculated value of ρ , i.e. on the correlation strength, a conclusion can be made about the existence or absence of synchronization if ρ is close to one or close to zero, respectively. We mention here that the notion of ‘correlation strength’ used in ecological studies is somewhat conventional [22]: the correlation is regarded as very strong for $0.8 \leq \rho \leq 1$, strong for $0.6 \leq \rho < 0.8$, moderate for $0.4 \leq \rho < 0.6$, weak for $0.2 \leq \rho < 0.4$ and very weak for $0 \leq \rho < 0.2$. We will use this verbal description of the correlation strength in our analysis below. Apparently, there is considerable difference in the population dynamics, and hence different implications, depending on whether the population fluctuations in different domains over a given area are synchronized (strongly correlated) or not synchronized (very weakly correlated). For instance, the existence of synchronization may result in population outbreaks or population falling to low numbers not in a single habitat but over vast areas of space, and hence may pose considerable problems for pest control or nature conservation.

We emphasize that, although the right-hand side of Eq. (2) does not contain the number of samples N explicitly, the correlation coefficient does depend on N because the values of average population density $\bar{u}_A^{(i)}$ and $\bar{u}_B^{(i)}$ depend on N . It is an inherent problem with expression (1) (as well as, in fact, with any other way to calculate \bar{u}) that its accuracy depends on N . A small number of samples can make the accuracy very low. Importantly, the number of samples required for a sufficiently accurate estimate is known to strongly depend on the properties of the population distribution [41, 43, 45]. For an approximately uniformly distributed population, a few samples can provide a very good estimate of the average density (ultimately, just one sample is enough if the distribution is exactly uniform). However, for a strongly heterogeneous or even ‘patchy’ distribution where the density exhibits large-amplitude fluctuations over space, a reasonable accuracy can only be achieved with a much larger number of nodes in the sampling grid, i.e. when all the peaks in the population distribution are somehow ‘resolved’ [40, 41]. On a coarse grid, i.e. where the number of nodes is not sufficiently large to resolve the details of patchy distribution, the evaluation accuracy can become poor. In particular, in the examples shown in Fig. 1, it is readily seen that the arithmetic average of the locally sampled population density would significantly underestimate the true value as the population peaks remain unresolved. The large numerical error of evaluating the average density on a coarse grid, i.e. the difference between the right-hand side in (1) and the true

value \bar{u} , is then carried on to the corresponding time series and therefore can significantly affect the calculated value of the correlation coefficient and hence the conclusions about the presence or absence of synchronization. In Sections 3 and 4, this heuristic inference will be confirmed by a detailed quantitative analysis of several case studies.

3 Synchronization ‘lost and found’: instructive example

Since synchronization in a system of multiple domains is usually considered pairwise, cf. Eq. (2), it is sufficient for our purposes to consider a system consisting of just two domains, A and B . In this section, we restrict our analysis to the one-dimensional case so that the domains are quantified by their lengths (rather than area) which we assume to be the same, $L_A = L_B = L$.

The idea of our analysis is to consider a system with some known, prescribed properties and to show how these properties may become obscure or distorted when the sampling grid is coarse. Specifically, we simulate two sequences of spatial population distributions, different in A and B , the corresponding total population sizes (or the average population densities) being certain known functions of time. These functions, $\omega_A(t)$ and $\omega_B(t)$ respectively, describe the ‘population dynamics’, i.e. how the population sizes in domains A and B are evolving with time. For the purposes of this section, we consider a strong test where these functions are identical, $\omega_A(t) = \omega_B(t) \equiv \omega(t)$. Thus, the population dynamics in A and B is perfectly synchronized and the theoretical value of the correlation coefficient (2) is exactly one. The question that we are going to consider is what can be the *empirical value* of the correlation coefficient for different value N of nodes in the spatial sampling grid.

The ways to generate ‘realistic’ heterogeneous population densities $u_A(x, t)$ and $u_B(x, t)$ are numerous; for instance, they can be obtained from a population dynamics model [30]. Here we use a simpler approach: we consider the population density being a superposition of normal distributions:

$$u_j(x, t) = \frac{\omega(t)}{p} \sum_{m=1}^p f_{j,t}^{(m)}(x), \quad j = A, B, \quad (4)$$

where $t = 1, 2, \dots, k$ are the moments when the census is taken (e.g. weekly, monthly or annually),

$$f_{j,t}^{(m)}(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp \left[-\frac{\left(x - \hat{x}_{j,t}^{(m)}\right)^2}{2\sigma^2} \right], \quad (5)$$

where parameter σ is the standard deviation and the peak location $\hat{x}_{j,t}^{(m)}$ are independent identically distributed random variables drawn from a certain probability distribution for every peak and every year. Parameter p has the meaning of the number of peaks in the distribution, although some of the peaks can merge or nearly merge when their location is close. The single-peak distribution ($p = 1$) can be regarded as a case of high population aggregation while the multi-peak case $p \gg 1$ corresponds to a case where the population is somehow distributed over the whole domain. Examples of function (4) are shown in Fig. 2.




Figure 2: An example of the superposition $u(x)$ of normal distributions for $p = 4$ (left) and $p = 8$ (right).

Integration of the population density (4) gives the population size. In the unbounded domain $-\infty < x < \infty$, the Gaussian distribution (5) is scaled to one, so that

$$\bar{M}(t) = \int_{-\infty}^{\infty} u_j(x, t) dx = \frac{\omega(t)}{p} \cdot p = \omega(t), \quad (6)$$

therefore ω is indeed the population size. In case the population (4) is considered in the bounded domain $0 < x < L$,

$$M(t) = \int_0^L u_j(x, t) dx < \omega(t), \quad (7)$$

because the tail of the distribution lies outside of the domain. Moreover, considering the location of each peak to be a random variable uniformly distributed over the domain, the deviation $|M - \omega|$ can be considerable (in the single-peak case it can be as large as $\omega/2$, i.e. 50% of the total). In order to reduce the effect of this random fluctuation in the population size, we consider the model where the location of each peak is drawn from a uniform probability distribution defined over a truncated domain $\epsilon < x < L - \epsilon$ where the auxiliary parameter ϵ is chosen sufficiently large compared to the standard deviation σ to provide $M \approx \bar{M} = \omega$.

With regard to the population dynamics, we begin with the simple hypothetical case where the population size grows linearly with time, i.e. $\omega(t) = Ct$ where C is a certain constant parameter. For this system, we consider the series of $k = 500$ censuses and, correspondingly, generate 500 population distributions for each domains A and B .

We then consider the sequence of sampling grids with the number of nodes N increasing from $N = 1$ to $N = N_{max}$. For every given N , the sampling grid is centered around the domain midpoint and the grid nodes (i.e. the location of the sampling points across space) are distributed uniformly over the domain with constant spacing Δx between the neighboring nodes:

$$x_i = i\Delta x, \quad i = 1, \dots, N, \quad (8)$$

where $\Delta x = L/(N + 1)$. For every given N , each of the generated population distributions is sampled on the corresponding grid and the average population density (1) is calculated. The

time course of values of the average population density obtained for a given N is then fed into Eqs. (2–3) to calculate the corresponding value of the correlation coefficient. Correspondingly, the correlation coefficient becomes a function of the number of grid nodes, $\rho = \rho(N)$, where for the sake of simplicity we now omit A , B and k from the notation for the correlation coefficient but emphasize its dependence on N .

Figure 3 shows the results obtained for a highly aggregated single-peak population distribution, i.e. Eq. (5) with $p = 1$ and $\sigma = 8.0$. The domain length is $L = 300$, truncation is done with $\epsilon = 40$. We observe that the truncation of the domain at the ends to generate the distribution (4) is indeed necessary to correctly describe the highly correlated dynamics between the two domains, with $\rho \approx 1$. In case the domain is not truncated, ρ does not approach one even for a very large N . In the rest of this section, we therefore stick to the case where the location each peak is a random variable uniformly distributed over the truncated domain $[\epsilon, L - \epsilon]$.

It is readily seen that correlation coefficient (2) depends on N quite strongly. In order to obtain its correct value $\rho \approx 1$, the sampling grid must contain a sufficiently large number of nodes, i.e. $N \geq 15$; see Fig. 3. For $N \geq 10$, the domains are strongly correlated ($\rho \geq 0.8$), which may be regarded as a good approximation to the actual situation of the almost perfect synchronization. However, for a number of nodes $N \leq 6$, the domains are only correlated weakly or very weakly ($\rho < 0.4$), which has little to do with reality: the synchronization is lost. We therefore conclude that, in case of an aggregated population distribution, synchronization cannot be seen unless the sampling grid contains a sufficiently large number of nodes.

Similar results are obtained in case the population spatial distribution is not highly aggregated but consists of several peaks or patches. Figure 4 shows the correlation coefficient $\rho(N)$ calculated in case of such a multi-peak distribution (4) with various number of peaks p . Although in this case the drop in the calculated value of ρ observed for small N is less dramatic, it still differs significantly from the true value. In particular, on a grid of $N = 3$ nodes we have $\rho = 0.42$ and $\rho = 0.6$ (instead of $\rho = 1$) for the number of peaks $p = 4$ and $p = 8$, respectively.




Figure 3: The correlation coefficient for the single peak case, see distribution (5), obtained in the full domain $[0, 300]$ (solid curve) and truncated domain $[40, 260]$ (dashed curve). The distribution parameter is $\sigma = 8.0$.




Figure 4: The correlation coefficient $\rho(N)$ for different number of peaks p in the distribution (4) in the case where the population dynamics is described by a linear function $\omega(t)$. Other parameters are the same as in Fig. 3.

The higher is the population aggregation, the more prominent becomes the dependence of the correlation coefficient on the number of nodes in the sampling grids. Figure 5 shows $\rho(N)$ in case of two different values of the standard deviation in the single-patch distribution, cf. Eq. (4) with $p = 1$, i.e. $\sigma = 8$ (dashed curve) and $\sigma = 3$ (solid curve). It is readily seen that in the latter case the true value $\rho \approx 1$ is not obtained until $N = 40$ or larger, and the domains do not appear to be strongly correlated unless $N \geq 28$. For $N \leq 20$, synchronization is lost as the domains appear to be correlated only weakly or very weakly.

A question arises here as to how the resolution of the sampling grid (i.e. the distance between the neighboring grid nodes) can be related to the spatial scale of the pattern in order to provide a reliable estimate of the correlation coefficient. A quantity known as the Nyquist frequency is often used in spatial ecology (cf. [19, 47]) to quantify spatial variability of the population density with the goal to determine the resolution of the sampling grid required to avoid any significant loss of information. Omitting mathematical details, the sampling strategy based on the Nyquist frequency




Figure 5: The case of a single peak distribution as given by Eq. (4) with $p = 1$: dashed curve for $\sigma = 8$, solid curve for $\sigma = 3$.

recommends to have at least two samples per population peak. At the first sight, it agrees well with our results shown above. Indeed, in the high aggregation case shown in Fig. 5, the correlation coefficient approaches one for $N \geq 15$ and $N \geq 42$ in case of $\sigma = 8$ and $\sigma = 3$, respectively. However, for a more complicated, multi-peak pattern the agreement is worse: inspection of Fig. 4 immediately reveals that the required sampling grid is almost twice coarser than the one based on the Nyquist frequency. The more complicated the spatial pattern is, the worse this disagreement becomes. In the next section (see also the last part of Section 6), we will show that in a more realistic case accounting for some details of the population dynamics the approach based on the Nyquist frequency can hardly be applied at all.

In conclusion to this section, we mention that the results shown in Figs. 3–5 as well as Fig. 6 below are obtained based on a single realization of a stochastic process (i.e. the random position of the population peak inside the domain). Another realization of the same process may lead to a somewhat different result. Generally speaking, for any given N , one should consider a distribution of values for $\rho(N)$ coming from different realizations, which could be quantified, for instance, by its median and the confidence interval. However, in the case of the above results, the lack of the ensemble of realizations is compensated by the length of the time series⁴: recall that $k = 500$. Results of complementary simulations (not shown here for sake of brevity) reveal that the confidence intervals for $\rho(N)$ shown in Figs. 3–6 are very small. For a smaller number of censuses, the effects of stochasticity may become more explicit. This issue will be further discussed at the end of Section 4.

3.1 Case of more complex dynamics

We have demonstrated that the information about synchronization can be lost if sampling data are collected over a coarse grid. However, the type of the population dynamics that we used – i.e. the population size being a linear function of time – is arguably a simple and rather special case. The question therefore arises as to whether our results on the synchronization loss on a coarse grid may also be a special case, or the situation remains qualitatively the same if the population dynamics is more complicated or more realistic.

In order to address this issue, we now consider a model where the population size $\omega(t)$ is given by the Ricker map:

$$\omega_{t+1} = r\omega_t e^{-\alpha\omega_t}, \quad (9)$$

where r and α are parameters. Note that Eq. (9) is a more realistic model than the simple linear increase used in the previous section; in particular, the Ricker map is widely used in fisheries [50, 60].

It is well-known that, depending on parameter values, model (9) can exhibit rich dynamics including multiperiodic oscillations and chaos [2]. We therefore use (9) in order to generate a sequence of values $\omega_1, \dots, \omega_k$ to simulate a ‘realistic’ dynamics (Fig. 6, left). This values are then

⁴The property known in the theory of complex systems as ergodicity states that, under certain conditions, the length of the observation time is equivalent to the number of realizations, e.g. see [5]

used in the same approach as in the previous section, i.e. first to generate a sequence of spatial distributions (4) and then to calculate the correlation coefficient for different number of the grid nodes N .

The results are shown in Fig. 6 (right). It is readily seen that the dependence of the correlation coefficient ρ on N possesses essentially the same features as for the simple linear population growth, i.e. the true value $\rho \approx 1$ is obtained only if N is sufficiently large. Similarly, the dependence on N is more prominent for a single peak distribution than for a multi-peak distribution, cf. cases $p = 1$, $p = 4$ and $p = 8$ in Fig. 6. We therefore conclude that the loss of information about synchronization (i.e. considerable decrease in the correlation strength) observed when the sampling data are collected on a coarse sampling grid is not case specific but takes place for the population dynamics with various properties ranging from very simple to very complicated.

4 Synchronization in different population models

As was discussed in the introduction, in ecological studies the information about population abundance such as the population size or the spatially average population density is usually deduced from data collected by spatially discrete sampling, i.e. by taking samples in the nodes of a certain spatial grid. In the previous section, we have demonstrated that sampling over a *coarse grid* can result in wrong conclusions about the population dynamics. When the sampling data are used to reveal the degree of correlation between population dynamics in different habitats, e.g. to reveal the presence or absence of synchronization, the correlation coefficient becomes a function of the number N of samples in a census, i.e. the number of nodes in the sampling grid. We have shown that, if N is small, the calculated correlation coefficient is likely to be small too (e.g. $\rho \approx 0.4$ or smaller) regardless of its actual value, even in case of the perfectly synchronized dynamics where the true value is $\rho \approx 1$. Therefore, when sampling is done over a coarse grid, the synchronization is likely to be lost.




Figure 6: (left) The time course of population size $\omega(t)$ simulated with model (9) for $\alpha = 2$ and $r = 19$; (right) the corresponding correlation coefficient $\rho(N)$ obtained for spatial distributions with different number of peaks p ; see Eq. (4).




Figure 7: (left) Probability density function of sample values in a census in case the spatial distribution is single-peaked (cf. Eq. (4) with $p = 1$) as described by the probability density function (10–11). (right) Frequency Q of sample values in case of a multi-peaked distribution (Eq. (4) with $p = 4$) obtained numerically.

The above results were, however, obtained in a rather idealized system. One limitation of the model used in Section 3 is that, in any given census, the statistical distribution of the values of the population density over the collection of samples is, in fact, predefined by the choice of the density profile as (4). To demonstrate this, let us consider the high aggregation case where the population density forms a single peak described by the Gaussian distribution (5). Consider the ultimate case where the sampling grid consists of a single node located in the center of the domain, $x_1 = L/2$. Using standard probability calculus, it is then straightforward to calculate the probability distribution function (pdf) $\phi(u)$ of the event that $u(x_1)$ takes the prescribed value u :

$$\phi(u) = 0 \quad \text{for} \quad u > \frac{\omega(t_1)}{\sigma\sqrt{2\pi}}, \quad (10)$$

$$\phi(u) = \frac{\sigma}{L\sqrt{2}} \left(\log \left[\frac{\omega(t_1)}{u\sigma\sqrt{2\pi}} \right] \right)^{-1/2} \frac{1}{u} \quad \text{for} \quad 0 < u \leq \frac{\omega(t)}{\sigma\sqrt{2\pi}}, \quad (11)$$

where t_1 is the time when the census is taken,

Probability distribution (10–11) is shown in Fig. 7 (left). For a more general case of a multi-peak distribution, i.e. Eq. (4) with $p > 1$, the analytical expression for the pdf is not available but it can be readily obtained by numerical simulations; an example is shown in Fig. 7 (right). Therefore, in both cases the pdf has a bimodal shape, this shape being more pronounced on the single peak case.

We mention here that the frequency distribution shown in Fig. 7 is not unrealistic: the distribution of sample data observed in the population census of some plant species has a similar shape [17]. Yet it gives only one possible case from a great multiplicity of various probability distribution functions that are used to describe sampling data collected for different species and under different ecological conditions [14, 63]. Questions therefore arise as to (i) how common is the situation where synchronization remains undetected on a coarse sampling grid and (ii) how the minimum number of samples sufficient to reveal synchronization may depend on the properties of the popu-

lation dynamics as reflected by the pdf of the sampling data. Indeed, as was discussed in Section 2, the accuracy of the estimate of the average population density (1) depends on the sample size N . However, the accuracy of the estimate depends also on the way how the sample values are distributed, because the rate of convergence of the arithmetic average to the true mean density is somewhat different for different probability distributions. Therefore, the same sample size N may be sufficient to reveal synchronization in one case, e.g. for one probability distribution of sample values, but insufficient in case of another probability distribution.

In order to address these issues, in this section we simulate sampling data using a variety of probability density functions; see Table 1. We assume that the distance between any two neighboring nodes of the sampling grid is large enough to exclude possible interference between them. Correspondingly, the population densities obtained at any two grid nodes in both domains A and B are independent identically distributed random variables drawn from a given pdf ϕ . This produces a certain spatial pattern of the population distribution which is somewhat different for different pdfs of the sampling data (e.g. having different variance); see Fig. 8. For a given ϕ , the procedure is repeated to generate the two time courses of the spatial patterns, i.e. in domains A and B. The correlation coefficient is then calculated basing on the time series of mean population density where the mean density is calculated as the arithmetic average over N samples taken in the corresponding census; see Eqs. (1–3). For the same sequence of the generated spatial patterns, this procedure is repeated on sampling grids with different number of nodes to obtain $\rho_\phi(N)$.

In order to place our analysis into the context of a real field study, we relate the problem of spatial sampling to the recent study on synchronization of *Tipula paludosa* in agricultural landscape [7]. The metapopulation of *T. paludosa* was monitored for fifteen years in thirty eight agricultural fields across South-West Scotland. In each of those fields, the *T. paludosa* population was subjected to annual census in winter, i.e. at the time when it is at the larvae stage and is mostly dwelling in the soil. In each field, twenty five soil cores were taken at randomly chosen locations, cf. [10]. The number of larvae in each soil core was counted. Given the known radius r of the core,

Name	Probability $P\{u = n\}$ or probability density $\phi(u)$	Distribution parameter(s)	Mean, \bar{u}
Poisson	$P\{u = n\} = \frac{\lambda^n}{n!} e^{-\lambda}$	λ	λ
Exponential	$\phi(u) = \lambda e^{-\lambda u}$	λ	$\frac{1}{\lambda}$
Gamma	$\phi(u) = \frac{1}{\lambda^m \Gamma(m)} u^{m-1} \exp\left(-\frac{u}{\lambda}\right)$	λ, m	$m\lambda$
Log-normal	$\phi(u) = \frac{1}{u\sigma\sqrt{2\pi}} \exp\left[-\frac{(\log u - \mu)^2}{2\sigma^2}\right]$	σ, μ	$\exp\left(\mu + \frac{\sigma^2}{2}\right)$
Power law	$\phi(u) = \frac{C}{(\delta+u)^m}, \quad C = (m-1)\delta^{(m-1)}$	$\delta, m \ (m > 1)$	$\frac{\delta}{m-2}, \ m > 2$

Table 1: *Population models to describe frequency of sample values u in a census.*




Figure 8: (colour online) Spatial population distribution obtained on a hypothetical rectangular sampling grid of $10 \times 10 = 100$ nodes with the grid step $\Delta x = \Delta y = 10$ for (left) Poisson distribution of the population density, (right) exponential distribution of the population density. In both cases, the average density is $\bar{u} = 10$. Note that different distributions of sampling data correspond to somewhat different spatial patterns, in particular the variance of the spatial distribution is 10 and 100 for the left and right panels, respectively.

the count n provides a reliable index of the local population abundance; in particular, it can be used to calculate the local population density u as $u = n/(\pi r^2)$. The arithmetic average of the counts was then calculated for each field and each census. The fifteen year courses of the population density obtained for each field were then fed pairwise into Eqs. (2–3) to calculate the correlation coefficient (3). It was observed that some of the fields are strongly synchronized, with $\rho \approx 0.8$ or larger; for details see [7].

For the purposes of our analysis, we choose two pairs of fields so that Pair 1 consists of fields A1 and B1 and Pair 2 consists of fields A2 and B2; see Table 2. Figure 9 shows the corresponding time courses of the spatially average population counts. It is readily seen that the population fluctuations are not independent; indeed, the calculated correlation coefficient (Table 2) shows that they are strongly correlated. The question that we are asking here is: how the conclusions about *T. paludosa* synchronization might have changed if the number of the samples (soil cores) would be much less than $N = 25$, could the synchronization still be seen? Similarly, if the number of samples would have been much larger than $N = 25$, would the results possibly reveal even stronger correlation between the fields?

Unfortunately, the raw data, i.e. the insect counts for each soil core, are not available. In each census, only the arithmetic average of the samples is available for each field. It is therefore not possible to work with the original data, in particular, it not possible to tell what was the probability distribution for the sampling data (i.e. larvae counts). It was shown in [7] that the average density⁵ is well described by a log-normal distribution but this does not necessarily mean that the individual

⁵more precisely, the residuals of the average density obtained after removing density dependence from the original data, see [7] for details.

Name (as used in the text)	Position (grid reference)	Max/min population count	Course average, μ (count)	Correlation coefficient, $\rho(A, B)$
A1	NS 111703	32/1	8.2	0.884
B1	NY 049748	22/0	4.73	
A2	NN 943236	8/0	2.33	
B2	NS 412331	27/0	6.6	

Table 2: *Two pairs of fields from [7] used in our correlation modelling.*

counts are distributed log-normally. In order to fill in for the missing information, we are going to replace the original sampling data with simulated data using different population models, i.e. different pdfs of the frequency of the sample values given in Table 1. For each year of the survey, we simulate N_{max} samples to reproduce the actual (observed) average:

$$\bar{u} = \frac{1}{N_{max}} \sum_{i=1}^{N_{max}} s_i^{(\phi)}, \quad (12)$$

where the superscript (ϕ) refers to one of the models in Table 1, $s_1, \dots, s_{N_{max}}$ are the samples generated according to a given pdf ϕ , and \bar{u} is the known average (see Fig. 9). N_{max} is chosen to be sufficiently large to ensure that the arithmetic mean approaches closely its theoretical limiting value. (In simulations shown below, we used $N_{max} = 100$.) We then use only a subset N of these simulated samples, $N \leq N_{max}$, to calculate the corresponding subset average:

$$\bar{u}_N^{(\phi)} = \frac{1}{N} \sum_{i=1}^N s_i^{(\phi)}. \quad (13)$$

Obviously, in a general case $\bar{u}_N^{(\phi)} \neq \bar{u}$, because the estimation on a coarser sampling grid is less




Figure 9: *The average population counts over time in the selected pairs of correlated fields (see Table 2), left: Pair 1, crosses for A1, diamonds for B1; right: Pair 2, crosses for A2, diamonds for B2.*

accurate. The average (13) is then used to calculate the correlation coefficient (2). By varying N , we reveal the dependence of ρ on the sample size N for any given pdf ϕ .

The results obtained for different population models as given by different pdfs in Table 1 are shown in Figs. 10–11 where the left and right columns correspond to Pair 1 and Pair 2, respectively. Note that, for any given pdf ϕ , for each year in the time course the parameter(s) of the pdf are chosen somewhat differently in order to agree with the current value of \bar{u} ; see the last column in Table 1. Given the random nature of the simulated samples and the relatively short length of the time courses (fifteen annual surveys, i.e. fifteen points), it is not surprising that the correlation coefficient ρ exhibits stochastic fluctuations. In order to decrease the effect of stochasticity and hence to make the general tendency clearer, for any given N the procedure was repeated ten times; the thick curve shows $\rho(N)$ averaged over those ten realizations. To show the range of possible values due to the inherent randomness of the system, the *confidence interval* is calculated: the dotted curves show the averaged value of ρ plus-minus the standard deviation calculated over the ten realizations.

It is readily seen from Figs. 10–11 that there is a clear difference between the case where the probability distribution has a maximum at some positive value $u > 0$ (as for the Poisson, gamma and log-normal distributions) and the case where the probability distribution has a maximum at $u = 0$ (as for the exponential and power law distributions). In the former case, see Fig. 10, a good estimate of the correlation coefficient (e.g. within ten percent of its true value shown by the dashed-and-dotted horizontal line) is typically obtained for a relatively small number of samples; in particular, just 3–4 samples per census can be sufficient for the Poisson and log-normal distributions and 9–10 in case of the Gamma distribution. The situation is different in the latter case; see Fig. 11. In the cases of the exponential distribution and the power law with $m = 3$, a reliable estimate of the true value of ρ is not obtained until the number of samples in a census is 18–20; see the top and middle rows in Fig. 11. The convergence of $\rho(N)$ to its true value is somewhat slower when the distribution of sample values is described by a power law with $m = 2$ where a reliable estimate is not obtained until the sample size $N \approx 25$; see the bottom-left panel in Fig. 11. Interestingly, some apparently minor details of the dynamics can affect the results too: the required sample size appears to be smaller in Pair 2 than in Pair 1 for all three cases shown in Fig. 11.

5 Ghost synchronization on a coarse sampling grid

We therefore have shown that, when the population density exhibits considerable variation over space and the data are collected on a coarse sampling grid, synchronization is often lost as the correlation coefficient is usually much smaller than its actual value. In this section, we are going to demonstrate that the opposite is also possible. Namely, we will show that sampling on a coarse grid may result in a ‘ghost synchronization’, i.e. in the situation where the correlation coefficient calculated from the coarse sampling data has a value close to one whilst the dynamics is actually anti-correlated.

We consider a simple system consisting of two 1D domains A and B where the population of




Figure 10: Correlation coefficient $\rho(N)$ (thick curve) calculated for different population models and parameters (as in Table 1): (top) Poisson distribution with $\lambda = \bar{u}$, (middle) gamma distribution with $m = 2$ and $\lambda = \frac{1}{2}\bar{u}$, (bottom) log-normal distribution with $\sigma = 0.45$ and $\mu = \log(\bar{u}) - 0.5 * \sigma^2$. Left column for Pair 1, right column for Pair 2. Dotted curves show the calculated value of ρ plus-minus standard deviation, the vertical distance between the curves thus being the confidence interval; the dashed-and-dotted line shows the true value of ρ .




Figure 11: Correlation coefficient $\rho(N)$ (thick curve) calculated for different population models and parameters (as in Table 1): (top) exponential distribution with $\lambda = 1/\bar{u}$, (middle) power law with $m = 3$ and $\delta = \bar{u}$, (bottom) power law with $m = 2$, in this case δ is chosen for \bar{u} to coincide with the median of the distribution. Left column for Pair 1, right column for Pair 2. Dotted curves show the calculated value of ρ plus-minus standard deviation, the vertical distance between the curves thus being the confidence interval; the dashed-and-dotted line shows the true value of ρ .

a certain species has a unimodal spatial distribution with the maximum at the domain boundary:

$$u_j(x, t) = \frac{\omega_j(t)}{\sigma\sqrt{2\pi}} \exp\left[-\frac{x^2}{2\sigma_j(t)^2}\right], \quad 0 \leq x \leq L_j, \quad j = A, B. \quad (14)$$

Note that, although the functional form (14) of the population distribution is specific, the case where the population density decays monotonously with the distance from the domain border is relatively common in ecosystems and agroecosystems; in particular, it may correspond to the invasion of a pest insect to a farm field from adjoining uncultivated areas [6].

We assume that the populations are affected by factors that are different in the two domains, so that the population size in domain A and domain B evolves differently with time (but preserving the shape of the spatial distribution). We consider the following hypothetical situation:

$$\omega_A(t) = \omega_0(t+1)^a, \quad \sigma_A(t) = \sigma_0, \quad \omega_B(t) = \omega_0 - \gamma t, \quad \sigma_B(t) = \sigma_0(t+1)^b, \quad (15)$$

where $t \geq 0$ and a, b, γ, ω_0 and σ_0 are parameters. Since $\omega_A(0) = \omega_B(0) = \omega_0$ and $\sigma_A(0) = \sigma_B(0) = \sigma_0$, the initial population distribution is the same in both domains. However, the dynamics is different: the maximum population density grows in domain A but decays in domain B while the width of the patch does not change in domain A but increases in domain B .

Let us consider the case where in both domains the initial distribution is aggregated in the vicinity of the habitat boundary, $\sigma_0 \ll \min\{L_A, L_B\}$. If, for the sake of simplicity, we restrict our analysis to the time when the tail of the distribution at the right-hand side of the domains is still thin, i.e. $\sigma_A(t) \ll L_A$ and $\sigma_B(t) \ll L_B$, then the population size in domains A and B is, respectively, $\frac{1}{2}\omega_A(t)$ and $\frac{1}{2}\omega_B(t)$. We then observe that the population size grows in domain A but decreases in domain B . Hence the dynamics is anti-correlated; the corresponding correlation coefficient must be negative, having a value close to -1 . However, the population density at a given location does not necessarily behave in the same way. In fact, it is readily seen from the properties of function (14) that there is a sub-domain where the population density actually tends to increase simultaneously in both domains. An example is shown in Fig. 12. One can expect that,




Figure 12: Population density at the location $x_1 = 10$ (as given by Eqs. (14–15) with parameters $a = 0.5$, $b = 0.25$, $\gamma = 0.0001$, $\omega_0 = 2$ and $\sigma_0 = 3$) in domain A (curve 1) and in domain B (curve 2).




Figure 13: *Dependence of the correlation coefficient on the number of nodes in the sampling grid for two different ways to refine the grid: (left) extra nodes are added towards the tail of the population spatial distribution, (right) extra nodes are added towards the maximum of the population spatial distribution; see the main text for more details.*

if the samples on the population density are taken in that sub-domain only, then the corresponding value of the correlation coefficient is going to be positive, possibly being close to one.

Having that said, it remains unclear how the correlation coefficient may depend on the number of samples and on their locations. In order to make a more quantitative insight into the properties of the system (14–15), we now use simulations, i.e. we generate a sequence of sampling grids with different number of nodes N and calculate the correlation coefficient $\rho(N)$ accordingly. To perform simulations, we use the following parameters: $L_A = L_B = 300$, $a = 0.5$, $b = 0.25$, $\gamma = 0.0001$, $\omega_0 = 2$ and $\sigma_0 = 3$.

The results are shown in Fig. 13 where the sampling grid is chosen differently in the left and right panels. In the case of Fig. 13 (left), in both domains A and B the additional nodes are added at the right, i.e. towards the tail of the spatial distribution, their location being defined as $x_i = x_1 + (i - 1)\Delta x$ where $2 \leq i \leq N$. The results shown in Fig. 13 (left) are obtained in case $\Delta x = x_1 = 10$. We therefore observe that ρ is not very sensitive to the sample size N ; the correlation coefficient changes just slightly from $\rho(N = 1) \approx 0.89$ to $\rho(N = 27) \approx 1$. Remarkably, these values has nothing to do with reality as the population dynamics described by Eqs. (14–15) is anti-correlated with $\rho \approx -1$.

In the case of Fig. 13 (right), in both domains A and B the additional nodes are added at the left, i.e. towards the center of the spatial distribution (14). The location of additional nodes is defined as $x_i = x_1 - (i - 1)\Delta x$, $2 \leq i \leq N$ (the results shown in Fig. 13-right are obtained for $x_1 = 10$ and $\Delta x = 1$). In this case, ρ strongly depends on N by exhibiting a monotonous decay from the false value $\rho(N = 1) \approx 0.89$ to the actual value $\rho(N = 10) \approx -1$.

We therefore conclude that, in order to obtain the correct value of the correlation coefficient, not only the number of nodes in the sampling grid is important but their location too, the latter being determined by the properties of the spatial pattern. In particular, in the case of system (14–15), samples collected in the area close to the maximum of the distribution are apparently more important as they bring more information than samples collected in the area at its tail. This

conclusion is further confirmed by the dependence $\rho(N)$ calculated on a grid with mixed properties (see Fig. 14) where the second node is placed in the ‘important’ range $0 < x < x_1$, namely at the location $x_2 = 0.5x_1$, but other additional nodes are added at the right of x_1 with the spatial step $\Delta x = 0.5x_1$. It is readily seen that, while the second node brings some essential information sufficient to change the value of the correlation coefficient from a completely false value $\rho \approx 0.89$ to a much more realistic $\rho \approx -0.85$, all other nodes added in the area towards the distribution tail do not improve the accuracy any further.




Figure 14: Correlation coefficient $\rho(N)$ in the case where the sampling grid is refined in a mixed way; see the main text for details.

6 Discussion and conclusions

In ecological and environmental studies as well as in other natural sciences and environmental engineering, it is often needed to estimate the population size of a given species or the total mass of a given substance based on local, spatially discrete data collected at the nodes of a certain sampling grid [9, 13, 54, 56, 62]. When the spatial distribution of a given population (or substance) exhibit a considerable variability in space, which is rather typical in ecology [18, 23, 27], the number of collected samples may not always be sufficient to resolve the details of the spatial configuration. Moreover, information about the spatial pattern of the population density distribution, e.g. the exact location of peaks or patches, usually is not known a priori (but see [1]); it is eventually obtained as a result of the analysis of the sampling data [26].

It therefore often happens that the number of samples in a census as well as the location of samples are chosen based on a guess or a certain rule of thumb. It can also be negatively affected by some external constraints, e.g. a limited budget. As a result, the sampling grid may appear to be coarse, i.e. not resolving the spatial population distribution in sufficient details. Estimation of the population size⁶ on a coarse grid would normally have low accuracy [11, 12, 41, 43, 45, 54]. When

⁶or the spatially average population density, which differs from the population size only by a factor of the area of the habitat which we assume can always be determined with sufficient precision.

the estimated population size is used as the input information for some further analysis, e.g. to assess the correlation strength between two habitats, this inaccuracy is likely to affect the results too. In this paper, we showed that the correlation coefficient ρ calculated based on sampling data collected over a coarse grid often has little to do with its true value. Even a very strong correlation (i.e. $0.8 \leq \rho \leq 1$), usually referred to as synchronization, can be “lost”, i.e. remain unseen, as the value obtained on a coarse grid is typically $\rho \sim 0.5$ or less, e.g. see Figs. 4–6 and 11. Moreover, we also showed that, when the location of nodes in the sampling grid is chosen inadequately, the opposite case is possible, i.e. the correlation coefficient calculated based on the sampling data is close to one whilst in reality there is no synchronization (see Section 5). Remarkably, this “ghost synchronization” can happen even if the sampling grid contains an apparently large number of nodes, e.g. see the left panel in Fig. 13.

Here we mention that synchronization has a variety of implications for ecology, agroecology and nature conservation, in particular because synchronization is one of the main dynamical mechanisms behind large scale population crashes [31, 57] as well as large scale outbreaks [28, 58]. Hence, the capability to detect the presence or absence of synchronization is crucial for planning, forecasting and decision making. Reliability of the calculated value of the correlation coefficient is therefore an issue of high practical importance.

Test of grid coarseness

An important question arises as to whether it may be possible, based on the available sampling data, to separate the cases where the calculated ρ is likely to be close to its true value from the cases where the calculated ρ is likely to be wrong. Based on the results of our analysis in Section 4, the following *test of grid coarseness* can be suggested. We do not normally know the spatial pattern but we can estimate, based on the collected data, what is the probability density function (pdf) of the sampling data. Once the pdf is revealed, additional data distributed accordingly to this pdf can easily be simulated to create a virtual sequence of sampling grids with different number of nodes N , and then $\rho(N)$ can be calculated following the procedure described in Section 4. Once $\rho(N)$ is available, its convergence to the large- N limit can be readily established and then it is straightforward to estimate from the shape of the graph how many nodes are needed. For instance, in case of the Poisson distribution (see the top row in Fig. 10), a sparse grid consisting of 4-5 nodes should be sufficient to obtain the true value of the correlation coefficient; however, in case of the exponential distribution the grid will only become sufficiently refined when the number of nodes is 30 or more, cf. the top-left panel in Fig. 11.

With this new understanding thus achieved, we are now going to briefly revisit some cases of synchronization reported in the literature with the purpose to assess whether the corresponding sampling grids were adequate or perhaps too coarse:

- Region-wide synchronization of *Tipula paludosa* in South-West Scotland was reported in [7]. The average population density in each farm field used in the study was estimated based on 25 samples (soil cores). Moreover, there was some evidence presented that the distribution of frequencies was well described by a log-normal distribution [7]. An inspection of

the bottom row of Fig. 10 reveals that $\rho(N)$ approaches the vicinity of its true value (with the 10% tolerance) when $N \geq 4$. We therefore conclude that the results reported in [7] are reliable.

- Synchronization of several Lepidoptera species was observed in central Appalachian deciduous forests [48]. The study included twelve plots, each of them having the same area of 200 ha. In each plot, the data were collected by a single light trap. Raimondo et al. [48] do not provide any information about the frequency distribution of their sampling data. However, in another study on Lepidoptera [16], it was shown that sampling data for at least some Lepidoptera species are well described by either negative binomial distribution or Poisson distribution [16]. If we assume that this result is transferrable between the two studies (which is a rather strong assumption, because the study [16] was done in another geographical region), then we can make use of the results shown in the top row of Fig. 10. It is then readily seen that a single sample (i.e. single trap) is very unlikely to provide an exact value of ρ as the true value does not even fall into the range of possible values (shown by the dotted curves). The estimate of the correlation coefficient obtained based on a single sample is likely to considerably underestimate its true value, with the deviation from the true value being about 20-25%. We therefore conclude that Lepidoptera species in Appalachian forests are likely to be correlated much stronger than it was observed in [48].
- Synchronization of carabid beetles due to the weather fluctuations (the phenomenon known as the Moran effect [32, 33, 53]) was observed in a study performed in a nature reserve in the Netherlands [3]. The study area covering a few square kilometers was split into a few zones and in each zone three pitfall traps were installed to sample the carabid beetles population. Baars and Van Dijk [3] did not provide any analysis of the frequency distribution of their sampling data (i.e. trap counts). Some relevant information is available from another study on carabid beetles performed in the same geographic region [25, 52]. Although Rossi et al. [52] did not do a formal fitting of the trap count data with a statistical model, the shape of the frequency histogram (e.g. see Figs. 3 and 6 in [52]) suggests that it is likely to be better described by a probability density function with the maximum at the origin and a relatively slow rate of decay at the tail. We therefore hypothesize that, from the cases analyzed in Section 4, either the exponential distribution or a power law distribution are best candidates. We should also mention here the apparent visual similarity between the qualitative properties of the field data on carabid beetles distribution shown in Fig. 1b and the simulated spatial pattern shown in Fig. 8, right. Making use of the results shown in Fig. 11, we readily observe that the sampling grid consisting of three nodes is coarse and is likely to considerably underestimate the actual strength of the correlation. This may be a reason why the correlation between different sites and/or different subpopulations reported by Baars & Van Dijk [3] was not as strong and widespread as it perhaps might intuitively be expected.

Concluding remarks

We therefore conclude that the properties of the sampling grid such as the total number of samples in a survey and their location must be decided based upon a rigorous argument rather than a guess or rule of thumb. Although this may sound as a trivial statement, in field studies focusing on revealing synchronization surprisingly little attention is paid to checking whether the number of samples (e.g. traps) is sufficient to provide a robust estimate of the population abundance. As just one example, here we cite Baars and Van Dijk [3]: “summed catches *were assumed* to represent the adult density around a series of pitfall trap”⁷. As we discussed it above, one rigorous argument could be based on the analysis of the sampling data frequencies which makes it possible to estimate the minimum required number of nodes in the grid. Where possible, this should also be combined with some a priori knowledge of typical properties of the spatial population distribution of the given species. For instance, this information can be obtained from relevant previous studies (e.g. as available from literature) or from a specially designed pilot study. Although it undoubtedly requires an extra effort, it seems to be a necessary step in order to make any conclusion on the presence or absence of synchronization reliable. In particular, in order to avoid the ‘ghost synchronization’, one should have some a priori knowledge of the population dynamics of the species. This should include not only the pattern of the population spatial distribution but also some information about the temporal scales of the dynamics. For instance, if the correlation coefficient in the model (14–15) is calculated using time courses obtained over much longer time, then the correct value $\rho(N) \approx -1$ can be obtained even for a relatively small N (for the sake of brevity, we do not show the results here).

One important message following from our study is that the spatial resolution of the sampling grid cannot be decided upon based on just one universal rule. The appropriate frequency of spatial sampling appears to be context-specific and hence depends of the focus of the study. For instance, if the focus is on revealing the details of population distribution across all scales of spatial variability, then the approach based on the Nyquist frequency is known to work well: in order to avoid information loss, it recommends to have at least two sampling nodes per peak in the smallest spatial scale involved [19, 47]. (We mention here that in the problem of ecological patterning relevant spatial scales range from the microscale of the size of an individual to the macroscale of the geographical and climatic variation [44], and hence the decision about the ‘minimum’ spatial scale may often be arguable.) However, if the focus is on the evaluation of the total population size, the requirements to the sampling grid can be much less restrictive as the resolution depends on the required accuracy [38, 39, 40]. Furthermore, in case sampling is needed not only across space but also over time, e.g. to reveal the presence or absence of synchronization, the sampling grid resolution strongly depends on the population dynamics of the sampled species so that the required number of nodes in a given spatial domain can differ by an order of magnitude, c.f. the top-left panel in Figs. 10 and 11.

Our study leaves a few open questions. Perhaps the most challenging one is about the ghost synchronization. In this paper we have identified only one case where this curious artefact can happen if samples are collected on a coarse grid, i.e. where the population density in the corresponding

⁷The italic is ours. NB & SP.

domains decreases monotonously away from a domain boundary, provided the properties of this density profile changes with time in a certain way. It remains unclear whether the ‘ghost synchronization’ is an exotic situation only happening under some specific conditions or it happens more commonly. This should become a focus of a separate study. Another highly practical issue is the effect of the environmental heterogeneity. Throughout this paper, we have assumed that all sampling locations over the given domain are equivalent, in particular assuming that the probability density function of the sampling data is the same at any location. In real ecosystems, this is not always the case. Further development of our approach to include the effects of spatial heterogeneity will be a focus of the future research.

Authors contributions

NP and SP contributed equally to conceiving and performing the study, NP and SP wrote the manuscript, NP and SP approve the publication of this manuscript in its current form.

Competing interests

We declare that we have no competing interests.

Data accessibility

All field data that are used or mentioned in our study are taken from the literature and hence freely available.

References

- [1] Alexander CJ, Holland JM, Winder L, Wooley C, Perry JN. 2005 Performance of sampling strategies in the presence of known spatial patterns. *Ann. Appl. Biol.* **146**, 361–370.
- [2] Allman ES, Rhodes JA. 2004 *Mathematical Models in Biology: an Introduction*. Cambridge, UK: CUP.
- [3] Baars MA, Van Dijk TS. 1984 Population dynamics of two carabid beetles at a Dutch heathland. I. Subpopulation fluctuations in relation to weather and dispersal. *J. Anim. Ecol.* **53**, 375–388.
- [4] Bakus GJ, Nishiyama G, Eduardo H *et al.* 2007 A comparison of some population density sampling techniques for biodiversity, conservation, and environmental impact studies. *Bio-div. Conserv.* **16**, 2445–2455.
- [5] Balescu R. 1975. *Equilibrium and Nonequilibrium Statistical Mechanics*. New York: John Wiley.
- [6] Bearup D, Petrovskaya N, Petrovskii SV. 2015 Some analytical and numerical approaches to understanding trap counts resulting from pest insect immigration. *Mathematical Biosciences* **263**, 143–160.
- [7] Bearup D, Petrovskii SV, Blackshaw R, Hastings A. 2013 The impact of terrain and weather conditions on the metapopulation of *Tipula paludosa* in South-Western Scotland: linking pattern to process. *American Naturalist* **182**, 393–409.
- [8] Bearup D, Benefer CM, Petrovskii SV, Blackshaw R. 2016 Revisiting Brownian motion as a description of animal movement: a comparison to experimental movement data. *Methods in Ecology and Evolution*, doi: 10.1111/2041-210X.12615 (in press).
- [9] Benjamin MM, Lawler DF. 2013 *Water Quality Engineering: Physical & Chemical Treatment Processes*. New York: John Wiley & Sons.
- [10] Blackshaw RP. 1983 The annual leatherjacket survey in Northern Ireland, 1965–1982, and some factors affecting populations. *Plant Path.* **32**, 345–349.
- [11] Boag B, Deeks L, Orr A, Neilson R. 2005 A spatio-temporal analysis of a New Zealand flatworm (*Arthurdendyus triangulatus*) population in western Scotland. *Ann. Appl. Biol.* **147**, 81–88.
- [12] Boag B, Mackenzie K, McNicol JW, Neilson R. 2010 Sampling for the New Zealand flatworm. *Proceedings Crop Protection in Northern Britain 2010*, 45–50.
- [13] Boissonnat J-D, Nullans S. 1996 *Reconstruction of Geological Structures from Heterogeneous and Sparse Data*. INRIA Report N3069.

- [14] Bolker BM, 2008 *Ecological Models and Data in R*. Princeton: Princeton University Press.
- [15] Byers JA. 1993 Simulation and equation models of insect population control by pheromone-based traps. *J. Chem. Ecol.* **19**, 1939–1956.
- [16] Clarke-Harris D, Fleischer SJ. 2003 Sequential sampling and biorational chemistries for management of Lepidopteran pests of vegetable amaranth in the Caribbean. *J. Econ. Entomol.* **96**(3), 798–804.
- [17] Damgaard C. 2009 On the distribution of plant abundance data. *Ecological Informatics* **4**, 76–82.
- [18] Davis MB, Calcote RR, Sugita S, Takahara H. 1998 Patchy invasion and the origin of a Hemlock-Hardwoods forest mosaic. *Ecology* **79**, 2641–2659.
- [19] Denny MW, Helmuth B, Leonard GH, Harley CDG, Luke J. H. Hunt LJH, Nelson EK. 2004. Quantifying scale in ecology: lessons from a wave-swept shore. *Ecological Monographs* **74**, 513–532.
- [20] Engler R, Guisan A, Rechsteiner L. 2004 An improved approach for predicting the distribution of rare and endangered species from occurrence and pseudo-absence data. *Journal of Applied Ecology* **41**, 263–274.
- [21] Eppstein MJ, Hawrysz DJ, Godavarty A, Sevick-Muraca EM. 2002 Three-dimensional, Bayesian image reconstruction from sparse and noisy data sets: Near-infrared fluorescence tomography. *PNAS* **99**, 9619–9624.
- [22] Evans JD. 1996 *Straightforward Statistics for the Behavioral Sciences*. Pacific Grove, CA: Brooks/Cole Publishing.
- [23] Grünbaum D. 2012 The logic of ecological patchiness. *Interface Focus* **2**, 150–155.
- [24] Hanski I, Woiwod IP. 1993 Spatial synchrony in the dynamics of moth and aphid populations. *J. Anim. Ecol.* **62**, 656–668.
- [25] Hengeveld R. 1979. The analysis of spatial patterns of some ground beetles (col. Carabidae). In: Cormack M, Ord JK (eds). *Spatial and temporal analysis in ecology*. International Cooperative Publishing House, Fairland MD, pp.333–346.
- [26] Holland JM, Perry JN, Winder L. 1999 The within-field spatial and temporal distribution of arthropods in winter wheat. *Bull. Entomol. Res.* **89**, 499–513.
- [27] Levin S. 1994 Patchiness in marine and terrestrial systems: from individuals to populations. *Phil. Trans. R. Soc. B* **343**, 99–103.
- [28] Liebhold A, Koenig WD, Bjornstad ON. 2004 Spatial synchrony in population dynamics. *Annu. Rev. Ecol. Evol. Syst.* **35**, 467–490.

- [29] Lyles D, Rosenstock TS, Hastings A, Brown PH. 2009 The role of large environmental noise in masting: general model and example from pistachio trees. *J. Theor. Biol.* **259**, 701-713.
- [30] Malchow H, Petrovskii SV, Venturino E. 2008 *Spatiotemporal Patterns in Ecology and Epidemiology: Theory, Models, and Simulations*. Boca Raton, FL: CRC Press.
- [31] Milne A, Laughlin R, Coggins RE. 1965 The 1955 and 1959 population crashes of the leather-jacket, *Tipula paludosa* Meigen, in Northumberland. *J. Anim. Ecol.* **34**, 529-534.
- [32] Moran PAP. 1953 The statistical analysis of the Canadian lynx cycle. I. Structure and prediction. *Aust. J. Zool.* **1**, 163-173.
- [33] Moran PAP. 1953 The statistical analysis of the Canadian lynx cycle II. Synchronization and meterology. *Austr. J. Zool.* **1**, 291-298.
- [34] Murray JD. 1989 *Mathematical Biology*. Berlin: Springer.
- [35] Noss RF. 1990 Indicators for monitoring biodiversity: a hierarchical approach. *Conservation Biology* **4**, 355-364.
- [36] Pedigo LP, Buntin GD (eds). 1994 *Handbook of Sampling Methods for Arthropods in Agriculture*. Boca Raton: CRC Press.
- [37] Peltonen M, Liebhold A, Bjrnstad ON, Williams DW. 2002 Variation in spatial synchrony among forest insect species: roles of regional stochasticity and dispersal. *Ecology* **83**, 3120–3129.
- [38] Petrovskaya NB. 2017 ‘Catch me if you can’: evaluating the population size in the presence of a spatial pattern. *Ecol. Compl.*, submitted.
- [39] Petrovskaya N, Embleton N. 2013. Evaluation of peak functions on ultra-coarse grids. *Proc. R. Soc. A* **469**, 20120665.
- [40] Petrovskaya NB, Embleton NL. 2014 Computational methods for accurate evaluation of pest insect population size. In: Godoy WAC, Ferreira CP (eds). *Ecological Modelling Applied to Entomology*. Berlin: Springer, pp.171–218.
- [41] Petrovskaya NB, Petrovskii SV. 2010 The coarse-grid problem in ecological monitoring. *Proc. R. Soc. A* **466**, 2933–2953.
- [42] Petrovskaya NB, Embleton NL, Petrovskii SV. 2013 Numerical study of pest population size at various diffusion rates. In: Lewis MA, Maini P, Petrovskii SV (eds). *Dispersal, Individual Movement, and Spatial Ecology: A Mathematical Perspective*. Springer Lecture Notes in Mathematics, Vol. 2071. Berlin: Springer, pp.355-386.
- [43] Petrovskaya NB, Petrovskii SV, Murchie AK. 2012 Challenges of ecological monitoring: estimating population abundance from sparse trap counts. *J. R. Soc. Interface* **9**, 420–435.

- [44] Petrovskii, S.V. 2016 Pattern, process, scale, and models sensitivity: a comment. *Physics of Life Reviews* **19**, 131–134.
- [45] Petrovskii SV, Petrovskaya N, Bearup D. 2014 Multiscale approach to pest insect monitoring: random walks, pattern formation, synchronization, and networks. *Physics of Life Reviews* **11**, 467–525.
- [46] Petrovskii SV, Bearup D, Ahmed DA, Blackshaw R. 2012 Estimating insect population density from trap counts. *Ecol. Compl.* **10**, 69–82.
- [47] Platt T, Denman KL. 1975. Spectral analysis in ecology. *Annual Review of Ecology and Systematics* **6**, 189–210.
- [48] Raimondo S, Liebhold AM, Strazanac JS, Butler L. 2004 Population synchrony within and among Lepidoptera species in relation to weather, phylogeny, and larval phenology. *Environ. Entomol.* **29**, 96–105.
- [49] Ranta E, Kaitala V, Lindstrom K, Helle E. 1997 Moran effect and synchrony in population dynamics. *Oikos* **78**, 136–142.
- [50] Ricker WE. 1954 Stock and recruitment. *Journal of the Fisheries Research Board of Canada* **11**(5), 559–623.
- [51] Rosenstock TS, Hastings A, Koenig WD, Lyles DJ, Brown PH. 2011 Testing Moran’s theorem in an agroecosystem. *Oikos* **120**, 1434–1440.
- [52] Rossi RE, Mulla DJ, Journel AG, Franz EH. 1992 Geostatistical tools for modeling and interpreting ecological spatial dependence. *Ecological Monographs* **62**, 277–314.
- [53] Royama T. 1992 *Analytical Population Dynamics*. Chapman & Hall.
- [54] Seber GA. 1982 *The Estimation of Animal Abundance and Related Parameters*. London: Charles Griffin.
- [55] Sudman S, Sirken MG, Cowan CD. 1988 Sampling rare and elusive populations. *Science* **240**, 991–996.
- [56] Sutherland WJ (ed.) 1996 *Ecological Census Techniques: A Handbook*. Cambridge: Cambridge University Press.
- [57] Sutcliffe OL, Thomas CD, Yates TJ, Greatorex-Davies JN. 1997 Correlated extinctions, colonizations and population fluctuations in a highly connected ringlet butterfly metapopulation. *Oecologia* **109**, 235–241.
- [58] Swetnam TW, Lynch AM. 1993 Multicentury, regional scale patterns of Western spruce budworm outbreaks. *Ecological Monographs* **63**, 399–424.

- [59] Taylor LR, Woiwod IP, Perry JN. 1978 The density-dependence of spatial behaviour and the rarity of randomness. *J. Anim. Ecol.* **47**, 383–406.
- [60] Toft JE, Burke JL, Carey MP et al. 2013 From mountains to sound: modelling the sensitivity of Dungeness crab and Pacific oyster to landsea interactions in Hood Canal, WA. *ICES J. Mar. Sci.* **71**(3), 725–738.
- [61] Williams DW, Liebhold AM. 1995 Influence of weather on the synchrony of gypsy moth (Lepidoptera: Lymantriidae) outbreaks in New England. *Environ. Entomol.* **24**, 987–995.
- [62] Wu Q, Xu H, Zou X. 2005 An effective method for 3D geological modeling with multi-source data integration. *Computers & Geosciences* **31**, 35–43.
- [63] Young LJ, Young J. 1998 *Statistical Ecology*. Berlin: Springer.