

An unexplored role for Peroxiredoxin in exercise-induced redox signalling?

Aldred, Sarah; Wadley, Alex; Coles, SJ

DOI:

[10.1016/j.redox.2015.10.003](https://doi.org/10.1016/j.redox.2015.10.003)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Aldred, S, Wadley, A & Coles, SJ 2016, 'An unexplored role for Peroxiredoxin in exercise-induced redox signalling?', *Redox Biology*, vol. 8, pp. 51-58. <https://doi.org/10.1016/j.redox.2015.10.003>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.



ELSEVIER

Contents lists available at ScienceDirect

Redox Biology

journal homepage: www.elsevier.com/locate/redox

Mini Review

An unexplored role for Peroxiredoxin in exercise-induced redox signalling?

Alex J. Wadley^{a,*}, Sarah Aldred^b, Steven J. Coles^a^a Institute of Science and the Environment, University of Worcester, Worcestershire WR2 6AJ, United Kingdom^b School of Sport, Exercise and Rehabilitation Sciences, University of Birmingham, B15 2TT, United Kingdom

ARTICLE INFO

Article history:

Received 29 September 2015

Received in revised form

13 October 2015

Accepted 19 October 2015

Available online 25 December 2015

Keywords:

Redox signalling

ROS

Exercise

Cysteine

Thiol

Peroxiredoxin

ABSTRACT

Peroxiredoxin (PRDX) is a ubiquitous oxidoreductase protein with a conserved ionised thiol that permits catalysis of hydrogen peroxide (H₂O₂) up to a million times faster than any thiol-containing signalling protein. The increased production of H₂O₂ within active tissues during exercise is thought to oxidise conserved cysteine thiols, which may in turn facilitate a wide variety of physiological adaptations. The precise mechanisms linking H₂O₂ with the oxidation of signalling thiol proteins (phosphates, kinases and transcription factors) are unclear due to these proteins' low reactivity with H₂O₂ relative to abundant thiol peroxidases such as PRDX. Recent work has shown that following exposure to H₂O₂ in vitro, the sulfenic acid of the PRDX cysteine can form mixed disulphides with transcription factors associated with cell survival. This implicates PRDX as an 'active' redox relay in transmitting the oxidising equivalent of H₂O₂ to downstream proteins. Furthermore, under oxidative stress, PRDX can form stable oxidised dimers that can be secreted into the extracellular space, potentially acting as an extracellular 'stress' signal. There is extensive literature assessing non-specific markers of oxidative stress in response to exercise, however the PRDX catalytic cycle may offer a more robust approach for measuring changes in redox balance following exercise. This review discusses studies assessing PRDX-mediated cellular signalling and integrates the recent advances in redox biology with investigations that have examined the role of PRDX during exercise in humans and animals. Future studies should explore the role of PRDX as a key regulator of peroxide mediated-signal transduction during exercise in humans.

© 2015 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Contents

1. Introduction	51
2. Cysteine thiol groups	52
3. Cysteine targets for hydrogen peroxide during exercise	52
4. Hydrogen Peroxide as a cellular signal: recent advances	52
5. Peroxiredoxin: an abundant and highly active peroxidase	53
6. Is PRDX a peroxide sensing protein during and following exercise?	54
7. Future Perspectives	55
8. Conclusion	56
References	56

1. Introduction

Reactive oxygen and nitrogen species (RONS) are known to mediate a range of signalling processes within mammalian tissues, with their production, interaction and removal by antioxidants all

* Correspondence to: Institute of Science and the Environment University of Worcester, St Johns Campus, Henwick Grove Worcester, WR2 6AJ, United Kingdom.
E-mail address: a.wadley@worc.ac.uk (A.J. Wadley).

critical to cell function. The acute production of RONS in response to exercise has been an active area of research over the last 20 years. The oxidising properties of RONS have been implicated with mediating skeletal muscle excitation-contraction coupling [1,2] (via regulation of calcium signalling [3]), enzyme release [4,5], as well as modulating post-exercise mitochondrial biogenesis [6] and cytoprotective gene expression (e.g. heat shock [7,8] and antioxidant proteins [8,9]). A range of cell types can produce RONS during exercise, including skeletal muscle, immune and endothelial cells [10,11]. The enzymes nicotinamide adenine dinucleotide phosphate- (NADPH) oxidase and nitric oxide synthase (NOS) [12,13] produce superoxide ($O_2^{\bullet -}$) and nitric oxide (NO^{\bullet}) respectively [14,15], with secondary oxidants such as peroxynitrite ($ONOO^-$) formed from the favourable reaction between $O_2^{\bullet -}$ and NO^{\bullet} [16,17], and hydrogen peroxide (H_2O_2), by the conversion of $O_2^{\bullet -}$ to H_2O_2 by the antioxidant enzyme superoxide dismutase (SOD) or spontaneous dismutation [18]. The role of RONS in redox-mediated exercise adaptation are far from understood, but evidence strongly implicates the reversible oxidation of conserved cysteine residues within various proteins [19,20]. The aim of this review is to discuss the role of H_2O_2 as a cellular signal, paying particular attention to the role of Peroxiredoxin (PRDX) in transmitting H_2O_2 signals via various thiol-mediated mechanisms. Given the growing body of evidence that point to H_2O_2 in mediating redox signalling pathways during exercise, we suggest that PRDX may be an important transducer of exercise-induced H_2O_2 levels. A combination of study models will be reviewed, ranging from monolayer cell culture experiments to the few studies that have investigated PRDX in response to exercise in humans.

2. Cysteine thiol groups

Cysteine is one of the least abundant amino acids within the primary protein structure and contains a terminal sulphhydryl ($-SH$) or 'thiol' group that is highly electronegative in nature. As a result, the thiol group of many solvent accessible cysteines is a prime target for RONS. Reversible oxidation of the cysteine thiol or thiolate anion ($-S^-$; deprotonated thiol) can form stable inter or intra-molecular disulphides which can govern a broad range of cellular events e.g. metabolism [21] and signal transduction [22]. In this context, 2-electron oxidants (H_2O_2 or $ONOO^-$) in particular have been shown to oxidise critical redox active proteins or low molecular weight cellular thiols such as glutathione (GSH) [23]. In many cases, cysteine oxidation can change the structure and/or function of a redox active protein, evoking altered cellular and physiological responses [24–26].

It is well established that H_2O_2 lends itself to cellular signalling more so than the aforementioned RONS. The 1-electron oxidation of thiols by NO^{\bullet} and $O_2^{\bullet -}$ occurs at a fairly slow rate [23,27,28], forming highly unstable thiyl radicals ($-S^{\bullet}$). Oxidants like $ONOO^-$ can react directly with thiolates via a 2-electron mechanism [29], however its rapid reaction rate with carbon dioxide yields 1-electron oxidant products (i.e. carbonate and nitrogen dioxide radicals) that also form thiyl radicals [30,31]. Although the thiyl radical pathway can form mixed disulphides, its mechanism within a signalling cascade per se is questioned due to the lack of specificity of both the oxidants in question (i.e. $O_2^{\bullet -}$, NO^{\bullet} and $ONOO^-$) and the thiyl radical itself [23]. Conversely, H_2O_2 is a small, uncharged and membrane permeable RONS that can oxidise protein thiolates to form the intermediate sulfenic acid ($-SOH$) that then rapidly resolves with other reduced cysteine residues ($-SH$ or $-S^-$), forming inter or intra-molecular disulphide bonds [32]. H_2O_2 is highly oxidising due to the presence of a peroxide bond ($O-O$), however its chemical reduction can be limited by its high activation energy [33,34]. As a result, this gives H_2O_2

enormous selectivity over its reactions with protein thiol/thiolates [33]. Redox sensitive protein targets such as phosphatases (i.e. PTP1B [35,36] and PTEN [37,38]), kinases (i.e. ATM [39]) and transcription factors (i.e. STAT3 [22], Nrf-2 [40] and NF- κ B [41]) have cysteine thiols that can be specifically oxidised by H_2O_2 , implicating perturbations in H_2O_2 levels during exercise with signal transduction.

3. Cysteine targets for hydrogen peroxide during exercise

Studies in animals have demonstrated that H_2O_2 can oxidise critical cysteine thiols that facilitate muscle contraction [2,42,43]. In a study by Andrade et al., 1998, 4 min of H_2O_2 exposure (300 μ M) coupled to electrical stimulation (50 Hz, 350 ms duration) markedly increased the force output of isolated mouse skeletal muscle fibres, relative to a stimulation only trial [1]. Increased force output was suggested to be mediated by the glutathionylation of critical thiols within skeletal muscle tissue that improved calcium channel sensitivity [2]. Interestingly, exceeding a particular ' H_2O_2 threshold' (i.e. 8 min of H_2O_2 exposure) and also quenching H_2O_2 with a potent reducing agent (e.g. 1 mM dithiothreitol) both depressed force output significantly [1]. This highlights a critical balance of H_2O_2 required to optimally modulate thiol oxidation in skeletal muscle, and how 'oxidative' and 'reductive' stress may limit force output. H_2O_2 may also have a role in controlling blood flow during exercise by altering NO production via endothelial NOS (eNOS). There is evidence of H_2O_2 -mediated increases in eNOS activity and expression following acute [44] and long term aerobic exercise in animals respectively [45]. In addition, there is some evidence for a role of H_2O_2 in mediating the vasodilation of gluteal muscle microvasculature during resistance-based leg press exercise in humans [46]. The effect of H_2O_2 on vascular perfusion during exercise is likely via protein kinase G, which is known to be redox-sensitive [47]. Finally, H_2O_2 may have an important role in post-exercise metabolic adaptation by increasing the expression of redox-sensitive and thiol-rich transcription factors such as PGC-1 α [48] and FOXO3a [49]. It must be noted that relative to the spatio-temporal specificity of established signalling pathways (calcium signalling, G-proteins and phosphorylation); it is unclear how a particular H_2O_2 molecule may exert thiol specificity during and following exercise [22]. Moreover, the affinities of the many signalling proteins for H_2O_2 are extremely low [33], despite clear changes in downstream transcriptional activation that ultimately provide cross-resistance to H_2O_2 following exercise [49].

4. Hydrogen Peroxide as a cellular signal: recent advances

A variety of hypotheses have been proposed to explain how H_2O_2 can act as an intracellular signal. These have primarily focussed on mechanisms that might explain transient and localised accumulation of H_2O_2 through the inactivation of glutathione peroxidase (GPx), catalase and PRDX, or at sites where these proteins are not present [32]. In this context, H_2O_2 generating enzymes such as NADPH oxidase may co-localise with lower reacting thiols (i.e. kinases and phosphatases) to generate 'hot spots' of H_2O_2 , that permits the proteins' oxidation and thus cellular signalling (Fig. 1) [50,51]. The PRDX family of proteins in particular have received a great deal of attention with regards to their high abundance and catalytic turnover of H_2O_2 [33]. Briefly, post-translational modifications (i.e. serine and threonine phosphorylation [52], glutathionylation [53], tyrosine nitration [54], acetylation [55] or s-nitrosylation [56]) on non-catalytic amino acids sites of PRDX or over-oxidation of the active site thiol [57] can




Fig. 1. A theoretical model of how PRDX floodgate signalling may transduce exercise-induced H_2O_2 signals in skeletal muscle. NADPH oxidase translocation and subsequent H_2O_2 production may generate ‘hotspots’ of H_2O_2 that facilitate the oxidation of thiol-containing signalling proteins (e.g. a phosphatase), alongside the simultaneous inactivation of the PRDX thiolate. There is evidence that H_2O_2 -mediated over oxidation as well as post-translational modifications such as acetylation, phosphorylation (serine and threonine), glutathionylation, s-nitrosylation and tyrosine nitration of non-catalytic PRDX amino acids can prevent nucleophilic attack of the PRDX thiolate on H_2O_2 . In this context, the formation of stable PRDX dimers may facilitate extracellular redox-signalling. *Notes:* Thicker arrows represent the dominant pathway during floodgate signalling; flat ended arrows indicate inhibition of the PRDX thiolate; dashed circle represents a ‘hot spot’ of hydrogen peroxide. S-S is a disulphide bond within the oxidised PRDX dimer. *Abbreviations:* SOD: superoxide dismutase; O_2^- : superoxide; H_2O_2 : hydrogen peroxide; PRDX: peroxiredoxin; H_2O : water; -SOH: sulfenic acid; S^- : thiolate.

drive the accumulation of H_2O_2 in specific cellular domains, depending on the active PRDX isoform [58]. These modifications can inactivate PRDX thiol activity and conceivably redirect the initial H_2O_2 signal to elicit a change in cell function via oxidation of signalling thiol proteins such as PTP1B, PTEN or ATM [26]. This would render PRDX as a negative regulator of H_2O_2 signalling in a mechanism known as the ‘floodgate model’ [59]. However, a recent paper by Sobotta et al. [22] eloquently demonstrated that PRDX has a ‘direct’ role in translating the oxidizing equivalent of H_2O_2 to the cysteine rich transcription factor STAT3. Following H_2O_2 exposure, the -SOH intermediate of PRDX-2 was shown to form a mixed disulphide with STAT3 directly, initiating its translocation to the nucleus. This implies that PRDX may act as an active ‘redox relay’ with respect to increased cellular H_2O_2 levels. The interplay between PRDX floodgate signalling and active redox relays are unclear, however the mechanisms are likely not mutually exclusive [60].

5. Peroxiredoxin: an abundant and highly active peroxidase

An array of factors can influence the capacity of a given redox active cysteine to reduce H_2O_2 by nucleophilic attack (Fig. 2). Accessibility of H_2O_2 to the catalytic thiol/thiolate motif and structural factors (e.g. adjacent neighbouring amino acids on the polypeptide chain) that affect electron density can alter the midpoint potential (E_m) of the thiol. Cysteine residues that have a lower E_m are more readily oxidised [24,61] and even alternative isoforms of the same redox active protein may contain cysteine residues with differing E_m and thus alternative propensities for oxidation [62]. The sensitivity of a redox active cysteine to oxidation is also, in part, determined by pH, which relates to the solution and microenvironment in which that cysteine resides.

Acid residues adjacent to a cysteine have been shown to alter sensitivity of that cysteine to oxidation presumably via thiol protonation (S^- to -SH) [63]. Whether a redox cysteine is protonated at physiological pH is determined by the pK_a of the thiol group [24]. All of these factors explain the broad range of sensitivities exhibited by thiol-based proteins to fluctuations in H_2O_2 concentrations. In particular, PRDX has a turnover of H_2O_2 up to a million times greater than protein phosphatases such as PTP1B and cdc25b in vitro [23,33].

PRDX is a ubiquitously expressed oxidoreductase protein (160–220 amino acids) located in the cytosolic (isoforms I, II and VI), endoplasmic reticulum (isoform IV) and mitochondrial (isoforms III and V) domains of the cell [64]. The nascent form of most PRDXs is the decamer form (I–IV), with PRDX V and VI unable to form oligomers. The ‘catalytic’ cysteine of all PRDXs (S^-) can convert H_2O_2 [65], ONOO^- [66] and other peroxide substrates [65] to H_2O via the oxidation of its conserved thiolate to a -SOH intermediate, before reacting with a ‘resolving’ cysteine thiol (-SH). In this regard, the S^- form acts as the ‘redox-sensor’ via nucleophilic attack and is the target for oxidation, whereas the -SH form is resolving in nature [67,68]. The mechanism of -SOH resolution determines the sub-classes of the PRDX family. These include typical-2 cysteine PRDX (I–IV), atypical-2 cysteine PRDX (V) or 1-cysteine (VI) PRDX, whereby mixed disulphides are formed through inter-molecular bonding with a neighbouring thiol (PRDX molecule or thiol-based protein), intra-molecular bonding with a native thiol or inter-molecular bonding with GSH [69] respectively. These disulphide bonds are reduced by the antioxidants TRX (I–IV), GSH-S-Transferase (V) and GSH (VI) in bioenergetically favoured reactions [69,70].

Other antioxidant enzymes such as catalase, GSH and GPx have prominent and defined roles in H_2O_2 catalysis [71]. Importantly, these enzymes likely work in synergy with PRDX to modulate the




Fig. 2. A simplified schematic of factors affecting protein thiol activity with H_2O_2 . The rate constants for selected thiol proteins are indicated [23,33]. The interactions between the 5 factors presented are complex and therefore scales are not quantified: (1) microenvironment, e.g. pH – a low pH will protonate the thiolate and potentially decrease reactivity; (2) structural arrangement – neighbouring amino acids that have a δ^- (aspartic acid/glutamic acid) charge will compete for electrons within the polypeptide chain, whereas δ^+ amino acids (arginine, lysine and histidine) will permit high electron density around the thiol/thiolate, thus altering thiol/thiolate sensitivity to oxidation. (3) *pKa* – a low *pKa* value will cause deprotonation of the thiol to form a negatively charged thiolate anion, which potentially increases reactivity; (4) mid-point potential – a lower mid-point potential will increase the capacity of the thiol/thiolate to become oxidised, thus increasing thiol/thiolate activity; (5) solvent accessibility – surface thiol/thiolates within the quaternary structure of a protein will allow more rapid nucleophilic attack on H_2O_2 , whereas embedded thiol/thiolates are less easily oxidised. Notes: the dotted grey boxes indicate electron density around the thiolate. Thick black arrow indicates a greater turnover of H_2O_2 relative to a thinner arrow.

overall peroxide signal [72]. Catalase is primarily located to peroxisomes where H_2O_2 is formed from the breakdown of various substrates by flavoprotein oxidases [71], and therefore has no role in the transfer of cytosolic to nucleic peroxide signals. GSH is a low molecular weight antioxidant with a relatively low turnover of H_2O_2 [33], and unlikely acts a direct ‘sensor’ for translating the peroxide signal for transcriptional activation.¹ GPx has a comparable rate constant to PRDX for H_2O_2 catalysis [33]; however the transient and highly reductive selenenic acid (Se^-) intermediate that is formed following GPx reaction with H_2O_2 rapidly resolves with a native amide group to form an intramolecular sulfenamide [73]. Therefore GPx does not act as a dimerising peroxide ‘sensor’ protein. It is the structure and high affinity that PRDX has for H_2O_2 to favourably form $-SOH$ intermediates and resolving dimers that may give this antioxidant protein a unique role in actively transducing peroxide signals into a dynamic biological response. Ubiquitous PRDX expression has been estimated to be far more abundant than both GPx and catalase [74] and the decamer conformation of typical 2-cysteine PRDXs can stabilise the active site and increase peroxidase activity [75]. Further, structural analysis of PRDX has revealed a hydrogen-bonding network surrounding the active site that favours peroxide substrate binding and catalysis [76,77]. A deprotonated catalytic thiolate (due to a low *pKa* value)

coupled with favoured polarisation of the peroxide bond (O–O) has led to estimates that PRDX reacts with up to 99% of cytosolic H_2O_2 [78,79].

6. Is PRDX a peroxide sensing protein during and following exercise?

The available evidence suggests that H_2O_2 plays a crucial role in mediating tissue function during exercise [2], as well as modulating post-exercise metabolic adaptation [49] via redox-sensitive pathways. Despite this, the direct reactivity of H_2O_2 with signalling protein thiols is known to be very low. Given PRDXs high abundance and turnover of cytosolic H_2O_2 , PRDX would likely favour the formation of $-SOH$ intermediates in response to heightened H_2O_2 production during exercise. Indeed, accumulating evidence suggests that cytosolic, rather than mitochondrial sources of RONS predominate during exercise [80,81]. Oxidation of PRDX may therefore act as a physiologically conserved mechanism for translating contractile signals from exercise-induced oxidants to downstream transcription factors, thus facilitating the adaptive response to exercise.

The targets of PRDX $-SOH$ intermediates are to date, largely unclear, with no studies exploring this in the context of exercise. It has been demonstrated *in vitro* that transcription factors associated with exercise adaptation, namely, STAT3 [22] and p38 [82] are responsive to this signalling pathway. Following H_2O_2

¹ It is important to note that glutathionylation is an important post-translational modification whereby GSH resolves with protein $-SOH$ or S^- directly (e.g. following oxidation by RONS) to modify protein function and thus cell signalling.

exposure, PRDX-2 can form mixed disulphides with STAT3 directly [22], whereas PRDX-1 resolves with ASK-1, subsequently increasing p38 phosphorylation [82]. It is well documented that STAT3 [83] and p38 [84] activation are linked with increased muscle anabolism and mitochondrial biogenesis following exercise respectively. It is important to note that PRDX has been shown to interact with various other signalling complexes (e.g. platelet-derived growth factors [85], receptor tyrosine kinases [52] and lipid phosphatases [37]) that are undoubtedly involved with signalling pathways following exercise. The translocation of PRDX to these receptors has been previously regarded as a mechanism to prevent localised protein oxidation via its reduced thiolate. However, advances in our understanding of how PRDX –SOH resolve with signalling proteins may question this viewpoint. Given that there is no known receptor for H₂O₂ binding, highly reactive and specific cysteine targets such as PRDX may offer a control point for managing H₂O₂ gradients with targeted precision in response to a bout of exercise.

Increased levels of oxidised PRDX (I–IV) dimers [86] and over-oxidised monomers [70,87,88] have been reported in immune cells and erythrocytes following exercise. Under cellular oxidative stress, the PRDX decamer can expose an oxidised cysteine that resolves with a neighbouring PRDX thiol to form a stable oxidised dimer. This state is known to be favoured following PRDX oxidation [89] and levels have been shown to increase following a single bout of ultra-endurance exercise (174 km, 30–44 h) in isolated PBMCs from well-trained male participants [87]. Recent work by Salzano et al. [53] has indicated that glutathionylation of non-catalytic cysteine residues may facilitate extracellular secretion of dimerised PRDX from immune cells, suggesting that this may act as an extracellular ‘stress’ signal. Interestingly, there is also evidence that skeletal muscle cells can also increase their secretion of PRDX in response to injury [90]. In this regard, extracellular PRDX may act in a paracrine or hormonal manner between cells under redox stress during exercise. Furthermore, PRDX-2 has been shown to bind to toll-like receptor-4 in immune cells, increasing inflammatory cytokine transcription via NF-κB (i.e. IL-1β) [91]. This indicates an additional aspect of extracellular communication under cellular redox stress, with immune cells known to target, infiltrate and repair skeletal muscle following exercise.

A unique feature of the PRDX catalytic cysteine is the capacity of the thiolate to react with a second and third H₂O₂ molecule, leading to sulfinic (–SO₂H) and sulfonic (SO₃H) acid oxidation states [92]. This ‘over-oxidation’ occurs at a rate too quickly for thiol ‘resolution’ and leads to the formation of over-oxidised monomers. Over-oxidised PRDX monomers have been reported during and following exercise in human peripheral blood mononuclear cells (PBMCs) [93,87] and erythrocytes respectively [88]. Formation of over-oxidised PRDX in PBMCs (I–IV isoforms) has been shown to be dependent on the intensity of exercise, with heightened peroxide concentrations during high intensity exercise (80% maximal oxygen consumption [$\dot{V}O_{2max}$] vs. 60% $\dot{V}O_{2max}$) likely exceeding the reduction power of TRX, the exclusive reductant of the PRDX (I–IV) disulphide [93]. As well as increased peroxide levels during exercise, these changes may also relate, in part, to reductions in cellular pH that might reduce the sensitivity of the TRX cysteine to oxidation. As introduced earlier, this ‘floodgate model’ mechanism (Fig. 1) may allow accumulation of H₂O₂ that permits oxidation of other signalling thiol proteins, for example PTP1B, which has a much lower *pKa* value than TRX [33]. Moreover, *in vitro* evidence suggests that TRX may then redirect its reducing power to transcription factors such as NF-κB [94] and AP-1 [57], eliciting changes in cell function in a different micro-environment (i.e. higher pH).

The research assessing the interplay between H₂O₂ and PRDX in response to acute exercise in humans is extremely limited, with

descriptive changes in PRDX expression and oxidation only previously monitored [93,87,95]. Mechanistic approaches in knockout mouse models have highlighted a clear and prominent role for mitochondrial PRDX-3 in controlling skeletal muscle force production [96] and mitochondrial homeostasis [97]. For example, absence of PRDX-3 caused deregulation of mitochondrial membrane potential and a faster rate of muscle fatigue in the extensor digitorum longus and soleus muscles of mice, possibly as a result of elevated peroxide concentrations [96]. A study by Kil et al., 2012 provided evidence to suggest that PRDX-III hyperoxidation has an important role in modulating steroid hormone production following physiological stress in mice [26], which will undoubtedly be important in an exercise context whereby steroidogenesis is elevated post-exercise [98]. These studies do not provide insight into the specific sources of RONS that mediate PRDX cysteine oxidation, nor the associated downstream signalling mechanisms. The only study to assess changes in PRDX following regular exercise in humans, reported an increase in erythrocyte PRDX-2 expression in overweight males after 3-months of aerobic exercise training (3 sessions per week at 75% of maximum heart rate; progressive increase in session duration (25–50 min) over the 3 month period) [95]. Increased PRDX expression following exercise indicates an important role for PRDX in cellular remodelling following exercise training, which likely occurs via the redox-sensitive transcription factor NF-κB (as recently demonstrated *in vitro* [72]). Interestingly, in a non-exercise context, PRDX has also been shown to have an active role in the progression of mitosis through selective phosphorylation of PRDX-1 during anaphase [99]. Phosphorylated and inactivated PRDX-1 allows the transient and localised accumulation of H₂O₂ that oxidises centrosome-bound phosphates, thus permitting cell proliferation. This highlights a complex interplay between thiol modifications and cellular phosphorylation in mediating cell growth, which may have applications to tissue remodelling (i.e. skeletal muscle) in response to exercise.

7. Future Perspectives

The redox environment within cells is a complex network of highly transient RONS that work in strict cooperation with cellular and dietary antioxidants. Despite rapid progression in our understanding of many aspects of redox signalling, it is clear that the analytical techniques currently available to monitor thiol modifications following exercise in humans are limited. Technologies for the evaluation of –SOH formation *in vitro* and *in vivo* have been developed, which include dimedone based reagents and conjugates [100]. These reagents ‘trap’ –SOH formation in real time, which can successfully be detected by mass-spectrometry [101,102]. The use of such reagents has real potential in the field of exercise physiology, particularly regarding the delineation of redox sensing and/or signalling pathways where PRDX and H₂O₂ are implicated. The evidence presented in this article highlights three primary functions of PRDX in managing H₂O₂ gradients: (1) an antioxidant with a very high affinity for H₂O₂, (2) an intracellular signal that transfers the oxidising equivalent of H₂O₂ to cysteine rich target proteins via its –SOH intermediate and (3) an extracellularly secreted protein that may permit local signalling. These signalling properties may well underpin a fundamental aspect of redox communication in response to exercise (Fig. 3).

At present, it is unclear how H₂O₂ acts as an intracellular signal *in vivo* and importantly, how widespread H₂O₂ ‘sensor’ proteins are within cells. A recent review by Cobley et al., [103] suggested that H₂O₂ contained to the cellular domain of production during exercise (by proteins such as PRDX) likely permits its actions as an intracellular signal, whereas distal diffusion permits non-specific




Fig. 3. A theoretical model of how PRDX could transduce exercise-induced H_2O_2 signals in active tissues during exercise. Note that for simplicity; the reaction of PRDX with $ONOO^-$ and other lipid peroxides has not been included. (1) The PRDX thiolate has a very high turnover of H_2O_2 relative to other signalling thiol proteins, giving this protein a prominent antioxidant role; (2) PRDX can also act as an intracellular and potentially an (3) extracellular signal through its sulfenic acid intermediate ($-SOH$) dissociating from the PRDX decamer after oxidation. The PRDX $-SOH$ can resolve with a signalling thiol protein directly or a neighbouring PRDX thiol that forms a dimer which can be secreted from the cell. Over-oxidation of PRDX may permit floodgate signalling via various mechanisms, including TRX translocation to the nucleus (*). Note that a decamer PRDX $-SOH$ can react with a neighbouring PRDX dimer to form a dodecameric structure that has been associated with protein chaperone activity [89]. Notes: Black arrows indicate the possible signalling properties of PRDX during exercise; dashed arrows indicate reduction reactions that modulate these pathways; light blue fill in box indicates oxidation of protein; purple fill in box indicates a reduced protein; red fill in box indicates other important antioxidant enzymes; green fill in box indicates the target thiol signalling protein; *over-oxidised PRDX monomers – these indirectly permit TRX to translocate to the nucleus and reduce transcription factors (floodgate model); S-S is a disulphide bond between oxidised PRDX cysteines. Abbreviations: PRDX: peroxiredoxin; TRX: thioredoxin; TRX-R: thioredoxin reductase; SRX: sulfiredoxin; SH: sulphur hydryl; $-SOH$: sulfenic acid; sulfinic acid; sulfonic acid; S-S: disulphide bond; NADPH: nicotinamide dinucleotide phosphate hydrogenase; SOD: superoxide dismutase; O_2^- : superoxide; H_2O_2 : hydrogen peroxide; FF: Fully Folded.

oxidation of biomolecules through reactions with free iron. At this stage, we cannot be certain that H_2O_2 acts solely as an intracellular signal. Indeed, H_2O_2 is a freely diffusible, uncharged molecule that may signal in an autocrine, paracrine or hormonal manner between cells during exercise [104]. This may also be spatially favoured given that membrane-bound NADPH oxidases are likely a source of O_2^- during exercise [105]. Similarly, the function of decamer PRDX, a product of high levels of peroxides such as H_2O_2 , needs further exploration as a possible extracellular redox signal.

The role of other 2-electron oxidants (i.e. $ONOO^-$) and even free radical species (i.e. 1-electron oxidants such as O_2^- and NO^*) in thiol-mediated redox signalling following exercise must not be discounted; particularly if the source of RONS is localised to the target protein [23]. Furthermore, other thiol modifications (i.e. s-nitrosylation and s-glutathionylation) may be responsible for mediating the adaptive response to exercise, with the role of reactive sulphur species (i.e. H_2S/S^-) in forming mixed persulphides emerging as a post-translational modification likely governing physiological effects [106].

8. Conclusion

PRDX is an abundant cellular protein with a diverse range of functions in mammalian cells, above and beyond its fundamental function as a thiol peroxidase. This review has drawn on a series of excellent advances in redox biology research to highlight the need to dissect the molecular mechanisms underpinning redox-mediated signal transduction in response to exercise. Future studies need to elucidate the precise role of PRDX and other peroxide sensors in transmitting H_2O_2 signals into dynamic biological responses during and following exercise.

References

- [1] F.H. Andrade, M.B. Reid, D.G. Allen, H. Westerblad, Effect of hydrogen peroxide and dithiothreitol on contractile function of single skeletal muscle fibres from the mouse, *J. Physiol.* 509 (1998) 565–575.
- [2] G.D. Lamb, H. Westerblad, Acute effects of reactive oxygen and nitrogen species on the contractile function of skeletal muscle, *J. Physiol.* 589 (2011) 2119–2127, <http://dx.doi.org/10.1113/jphysiol.2010.199059>.
- [3] P. Donoso, C. Hidalgo, G. Sánchez, Posttranslational modifications of cardiac ryanodine receptors/calcium release channels by reactive oxygen species

- (ROS) and reactive nitrogen species (RNS), *Syst. Biol. Free Radic. Antioxid.* 46 (2014) 1031–1046.
- [4] M.J. Jackson, D.A. Jones, R.H. Edwards, Vitamin E and skeletal muscle, *Ciba Found. Symp.* 101 (1983) 224–239.
 - [5] M.J. Jackson, M. Khassaf, A. Vasilaki, et al., Vitamin E and the oxidative stress of exercise, *Ann. N. Y. Acad. Sci.* 1031 (2004) 158–168, <http://dx.doi.org/10.1196/annals.1331.015>.
 - [6] M.C. Gomez-Cabrera, E. Domenech, M. Romagnoli, et al., Oral administration of vitamin C decreases muscle mitochondrial biogenesis and hampers training-induced adaptations in endurance performance, *Am. J. Clin. Nutr.* 87 (2008) 142–149.
 - [7] D.C. Salo, C.M. Donovan, K.J.A. Davies, HSP70 and other possible heat shock or oxidative stress proteins are induced in skeletal muscle, heart, and liver during exercise, *Free Radic. Biol. Med.* 11 (1991) 239–246.
 - [8] M. Khassaf, R.B. Child, A. McArdle, et al., Time course of responses of human skeletal muscle to oxidative stress induced by nondamaging exercise, *J. Appl. Physiol.* 90 (2001) 1031–1035.
 - [9] M. Khassaf, A. McArdle, C. Esanu, et al., Effect of vitamin C supplements on antioxidant defence and stress proteins in human lymphocytes and skeletal muscle, *J. Physiol.* 549 (2003) 645–652.
 - [10] S.K. Powers, E.E. Talbert, P.J. Adhithetty, Reactive oxygen and nitrogen species as intracellular signals in skeletal muscle, *J. Physiol.* 589 (2011) 2129–2138, <http://dx.doi.org/10.1113/jphysiol.2010.201327>.
 - [11] M.J. Jackson, D. Pye, J. Palomero, The production of reactive oxygen and nitrogen species by skeletal muscle, *J. Appl. Physiol.* 102 (2007) 1664–1670, <http://dx.doi.org/10.1152/jappphysiol.01102.2006> [pii].
 - [12] J. St-Pierre, J.A. Buckingham, S.J. Roebuck, M.D. Brand, Topology of superoxide production from different sites in the mitochondrial electron transport chain, *J. Biol. Chem.* 277 (2002) 44784–44790, <http://dx.doi.org/10.1074/jbc.M207217200>.
 - [13] S. Di Meo, P. Venditti, Mitochondria in exercise-induced oxidative stress, *Biol. Signals Recept.* 10 (2001) 125–140.
 - [14] F. McArdle, D.M. Pattwell, A. Vasilaki, et al., Intracellular generation of reactive oxygen species by contracting skeletal muscle cells, *Free Radic. Biol. Med.* 39 (2005) 651–657, <http://dx.doi.org/10.1016/j.freeradbiomed.2005.04.010>.
 - [15] T.W. Balon, J.L. Nadler, Nitric oxide release is present from incubated skeletal muscle preparations, *J. Appl. Physiol.* 77 (1994) 2519–2521.
 - [16] D.M. Pattwell, A. McArdle, J.E. Morgan, et al., Release of reactive oxygen and nitrogen species from contracting skeletal muscle cells, *Free Radic. Biol. Med.* 37 (2004) 1064–1072.
 - [17] J.S. Beckman, W.H. Koppenol, Nitric oxide, superoxide, and peroxynitrite: the good, the bad, and ugly, *Am. J. Physiol.* 271 (1996) C1424–C1437.
 - [18] L.R. Silveira, C.J. Lucia Pereira-Da-Silva, Y. Hellsten, Formation of hydrogen peroxide and nitric oxide in rat skeletal muscle cells during contractions, *Free Radic. Biol. Med.* 35 (2003) 455–464.
 - [19] B. McDonagh, G.K. Sakellariou, N.T. Smith, et al., Differential cysteine labeling and global label-free proteomics reveals an altered metabolic state in skeletal muscle aging, *J. Proteome Res.* 13 (2014) 5008–5021.
 - [20] J.N. Cobley, H. McHardy, J.P. Morton, et al., Influence of vitamin C and vitamin E on redox signalling: implications for exercise adaptations, *Free Radic. Biol. Med.* 84 (2015) 65–76, <http://dx.doi.org/10.1016/j.freeradbiomed.2015.03.018>.
 - [21] V.L. Kolossov, J.N. Beaudoin, N. Prabhu Ponnuraj, Thiol-based antioxidants elicit mitochondrial oxidation via respiratory complex III, *Am. J. Physiol. Cell Physiol.* 309 (2) (2015) C81–91.
 - [22] M.C. Sobotta, W. Liou, S. Stöcker, et al., Peroxiredoxin-2 and STAT3 form a redox relay for H₂O₂ signaling, *Nat. Chem. Biol.* 11 (2014) 64–70, <http://dx.doi.org/10.1038/nchembio.1695>.
 - [23] C.C. Winterbourn, Are free radicals involved in thiol-based redox signaling? *Free Radic. Biol. Med.* 80 (2015) 164–170, <http://dx.doi.org/10.1016/j.freeradbiomed.2014.08.017>.
 - [24] L.B. Poole, The basics of thiols and cysteines in redox biology and chemistry, *Free Radic. Biol. Med.* 80 (2015) 148–157, <http://dx.doi.org/10.1016/j.freeradbiomed.2014.11.013>.
 - [25] C. Berndt, C.H. Lillig, A. Holmgren, Thiol-based mechanisms of the thiorodoxin and glutaredoxin systems: implications for diseases in the cardiovascular system, *Am. J. Physiol. Heart Circ. Physiol.* 292 (2007) H1227–H1236, <http://dx.doi.org/10.1152/ajpheart.01162.2006> [pii].
 - [26] I.S. Kil, S.K. Lee, K.W. Ryu, et al., Feedback control of adrenal steroidogenesis via H₂O₂-dependent, reversible inactivation of peroxiredoxin III in mitochondria, *Mol. Cell* 46 (2012) 584–594, <http://dx.doi.org/10.1016/j.molcel.2012.05.030>.
 - [27] C.M. Jones, A. Lawrence, P. Wardman, M.J. Burkitt, Electron paramagnetic resonance spin trapping investigation into the kinetics of glutathione oxidation by the superoxide radical: re-evaluation of the rate constant, *Free Radic. Biol. Med.* 32 (2002) 982–990.
 - [28] J.S. Stampler, O. Jaraki, J. Osborne, et al., Nitric oxide circulates in mammalian plasma primarily as an S-nitroso adduct of serum albumin, *Proc. Natl. Acad. Sci. USA* 89 (1992) 7674–7677.
 - [29] C. Quijano, B. Alvarez, R.M. Gatti, et al., Pathways of peroxynitrite oxidation of thiol groups, *Biochem. J.* 322 (1997) 167–173.
 - [30] S. Carballeda, S. Bartsagha, R. Rafael, Kinetic and mechanistic considerations to assess the biological fate of peroxynitrite, *Biochim. Biophys. Acta: Gen. Subj.* 1840 (2014) 768–780.
 - [31] G. Ferrer-Sueta, R. Radi, Chemical biology of peroxynitrite: kinetics, diffusion, and radicals, *ACS Chem. Biol.* 4 (2009) 161–177.
 - [32] C.C. Winterbourn, M.B. Hampton, Signaling via a peroxiredoxin sensor, *Nat. Publ. Gr.* 11 (2014) 5–6, <http://dx.doi.org/10.1038/nchembio.1722>.
 - [33] C.C. Winterbourn, The biological chemistry of hydrogen peroxide, 1st ed. *Methods in Enzymology*, 2013. Doi: 10.1016/B978-0-12-405881-1.00001-X.
 - [34] A. Zeida, R. Babbush, M.C.G. Lebrero, et al., Molecular basis of the mechanism of thiol oxidation by hydrogen peroxide in aqueous solution: challenging the SN2 paradigm, *Chem. Res. Toxicol.* 25 (2012) 741–746.
 - [35] T.C. Meng, D. Buckley, S. Galic, et al., Regulation of insulin signaling through reversible oxidation of the protein-tyrosine phosphatases TC45 and PTP1B, *J. Biol. Chem.* 279 (2004) 37716–37725, <http://dx.doi.org/10.1074/jbc.M404606200>.
 - [36] S.R. Lee, K.S. Yang, J. Kwon, et al., Reversible inactivation of the tumor suppressor PTEN by H₂O₂, *J. Biol. Chem.* 277 (2002) 20336–20342.
 - [37] J. Cao, J. Schulte, A. Knight, et al., Prdx1 inhibits tumorigenesis via regulating PTEN/AKT activity, *EMBO J.* 28 (2009) 1505–1517, <http://dx.doi.org/10.1038/emboj.2009.101>.
 - [38] S.R. Lee, K.S. Kwon, S.R. Ki, S.G. Rhee, Reversible inactivation of protein-tyrosine phosphatase 1B in A431 cells stimulated with epidermal growth factor, *J. Biol. Chem.* 273 (1998) 15366–15372.
 - [39] Z. Guo, S. Kozlov, M.F. Lavin, et al., ATM activation by oxidative stress, *Science* 80–330AD (2010) 517–521.
 - [40] X. He, Q. Ma, NRF2 cysteine residues are critical for oxidant/electrophile-ubiquitination-proteasomal degradation, and transcription activation, *76 (2009) 1265–1278*, <http://dx.doi.org/10.1124/mol.109.058453.The>.
 - [41] G. Gloire, S. Legrand-Poels, J. Piette, NF-kappaB activation by reactive oxygen species: fifteen years later, *Biochem. Pharmacol.* 72 (2006) 1493–1505.
 - [42] L.F. Ferreira, M.B. Reid, Muscle-derived ROS and thiol regulation in muscle fatigue, *J. Appl. Physiol.* 104 (2008) 853–860, <http://dx.doi.org/10.1152/jappphysiol.00953.2007>.
 - [43] M. Reid, Free radicals and muscle fatigue: of ROS, canaries, and the IOC, *Free Radic. Biol. Med.* 44 (2008) 169–179.
 - [44] L.Y. Tanaka, L.R.G. Bechara, A.M. dos Santos, et al., Exercise improves endothelial function: a local analysis of production of nitric oxide and reactive oxygen species, *Nitric Oxide* 45 (2015) 7–14, <http://dx.doi.org/10.1016/j.niox.2015.01.003>.
 - [45] N. Lauer, T. Suvorava, U. Rütger, et al., Critical involvement of hydrogen peroxide in exercise-induced up-regulation of endothelial NO synthase, *Cardiovasc. Res.* 65 (2005) 254–262.
 - [46] M.J. Durand, K. Dharmashankar, J.-T. Bian, et al., Acute exertion elicits a H₂O₂-dependent vasodilator mechanism in the microvasculature of exercise-trained but not sedentary adults, *Hypertension* 65 (2015) 140–145.
 - [47] B.H. Neo, S. Kandhi, M.S. Wolin, Roles for soluble guanylate cyclase and a thiol oxidation-elicited subunit dimerization of protein kinase G in pulmonary artery relaxation to hydrogen peroxide, *Am. J. Physiol. Heart Circ. Physiol.* 299 (2010) H1235–H1241, <http://dx.doi.org/10.1152/ajpheart.00513.2010>.
 - [48] L.R. Silveira, H. Pilegaard, K. Kusuhara, et al., The contraction induced increase in gene expression of peroxisome proliferator-activated receptor (PPAR)-γ coactivator 1α (PGC-1α), mitochondrial uncoupling protein 3 (UCP3) and hexokinase II (HKII) in primary rat skeletal muscle cells is dependent on re, *Biochim. Biophys. Acta: Mol. Cell Res.* 1763 (2006) 969–976, <http://dx.doi.org/10.1016/j.bbamcr.2006.06.010>.
 - [49] P. Wang, C.G. Li, Z. Qi, et al., Acute exercise induced mitochondrial H₂O₂ production in mouse skeletal muscle: association with p^{66shc} and FOXO3a signaling and antioxidant enzymes, *Oxid. Med. Cell. Longev.* (2014) 1–10.
 - [50] C. Chen, T. Cheng, H. Lin, et al., Reactive oxygen species generation is involved in epidermal growth factor receptor transactivation through the transient oxidization of Src homology 2-containing tyrosine phosphatase in endothelin-1 signaling pathway in rat cardiac fibroblasts, *Mol. Pharmacol.* 69 (2006) 1347–1355, <http://dx.doi.org/10.1124/mol.105.017558.by>.
 - [51] M. Ushio-Fukai, Compartmentalization of redox signaling through NADPH oxidase-derived ROS, *Antioxid. Redox Signal.* 11 (2009) 1289–1299.
 - [52] H.A. Woo, S.H. Yim, D.H. Shin, et al., Inactivation of peroxiredoxin I by phosphorylation allows localized H₂O₂ accumulation for cell signaling, *Cell* 140 (2010) 517–528, <http://dx.doi.org/10.1016/j.cell.2010.01.009>.
 - [53] S. Salzano, P. Checconi, E.-M. Hanschmann, et al., Linkage of inflammation and oxidative stress via release of glutathionylated peroxiredoxin-2, which acts as a danger signal, *Proc. Natl. Acad. Sci. USA* 111 (2014) 12157–12162, <http://dx.doi.org/10.1073/pnas.1401712111>.
 - [54] L.M. Randall, B. Manta, M. Hugo, et al., Nitration transforms a sensitive peroxiredoxin 2 into a more active and robust peroxidase, *J. Biol. Chem.* 289 (2014) 15536–15543, <http://dx.doi.org/10.1074/jbc.M113.539213>.
 - [55] Y. Pan, J.-H. Jin, Y. Yu, J. Wang, Significant enhancement of hPrx1 chaperone activity through lysine acetylation, *Chembiochem* 15 (2014) 1773–1776, <http://dx.doi.org/10.1002/cbic.201402164>.
 - [56] J. Fang, T. Nakamura, D.-H. Cho, et al., S-nitrosylation of peroxiredoxin 2 promotes oxidative stress-induced neuronal cell death in Parkinson's disease, *Proc. Natl. Acad. Sci. USA* 104 (2007) 18742–18747, <http://dx.doi.org/10.1073/pnas.0705904104>.
 - [57] A.M. Day, J.D. Brown, S.R. Taylor, et al., Inactivation of a peroxiredoxin by hydrogen peroxide is critical for thioredoxin-mediated repair of oxidized proteins and cell survival, *Mol. Cell* 45 (2012) 398–408, <http://dx.doi.org/10.1016/j.molcel.2011.11.027>.
 - [58] A. Perkins, L.B. Poole, P.A. Karplus, Tuning of peroxiredoxin catalysis for

- various physiological roles, *Biochemistry* 53 (2014) 7693–7705.
- [59] Z.A. Wood, L.B. Poole, P.A. Karplus, Peroxiredoxin evolution and the regulation of hydrogen peroxide signaling, *Science* 300 (2003) 650–653, <http://dx.doi.org/10.1126/science.1080405>.
- [60] A. Perkins, K.J. Nelson, D. Parsonage, et al., Peroxiredoxins: guardians against oxidative stress and modulators of peroxide signaling, *Trends Biochem. Sci.* 40 (2015) 435–445, <http://dx.doi.org/10.1016/j.tibs.2015.05.001>.
- [61] F.Q. Schafer, G.R. Buettner, Redox environment of the cell as viewed through the redox state of the glutathione disulfide/glutathione couple, *Free Radic. Biol. Med.* 30 (2001) 1191–1212.
- [62] S.J. Coles, J.T. Hancock, M.E. Conway, Differential redox potential between the human cytosolic and mitochondrial branched-chain aminotransferase, *Acta Biochim. Biophys. Sin.* 44 (2012) 172–176, <http://dx.doi.org/10.1093/abbs/gmr103>.Advance.
- [63] A.T.P. Carvalho, P.A. Fernandes, M.J. Ramos, Determination of the DeltapKa between the active site cysteines of thioredoxin and DsbA, *J. Comput. Chem.* 27 (2006) 966–975, <http://dx.doi.org/10.1002/jcc.20404>.
- [64] Z.A. Wood, L.B. Poole, P.A. Karplus, Peroxiredoxin evolution and the regulation of hydrogen peroxide signaling, *Science* 300 (2003) 650–653, <http://dx.doi.org/10.1126/science.1080405>.
- [65] Z.A. Wood, E. Schröder, J.R. Harris, L.B. Poole, Structure, mechanism and regulation of peroxiredoxins, *Trends Biochem. Sci.* 28 (2003) 32–40, [http://dx.doi.org/10.1016/S0968-0004\(02\)00003-8](http://dx.doi.org/10.1016/S0968-0004(02)00003-8).
- [66] R. Bryk, P. Griffin, C. Nathan, Peroxynitrite reductase activity of bacterial peroxiredoxins, *Nature* 407 (2000) 211–215.
- [67] N. Fujiwara, T. Fujii, J. Fujii, N. Taniguchi, Roles of N-terminal active cysteines and C-terminal cysteine-selenocysteine in the catalytic mechanism of mammalian thioredoxin reductase, *J. Biochem.* 129 (2001) 803–812.
- [68] C. Klomsiri, P.A. Karplus, L.B. Poole, Cysteine-based redox switches in enzymes, *Antioxid. Redox Signal.* 14 (2011) 1065–1077, <http://dx.doi.org/10.1089/ars.2010.3376>.
- [69] J.R. Pedrajas, B. McDonagh, F. Hernández-Torres, et al., Glutathione is the resolving thiol for thioredoxin peroxidase activity of 1-cys peroxiredoxin without being consumed during the catalytic cycle, *Antioxid. Redox Signal.* (2015), in press.
- [70] A.J. Wadley, Y.W. Chen, S.J. Bennett, et al., Monitoring changes in thioredoxin and over-oxidised peroxiredoxin in response to exercise in humans, *Free Radic. Res.* 49 (2015) 290–298, <http://dx.doi.org/10.3109/10715762.2014.1000890>.
- [71] N.A. Bonekamp, A. Völkl, H.D. Fahimi, M. Schrader, Reactive oxygen species and peroxisomes: Struggling for balance, *BioFactors* 35 (2009) 346–355, <http://dx.doi.org/10.1002/biof.48>.
- [72] H. Won, S. Lim, M. Jang, et al., Peroxiredoxin-2 upregulated by NF-κB attenuates oxidative stress during the differentiation of muscle-derived C2C12 cells, *Antioxid. Redox Signal.* 16 (2012) 245–261.
- [73] H.S. Marinho, C. Real, L. Cyrne, et al., Hydrogen peroxide sensing, signaling and regulation of transcription factors, *Redox Biol.* 2 (2014) 535–562, <http://dx.doi.org/10.1016/j.redox.2014.02.006>.
- [74] S. Ghaemmaghami, W. Huh, K. Bower, et al., Global analysis of protein expression in yeast, *Nature* 425 (2003) 737–741.
- [75] A. Perkins, K.J. Nelson, D. Parsonage, et al., Peroxiredoxins: guardians against oxidative stress and modulators of peroxide signaling, *Trends Biochem. Sci.* 40 (8) (2015) 1–11, <http://dx.doi.org/10.1016/j.tibs.2015.05.001>.
- [76] D. Parsonage, K.J. Nelson, G. Ferrer-sueta, S. Alley, Dissecting peroxiredoxin catalysis: separating binding, peroxidation, and resolution for a bacterial AhpC, *Biochemistry* 54 (2015) 1567–1575.
- [77] A. Hall, D. Parsonage, L.B. Poole, P.A. Karplus, Structural evidence that peroxiredoxin catalytic power is based on transition state stabilization, *J. Mol. Biol.* 402 (2010) 194–209.
- [78] P.A. Karplus, A primer on peroxiredoxin biochemistry, *Free Radic. Biol. Med.* 80 (2015) 183–190, <http://dx.doi.org/10.1016/j.freeradbiomed.2014.10.009>.
- [79] N.J. Adimora, D.P. Jones, M.L. Kemp, A model of redox kinetics implicates the thiol proteome in cellular hydrogen peroxide responses, *Antioxid. Redox Signal.* 13 (2010) 731–743.
- [80] R.L.S. Goncalves, C.L. Quinlan, I.V. Perevoshchikova, et al., Sites of superoxide and hydrogen peroxide production by muscle mitochondria assessed ex vivo under conditions mimicking rest and exercise, *J. Biol. Chem.* 290 (2015) 209–227, <http://dx.doi.org/10.1074/jbc.M114.619072>.
- [81] T. Pearson, T. Kabayo, R. Ng, et al., Skeletal muscle contractions induce acute changes in cytosolic superoxide, but slower responses in mitochondrial superoxide and cellular hydrogen peroxide, *Plos One* 9 (2014) e96378, <http://dx.doi.org/10.1371/journal.pone.0096378>.
- [82] R.M. Jarvis, S.M. Hughes, E.C. Ledgerwood, Peroxiredoxin 1 functions as a signal peroxidase to receive, transduce, and transmit peroxide signals in mammalian cells, *Free Radic. Biol. Med.* 53 (2012) 1522–1530.
- [83] M.K. Trenerry, K.A. Carey, A.C. Ward, D. Cameron-Smith, STAT3 signaling is activated in human skeletal muscle following acute resistance exercise, *J. Appl. Physiol.* 102 (2007) 1483–1489, <http://dx.doi.org/10.1152/jappphysiol.01147.2006>.
- [84] D.C. Wright, D.H. Han, P.M. Garcia-Roves, et al., Exercise-induced mitochondrial biogenesis begins before the increase in muscle PGC-1α expression, *J. Biol. Chem.* 282 (2007) 194–199, <http://dx.doi.org/10.1074/jbc.M606116200>.
- [85] M.H. Choi, I.K. Lee, G.W. Kim, et al., Regulation of PDGF signalling and vascular remodelling by peroxiredoxin II, *Nature* 435 (2005) 347–353.
- [86] J.E. Turner, J.A. Bosch, M.T. Drayton, S. Aldred, Assessment of oxidative stress in lymphocytes with exercise, *J. Appl. Physiol.* 111 (2011) 206–211.
- [87] J.E. Turner, S.J. Bennett, J.P. Campbell, et al., The antioxidant enzyme peroxiredoxin-2 is depleted in lymphocytes seven days after ultra-endurance exercise, *Free Radic. Res.* 47 (2013) 821–828.
- [88] C. Brinkmann, J. Blossfeld, M. Pesch, et al., Lipid-peroxidation and peroxiredoxin-overoxidation in the erythrocytes of non-insulin-dependent type 2 diabetic men during acute exercise, *Eur. J. Appl. Physiol.* 112 (2012) 2277–2287, <http://dx.doi.org/10.1007/s00421-011-2203-x>.
- [89] Z. a Wood, L.B. Poole, R.R. Hantgan, P.A. Karplus, Dimers to doughnuts: redox-sensitive oligomerization of 2-cysteine peroxiredoxins, *Biochemistry* 41 (2002) 5493–5504, <http://dx.doi.org/10.1021/bi012173m>.
- [90] Y. Manabe, M. Takagi, M. Nakamura-Yamada, et al., Redox proteins are constitutively secreted by skeletal muscle, *J. Physiol. Sci.* 64 (2014) 401–409.
- [91] J.R. Riddell, X.Y. Wang, H. Minderman, S.O. Gollnick, Peroxiredoxin 1 stimulates secretion of proinflammatory cytokines by binding to TLR4, *J. Immunol.* 184 (2010) 1022–1030, <http://dx.doi.org/10.4049/jimmunol.0901945>.
- [92] T. Rabilloud, M. Heller, F. Gasnier, et al., Proteomics analysis of cellular response to oxidative stress. Evidence for in vivo overoxidation of peroxiredoxins at their active site, *J. Biol. Chem.* 277 (2002) 19396–19401, <http://dx.doi.org/10.1074/jbc.M106585200>.
- [93] A.J. Wadley, Y.W. Chen, S.J. Bennett, et al., Monitoring changes in thioredoxin and over-oxidised peroxiredoxin in response to exercise in humans, *Free Radic. Res.* 49 (2015) 290–298, <http://dx.doi.org/10.1080/09650790802667527>.
- [94] H. Schenk, M. Klein, W. Erdbrügger, et al., Distinct effects of thioredoxin and antioxidants on the activation of transcription factors NF-κB and AP-1, *Proc. Natl. Acad. Sci. USA* 91 (1994) 1672–1676.
- [95] D.A. Moghaddam, A. Heber, D. Capin, et al., Training increases peroxiredoxin 2 contents in the erythrocytes of overweight/obese men suffering from type 2 diabetes, *Wien. Med. Wochenschr.* 161 (2011) 511–518, <http://dx.doi.org/10.1007/s10354-011-0037-0>.
- [96] K.P. Lee, Y.J. Shin, S.C. Cho, et al., Peroxiredoxin 3 has a crucial role in the contractile function of skeletal muscle by regulating mitochondrial homeostasis, *Free Radic. Biol. Med.* 77 (2014) 298–306.
- [97] T.-S. Chang, C.-S. Cho, S. Park, et al., Peroxiredoxin III, a mitochondrion-specific peroxidase, regulates apoptotic signaling by mitochondria, *J. Biol. Chem.* 279 (2004) 41975–41984.
- [98] J.L. Vingren, W.J. Kraemer, D.L. Hatfield, et al., Effect of resistance exercise on muscle steroidogenesis, *J. Appl. Physiol.* 105 (2008) 1754–1760, <http://dx.doi.org/10.1152/jappphysiol.91235.2008>.
- [99] J.M. Lim, K.S. Lee, H. Woo, Control of the pericentrosomal H₂O₂ level by peroxiredoxin I is critical for mitotic progression, *J. Cell Biol.* 210 (2015) 23–33, <http://dx.doi.org/10.1083/jcb.201412068>.
- [100] L.B. Poole, C. Klomsiri, S.A. Knaggs, et al., Fluorescent and affinity-based tools to detect cysteine sulfenic acid formation in proteins, *Bioconjug Chem.* 18 (2007) 2004–2017, <http://dx.doi.org/10.1021/bc700257a>.
- [101] S.M. Hutson, L.B. Poole, S. Coles, M.E. Conway, Redox regulation and trapping sulfenic acid in the peroxide-sensitive human mitochondrial branched chain aminotransferase, *Methods Mol. Biol.* 476 (2008) 139–152.
- [102] M.E. Conway, S.J. Coles, M.M. Islam, S.M. Hutson, Regulatory control of human cytosolic branched-chain aminotransferase by oxidation and S-glutathionylation and its interactions with redox sensitive neuronal proteins, *Biochemistry* 47 (2008) 5465–5479, <http://dx.doi.org/10.1021/bi800303h>.
- [103] J.N. Cobley, N.V. Margaritelis, J.P. Morton, et al., The basic chemistry of exercise-induced DNA oxidation: oxidative damage, redox signaling, and their interplay, *Front. Physiol.* 6 (2015) 1–8, <http://dx.doi.org/10.3389/fphys.2015.00182>.
- [104] L.Y. Tanaka, L.R. Bechara, A.M. dos Santos, C.P. Jordão, L.G. de Sousa, et al., Exercise improves endothelial function: a local analysis of production of nitric oxide and reactive oxygen species, *Nitric Oxide* 45 (2015) 7–14.
- [105] M.J. Jackson, Redox regulation of adaptive responses in skeletal muscle to contractile activity, *Free Radic. Biol. Med.* 47 (2009) 1267–1275, <http://dx.doi.org/10.1016/j.freeradbiomed.2009.09.005>, S0891-5849(09)00531-0 [pii].
- [106] E. Cuevasanta, M. Lange, J. Bonanata, et al., Reaction of hydrogen sulfide with disulfide and sulfenic acid to form the strongly nucleophilic persulfide, *J. Biol. Chem.* 290 (2015) 26866–26880, <http://dx.doi.org/10.1074/jbc.M115.672816>.