

The Impact of Alemtuzumab Scheduling on Graft Versus Host Disease Following Unrelated Donor Fludarabine and Melphalan Allografts

Green, Kile; Pearce, Kim; Sellar, Rob S; Jardine, Laura; Nicolson, Phillip; Nagra, Sandeep; Bigley, Venetia; Jackson, Graham; Dickinson, Anne M; Thomson, Kirsty; Mackinnon, Stephen; Craddock, Charles; Peggs, Karl S; Collin, Matthew

DOI:

[10.1016/j.bbmt.2017.02.007](https://doi.org/10.1016/j.bbmt.2017.02.007)
[10.1016/j.bbmt.2017.02.007](https://doi.org/10.1016/j.bbmt.2017.02.007)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Green, K, Pearce, K, Sellar, RS, Jardine, L, Nicolson, P, Nagra, S, Bigley, V, Jackson, G, Dickinson, AM, Thomson, K, Mackinnon, S, Craddock, C, Peggs, KS & Collin, M 2017, 'The Impact of Alemtuzumab Scheduling on Graft Versus Host Disease Following Unrelated Donor Fludarabine and Melphalan Allografts', *Biology of Blood and Marrow Transplantation*, vol. 23, no. 5, pp. 805-812. <https://doi.org/10.1016/j.bbmt.2017.02.007>, <https://doi.org/10.1016/j.bbmt.2017.02.007>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Accepted Manuscript

Title: The Impact of Alemtuzumab Scheduling on Graft Versus Host Disease Following Unrelated Donor Fludarabine and Melphalan Allografts

Author: Kile Green, Kim Pearce, Rob S Sellar, Laura Jardine, Phillip LR Nicolson, Sandeep Nagra, Venetia Bigley, Graham Jackson, Anne M Dickinson, Kirsty Thomson, Stephen Mackinnon, Charles Craddock, Karl S Peggs, Matthew Collin


PII: S1083-8791(17)30276-8
DOI: <http://dx.doi.org/doi: 10.1016/j.bbmt.2017.02.007>
Reference: YBBMT 54580

To appear in: *Biology of Blood and Marrow Transplantation*

Received date: 7-10-2016
Accepted date: 8-2-2017

Please cite this article as: Kile Green, Kim Pearce, Rob S Sellar, Laura Jardine, Phillip LR Nicolson, Sandeep Nagra, Venetia Bigley, Graham Jackson, Anne M Dickinson, Kirsty Thomson, Stephen Mackinnon, Charles Craddock, Karl S Peggs, Matthew Collin, The Impact of Alemtuzumab Scheduling on Graft Versus Host Disease Following Unrelated Donor Fludarabine and Melphalan Allografts, *Biology of Blood and Marrow Transplantation* (2017), <http://dx.doi.org/doi: 10.1016/j.bbmt.2017.02.007>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



The impact of alemtuzumab scheduling on graft versus host disease following unrelated donor fludarabine and melphalan allografts

Kile Green 1
Kim Pearce 1
Rob S Sellar 2,3
Laura Jardine 1,4
Phillip LR Nicolson 5
Sandeep Nagra 6
Venetia Bigley 1,4
Graham Jackson 4,7
Anne M Dickinson 1
Kirsty Thomson 2,3
Stephen Mackinnon 2,3
Charles Craddock 5,6
Karl S Peggs 2,3
Matthew Collin 1,4

1. Institute of Cellular Medicine, Newcastle University, Newcastle upon Tyne, United Kingdom
2. Cancer Institute, University College London, London, United Kingdom
3. Department of Haematology, University College London Hospitals NHS Foundation Trust, London, United Kingdom
4. Northern Centre for Bone Marrow Transplantation, Newcastle upon Tyne Hospitals NHS Foundation Trust, Newcastle upon Tyne, United Kingdom
5. School of Cancer Sciences, University of Birmingham, Birmingham, United Kingdom
6. Centre for Clinical Haematology, University Hospitals Birmingham NHS Foundation Trust, Birmingham, United Kingdom
7. Northern Institute for Cancer Research, Newcastle University, Newcastle Upon Tyne, United Kingdom

Text	3793 (inc figure legends)
Abstract	202
Figures/Tables	6/2
References	36

Conflict of interest disclosures


The authors have no conflict of interest to declare.

Short title

Alemtuzumab scheduling in unrelated donor allografts

Accepted Manuscript

Graphical Abstract



Highlights:

- Alemtuzumab dosing in unrelated reduced intensity transplantation is undefined.
- 60mg Alemtuzumab provides equivalent control of GVHD compared with 100mg
- 50mg Alemtuzumab is associated with threefold increased risk of severe acute GVHD.

Abstract

Alemtuzumab conditioning is highly effective at reducing the incidence of acute and chronic graft versus host disease (GVHD) in reduced intensity fludarabine and melphalan transplantation with ciclosporin monotherapy. Less frequent and lower dose scheduling may be used with sibling donors but an optimal regimen for matched unrelated donors has not been defined. In this retrospective observational

study of 313 patients, the incidence and severity of GVHD was compared in patients receiving three different dose schedules: the standard 100mg regimen (20mg on day -7 to -3), 60mg (30mg day -4 and -2) or 50mg (10mg on day -7 to -3). Patients treated with 100mg, 60mg or 50mg developed acute GVHD grade I-IV with an incidence of 74%, 65% and 64%, respectively, while 36%, 32% and 41% developed chronic GVHD. An excess of severe acute grade III/IV GVHD was observed in the 50mg cohort (15% vs. 2-6%; $p = 0.016$). The relative risk of severe acute grade GVHD remained more than three-fold higher in the 50mg cohort, compared with 100mg, after adjustment for differences in HLA match, age, gender mismatch, CMV risk and diagnosis ($p = 0.030$). The findings indicate that the 60mg alemtuzumab schedule was comparable to 100mg but more attenuated schedules may increase the risk of severe grade GVHD.

Keywords

Alemtuzumab
T cell depletion
Graft versus host disease
Conditioning regimens

Accepted Manuscript

Introduction

Alemtuzumab (humanized anti-CD52 antibody) is highly effective at reducing the incidence of acute and chronic GVHD in the setting of reduced intensity transplantation with fludarabine and melphalan (1-7). When delivered to recipients during conditioning therapy, it effects *in vivo* depletion of both recipient and donor T cells, NK cells, B cells, monocytes and dendritic cells, owing persistence in the recipient with a half life of 8 days (8-11). Freedom from GVHD is associated with partial chimerism of donor T cells but this may be corrected with donor lymphocyte infusions to deliver good overall survival with minimal long term morbidity (7,12-14).

The original fludarabine, melphalan and alemtuzumab regimen used an empiric alemtuzumab schedule of 100mg comprising five 20mg doses given on consecutive days between day -7 and -3. This regimen is effective at abrogating GVHD in mixed cohorts of matched related and unrelated donor transplants used to treat patients with both myeloid and lymphoid malignancy (1-7). It has also been noted that GVHD is well controlled in PBSC and BM grafts from unrelated donors (15) and that a degree of antigen mismatching is well tolerated (16). Similar schedules of alemtuzumab have also been used with other fludarabine-based reduced intensity protocols with equivalent efficacy (14).

Excessive T cell depletion may be associated with increased relapse and risk of infection (11,17-19) and several groups have shown that dose reduction is possible in unrelated donor transplantation. A number of schedules with doses of between 50-100mg, administered over 2-5 days, have been tested (20-22) and it is reported that as little as 10mg reduces the burden of GVHD (23). A phased dose de-escalation study in sibling transplants concluded that a single dose of 30mg on day -1 was sufficient to reduce GVHD to a similar level as 100mg of alemtuzumab (24) but a comparable study has not been performed using only unrelated donors in a common protocol. Retrospective comparison of 30mg and 60mg dosing in sibling and unrelated donor transplants, respectively, indicated that the unrelated cohort still experienced more GVHD and had higher donor T cell chimerism (25).

Owing to the long *in vivo* half-life of alemtuzumab, the total dose and scheduling both have the potential to modify GVHD risk substantially (11) but there is no consensus about an optimal regimen in fludarabine-melphalan unrelated donor transplantation. Here we report a retrospective observational study in which we compared three commonly used protocols. Reduction and compression of the alemtuzumab schedule to two 30mg doses on day -4 and -2 was comparable to 100mg between days -7 to -3 but patients receiving 50mg alemtuzumab between day -7 to -3 were at greater risk of severe acute grade III/IV GVHD.

Methods

Patients and donors

Data were collated from three UK transplant centers: University College Hospital/Royal Free Hospital, University College London Hospitals NHS Foundation

Trust, London; Northern Centre for Bone Marrow Transplantation, Newcastle upon Tyne Hospitals NHS Foundation Trust, Newcastle upon Tyne; and, Queen Elizabeth Hospital, University Hospitals Birmingham NHS Foundation Trust, Birmingham. Sequential patients transplanted between January 2007 and December 2011 were included. Patients were over the age of 18 years at transplantation, had a hematological malignancy at any stage and were transplanted with an unrelated donor with at least 8/10 HLA four digit allele matching. The number of transplants with less than 10/10 matching was too small to discern differences between A, B, C, or DQ antigen mismatches. The major stem cell source was PBSC at all centers. All patients gave consent for their clinical data to be reported anonymously according to local ethical approval. Clinical data were collected by transplant center data managers and analyzed by KG, KP and MC.

Conditioning and alemtuzumab dosing

All patients were conditioned with five doses of fludarabine $30\text{mg}/\text{m}^2$ daily days -7 to -3 and melphalan $140\text{mg}/\text{m}^2$ day -2. GVHD prophylaxis consisted of ciclosporin monotherapy starting at $3\text{mg}/\text{kg}/\text{day}$ adjusting to an initial level of at least $200\text{ng}/\text{ml}$. Each center delivered a different alemtuzumab schedule. London patients received 100mg in five 20mg daily doses given consecutively from day -7 to day -3. Newcastle patients received 60mg as two 30mg doses on day -4 and day -2 and Birmingham patients received 50mg in five 10mg doses given consecutively from day -7 to day -3. Donor lymphocyte infusions (DLI) were given for mixed chimerism in all centers according to three monthly escalating schedules starting at 6 months post transplantation with a dose of 1×10^6 CD3+ cells per kg. As the DLI schedule did not start until 6 months, it did not influence the incidence or severity of acute GVHD.

Study endpoints and supportive care

The primary endpoints of the study were the raw incidence and severity of acute GVHD according to Seattle grading and the cumulative incidence of chronic GVHD. Secondary endpoints included donor T cell chimerism, CMV reactivation, non-relapse mortality (NRM), relapse and overall survival. Patients were evaluable for acute GVHD if they survived until engraftment and for chronic GVHD if they lived more than 100 days. CMV reactivation was defined as more than 2 sequential blood PCR results $> 2 \times 10^3$ copies/ml or requiring anti-CMV therapy with ganciclovir, valganciclovir or foscarnet. Anti-viral prophylaxis was maintained with acyclovir 200mg BD PO in the 100mg cohort or 200mg TDS PO in the 60mg and 50mg cohorts.

Statistical analysis

Analysis was performed according to EBMT guidelines (26-29), on consecutive patients transplanted between January 2007 and December 2011. Chi-square, Mann-Witney tests and Kaplan-Meier survival curves were plotted with GraphPad Prism 6 and cumulative incidence analysis with competing risks was performed with the `cmprsk` function of the open-source software R. Kaplan-Meier curves were compared with log-rank (Mantel-Cox) tests while cumulative incidence curves were compared with Gray's test. Patients were censored at last follow up. Multivariate

analysis was performed in SPSS using binary logistic regression (grade III/IV vs other)

Results

Conditioning protocols used in this three-center retrospective observational study are summarized in **Figure 1**. Both the scheduling and total dose of Alemtuzumab was different between each cohort.

Patient characteristics

Consecutive patients recruited at each center were included in the study. Several differences were noted in the patient characteristics reflecting local practice and referral bias, summarized in **Table 1**. Patients in each cohort were matched for donor cell source (90-97% receiving PBSC), HLA matching (69-83% at least 10/10), disease status and follow up. There were however significant differences in age, the proportion of female to male transplants, the frequency of high risk CMV positive recipients and the distribution of disease between acute myeloid leukemia and lymphoproliferative disease. The 50mg cohort had the highest rate of high risk CMV serostatus (negative into positive; 25.23%; $p = 0.0058$) and patients were more likely to have a diagnosis of AML or MDS (62.62%; $p = 0.0005$). Patients receiving 60mg of alemtuzumab were younger (median age 45; $p = 0.0051$) and had no female to male transplants ($p = 0.009$). Patients receiving 100mg had the highest rate of female to male transplants (21.82%; $p = 0.009$) and the most patients with lymphoproliferative disease (48.51%; $p = 0.0005$).

Acute GVHD

There were no differences in the overall incidence of acute GVHD between the three cohorts of patients with 64-74% of patients experiencing at least grade I GVHD and 32-40% of patients experiencing clinically significant grade II-IV GVHD in all cohorts (**Figure 2A**). However there distribution of acute GVHD grades was skewed ($p = 0.0124$ for all grades), with an excess of grade III/IV acute GVHD in the 50mg cohort (15% compared with 2-6%; $p = 0.0161$ for grades III-IV vs. grades 0-II; Chi-square tests). It was not possible to calculate the cumulative incidence of acute GVHD from the available data, so early relapse death remains a potential competing risk. However, the relapse risk within 100 days was consistently low at 5.6%, 1.4% and 6.7% respectively for the 50mg, 60mg and 100mg cohorts. HLA matching was considered an important potential confounding variable by subgroup analysis. The 50mg and 100mg cohorts had a slighter higher percentage of <10/10 HLA-matched donors than the 60mg cohort but this was not a significant baseline difference and

<10/10 matched donors did not experience more GVHD, within each cohort. **(Figure 2B)**.

The control of severe grade GVHD in the <10/10 patients was not improved by higher dosing of alemtuzumab. In fact, less severe GVHD was observed in the 50mg and 60mg cohorts who received a <10/10 antigen matched donor compared with 10/10 matched donors, presumably due to confounding factors in the selection of lower risk patients or anticipatory adjustment of post-transplant immunosuppression. The number of patients with <10/10 mismatching was not sufficient to stratify the degree of mismatch further.

Multivariate analysis

Although an apparent excess of grade III/IV acute GVHD was observed in the 50mg cohort, several variables that might affect the incidence of GVHD or were non-randomly distributed between the cohorts. A multivariate analysis was therefore performed to analyse the variables that might predict the bivariate outcome grade III/IV GVHD vs all other grades. The covariates considered were Alemtuzumab schedule 100mg, 60mg or 50mg; >10/10 or 10/10 HLA matching; age group <40, 41-60 and >60; female into male or other; CMV negative into positive or other; and, diagnosis MDS/AML or NHL. In a general linear model, older age, female into male, CMV high risk and NHL were associated with increased but non-significant risk of grade III/IV acute GVHD. Only alemtuzumab schedule was significantly predictive of grade III/IV GVHD conferring more than a three-fold increased risk, which remained after adjustment for the other variables **(Table 2)**.

Chronic GVHD

The cumulative incidence of chronic GVHD was similar in all cohorts ranging from 33% in the 100mg and 32% in the 60mg cohort, to 41% in the 50mg cohort at two years ($p = 0.6248$) **(Figure 3A)**. Although there was a trend for more chronic GVHD in the 50mg cohort, the number of patients with extensive chronic GVHD was not increased between the cohorts ($p = 0.1612$) **(Figure 3B)**.

Engraftment and donor chimerism

In keeping with previous reports, the rate of engraftment and achievement of complete donor myeloid chimerism was more than 95% in all cohorts (not shown). The level of donor T cell chimerism was however very variable **(Figure 4A)**. Analysis was constrained by the reporting donor chimerism in the 100mg cohort in bands of 0-49; 50-94 and ≥ 95 , so data from the other cohorts were transformed to

match. Only just over half of all patients achieved more than 95% donor chimerism with the lowest levels of donor T cell chimerism observed in the 60mg cohort ($p = 0.0460$). This may reflect a lower proportion of mismatched donors or more liberal use of post-transplant immunosuppression in this cohort. Continuously distributed data were available for the 50mg and 60mg cohorts and showed a significant skewing between cohorts with lower median T cell chimerism in the 60mg cohort (85% vs. 96%; $p =$ **(Figure 4B)**).

CMV reactivation

The risk of CMV reactivation is high with alemtuzumab containing regimens. More than 80% of patients in all cohorts with high risk serostatus (positive recipient) had a positive CMV PCR result after transplant. Intermediate status (positive donor into negative recipient) reactivation was similar in all cohorts. Within the high risk serostatus groups, significantly higher rates of reactivation were observed in the 100mg cohort ($p = 0.0267$) **(Figure 5)**.

Mortality and survival

No significant differences were observed in the cumulative incidence of non-relapse mortality at 2 years between 100mg, 60mg and 50mg cohorts (16%, 16%, and 24%, respectively; $p = 0.3232$) **(Figure 6A)** or relapse rates at 2 years (36%, 24%, and 29%, respectively; $p = 0.3454$) **(Figure 6B)**. The overall survival of each cohort was not significantly different and is shown for the major disease groups AML/MDS and NHL **(Figure 6C)**. Trends for lower survival in the 50mg cohort were observed in both relapse and non-relapse mortality. There were no differences within each cohort when survival was stratified according to diagnosis or disease risk (not shown).

Discussion

The use of T cell depletion remains controversial in all forms of allogeneic bone marrow transplantation but particularly in the setting of reduced intensity conditioning, where it is argued that graft versus tumor effects need to be stronger to compensate for attenuated cytoreduction(19,30). However, even with minimal conditioning, T replete transplants still incur a similar burden of graft versus host disease to full intensity transplants albeit with delayed kinetics(31). The prevalence of GVHD with reduced intensity conditioning is further amplified by the increasing age of eligible patients and widespread use of peripheral blood stem cells. For these reasons, many UK centers have developed reduced intensity protocols using alemtuzumab to effect in vivo T cell depletion and reduce acute GVHD(1-7). This

approach is supported by a retrospective comparison of T depletion strategies with reduced intensity regimens showing superior control of acute and chronic GVHD and a trend for greater overall survival with alemtuzumab-containing regimens, despite higher relapse rates (19). These data underscore the need to define optimal dosing regimens that balance the risks of toxicity and relapse.

Several cohort studies have compared protocols with and without alemtuzumab and arrived at the conclusion that overall survival is similar and survivorship quality is enhanced even though there are higher rates of relapse and CMV reactivation in patients given alemtuzumab. In sibling transplants for lymphoproliferative disease, acute and chronic GVHD was reduced by alemtuzumab with no impact on overall survival (32). In a comparison of patients with AML, alemtuzumab reduced the incidence of extensive chronic GVHD from 47% to 4% and was associated with slightly superior survival than those not receiving alemtuzumab (5). This study included sibling and matched unrelated donor transplants given a range of alemtuzumab schedules from 30mg to 100mg. Another more controlled study of sibling and unrelated transplants for AML showed that 100mg alemtuzumab reduced the incidence of acute grade II-IV GVHD from 58.1% to 23.3% and chronic GVHD from 78.4% to 16.0% with no difference in overall survival(33). Increases in the rate of CMV reactivation with alemtuzumab do not appear to impact on survival with modern pre-emptive antiviral therapy (34). Although intervention with DLI may be required to achieve an optimal outcome in alemtuzumab-containing transplants, recipients of DLI appear to enjoy a favorable outcome in both lymphoid and myeloid malignancy (14,35).

The scheduling of alemtuzumab has a significant impact on plasma concentration on the day of transplant(11). We have previously demonstrated that the 60mg regimen results in a similar level of alemtuzumab to the 100mg regimen at 5-6 μ g/ml on day +1 (25), thus it is not surprising that comparable control of GVHD was observed. It would be erroneous to conclude that 60mg was a safe threshold dose without reference to the timing of administration. The excess of severe acute GVHD in the 50mg cohort probably reflects earlier scheduling as much as a lower total dose, resulting in a substantially lower in vivo concentration. Although Alemtuzumab levels have not been measured in this regimen, the continuously distributed T cell chimerism at day 100 is consistent with a much lower level of donor T cell depletion in the 50mg cohort, compared with the 60mg cohort. Even later scheduling of two 30mg doses to day -2 to -1 has been proposed and is likely to deliver alemtuzumab concentrations that exceed the original 100mg regimen(36). The 100mg dose of alemtuzumab may be used with <10/10 antigen matched donors(16) and it appeared here that reduction to 60mg or 50mg did not significantly increase GVHD, relative to patients with 10/10 matched donors given the same prophylaxis.

These results indicate that fludarabine-melphalan reduced intensity transplantation is feasible with or without alemtuzumab but the question remains whether it is possible to define an optimal schedule of alemtuzumab that achieves good control of GVHD and survivorship without excessive detriment to immune reconstitution. A phase II

study in sibling transplants reported that a significant dose reduction to a single 30mg vial at day -1 almost completely ablated acute grade II-IV GVHD and extensive chronic GVHD (both 4%). Another dose reduction study using a variety of fludarabine-based regimens showed that lower doses were also feasible in mixed sibling and unrelated donors given a variety of conditioning regimens, but there was insufficient power to discriminate between different dose levels (23)

In unrelated donor transplants, there does not appear to be a simple linear relationship between alemtuzumab schedule and GVHD risk. At the highest alemtuzumab dose of 100mg, the rate of grade II-IV GVHD in our study was approximately 10-fold higher (40%) than sibling transplants exposed to 30mg of alemtuzumab (4%), although the majority was grade II, and remained relatively constant with 60mg and 50mg dosing. This is consistent with a previous report of unrelated donors transplants performed with 100mg of alemtuzumab(16). A recent study comparing the 60mg schedule for unrelated donors with 30mg for sibling transplants showed similarly high rates of grade II-IV GVHD (44%) and donor T cell chimerism in the unrelated transplants(25). Reducing alemtuzumab to 40mg or less did not appear to increase the rate of grade II-IV acute GVHD in a series of small cohorts (23). However, below 60mg, there may be a trend towards increasing severe grade III/IV acute GVHD. In our study, patients receiving 50mg experienced 15% grade III/IV acute GVHD compared with 2-6% in the 60mg and 100mg cohorts. This bias was also observed in the 10/10 matched donors. An even higher rate of 23.8% severe acute grade III/IV GVHD was reported in unrelated donor transplants receiving 40mg of alemtuzumab although this did not further increase when alemtuzumab dose was reduced to 20mg and 10mg (23).

The main limitation of this study is the difficulty in controlling center bias. The increase in grade III/IV acute GVHD in the 50mg cohort remained more than three-fold after adjustment for the major covariables that might have influenced GVHD, but other factors such as disease risk index and performance status could not be completely controlled for. Stratification according to disease risk was not possible given the many different indications that were included in the analysis and Hematopoietic Cell Transplantation Comorbidity Index data were not routinely reported for patients transplanted over this period. Less tangible bias may also have been introduced by variations in clinical practice such as the use of post-transplant immunosuppression. Although centers reported similar operating procedures governing the withdrawal of ciclosporin, ciclosporin levels were not collected as primary data. Similarly, detailed donor lymphocyte infusion records were not available. All three centers reported a common escalating DLI regimen but we could not audit this directly. Similar rates of chronic GVHD were observed, regardless of any institutional variations in practice that may have occurred in relation to the use of DLI.

In conclusion, this study suggests that it is feasible to use two 30mg doses of alemtuzumab in fludarabine-melphalan unrelated donor transplants without significantly increasing the risk of GVHD or impairing outcome compared with the

original 100mg dose. Given at day -4 and -2, this results in similar plasma levels compared with five 20mg doses (25) and is cheaper, assuming a 30mg vial size. Further dose reduction may be feasible with later scheduling (day -2 to -1) and ideally, should be studied prospectively with in vivo alemtuzumab levels and accurate T cell chimerism, in addition to clinical outcome data. The current 50mg regimen in use in the UK was associated with an increased risk of severe grade III/IV acute GVHD, although confounding factors may have increased the baseline risk in this cohort. Overall, the use of T cell depletion remains subject to historical precedent and individual center bias regarding the acceptable burden of GVHD in transplant survivors.

Acknowledgements

The authors thank patients who participated and the transplant teams who provided their clinical care. The research was supported by Bright Red and the British Society for Haematology.

Authorship contributions

Kile Green analyzed data and produced figures
Kim Pearce performed statistical analysis
Rob S Sellar provided clinical data
Laura Jardine analyzed data and produced figures
Phillip LR Nicolson provided clinical data
Sandeep Nagra provided clinical data
Venetia Bigley provided clinical data
Graham Jackson provided clinical data
Anne M Dickinson provided clinical data
Kirsty Thomson provided clinical data
Stephen Mackinnon designed the study
Charles Craddock designed the study
Karl S Peggs designed the study and wrote the paper
Matthew Collin designed the study and wrote the paper

References

1. Kottaridis PD, Milligan DW, Chopra R et al. In vivo CAMPATH-1H prevents graft-versus-host disease following nonmyeloablative stem cell transplantation. *Blood*. 2000;96 (7):2419-2425.
2. Chakraverty R, Peggs K, Chopra R et al. Limiting transplantation-related mortality following unrelated donor stem cell transplantation by using a nonmyeloablative conditioning regimen. *Blood*. 2002;99 (3):1071-1078.
3. Tauro S, Craddock C, Peggs K et al. Allogeneic stem-cell transplantation using a reduced-intensity conditioning regimen has the capacity to produce durable remissions and long-term disease-free survival in patients with high-risk acute myeloid leukemia and myelodysplasia. *J Clin Oncol*. 2005;23 (36):9387-9393.
4. Delgado J, Thomson K, Russell N et al. Results of alemtuzumab-based reduced-intensity allogeneic transplantation for chronic lymphocytic leukemia: a British Society of Blood and Marrow Transplantation Study. *Blood*. 2006;107 (4):1724-1730.
5. Malladi RK, Peniket AJ, Littlewood TJ et al. Alemtuzumab markedly reduces chronic GVHD without affecting overall survival in reduced-intensity conditioning sibling allo-SCT for adults with AML. *Bone Marrow Transplant*. 2009;43 (9):709-715.
6. Thomson KJ, Morris EC, Bloor A et al. Favorable long-term survival after reduced-intensity allogeneic transplantation for multiple-relapse aggressive non-Hodgkin's lymphoma. *J Clin Oncol*. 2009;27 (3):426-432.
7. Thomson KJ, Morris EC, Milligan D et al. T-cell-depleted reduced-intensity transplantation followed by donor leukocyte infusions to promote graft-versus-lymphoma activity results in excellent long-term survival in patients with multiply relapsed follicular lymphoma. *J Clin Oncol*. 2010;28 (23):3695-3700.
8. Klanginsirikul P, Carter GI, Byrne JL, Hale G, Russell NH. Campath-1G causes rapid depletion of circulating host dendritic cells (DCs) before allogeneic transplantation but does not delay donor DC reconstitution. *Blood*. 2002;99 (7):2586-2591.
9. Buggins AG, Mufti GJ, Salisbury J et al. Peripheral blood but not tissue dendritic cells express CD52 and are depleted by treatment with alemtuzumab. *Blood*. 2002;100 (5):1715-1720.
10. Ratzinger G, Reagan JL, Heller G, Busam KJ, Young JW. Differential CD52 expression by distinct myeloid dendritic cell subsets: implications for alemtuzumab activity at the level of antigen presentation in allogeneic graft-host interactions in transplantation. *Blood*. 2003;101 (4):1422-1429.
11. Morris EC, Rebello P, Thomson KJ et al. Pharmacokinetics of alemtuzumab used for in vivo and in vitro T-cell depletion in allogeneic transplantations: relevance for early adoptive immunotherapy and infectious complications. *Blood*. 2003;102 (1):404-406.

12. Peggs KS, Thomson K, Hart DP et al. Dose-escalated donor lymphocyte infusions following reduced intensity transplantation: toxicity, chimerism, and disease responses. *Blood*. 2004;103 (4):1548-1556.
13. Bloor AJ, Thomson K, Chowdhry N et al. High response rate to donor lymphocyte infusion after allogeneic stem cell transplantation for indolent non-Hodgkin lymphoma. *Biol Blood Marrow Transplant*. 2008;14 (1):50-58.
14. Potter VT, Krishnamurthy P, Barber LD et al. Long-term outcomes of alemtuzumab-based reduced-intensity conditioned hematopoietic stem cell transplantation for myelodysplastic syndrome and acute myelogenous leukemia secondary to myelodysplastic syndrome. *Biol Blood Marrow Transplant*. 2014;20 (1):111-117.
15. Shaw BE, Apperley JF, Russell NH et al. Unrelated donor peripheral blood stem cell transplants incorporating pre-transplant in-vivo alemtuzumab are not associated with any increased risk of significant acute or chronic graft-versus-host disease. *Br J Haematol*. 2011;153 (2):244-252.
16. Mead AJ, Thomson KJ, Morris EC et al. HLA-mismatched unrelated donors are a viable alternate graft source for allogeneic transplantation following alemtuzumab-based reduced-intensity conditioning. *Blood*. 2010;115 (25):5147-5153.
17. Chakrabarti S, Avivi I, Mackinnon S et al. Respiratory virus infections in transplant recipients after reduced-intensity conditioning with Campath-1H: high incidence but low mortality. *Br J Haematol*. 2002;119 (4):1125-1132.
18. Doderio A, Carrabba M, Milani R et al. Reduced-intensity conditioning containing low-dose alemtuzumab before allogeneic peripheral blood stem cell transplantation: graft-versus-host disease is decreased but T-cell reconstitution is delayed. *Exp Hematol*. 2005;33 (8):920-927.
19. Soiffer RJ, Lerademacher J, Ho V et al. Impact of immune modulation with anti-T-cell antibodies on the outcome of reduced-intensity allogeneic hematopoietic stem cell transplantation for hematologic malignancies. *Blood*. 2011;117 (25):6963-6970.
20. Faulkner RD, Craddock C, Byrne JL et al. BEAM-alemtuzumab reduced-intensity allogeneic stem cell transplantation for lymphoproliferative diseases: GVHD, toxicity, and survival in 65 patients. *Blood*. 2004;103 (2):428-434.
21. Tholouli E, Liakopoulou E, Greenfield HM et al. Outcomes following 50 mg versus 100 mg alemtuzumab in reduced-intensity conditioning stem cell transplants for acute myeloid leukaemia and poor risk myelodysplasia. *Br J Haematol*. 2008;142 (2):318-320.
22. Ingram W, Devereux S, Das-Gupta EP et al. Outcome of BEAM-autologous and BEAM-alemtuzumab allogeneic transplantation in relapsed advanced stage follicular lymphoma. *Br J Haematol*. 2008;141 (2):235-243.
23. Bertz H, Spyridonidis A, Wasch R, Grulich C, Egger M, Finke J. A novel GVHD-prophylaxis with low-dose alemtuzumab in allogeneic sibling or unrelated

donor hematopoietic cell transplantation: the feasibility of deescalation. *Biol Blood Marrow Transplant*. 2009;15 (12):1563-1570.

24. Chakraverty R, Orti G, Roughton M et al. Impact of in vivo alemtuzumab dose before reduced intensity conditioning and HLA-identical sibling stem cell transplantation: pharmacokinetics, GVHD, and immune reconstitution. *Blood*. 2010;116 (16):3080-3088.

25. Jardine L, Publicover A, Bigley V et al. A comparative study of reduced dose alemtuzumab in matched unrelated donor and related donor reduced intensity transplants. *Br J Haematol*. 2015;168 (6):874-881.

26. Klein JP, Rizzo JD, Zhang MJ, Keiding N. Statistical methods for the analysis and presentation of the results of bone marrow transplants. Part I: unadjusted analysis. *Bone Marrow Transplant*. 2001;28 (10):909-915.

27. Klein JP, Rizzo JD, Zhang MJ, Keiding N. Statistical methods for the analysis and presentation of the results of bone marrow transplants. Part 2: Regression modeling. *Bone Marrow Transplant*. 2001;28 (11):1001-1011.

28. Fine JP, Gray RJ. A Proportional Hazards Model for the Subdistribution of a Competing Risk. *Journal of the American Statistical Association*. 1999;94 (446):496-509.

29. Scrucca L, Santucci A, Aversa F. Regression modeling of competing risk using R: an in depth guide for clinicians. *Bone Marrow Transplant*. 2010;45 (9):1388-1395.

30. McSweeney PA, Niederwieser D, Shizuru JA et al. Hematopoietic cell transplantation in older patients with hematologic malignancies: replacing high-dose cytotoxic therapy with graft-versus-tumor effects. *Blood*. 2001;97 (11):3390-3400.

31. Mielcarek M, Martin PJ, Leisenring W et al. Graft-versus-host disease after nonmyeloablative versus conventional hematopoietic stem cell transplantation. *Blood*. 2003;102 (2):756-762.

32. Perez-Simon JA, Kottaridis PD, Martino R et al. Nonmyeloablative transplantation with or without alemtuzumab: comparison between 2 prospective studies in patients with lymphoproliferative disorders. *Blood*. 2002;100 (9):3121-3127.

33. van Besien K, Kunavakkam R, Rondon G et al. Fludarabine-melphalan conditioning for AML and MDS: alemtuzumab reduces acute and chronic GVHD without affecting long-term outcomes. *Biol Blood Marrow Transplant*. 2009;15 (5):610-617.

34. Peggs KS, Preiser W, Kottaridis PD et al. Extended routine polymerase chain reaction surveillance and pre-emptive antiviral therapy for cytomegalovirus after allogeneic transplantation. *Br J Haematol*. 2000;111 (3):782-790.


35. Peggs KS, Sureda A, Qian W et al. Reduced-intensity conditioning for allogeneic haematopoietic stem cell transplantation in relapsed and refractory

Hodgkin lymphoma: impact of alemtuzumab and donor lymphocyte infusions on long-term outcomes. *Br J Haematol.* 2007;139 (1):70-80.

36. Okasha D, Kirkwood A, Copland M et al. Fludarabine, Melphalan and Alemtuzumab Conditioned Reduced Intensity (RIC) Allogeneic Hematopoietic Cell Transplantation for Adults Aged >40 Years with De Novo Acute Lymphoblastic Leukemia: A Prospective Study from the UKALL14 Trial ([ISRCTN 66541317](#)). *Blood.* 2015; epub December.

Accepted Manuscript

Figure 1

**Figure 1. Preparative regimens**

Outline of preparative regimens for the three patient cohorts.

Figure 2




Figure 3




Figure 4





Figure 4. Chimerism

A. Donor T cell chimerism at 100 days. Bins of 0-50; 50-95 and 95-100 were selected based on the discontinuous data available for the 100mg cohort. A significant difference was detected owing to lower T cell chimerism in the 60mg cohort.

B. Comparison of donor T cell chimerism at 100 days for patients in 50mg and 60mg cohorts showing skewing towards lower median T cell chimerism in the 60mg cohort (Mann-Witney test).

Figure 5




* $p = 0.0267$

Figure 5. CMV reactivation

The proportion of patients in each cohort requiring treatment for CMV reactivation, according to CMV risk status. A higher risk of reactivation was detected for high risk CMV positive recipients in the 100mg cohort (*).

Figure 6

**Figure 6. Outcome**

A. Cumulative incidence of NRM with relapse as a competing factor showing no significant difference between the cohorts by Gray's test.

B. Cumulative incidence of relapse with NRM as a competing factor showing no significant difference between the cohorts by Gray's test.

C. Overall survival for patients with i) AML/MDS or ii) NHL by cohort. Kaplan-Meier curves were compared by Log Rank (Mantel-Cox) test.

Table 1. Patient Characteristics

Summary of patient demographics, donor source, HLA matching and gender, CMV serostatus, disease and disease status at transplantation. Differences between cohorts were compared with t-tests or contingency analysis and p values were derived. Significantly outlying values by 2x2 Chi-square tests are indicated in bold in the table.

	50mg (n =107)	60mg (n =72)	100mg (n =134)	P-value
Age at transplantation				
Median (range)	56 (24-71)	45 (22-67)	50 (21-67)	0.0051
Graft				
Peripheral blood	97.20%	90.28%	97.47%	0.0551
Bone marrow	2.80%	9.72%	2.53%	
HLA status				
10/10	69.16%	83.33%	70.90%	0.1002
<10/10	30.84%	16.67%	29.10%	
Gender (donor/recipient)				
Female to male	5.61%	0.00%	21.82%	0.009
CMV (donor/recipient)				
- / positive	42.99%	38.89%	40.30%	0.018
positive / negative	25.23%	12.50%	11.94%	
negative / negative	31.78%	48.61%	47.76%	
Disease indication				
ALL	3.74%	1.39%	3.73%	0.0005
AML/MDS	62.62%	50.00%	32.84%	
Lymphoproliferative disorders	25.23%	44.44%	48.51%	
Other	8.41%	4.17%	14.93%	
Status at transplantation				
Untreated	5.61%	2.78%	3.73%	0.0909
Partial remission	10.28%	6.94%	17.91%	
Complete remission	74.77%	81.94%	55.22%	
Relapse/progression	5.61%	4.17%	6.72%	
Other	3.74%	4.17%	16.42%	
Follow up period				
Median follow up (months)	18.12	21.63	19.12	0.3339

Table 2. Multivariate analysis of the risk of grade III/IV acute GVHD

Effect of covariables on the incidence of grade III/IV GVHD. Odds ratios are given for the first variable versus the second as baseline.

Variable	Univariate Odds Ratio of GVHD III/IV	p value	Multivariate Odds Ratio of GVHD III/IV	p value
Alemtuzumab 60mg vs 100mg	1.067	0.920	0.837	0.792
Alemtuzumab 50 mg vs 100mg	3.190	0.014	3.024	0.030
<u><10/10 vs 10/10 match</u>	<u>0.933</u>	<u>0.888</u>	<u>0.881</u>	<u>0.803</u>
Age 41-60 vs <40	0.847	0.783	0.638	0.458
Age >60 vs <40	0.607	0.285	1.144	0.858
F into M vs other	0.340	0.300	0.342	0.318
CMV neg pos vs other	1.491	0.415	1.423	0.499
NHL vs MDS/AML	1.393	0.620	2.086	0.162