

Identifying the causal mechanisms of the quiet eye

Gonzales, Claudia; Causer, Joseph; Miall, Rowland; Grey, Michael; Humphreys, GW; williams, AM

DOI:

[10.1080/17461391.2015.1075595](https://doi.org/10.1080/17461391.2015.1075595)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Gonzales, C, Causer, J, Miall, R, Grey, M, Humphreys, GW & williams, AM 2017, 'Identifying the causal mechanisms of the quiet eye', *European Journal of Sport Science*, vol. 17, pp. 74. <https://doi.org/10.1080/17461391.2015.1075595>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.



Identifying the causal mechanisms of the quiet eye

Journal:	<i>European Journal of Sports Science</i>
Manuscript ID:	TEJS-2015-0109
Manuscript Type:	Review
Keywords:	Behavior, Measurement, Neuroscience, Performance

SCHOLARONE™
Manuscripts

Review Only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Short title: functional mechanisms of the quiet eye

Identifying the causal mechanisms of the quiet eye

For Peer Review Only

Abstract

Investigations into the gaze strategies employed by athletes have determined that longer quiet eye (QE) durations (QED) are characteristic of skilled compared to less-skilled performers. However, the cognitive mechanisms of the QE and, specifically, how the QED affects performance, are not yet fully understood. In this review we integrate research that has examined the functional mechanism underlying QE and based on these observations, discuss the underlying neural networks that may be involved. We also highlight the limitations surrounding QE measurement and its definition and propose future research directions to address these shortcomings. Investigations into the behavioural and neural mechanisms of QE will aid understanding of the perceptual and cognitive processes underlying expert performance and the factors that change as expertise develops. This research has important implications for the development of expertise in sports as well as in other fields including medicine and amongst clinical populations to develop and enhance goal-directed action.

Key words: gaze, programming, attention, expertise

The Quiet Eye

In recent years, attention has been devoted to the investigation of gaze behaviour to identify the perceptual and cognitive factors involved in expert performance across different sports, as well as in other domains. Vickers (Vickers, 1992) highlighted distinct gaze patterns that differentiate expert and novice golfers while performing putts and identified that experts kept a steady fixation at a specific location before ball contact. This steady fixation just before movement initiation was later identified in basketball players and termed “quiet eye” (QE; Vickers, 1996).

The QE corresponds to the final fixation within 1–3 degrees of visual angle and a duration of at least 100 ms prior to a movement. Longer quiet eye durations (QED) are exhibited by experts compared to non-experts, while within-participant analyses also show that they are characteristic of successful compared to unsuccessful attempts, (Vickers & Williams, 2007; Vickers, 1996). In addition, with the use of video-based mobile eye tracking systems, these findings have been replicated across various types of aiming and interceptive sports including shooting (Causer, Bennett, Holmes, Janelle, & Williams, 2010), darts (Rienhoff et al., 2013), billiards (Williams, Singer, & Frehlich, 2002), table tennis (Rodrigues, Vickers, & Williams, 2002), and football (Piras & Vickers, 2011).

A number of researchers have successfully used the QE as a training tool to improve performance in different targeting sports (Causer et al., 2011; Moore, Vine, Cooke, Ring, & Wilson, 2012; Vine & Wilson, 2011) and recently, outside of the sporting area such as when training surgical skills (Causer, Harvey, Snelgrove, Arsenault, & Vickers, 2014). In these studies, QE training aimed at increasing QED (where to look and for how long), results in enhanced performance linked to relative increases in QED (for a detailed review, see Vine,

1
2
3 Moore, & Wilson, 2014). However, whether or not the duration of the QE *per se* causes these
4
5 improvements in performance and how these benefits come into place are still subjects of
6
7 interest. This research has highlighted the need to better understand the underlying
8
9 mechanisms of QE and, in particular, investigate the beneficial effects of the QE on
10
11 performance to implement effective training protocols (Behan & Wilson, 2008).
12
13

14
15 Much of the research on QE has focused on identifying distinct QED effects amongst expert
16
17 and novice performers and in evaluating the training of QE as a performance-enhancing
18
19 technique. A review that focuses on the latter and on learning under anxiety can be found in
20
21 Vine et al. (2014). Vine et al. propose three potential explanations (attention control, focus
22
23 of attention and response planning) of how QE may benefit performance and expedite
24
25 learning. However, they only present one study in which the QE was directly manipulated
26
27 and questions about the causality of the QED on performance still remain. In addition,
28
29 limitations in the QE methodology have not yet been adequately addressed. Thus, in this
30
31 paper, we review current research aimed at investigating the underlying mechanisms of the
32
33 QE and the effects of directly manipulating it. Moreover, based on this current research, we
34
35 propose potential neural networks that may be involved in QE. These notions may provide
36
37 further insight into the facilitatory effects of longer QED on performance. In addition, we
38
39 aim to highlight limitations surrounding the QE definition and measurement techniques, as
40
41 well as the potential impact this may have on the interpretation of current literature.
42
43 Finally, we propose future research directions to better understand the critical processes
44
45 involved in QE.
46
47
48
49
50
51
52
53
54

55 **The programming hypothesis**

56
57
58
59
60

1
2
3 The QE is suggested to facilitate information processing and its duration reflects the time
4 needed to program and fine-tune a response (Vickers, 2009). Long QEDs are suggested to
5 extend the critical motor preparation period that consists of response selection and the fine
6 tuning of movement parameters for motor programming (Moore et al., 2012). Williams et
7 al. (2002) manipulated the complexity of a billiards shot (near versus far) as well as the
8 time needed to complete a specific shot and demonstrated that QEDs were longer in skilled
9 performers and in successful compared with unsuccessful shots. These authors further
10 demonstrated longer QED with increasing levels of task complexity and when increasing the
11 time needed for motor programming, in skilled and less-skilled billiard players. This
12 evidence supports the programming hypothesis, as longer QEDs were observed in more
13 complex tasks that required more information processing and longer programming times
14 (Vickers, 2011).
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29

30 Horn, Okumura, Alexander, Gardin, and Sylvester (2012) investigated QED and the effects of
31 variable versus blocked practice in a dart-throwing task. They hypothesized that random
32 target changes (to different targets at similar distances) would produce longer QED due to
33 the increased programming demands of having to parameterize the response from one trial
34 to the next, compared to the consistency of the blocked target presentation (same target
35 and distance). Although performance (accuracy) differences between the practice groups
36 were not apparent, longer QEDs were indeed observed during random compared to blocked
37 practice and likely reflected the heightened cognitive effort during random practice to
38 compensate for the additional task demands (increased on-line movement
39 parameterization) (Figure 1). Horn et al. (2012) concluded that the longer QED was a
40 functional element of programming demand rather than a by-product of successful aiming.
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

-Insert Figure1-

As with previous research, however, the QED effects on performance found in Williams et al.'s (2002) and Horn et al.'s (2012) studies were shown as a correlation of performance and did not reveal a causal relationship. In a recent study, Klostermann, Kredel and Hossner (2013) investigated the performance-enhancing effects of experimentally manipulated QED by varying response selection and stimulus identification demands. In their experiments, participants took part in an externally paced throwing task in which the onset of the last fixation and the amount of information to be processed were manipulated by presenting the target at different timings (short and long presentations) and locations (random and predictive) during movement unfolding. They showed that the facilitatory effects of longer QED on performance were apparent under a high information-processing load; however, QED effects seemed to disappear with increased predictability of the target's location and decreased task demands (Figure 2). Klostermann et al. (2013) argued that the predictability of the target might have facilitated relevant information processing that was not required during the QE period; consequently, QED effects were "dispensable" with high predictability and low task demands. This finding suggests that on-line control may have an important role for QE during aiming tasks (cf., Horn et al., 2012), highlighting the need to investigate these mechanisms while maintaining information load constant, as proposed by Klostermann et al. (2013). These data, together with Williams et al.'s (2002) and Horn et al.'s (Horn et al., 2012) findings, provide evidence for functional links between information processing and the effects of prolonged QED on performance. However, the precise

1
2
3 information that is important remains unclear; furthermore, additional mechanisms could
4
5 be involved, including the effective allocation of attention for information extraction.
6
7
8
9

10
11
12 **-Insert Figure2-**
13

14 15 16 17 18 **QED and attentional focus** 19

20
21
22
23
24 The allocation of visual attention is influenced by two control processes which include a
25
26 volitional (top-down) and a stimulus-driven (bottom-up) network (Corbetta & Shulman,
27
28 2002). Explanations of the QE have integrated these processes and the QE has been
29
30 suggested to aid the allocation of attention to relevant cues, while suppressing responses
31
32 from other stimuli (top-down control) (Vickers, 1996). It has also been suggested that, in
33
34 particular, external allocation of attention is responsible for the beneficial effects of the QED
35
36 on performance, as shown in QE training studies. Moore et al. (2012) showed that a QE
37
38 trained group developed longer QED and better performance compared to a technical
39
40 trained group during a golf putting task. The authors suggested that the benefits of the QED
41
42 derive from better response programming, yet the role of attentional focus (external focus)
43
44 was also highlighted, which possibly promotes longer QED, or is enhanced by the longer
45
46 QED (Moore et al., 2012; Vine & Wilson, 2011).
47
48
49

50
51 The facilitatory effects of an external focus of attention on motor performance and skill
52
53 acquisition have been previously described (for a review, see Wulf, 2007). It has been
54
55 suggested that an internal or an external focus of attention disrupts or facilitates automatic
56
57
58
59
60

1
2
3 control mechanisms respectively, and that the utilisation of these automatic processes
4 results in decreased attentional demands, smaller movement adjustments and faster
5 learning (Wulf, Shea, & Lewthwaite, 2010). Klostermann et al. (2014) also examined the
6 links between QED and focus of attention. Their results replicated previous findings, in that
7 instructed external focus resulted in better performance and experts exhibited longer QEDs.
8 However, their correlation-based findings did not provide support for a performance-
9 enhancing effect of the longer QED with an external focus of attention. It is worth noting
10 that their instructions varied from previous research in order to exclude the impact of
11 directing the participant's gaze behaviour, and the authors themselves expressed caution
12 when interpreting these results. Klostermann et al. (2014) suggested an alternate
13 explanation for their findings related to an inhibition hypothesis. It follows that the positive
14 effects of longer QED are explained by the need to inhibit alternate movement variables to
15 allow for the effective parameterisation of a single movement. Further research is needed to
16 replicate these findings to provide support for this hypothesis.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33

34
35 It is possible that some aspects of external attention allocation are important for QED;
36 however, to date the potential explanations with respect to attention allocation (i.e.,
37 movement automaticity or "constrained control theory" and a working memory load
38 hypothesis) have not been experimentally related to QE, and more research is needed to
39 probe both phenomena. In addition, given that gaze is typically directed in most QE training
40 protocols, the effects of focus of attention may be influenced by these instructions as
41 suggested by Klostermann et al. (2014). Moore et al. (2012) conducted a QE training study
42 in golf and only included instructions related to gaze control for the QE training group
43 compared to a technical trained group. They found that the QE trained group exhibited
44 more expert-like putting kinematics indirectly associated with an external focus of attention
45 (also see Causer et al. 2011). Moore et al. (2012) suggested that future research should
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 implement more than one group to manipulate focus of attention and further understand
4
5 the benefits of QE training.
6
7
8
9

10
11 The current QE research is mainly focused on identifying the functional links that explain
12 the effects of performance-enhancing QEDs. To date, the hypothesis based on the
13 information processing load of motor programming (Williams et al., 2002) seems to be a
14 suitable explanation for the QE phenomenon. Given that the QED effects have been
15 successfully observed under experimentally controlled conditions (Klostermann et al.,
16 2013; Williams et al., 2002) and during virtual tasks (see Behan & Wilson, 2008), it seems
17 promising that the mechanisms explaining the benefits of the QE can be further explored in
18 a controlled environment. From the studies included in this review, the QED effects on
19 performance are not clear-cut, in that longer QEDs do not always correlate with better
20 performance, as in Horn et al. (2012). The relative contribution of the QED on performance
21 may be addressed through these controlled QED manipulations to identify why, how and
22 when the QED effects manifest during targeting tasks. We further propose the investigation
23 of the neural mechanisms that are involved in QE using these controlled manipulations that
24 replicate the QE effects obtained in previous field research.
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

48 **Functional mechanisms of QE: psychophysiological evidence**

49
50
51
52

53 Visually-guided tasks such as targeting tasks require the accurate visual acquisition of a
54 target, the integration of visual and proprioceptive (afferent) signals for motor
55 programming and response selection to generate a motor response (Balslev, Miall, & Cole,
56
57
58
59
60

1
2
3 2007; Emma Gowen & Miall, 2006). The frontoparietal networks, particularly the posterior
4 parietal cortex (PPC), are suggested to be involved in the integration of signals related to
5 eye and limb movements (van Donkelaar & Staub, 2000). During these processes, internal
6 representations of eyes, limbs, and the target are formulated together with predictions of
7 the sensory consequences of the motor response (Wolpert, Ghahramani, & Jordan, 1995),
8 and may also engage the cerebellum in this predictive process (Miall & King, 2008).
9 Similarly, the allocation of visuospatial attention includes areas within the frontal and
10 parietal lobes (Corbetta et al., 1998). It is evident that a tight coupling between visual
11 attention orientation and the processing of the motor components of the task at hand is
12 important for performance (Behan & Wilson, 2008).
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27

28 *Attention, eye movements and QE*

29
30
31
32
33

34 The networks activated during the allocation of spatial attention include the frontal eye
35 fields (FEF), dorsolateral prefrontal cortex (DLPC) and PPC; areas that have also been
36 individually associated with the control of eye movements. There is evidence supporting the
37 strong links between attention and eye movements (Moore & Fallah, 2001, 2004; Rizzolatti,
38 Riggio, Dascola, & Umiltá, 1987). Furthermore, it has been shown that attention allocated to
39 a fixation point results in a “suppression” (reduced saccade amplitude and velocity) of the
40 oculomotor system (Goldberg, Bushnell, & Bruce, 1986), suggesting that actively fixating on
41 a location results in attention allocated to that point and not to a peripheral location. Thus,
42 there seems to be extensive overlap between programming and attention networks as well
43 as with areas involved in oculomotor control.
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 The definition of the QE implies the suppression of large eye movements between 1—3
4 degrees of visual angle and an enhanced ability to fixate on relevant cues and suppress
5 responses to irrelevant stimuli for a more efficient extraction of information. However,
6 given the definition, a fixation of 100 ms may not be long enough to allow the shift of
7 conscious attention, for example, saccade reaction times take about 200 to 250 ms (Saslow,
8 1967) after a stimulus is perceived and also, studies involving neuroimaging have shown
9 perceptual processing of visual stimuli occurring > 100 ms (Pins & Ffytche, 2003). It is also
10 important to note that the term “fixation” at 1-3 degrees of visual angle is determined by the
11 technological limitations of current gaze trackers and may encompass other types of small
12 eye movements within this definition. These eye movements may be used as a favourable
13 strategy of a given task; for example smooth pursuit in golf putting, saccades during
14 basketball throwing. FEF involvement is likely candidate given the close integration
15 between the frontoparietal network and eye movement (fixation, saccade and pursuit)
16 networks (Corbetta et al., 1998).
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34

35 With respect to the “quiet eye” terminology, it is worth noting that the eyes are seldom
36 “quiet” in that small eye movements occur during visual fixation. Fixational eye movements
37 are essential for natural vision (McCamy et al., 2012) and include small (low velocity) drifts
38 as well as (high velocity) microsaccades (< 1 degree occurring 1-2 per second) that
39 contribute to fixation stability and prevent perceptual fading of a stationary object; however
40 the precise role of microsaccades in long fixations remains controversial (for review see
41 Collewijn & Kowler, 2008). A recent study investigating fixational eye movements (drifts
42 and microsaccades) during distinct fading conditions, supports previous notions that
43 microsaccades prevent and correct visual fading during prolonged fixation (McCamy et al.,
44 2012). Furthermore, the size and frequency of the microsaccades determined the
45 effectiveness of counteracting foveal and peripheral fading (McCamy et al., 2012).
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Microsaccades can be suppressed during fine visual tasks, suggesting they may be modulated by attention, similar to larger saccades. For example, a higher rate of microsaccades to one side may indicate parafoveal information processing during visual fixation, as indicated by (Engbert & Kliegl, 2003), who investigated microsaccades during covert attention. Microsaccades during fixation appear functionally related to saccadic intrusions, which are also influenced by the shift of attention (Gowen, Abadi, Poliakoff, Hansen, & Miall, 2007). Thus, the amplitude and frequency of eye movements may provide important insights into the links between oculomotor control, visual perception and the allocation of attention in expert performance, which may be explored within the QE. Very small amplitude oculomotor noise is unlikely to disrupt vision; however, even moderate head motion requires the involvement of oculomotor compensatory mechanisms, such as the vestibulo-ocular response, optokinetic reflexes, smooth pursuit or saccades. This again suggests that other gaze behaviours may be occurring in the measured QE.

Motor programming, response selection and inhibition networks

42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

It is suggested that the ability to maintain attention and ignore/suppress external stimuli, also found to be frontally controlled (FEF and DLPFC, DeSouza, Menon, & Everling, 2003) and a characteristic observed in expert performers, aids the programming and execution of a motor response. This selective process is therefore important during the aiming period in a targeting task. Vickers (1992, 1996) supported this suppression hypothesis (also see Klostermann et al., 2014) after observations of a QE offset during movement initiation, that is, eye movements outside of the QE threshold. Vickers (1996) suggested that this suppression behaviour prevents the interference of the movement plan from online visual

1
2
3 feedback. Given that QED effects have been recently associated with more complex tasks
4 that require on-line control (such as during variable practice and the presentation of
5 random targets), it may be that feedback and continuous monitoring (afferent) signals,
6 including visual and proprioceptive information, are integrated during the QED. This finding
7 suggests interactions between top-down and bottom-up control networks (or dorsal and
8 ventral; Corbetta & Shulman, 2002) during target selection and computations for movement
9 parameterization during the QE in goal-directed actions.
10
11

12
13
14
15
16
17
18
19
20 The relative contributions of inhibition and on-line feedback integration may be addressed
21 by investigating the QE's temporal components (its onset and offset) during tasks with
22 distinct requirements for these processes. For example, in Klosterman et al.'s (2013)
23 experiments it was early QE onset that seemed to be important for accurate throwing, while
24 findings from Vine et al.'s (2013) experiment using golf putting, suggested it was the late
25 components (offset) of the QE that seem to affect performance, possibly due to higher
26 demands of on-line control functions throughout the movement. In addition, examining
27 these temporal components in QE may indicate a more efficient allocation of the timings
28 that correspond to information extraction, motor programming and movement execution.
29 These studies suggest that an important aspect of QE measurement is not only the duration
30 but also the actual timing relative to the movement.
31
32
33
34
35
36
37
38
39
40
41
42
43
44

45 Finding the optimal balance between the timings when information extraction, motor
46 programming and movement execution is particularly important for time-constrained
47 motor actions, which are encountered in many sporting contexts. The time spent preparing
48 a movement will facilitate the development of appropriate actions and will often help to
49 minimize errors (Battaglia & Schrater, 2007). Expert performers have earlier QE onsets
50 which result in longer QED, suggesting that they are able to find the relevant visual
51
52
53
54
55
56
57
58
59
60

1
2
3 information earlier and program the movement for longer irrespective of task constraints
4 (Vickers, 2009). The first neurophysiological evidence of the QE's role in motor planning is
5 reported by Mann et al. (2011), who investigated the Bereitschaftspotential (BP) during the
6 QE period of expert and near-expert golfers. Their results showed increased cortical
7 activation in right-central regions in experts compared to non-experts with an enhanced BP
8 peak and greater negativity, which is suggested to be indicative of movement preparation.
9

10
11 The intraparietal sulcus and the premotor cortex (within frontoparietal networks) have
12 been systematically shown to be involved in the programming of actions (Rushworth,
13 Johansen-Berg, Göbel, & Devlin, 2003) and in particular, the pre-supplementary motor area
14 (pre-SMA) has been associated with action selection (Mars, Piekema, Coles, Hulstijn, & Toni,
15 2007). The anterior cingulate cortex (ACC) receives input from the prefrontal cortex, also
16 playing a role in action selection (Halsband & Lange, 2006). The cingulate cortex has also
17 been associated with emotion regulation during goal-directed actions by manipulating goal
18 outcomes in various tasks (Rolls, 2014). Emotion-related decision-making has been
19 suggested to be an important aspect of the QE mechanisms. More specifically, a longer QED
20 aids emotional control by enabling the individual to get into a suitable level of arousal to
21 complete the task. This seems important in aiming tasks where a low arousal level is related
22 to better performance and a longer QE may enable the individual to 'quieten' the body. For
23 example, Vickers and Williams (2007) tested gaze control of biathletes under low and high
24 anxiety conditions and found that longer QEDs were indirectly related to better
25 performance under pressure situations and coincided with a reduced heart rate and
26 reduced physiological arousal. Furthermore, the QE trained group in Moore et al.'s (2012)
27 study, mentioned above, also showed a deceleration of heart rate prior to the golf putt
28 combined with longer QED and improved putt-kinematics, compared to the technical
29 trained group. However, phasic heart rate did not change during elevated anxiety
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 conditions in either training group, suggesting that this “quieting” of the body may explain
4 some of the benefits of QE training, but do not explain its effects during heightened anxiety
5 situations.
6
7
8

9
10 The involvement of these networks reflects higher order cognitive processing presumably
11 occurring during the QED. Studying the neural networks of a phenomenon observed in the
12 sporting field is no easy task. However, researchers have reported promising results, such
13 as the study by Wright et al. (2011), in which fMRI was used to investigate the neural
14 mechanisms associated with anticipation in badminton players of varying skill levels. The
15 authors were able to identify distinct levels of activation in task-relevant areas in experts
16 compared to novices and were able to correlate the activation of specific areas to show their
17 influence on performance (also see Heinen, Rowland, Lee, & Wade, 2006).
18
19
20
21
22
23
24
25
26
27
28
29
30
31

32 In summary, the neural networks and mechanisms responsible for the QE effects are not yet
33 fully understood. Skilled movements have been suggested to involve the
34 dorsofrontoparietal networks, associated with the allocation of attention to relevant stimuli
35 and the suppression of distractors in the environment (Corbetta & Shulman, 2002;
36 Mevorach, Hodson, Allen, Shalev, & Humphreys, 2010; Mevorach, Humphreys, & Shalev,
37 2006); however, the extent to which these networks are involved in QE needs to be
38 investigated. Current research shows that QED is a function of task complexity, a finding
39 that agrees with the current proposed response programming theory of QE. It is further
40 suggested that the QE aids the maintenance of goal-directed (dorsal or top-down attention
41 system) attention control, perhaps via superior gaze control and the effective allocation of
42 visuospatial attention, while “suppressing” a stimulus-driven attentional system (ventral or
43 bottom-up system) (Vickers, 2009). These processes then allow for the motor programming
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 to take place during the QED. Thus, the programming hypothesis may be simplifying a more
4
5 complex sequence of events that occur during the QE of a targeting task. Understanding the
6
7 mechanisms of QE will aid the formulation of training programmes, which are often based
8
9 on accepted practice rather than on procedures that try to optimise the critical processes
10
11 involved (Causer et al., 2011).
12
13

14 15 16 17 18 **Future directions and recommendations** 19

20
21
22
23
24 There is still much to be addressed when attempting to explain the links between gaze and
25
26 performance and the beneficial effects of a prolonged QED. The formulation of controlled
27
28 manipulations of QE and the investigation of QE neural networks using neuroimaging and
29
30 event related potentials (ERP) may shed some light on these links (e.g., Mann et al., 2011).
31
32 Furthermore, with current neurophysiological techniques such as transcranial magnetic
33
34 stimulation (TMS), causal relationships between anatomical regions and behaviour can be
35
36 established, in addition to determining the extent and timing of their involvement.
37
38 Neuroimaging reveals activation of specific neural networks during a performed task but
39
40 does not determine the functional role of these regions in the observed behaviour. TMS is
41
42 often used in conjunction with fMRI to determine the links between brain and behaviour by
43
44 altering the activity of a specific region and examining the affects on behaviour. For
45
46 example, Mevorach et al. (2010) combined fMRI and TMS techniques to investigate the role
47
48 of different brain regions in attention control during a task that required the suppression of
49
50 a response to stimuli. They found that TMS applied to the early visual areas led to reduced
51
52 responses to distractors, which provided evidence for the functional role of these regions in
53
54 attending or suppressing responses to salient stimuli. Similarly, the effects of QED may be
55
56
57
58
59
60

1
2
3 modulated by attention control regions, which may be determined by implementing these
4 techniques. The combination of TMS and fMRI techniques has been proven to be robust in
5
6 techniques. The combination of TMS and fMRI techniques has been proven to be robust in
7
8 examining functional mechanisms and neural networks involved in behaviour compared to
9
10 the less robust ERP measures obtained by electroencephalography (EEG). As such, caution
11
12 needs to be taken when inferring brain activation of specific areas using EEG methods given
13
14 the poor spatial resolution; however, EEG is well suited to test the timing of neural activity
15
16 in a given task. With new technology emerging, it is now possible to measure
17
18 electrophysiological responses during a golf-putting task, for example (see Mann et al.,
19
20 2011), although regional information may not be robust using this method. Thus, combining
21
22 these techniques (e.g., fMRI with EEG and/or TMS) may be of great benefit when trying to
23
24 explain spatially and timing dependant networks involved in QE.
25
26

27
28 The relative contributions of the neural networks for attention control, programming and
29
30 on-line control (functions presumably involved in QE) may depend on specific task-goal
31
32 demands. For example, golf putting requires more dynamic on-line control functions
33
34 compared to shooting a target, or shooting a moving target may require a balance between
35
36 anticipation (pre-planning) and inhibition mechanisms afforded by increased attention
37
38 control. Thus, generalisations of the functional mechanisms involved during the QE and how
39
40 they influence performance should be characterised based on task demands.
41
42

43
44 A key issue is that the definition of the QE period is restricted by the sensitivity of the
45
46 measurement systems that have been employed, which in turn impacts the interpretation of
47
48 the results. Future research should address the arbitrarily defined threshold for
49
50 determining QED and systematically evaluate whether performance and skill differences
51
52 still hold at varying thresholds (e.g., 1 vs. 2 or 2 vs. 3 degrees of visual angle), thus
53
54 examining the importance of the amplitude of eye movement during the QE. Research must
55
56
57
58
59
60

1
2
3 determine whether these eye movement differences are task or skill-related (i.e., experts
4 may have better oculomotor control) and whether alternate gaze strategies other than
5 fixations may come into play. Thus, caution should be taken when relying on fixation-
6 attention mechanisms to explain the QE phenomenon, since other gaze strategies may be
7 taking place within the QE, and distinct cognitive processes may be encompassed by a single
8 QE definition or threshold. High-resolution eye trackers (under 0.1 degrees spatial
9 resolution and sampling > 500 Hz) have limited use in real world sport settings, but may be
10 used in laboratory-based tasks in which QED effects on performance are obtained during an
11 aiming or interceptive task to better identify differences in oculomotor control and the
12 amount and/or type (pursuit, saccades, microsaccades) of eye movement related to
13 attention/inhibition mechanisms. For example: observed high frequency of microsaccades
14 versus a higher rate of larger saccades would indicate an enhanced fixation ability of the
15 former compared to the latter, which reveals disruptions in visuospatial attention. The use
16 of these high-resolution eye trackers may also address discrepancies between different gaze
17 measuring and analyses techniques for QE. QEDs have been reported using mobile video-
18 based gaze trackers of different resolutions (e.g., 30 – 60 Hz), or using video cameras only,
19 and one paper reported QED using electro-oculography (EOG; see Mann et al., 2011). The
20 sensitivity of one technique over another may indicate that some aspects of QE are
21 overlooked; for instance high variability between participants may hinder the effects of QED
22 on hit or miss trials or QED differences between expert and novice groups. Different
23 techniques may also underestimate or overestimate the duration of the QE, which has
24 implications when attempting to examine the mechanisms involved, particularly when
25 breaking the QED down into its temporal components (onset and offset).

26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55 In this review we also highlight the need to further examine the links between QE and
56 attention and address whether it is simply gaze control or gaze control and the specific
57
58
59
60

1
2
3 allocation of attention that matters in QE. Experiments similar to Klostermann et al. (2014),
4 in which gaze is not influenced by instructions but QE is manipulated may probe into these
5 attention allocation mechanisms. We also suggest that training protocols need to implement
6 approaches that go beyond the comparisons between elite and sub-elite individuals, and
7 assess whether experts differ a priori from novices. These experimental approaches may
8 include perceptual and cognitive manipulations to examine the ability to control gaze and
9 inhibit intrusive responses when aiming in a crowded environment or during high anxiety
10 situations. The response to these manipulations may change pre- to post-QE training
11 within, and between skill groups, demonstrating not only that these mechanisms are
12 important for QE but that they can be learned and adapted.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28

29 **Conclusions**

30
31
32
33
34
35 The QE appears to be an important measure of perceptual-cognitive expertise. In this
36 review we have highlighted current research that has focused on the functional mechanisms
37 of QE, involving direct manipulations and probes of the information programming theory.
38 However, there are strong links between eye movement, attention and action programming
39 networks, which may prove to be important during QE and thus, the relative contributions
40 of these still need to be addressed. The importance of understanding the mechanisms of the
41 QE will provide further knowledge of the behavioural and neural mechanisms of
42 performance-enhancing strategies used by expert performers, used to improve goal-
43 directed movements not only in sports but in other fields, such as surgery and amongst
44 clinical populations such as children with developmental coordination disorders (DCD) and
45 stroke survivors.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- Balslev, D., Miall, R. C., & Cole, J. (2007). Proprioceptive deafferentation slows down the processing of visual hand feedback. *Journal of Vision*, 7(5), 12.1–7.
doi:10.1167/7.5.12
- Battaglia, P. W., & Schrater, P. R. (2007). Humans trade off viewing time and movement duration to improve visuomotor accuracy in a fast reaching task. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 27(26), 6984–6994.
doi:10.1523/JNEUROSCI.1309-07.2007
- Behan, M., & Wilson, M. (2008). State anxiety and visual attention: the role of the quiet eye period in aiming to a far target. *Journal of Sports Sciences*, 26(2), 207–215.
doi:10.1080/02640410701446919
- Causser, J., Bennett, S. J., Holmes, P. S., Janelle, C. M., & Williams, A. M. (2010). Quiet eye duration and gun motion in elite shotgun shooting. *Medicine and Science in Sports and Exercise*, 42(8), 1599–1608. doi:10.1249/MSS.0b013e3181d1b059
- Causser, J., Harvey, A., Snelgrove, R., Arsenault, G., & Vickers, J. N. (2014). Quiet eye training improves surgical knot tying more than traditional technical training: a randomized controlled study. *American Journal of Surgery*, 208(2), 171–177.
doi:10.1016/j.amjsurg.2013.12.042
- Causser, J., Holmes, P. S., & Williams, A. M. (2011). Quiet eye training in a visuomotor control task. *Medicine and Science in Sports and Exercise*, 43(6), 1042–1049.
doi:10.1249/MSS.0b013e3182035de6
- Collewijn, H., & Kowler, E. (2008). The significance of microsaccades for vision and oculomotor control. *Journal of Vision*, 8(14), 20.1–21. doi:10.1167/8.14.20

- 1
2
3 Corbetta, M., Akbudak, E., Conturo, T. E., Snyder, A. Z., Ollinger, J. M., Drury, H. A., ... Shulman,
4
5 G. L. (1998). A common network of functional areas for attention and eye
6
7 movements. *Neuron*, *21*(4), 761–773.
8
9
10 Corbetta, M., & Shulman, G. L. (2002). Control of goal-directed and stimulus-driven attention
11
12 in the brain. *Nature Reviews. Neuroscience*, *3*(3), 201–215. doi:10.1038/nrn755
13
14 DeSouza, J. F. X., Menon, R. S., & Everling, S. (2003). Preparatory set associated with pro-
15
16 saccades and anti-saccades in humans investigated with event-related FMRI. *Journal*
17
18 *of Neurophysiology*, *89*(2), 1016–1023. doi:10.1152/jn.00562.2002
19
20
21 Engbert, R., & Kliegl, R. (2003). Microsaccades uncover the orientation of covert attention.
22
23 *Vision Research*, *43*(9), 1035–1045.
24
25
26 Goldberg, M. E., Bushnell, M. C., & Bruce, C. J. (1986). The effect of attentive fixation on eye
27
28 movements evoked by electrical stimulation of the frontal eye fields. *Experimental*
29
30 *Brain Research*, *61*(3), 579–584.
31
32
33 Gowen, E., Abadi, R. V., Poliakoff, E., Hansen, P. C., & Miall, R. C. (2007). Modulation of
34
35 saccadic intrusions by exogenous and endogenous attention. *Brain Research*, *1141*,
36
37 154–167. doi:10.1016/j.brainres.2007.01.047
38
39
40 Gowen, E., & Miall, R. C. (2006). Eye-hand interactions in tracing and drawing tasks. *Human*
41
42 *Movement Science*, *25*(4-5), 568–585. doi:10.1016/j.humov.2006.06.005
43
44
45 Halsband, U., & Lange, R. K. (2006). Motor learning in man: a review of functional and
46
47 clinical studies. *Journal of Physiology, Paris*, *99*(4-6), 414–424.
48
49 doi:10.1016/j.jphysparis.2006.03.007
50
51
52 Heinen, S. J., Rowland, J., Lee, B.-T., & Wade, A. R. (2006). An oculomotor decision process
53
54 revealed by functional magnetic resonance imaging. *The Journal of Neuroscience:*
55
56 *The Official Journal of the Society for Neuroscience*, *26*(52), 13515–13522.
57
58
59
60 doi:10.1523/JNEUROSCI.4243-06.2006

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- Horn, R. R., Okumura, M. S., Alexander, M. G. F., Gardin, F. A., & Sylvester, C. T. (2012). Quiet eye duration is responsive to variability of practice and to the axis of target changes. *Research Quarterly for Exercise and Sport*, *83*(2), 204–211. doi:10.1080/02701367.2012.10599851
- Klostermann, A., Kredel, R., & Hossner, E. J. (2014). On the interaction of attentional focus and gaze: the quiet eye inhibits focus-related performance decrements. *Journal of Sport & Exercise Psychology*, *36*(4), 392–400. doi:10.1123/jsep.2013-0273
- Klostermann, A., Kredel, R., & Hossner, E.-J. (2013). The “quiet eye” and motor performance: task demands matter! *Journal of Experimental Psychology. Human Perception and Performance*, *39*(5), 1270–1278. doi:10.1037/a0031499
- Mann, D. T. Y., Coombes, S. A., Mousseau, M. B., & Janelle, C. M. (2011). Quiet eye and the Bereitschaftspotential: visuomotor mechanisms of expert motor performance. *Cognitive Processing*, *12*(3), 223–234. doi:10.1007/s10339-011-0398-8
- Mars, R. B., Piekema, C., Coles, M. G. H., Hulstijn, W., & Toni, I. (2007). On the programming and reprogramming of actions. *Cerebral Cortex (New York, N.Y.: 1991)*, *17*(12), 2972–2979. doi:10.1093/cercor/bhm022
- McCamy, M. B., Otero-Millan, J., Macknik, S. L., Yang, Y., Troncoso, X. G., Baer, S. M., ... Martinez-Conde, S. (2012). Microsaccadic efficacy and contribution to foveal and peripheral vision. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *32*(27), 9194–9204. doi:10.1523/JNEUROSCI.0515-12.2012
- Mevorach, C., Hodsoll, J., Allen, H., Shalev, L., & Humphreys, G. (2010). Ignoring the elephant in the room: a neural circuit to downregulate salience. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *30*(17), 6072–6079. doi:10.1523/JNEUROSCI.0241-10.2010

- 1
2
3 Mevorach, C., Humphreys, G. W., & Shalev, L. (2006). Opposite biases in salience-based
4
5 selection for the left and right posterior parietal cortex. *Nature Neuroscience*, 9(6),
6
7 740–742. doi:10.1038/nn1709
8
9
10 Miall, R. C., & King, D. (2008). State estimation in the cerebellum. *Cerebellum (London,*
11
12 *England)*, 7(4), 572–576. doi:10.1007/s12311-008-0072-6
13
14 Moore, L. J., Vine, S. J., Cooke, A., Ring, C., & Wilson, M. R. (2012). Quiet eye training expedites
15
16 motor learning and aids performance under heightened anxiety: the roles of
17
18 response programming and external attention. *Psychophysiology*, 49(7), 1005–1015.
19
20 doi:10.1111/j.1469-8986.2012.01379.x
21
22
23 Moore, T., & Fallah, M. (2001). Control of eye movements and spatial attention. *Proceedings*
24
25 *of the National Academy of Sciences of the United States of America*, 98(3), 1273–
26
27 1276. doi:10.1073/pnas.021549498
28
29
30 Moore, T., & Fallah, M. (2004). Microstimulation of the frontal eye field and its effects on
31
32 covert spatial attention. *Journal of Neurophysiology*, 91(1), 152–162.
33
34 doi:10.1152/jn.00741.2002
35
36
37 Pins, D., & Ffytche, D. (2003). The neural correlates of conscious vision. *Cerebral Cortex*
38
39 *(New York, N.Y.: 1991)*, 13(5), 461–474.
40
41 Piras, A., & Vickers, J. N. (2011). The effect of fixation transitions on quiet eye duration and
42
43 performance in the soccer penalty kick: instep versus inside kicks. *Cognitive*
44
45 *Processing*, 12(3), 245–255. doi:10.1007/s10339-011-0406-z
46
47
48 Rienhoff, R., Hopwood, M. J., Fischer, L., Strauss, B., Baker, J., & Schorer, J. (2013). Transfer of
49
50 motor and perceptual skills from basketball to darts. *Frontiers in Psychology*, 4, 593.
51
52 doi:10.3389/fpsyg.2013.00593
53
54
55
56
57
58
59
60

- 1
2
3 Rizzolatti, G., Riggio, L., Dascola, I., & Umiltá, C. (1987). Reorienting attention across the
4
5 horizontal and vertical meridians: evidence in favor of a premotor theory of
6
7 attention. *Neuropsychologia*, *25*(1A), 31–40.
8
9
- 10 Rodrigues, S. T., Vickers, J. N., & Williams, A. M. (2002). Head, eye and arm coordination in
11
12 table tennis. *Journal of Sports Sciences*, *20*(3), 187–200.
13
14 doi:10.1080/026404102317284754
15
- 16 Rolls, E. T. (2014). Emotion and decision-making explained: a précis. *Cortex; a Journal*
17
18 *Devoted to the Study of the Nervous System and Behavior*, *59*, 185–193.
19
20 doi:10.1016/j.cortex.2014.01.020
21
22
- 23 Rushworth, M. F. S., Johansen-Berg, H., Göbel, S. M., & Devlin, J. T. (2003). The left parietal
24
25 and premotor cortices: motor attention and selection. *NeuroImage*, *20 Suppl 1*, S89–
26
27 100.
28
29
- 30 Saslow, M. G. (1967). Effects of components of displacement-step stimuli upon latency for
31
32 saccadic eye movement. *Journal of the Optical Society of America*, *57*(8), 1024–9.
33
- 34 Van Donkelaar, P., & Staub, J. (2000). Eye-hand coordination to visual versus remembered
35
36 targets. *Experimental Brain Research*, *133*(3), 414–418.
37
- 38 Vickers, J. N. (1992). Gaze control in putting. *Perception*, *21*(1), 117–132.
39
- 40 Vickers, J. N. (1996). Visual control when aiming at a far target. *Journal of Experimental*
41
42 *Psychology. Human Perception and Performance*, *22*(2), 342–354.
43
44
- 45 Vickers, J. N. (2009). Advances in coupling perception and action: the quiet eye as a
46
47 bidirectional link between gaze, attention, and action. *Progress in Brain Research*,
48
49 *174*, 279–288. doi:10.1016/S0079-6123(09)01322-3
50
51
- 52 Vickers, J. N. (2011). Mind over muscle: the role of gaze control, spatial cognition, and the
53
54 quiet eye in motor expertise. *Cognitive Processing*, *12*(3), 219–222.
55
56 doi:10.1007/s10339-011-0411-2
57
58
59
60

- 1
2
3 Vickers, J. N., & Williams, A. M. (2007). Performing under pressure: the effects of
4
5 physiological arousal, cognitive anxiety, and gaze control in biathlon. *Journal of*
6
7 *Motor Behavior*, 39(5), 381–394. doi:10.3200/JMBR.39.5.381-394
8
9
10 Vine, S. J., Lee, D., Moore, L. J., & Wilson, M. R. (2013). Quiet eye and choking: online control
11
12 breaks down at the point of performance failure. *Medicine and Science in Sports and*
13
14 *Exercise*, 45(10), 1988–1994. doi:10.1249/MSS.0b013e31829406c7
15
16
17 Vine, S. J., Moore, L. J., & Wilson, M. R. (2014). Quiet eye training: the acquisition, refinement
18
19 and resilient performance of targeting skills. *European Journal of Sport Science*, 14
20
21 *Suppl 1*, S235–242. doi:10.1080/17461391.2012.683815
22
23
24 Vine, S. J., & Wilson, M. R. (2011). The influence of quiet eye training and pressure on
25
26 attention and visuo-motor control. *Acta Psychologica*, 136(3), 340–346.
27
28 doi:10.1016/j.actpsy.2010.12.008
29
30 Williams, A. M., Singer, R. N., & Frehlich, S. G. (2002). Quiet eye duration, expertise, and task
31
32 complexity in near and far aiming tasks. *Journal of Motor Behavior*, 34(2), 197–207.
33
34 doi:10.1080/00222890209601941
35
36
37 Wolpert, D. M., Ghahramani, Z., & Jordan, M. I. (1995). An internal model for sensorimotor
38
39 integration. *Science (New York, N.Y.)*, 269(5232), 1880–1882.
40
41 Wright, M. J., Bishop, D. T., Jackson, R. C., & Abernethy, B. (2011). Cortical fMRI activation to
42
43 opponents' body kinematics in sport-related anticipation: expert-novice differences
44
45 with normal and point-light video. *Neuroscience Letters*, 500(3), 216–221.
46
47 doi:10.1016/j.neulet.2011.06.045
48
49
50 Wulf, G. (2007). Attentional focus and motor learning: A review of 10 years of research. *E-*
51
52 *Journal Bewegung Und Training*, 1(2-3), 1–11.
53
54
55
56
57
58
59
60

1
2
3 Wulf, G., Shea, C., & Lewthwaite, R. (2010). Motor skill learning and performance: a review
4
5 of influential factors. *Medical Education*, 44(1), 75–84. doi:10.1111/j.1365-
6
7 2923.2009.03421.x
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review Only

Figure Captions

Figure1. The graph displays the mean QED and standard deviations (SD) of random and blocked practice (RP and BP, respectively) groups in targets aimed in the horizontal and vertical axes (HA and VA, respectively) during a dart-throwing task from Horn, Okumura, Alexander, Gardin, and Sylvester (2012). The random practice group exhibited longer QED compared to participants trained under blocked practice conditions. Despite this QED differences, the authors reported no significant correlations between accuracy of the throw and QED.

Figure2. The figure displays the externally-paced throwing task (A) implemented by Klostermann et al., (2013). Figure (B) shows the mean and SD radial error (measure of performance) and (C) displays the mean and SD QE onset and offset during the low task demands and high task demands conditions (LTD and HTD, respectively) in short and long target presentations (SP and LP, respectively). Statistically significant findings are highlighted ($p < 0.05$ * and **). Complexity was manipulated by implementing a random and a predictive target presentation during movement unfolding. Their results showed earlier QE onsets, which resulted in longer QED, during more complex tasks and during longer compared to short target presentations. A significant interaction was found between task demands and target duration presentation which revealed that participants threw more accurately only in high task demands conditions. Adapted from Klostermann, Kredel, and Hossner (2013).

Unable to Convert Image

The dimensions of this image (in pixels) are too large to be converted. For this image to convert, the total number of pixels (height x width) must be less than 40,000,000 (40 megapixels).

Figure1. The graph displays the mean QED and standard deviations (SD) of random and blocked practice (RP and BP, respectively) groups in targets aimed in the horizontal and vertical axes (HA and VA, respectively) during a dart-throwing task from Horn, Okumura, Alexander, Gardin, and Sylvester (2012). The random practice group exhibited longer QED compared to participants trained under blocked practice conditions. Despite this QED differences, the authors reported no significant correlations between accuracy of the throw and QED.

Unable to Convert Image

The dimensions of this image (in pixels) are too large to be converted. For this image to convert, the total number of pixels (height x width) must be less than 40,000,000 (40 megapixels).

Figure 2. The figure displays the externally-paced throwing task (A) implemented by Klostermann et al., (2013). Figure (B) shows the mean and SD radial error (measure of performance) and (C) displays the mean and SD QE onset and offset during the low task demands and high task demands conditions (LTD and HTD, respectively) in short and long target presentations (SP and LP, respectively). Statistically significant findings are highlighted ($p < 0.05$ * and **). Complexity was manipulated by implementing a random and a predictive target presentation during movement unfolding. Their results showed earlier QE onsets, which resulted in longer QED, during more complex tasks and during longer compared to short target presentations. A significant interaction was found between task demands and target duration presentation which revealed that participants threw more accurately only in high task demands conditions. Adapted from Klostermann, Kredel, and Hossner (2013).

Review Only