

Vacuum-UV negative photoion spectroscopy of CF₃Cl, CF₃Br, and CF₃I

Simpson, Matthew; Tuckett, Richard; Dunn, KF; Hunniford, CA; Latimer, CJ

DOI:

[10.1063/1.3137103](https://doi.org/10.1063/1.3137103)

Citation for published version (Harvard):

Simpson, M, Tuckett, R, Dunn, KF, Hunniford, CA & Latimer, CJ 2009, 'Vacuum-UV negative photoion spectroscopy of CF₃Cl, CF₃Br, and CF₃I', *Journal of Chemical Physics*, vol. 130, no. 19, pp. 194302-1 - 194302-11. <https://doi.org/10.1063/1.3137103>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Vacuum-UV negative photoion spectroscopy of CF₃Cl, CF₃Br, and CF₃IM. J. Simpson,¹ R. P. Tuckett,^{1,a)} K. F. Dunn,² C. A. Hunniford,² and C. J. Latimer²¹*School of Chemistry, University of Birmingham, Edgbaston, Birmingham B15 2TT, United Kingdom*²*Department of Physics and Astronomy, Queen's University Belfast, Belfast BT7 1NN, United Kingdom*

(Received 12 February 2009; accepted 26 April 2009; published online 19 May 2009)

Using synchrotron radiation, negative ions are detected by mass spectrometry following vacuum-UV photoexcitation of trifluorochloromethane (CF₃Cl), trifluorobromomethane (CF₃Br), and trifluoroiodomethane (CF₃I). The anions F⁻, X⁻, F₂⁻, FX⁻, CF⁻, CF₂⁻, and CF₃⁻ are observed from all three molecules, where X=Cl, Br, or I, and their ion yields recorded in the range of 8–35 eV. With the exception of Br⁻ and I⁻, the anions observed show a linear dependence of signal with pressure, showing that they arise from unimolecular ion-pair dissociation. Dissociative electron attachment, following photoionization of CF₃Br and CF₃I as the source of low-energy electrons, is shown to dominate the observed Br⁻ and I⁻ signals, respectively. Cross sections for ion-pair formation are put onto an absolute scale by calibrating the signal strengths with those of F⁻ from both SF₆ and CF₄. These anion cross sections are normalized to vacuum-UV absorption cross sections, where available, and the resulting quantum yields are reported. Anion appearance energies are used to calculate upper limits to 298 K bond dissociation energies for $D^0(\text{CF}_3\text{-X})$, which are consistent with literature values. We report new data for $D^0(\text{CF}_2\text{I}^+\text{-F}) \leq 2.7 \pm 0.2$ eV and $\Delta_f H_{298}^0(\text{CF}_2\text{I}^+) \leq (598 \pm 22)$ kJ mol⁻¹. No ion-pair formation is observed below the ionization energy of the parent molecule for CF₃Cl and CF₃Br, and only weak signals (in both I⁻ and F⁻) are detected for CF₃I. These observations suggest that neutral photodissociation is the dominant exit channel to Rydberg state photoexcitation at these lower energies. © 2009 American Institute of Physics. [DOI: 10.1063/1.3137103]

I. INTRODUCTION

Ion-pair formation from a molecule is a unimolecular dissociation reaction in which two of the fragments produced are ionic; a cation-anion pair is formed. It is one of many ways in which a molecule releases energy following photoexcitation. Photoexcited states, usually Rydberg in character, may predissociate into ion pairs. This *indirect* mechanism is more favorable than *direct* ion-pair photodissociation based on Frank–Condon arguments and experimental results.¹ The formation and detection of ion pairs, therefore, can provide information on the electronic structure of a molecule and the decay dynamics of excited states. Our interest in the CF₃X series of substituted methanes, where X=Cl, Br, or I, is primarily fundamental—to compare the data and see the effects and resulting trends of changing substituent X. The interest in these molecules, however, is also environmental as CF₃Cl, CF₃Br, and CF₃I are all greenhouse gases and potential ozone depleters. The use of these molecules in industrial applications has inevitably led to their release into the atmosphere. For example, CF₃Cl (CFC-13) was used as a refrigerant and CF₃Br (Halon 1301) as a fire suppressor, but both are now banned in accordance with the Montreal Protocol.² CF₃I is considered less environmentally unfriendly than CF₃Cl or CF₃Br, and it is expected to have a relatively low atmospheric lifetime due to the weak C–I bond.³ This prop-

erty increases the potential for CF₃I applications, for example, as a plasma etching gas⁴ and as a possible replacement for CF₃Br in fire extinguishing systems.⁵

This series of CF₃X molecules have C_{3v} symmetry, and the main effect of a change in the substituent X is the elongation and subsequent weakening of the C–X bond. The effect on the overall electronic structure of the molecule on changing X is not dramatic since the orbitals of the X atom show little mixing with the CF₃ orbitals. The evidence for this property is best observed from photoelectron spectroscopy, where HeI, HeII, and threshold photoelectron spectra (TPES) have been reported for CF₃Cl, CF₃Br, and CF₃I.^{6–11} Bands observed in the spectra from ionization of an X lone pair or a C–X bonding electron shift to lower energy as X gets larger. However, bands observed from ionization of a F lone pair or a C–F bonding electron are very similar in energy for CF₃Cl, CF₃Br, and CF₃I. Absorption data on CF₃Cl have been well studied by photoabsorption spectroscopy^{12,13} and electron energy loss spectroscopy (EELS).^{14,15} More recent absorption¹⁶ and EELS (Ref. 17) studies compare data for all three CF₃X molecules. While most of this work is restricted to energies of <15 eV, absorption data for CF₃Cl are reported up to 25 eV (Refs. 13 and 18) and for CF₃Br up to 30 eV.¹⁸ Vacuum-UV (VUV) fluorescence spectroscopy has also been studied for CF₃X molecules, where X=F, H, Cl, and Br (Ref. 19) and where X=F, H, Cl, Br, and I (Ref. 18).

In this paper we report data on the negative ions formed following VUV photoexcitation of CF₃Cl, CF₃Br, and CF₃I, and ion yields have been recorded as a function of photon energy in the range of 8–35 eV using synchrotron radiation.

^{a)} Author to whom correspondence should be addressed. Telephone: +44 121 414 4425. Fax: +44 121 414 4403. Electronic mail: r.p.tuckett@bham.ac.uk.

Absolute cross sections for anions attributed to ion-pair formation have been evaluated using the negative ion data of CF_4 and SF_6 reported by Mitsuke *et al.*,^{20,21} and quantum yields have been calculated from photoabsorption data.^{16,18} The VUV photoion-pair formation of CF_3Cl was studied previously using a quadrupole mass analyzer by Schenk *et al.*,²² but to our knowledge this is the first report of ion-pair production following photoexcitation of CF_3Br and CF_3I .

II. EXPERIMENTAL

The tunable VUV radiation was provided by a 1 m Wadsworth monochromator on Beamline 3.1 at the UK Daresbury Synchrotron Radiation Source (SRS). This beamline is optimized for high flux in the 8–35 eV region of the electromagnetic spectrum.²³ All the spectra were recorded with a modest resolution of 0.6 nm. The experimental apparatus used for the detection of negative ions has been described in detail elsewhere,²⁴ and only a brief description is provided here. The gas under study is injected via a needle generating a directed jet, which bisects orthogonally the incident photon beam. The crossing point, which dictates the center of the interaction region, is positioned in the middle of two grids on the third orthogonal axis. A potential difference across the grids sweeps negative ions along this axis toward a three-element electrostatic lens for focusing, and into a Hitachi Analytical HAL IV triple quadrupole mass spectrometer (QMS) for mass selection. Detection is achieved by a channeltron electron multiplier. Sensitivity is considerably enhanced by differential pumping, which reduces the number of free electrons and secondary collisions in the QMS. The relative photon flux is measured using a sodium salicylate window and visible photomultiplier tube combination. The apparatus and QMS, connected via a 1 mm diameter aperture, are pumped separately by turbo pumps, which are backed by a common rotary pump, and the base pressure of the apparatus is $\sim 10^{-7}$ mbar. With sample gas running, the typical pressure in the chamber is $\sim 10^{-5}$ mbar. The pressure inside the chamber was measured using an ionization gauge, the sensitivity of which to CF_3Cl , CF_3Br , and CF_3I is calibrated in a separate experiment relative to N_2 gas using a capacitance manometer. Detected anion signals are initially recorded as a function of sample gas pressure over the range $(0.5\text{--}5.0) \times 10^{-5}$ mbar. Anions that show a *linear* dependence of signal with pressure most likely arise from unimolecular dissociation and are attributed to ion-pair formation. Anions that show a *nonlinear* dependence with pressure cannot be assigned as ion-pair products, and their signal is most likely influenced by secondary processes. For all anions produced from CF_3Cl , CF_3Br , and CF_3I , ion yields were recorded from 8 to 35 eV. For all scans presented below 11.8 eV (or 105 nm) a LiF window has been inserted to eliminate higher-order radiation. Gas samples were obtained from Apollo Scientific with a quoted purity of >99% and were used without further purification.

The ion yields are presented as anion cross sections σ in units of cm^2 . The method for obtaining these absolute measurements is identical to that from another recent ion-pair study and is described in detail elsewhere.²⁵ In summary, the

anion signal strengths (in counts s^{-1}) are normalized to relative photon flux, gas pressure, ring current, and relative mass sensitivity of the quadrupole. The F^- signals from both CF_4 and SF_6 are also recorded and normalized as described above. The corrected signal for F^- from SF_6 is then normalized to the known cross section at 14.3 eV, $(7 \pm 2) \times 10^{-21} \text{ cm}^2$ (Ref. 21). Likewise, the corrected signal for F^- from CF_4 is normalized to its value at 13.9 eV, $(1.25 \pm 0.25) \times 10^{-21} \text{ cm}^2$ (Ref. 20). A multiplication factor k is obtained, which converts the arbitrary normalized signals into the quoted absolute values. In theory, the values $k(\text{F}^-/\text{SF}_6)$ and $k(\text{F}^-/\text{CF}_4)$ should then be equal, but in fact they differ by a factor of 1.6. Given the number of corrections made to the anion signals, this difference seems a reasonable representation of experimental error. An average of the two k values is then used to determine absolute cross sections for the CF_3X anion signals. We comment that while these values of anion cross sections probably have an error as high as $\pm 50\%$ – 100% , such *absolute* measurements are notoriously difficult to make and prone to errors, which are often underestimated in the literature. These corrections are not made to anion signals which show a nonlinear dependence on pressure (i.e., which are *not* formed by ion-pair formation), because one of the requirements is to correct for gas pressure.

III. THERMOCHEMISTRY: GENERAL COMMENTS

Our work also determines appearance energies (AE) at 298 K for many fragment anions from CF_3Cl , CF_3Br , and CF_3I , and we compare these values with those calculated from thermochemical data. Berkowitz¹ noted that for many polyatomic molecules, a calculated threshold energy provides a lower limit to the experimental AE of an anion when suitable assumptions are made about the nature of the accompanying cation and/or neutral fragments. However, usually there is equality in these two values, although *energy* and *enthalpy* are often indistinguishable words. In comparing our experimental AE values of anions with calculated enthalpies of appropriate dissociation reactions, we make two assumptions that are justified at the relatively modest resolution of our experiment, approximately 0.1–0.2 eV. First, although it is not accurate to equate an AE_{298} to the enthalpy of the corresponding unimolecular reaction at 298 K because of thermal effects,²⁶ the corrections needed to the AE_{298} values are typically only 0.05–0.15 eV, and we feel justified in ignoring them. Second, the effects of entropy are disregarded in our calculations, even though all unimolecular reactions involve $\Delta n > 0$, where Δn is the number of product species minus the number of reactant species. Thus $\Delta_r S_{298}^\circ$ will be positive, and $\Delta_r G_{298}^\circ$ for the unimolecular reactions will be more negative than the calculated $\Delta_r H_{298}^\circ$ values.

We use enthalpies of formation $\Delta_f H_{298}^\circ$ to calculate dissociation enthalpies. The majority of these values is obtained from the JANAF tables.²⁷ Data obtained from other sources are listed below in kJ mol^{-1} . The parent molecule $\Delta_f H_{298}^\circ$ values for CF_3Cl , CF_3Br , and CF_3I are taken as -709 , -650 , and -586 , respectively, from the work of Ruscic *et al.*²⁸ In calculations for F^- , we use the value of -249 using

the electron affinity (EA) reported by Blondel *et al.*²⁹ and for F₂⁻, the value of -301, which uses the EA reported by Artau *et al.*³⁰ The values used for Cl⁻, Br⁻, and I⁻ are -227, -213, and -188, respectively, which use experimental EAs from a recent review paper.³¹ For CF⁻ we use a value of -63 using the EA(CF)=3.3±1.1 eV (reported as a lower limit)³² and for CF₂⁻ a value of -199, which uses the EA(CF₂)=0.179±0.005 eV.³¹ For CF₃⁻, we calculate $\Delta_f H_{298}^0 = -642$ kJ mol⁻¹ (Refs. 28 and 33). For CF₃⁺ we take the value of +406 reported by Garcia *et al.*³⁴ For CCl⁺ and CFCI⁺ we use +1311 and +1101, respectively.²² For CF₂Cl⁺ we use the value of +526 from Creasey *et al.*,¹⁰ but note that this uses a 0 K heat of formation of CF₂Cl. For $\Delta_f H_{298}^0$ (CF₂Br⁺), we use the upper-limit value of 570 kJ mol⁻¹ quoted by Secombe *et al.*³⁵

IV. RESULTS AND DISCUSSION

The negative ion mass spectra for the three CF₃X (X=Cl, Br, I) molecules recorded with white light at 0 nm all show the presence of the same seven anions; F⁻, X⁻, F₂⁻, FX⁻, CF⁻, CF₂⁻, and CF₃⁻. F⁻ and X⁻ are always the strongest signals. The remaining five anions were detected just above the sensitivity of the apparatus, the signals being approximately ≤2% of that of the dominant anion (F⁻ or X⁻). It was observed that the X⁻ relative signal strengths increased with increasing mass and size of X; Cl⁻=18%, Br⁻=37%, and I⁻=100% from CF₃Cl, CF₃Br, and CF₃I, respectively, of the strongest anion signal (F⁻ from CF₃Cl and CF₃Br, I⁻ from CF₃I). Of all the anions, only FI⁻ was too weak to record as a function of photon energy. Negative ion yields for all other anions are presented below.

To our knowledge this is the first report of ion-pair formation from CF₃Br and CF₃I. Similar experiments on CF₃Cl, however, have been reported in the literature.^{22,36} Of particular relevance to our study is the work of Schenk *et al.*,²² who also investigated the valence region of CF₃Cl with VUV synchrotron radiation, and comparisons between the two sets of results are detailed in the discussion below. In summary, Schenk *et al.*²² were only able to detect F⁻, Cl⁻, and CF₃⁻. CF₃⁻ was detected with low intensity and an ion yield was not recorded. The F⁻ and Cl⁻ ion yields are in excellent agreement with the results presented here.

A. F⁻ from CF₃Cl, CF₃Br, and CF₃I

The F⁻ ion yields from CF₃Cl, CF₃Br, and CF₃I are presented in Fig. 1 in the photon energy range of 8–32 eV. For comparative purposes Fig. 1 also includes the total photoabsorption spectrum,¹⁸ TPES (Ref. 10), and total fluorescence yield¹⁸ for CF₃Cl and CF₃Br, and the TPES (Ref. 11) and total fluorescence yield¹⁸ for CF₃I. The corresponding numerical data from the F⁻ ion yields are presented in Table I. The small rise in signal at 12 eV seen in the F⁻ ion yields from CF₃Cl and CF₃Br is considered to result from second-order radiation and is exaggerated by normalization to photon flux, which is low at this energy. In all three cases the F⁻ signal showed a linear rise with gas pressure, indicating that F⁻ ions are formed by unimolecular ion-pair dissociation.




FIG. 1. (Color online) Cross sections for F⁻ production following photoexcitation of (a) CF₃Cl and (b) CF₃Br between 12 and 32 eV. The total photoabsorption spectra (Ref. 18), TPES (Ref. 10), and total fluorescence yields (Ref. 18) for CF₃Cl and CF₃Br are included for comparative purposes. (c) Cross section for F⁻ production following photoexcitation of CF₃I between 8 and 32 eV. The TPES (Ref. 11, with permission from the authors) and total fluorescence yield (Ref. 18) are included for comparative purposes. All F⁻ ion yields were recorded with a step size of 0.1 eV and a wavelength resolution of 6 Å. This resolution is equivalent to 0.2 eV at 20.0 eV.

1. Onsets and thermochemistry

The F⁻ ion yield from CF₃Cl shows a gradual onset. The first indication of a rise in signal above the background is at 16.0±0.2 eV (Fig. 1, Table I). In the earlier work of Schenk *et al.*²² the F⁻ ion yield from CF₃Cl was reported with a wavelength resolution of 2 Å. They reported the onset of F⁻ ions to be 15.9±0.3 eV, correlating this onset to reaction (1) using thermochemical calculations

TABLE I. AEs, cross sections, and quantum yields for anions observed from photoexcitation of CF₃Cl, CF₃Br, and CF₃I.

Molecule [AIE ^a (eV)]	Anion	AE ^b (eV)	Cross section maximum ^c (cm ²)	Energy ^d (eV)	Quantum yield ^e
CF ₃ Cl [12.4]	F ⁻	16.0	1.5 × 10 ⁻²⁰	21.0	1.8 × 10 ⁻⁴
	Cl ⁻	16.1	2.3 × 10 ⁻²¹	20.9	2.9 × 10 ⁻⁵
	F ₂ ⁻	~21 ^f	6.8 × 10 ⁻²³	22.7	8.5 × 10 ⁻⁷
	FCl ⁻	~18 ^f	6.5 × 10 ⁻²³	20.8	8.0 × 10 ⁻⁷
	CF ⁻	25.5 ^g	1.6 × 10 ⁻²²	27.3	... ^h
	CF ₂ ⁻	20.2	1.5 × 10 ⁻²²	21.3	1.8 × 10 ⁻⁶
	CF ₃ ⁻	15.5	2.8 × 10 ⁻²²	18.1	3.5 × 10 ⁻⁶
CF ₃ Br [11.5]	F ⁻	14.7	9.7 × 10 ⁻²¹	19.6	1.2 × 10 ⁻⁴
	Br ⁻	15.1	... ⁱ ⁱ
	F ₂ ⁻	~19 ^f	2.8 × 10 ⁻²²	20.4	3.4 × 10 ⁻⁶
	FBr ⁻	~18 ^f	5.5 × 10 ⁻²²	20.4	6.6 × 10 ⁻⁶
	CF ⁻	23.6	3.4 × 10 ⁻²²	25.6	5.2 × 10 ⁻⁶
	CF ₂ ⁻	18.2	4.9 × 10 ⁻²²	19.5	5.8 × 10 ⁻⁶
	CF ₃ ⁻	13.6	2.5 × 10 ⁻²²	14.8	4.0 × 10 ⁻⁶
CF ₃ I [10.4]	F ⁻	9.7	1.1 × 10 ⁻²⁰	20.4	... ^j
	I ⁻	8.8	... ⁱ ⁱ
	F ₂ ⁻	~17 ^f	8.5 × 10 ⁻²³	20.1	... ^j
	CF ⁻	21.6	1.1 × 10 ⁻²²	23.6	... ^j
	CF ₂ ⁻	16.0	4.6 × 10 ⁻²²	16.8	... ^j
	CF ₃ ⁻	11.0	5.7 × 10 ⁻²²	12.7	... ^j

^aAdiabatic IE for CF₃Cl (Ref. 10), CF₃Br (Ref. 10), and CF₃I (Ref. 41).

^bObserved AE from this work. We estimate the error in the reported values to be ±0.2 eV, based on the resolution and step size used to record the ion yields.

^cCross section for anion production following photoexcitation of the parent molecule.

^dEnergy of peak maximum at which cross section and quantum yield measurements are taken.

^eQuantum yields for anion production obtained by dividing cross sections for anions (column 4) by total photoabsorption cross sections. The latter values are given for CF₃Cl and CF₃Br (Ref. 18).

^fCannot state AE with any confidence due to poor signal/noise.

^gThere is some ambiguity surrounding the mass of anions detected contributing to the CF⁻ ion yield from CF₃Cl. The signal observed in the range of 16–25 eV is thought to arise from Cl⁻ ions (see text), and the value of 25.5 eV represents our interpretation of the true onset to CF⁻ ions.


^hQuantum yield is not calculated because absolute photoabsorption data for CF₃Cl is not available at this energy.

ⁱThe Br⁻ and I⁻ ion yields are significantly influenced by anions arising from dissociative electron attachment and cross sections, and hence quantum yields cannot be defined.

^jQuantum yields cannot be calculated at this photon energy because the available absolute photoabsorption data for CF₃I is limited to photon energies <12 eV.



Schenk *et al.*²² also reported second (16.8 ± 0.1 eV), third (18.2 ± 0.1 eV), and fourth (20.0 ± 0.1 eV) onsets corresponding to the dissociation reactions (2)–(4), respectively,



Our thermochemical analysis, as outlined in Sec. III, agrees with all these assignments. However, the lack of well-defined onsets and features in the F⁻ ion yield from CF₃Cl, combined with the number of different dissociation channels possible, does not allow these assignments to be made with confidence. For example, the calculated dissociation enthalpies

for producing the ion pairs F⁻/CFCl⁺ (+F) [reaction (2)] and F⁻/Cl⁺ (+CF₂) are 17.0 and 17.1 eV, respectively. Not only are both these values *higher*, and not lower, in energy than the second onset, but from this analysis alone both are equally valid assignments.

The F⁻ ion yield from CF₃Br shows the first onset at 14.7 ± 0.2 eV (Fig. 1, Table I), which correlates best to the dissociation enthalpy of 14.9 eV calculated for reaction (5),



For the same reasons as discussed above in the thermochemical analysis of F⁻ from CF₃Cl, even tentative assignments of other unimolecular dissociation reactions to onsets of features in the F⁻ ion yield from CF₃Br are not suggested here.

Assignments of dissociation processes to onsets in the F⁻ ion yield from CF₃I can be made more confidently; calculated thresholds for reactions (6)–(9) coincide with local minima and hence with onsets to features in the ion yield [Fig. 1(c)],



The calculated enthalpies for reactions (6)–(9) are 14.2, 14.3, 15.7, and 18.5 eV, respectively. It is likely that features in the ion yield, which occur just after these values, represent the “turning on” of the newly available dissociation channel(s). In addition, the sharp onset observed at 12.7 ± 0.2 eV [Fig. 1(c)] can be correlated with formation of the F[−]/I⁺ (+CF₂) ion pair—although this assignment is made more tentatively since the calculated enthalpy is 13.2 eV, 0.5 eV above this onset.

The lowest energy ion-pair reaction, which yields F[−], must be



Lack of reliable information for $\Delta_f H^\circ(\text{CF}_2\text{I}^+)$ prevented a dissociation enthalpy for CF₃I in reaction (10) to be calculated. For CF₃Cl and CF₃Br the calculated thresholds for this reaction are 10.2 and ≤ 10.1 eV, respectively. In both cases these calculated dissociation enthalpies are significantly below the experimentally observed AE of F[−] ions; the AEs are 16.0 and 14.7 eV for F[−] from CF₃Cl and CF₃Br, respectively (Fig. 1, Table I). There is therefore no evidence from this thermochemical analysis that F[−] ions produced from CF₃Cl and CF₃Br arise via reaction (10). The AE for F[−] from CF₃I, however, is much lower at 9.7 eV (Fig. 1, Table I). Even though a threshold energy could not be calculated for reaction (10) when X=I, it is the only ion-pair channel forming F[−] from CF₃I that is likely to occur at energies below approximately 13 eV. The peak at 9.8 eV in the F[−] ion yield from CF₃I, albeit very weak, must therefore arise from reaction (10).

2. Discussion of the F[−] spectra

The photoabsorption spectra of CF₃Cl [Fig. 1(a)] and CF₃Br [Fig. 1(b)] (Ref. 18) extend over the energy range where F[−] ions are observed from the two molecules. Figure 1 does not include a photoabsorption spectrum for CF₃I, and published data in the energy range of interest (up to 25 eV) are limited.

The peak centered at 16.32 eV in the CF₃Cl absorption spectrum has been assigned as a transition to a 3s Rydberg orbital converging on the fifth excited valence state of CF₃Cl⁺ (\tilde{E}^2A_1).¹⁸ From EELS of CF₃Cl, King and McConkey assigned observed features at 16.29, 17.1, and 18.2 eV as transitions to 3s, 3p, and 3d Rydberg orbitals, respectively, all converging to CF₃Cl⁺ (\tilde{E}^2A_1).¹⁴ These features occur in the same energy range where the gradual onset of F[−] ions from CF₃Cl is observed. The cross section for F[−] ions in this energy range is relatively small (6×10^{-22} cm² at 17.6 eV) and well-defined peaks are not observed. As a result, and given the tentative nature of the assignments made from the photoabsorption and EEL spectra, we consider as-

signing the same transitions to the F[−] ion yield as speculative. The one peak we do observe at 21.0 eV has not been clearly observed in the absorption spectrum.¹⁸ It may correspond to a Rydberg state of CF₃Cl converging on either the \tilde{F}^2E or \tilde{G}^2A_1 state of the parent ion. The above discussion assumes the formation mechanism is predissociative, yet direct excitation to the ion-pair state should not be discounted. The gradual onset and small cross section indicate weak Frank–Condon overlap, and therefore direct ion-pair formation is plausible. If this is the case, the AE of F[−] ions may exceed the thermochemical ion-pair dissociation threshold by a greater amount than that from a predissociation mechanism where these two energies are more likely to be similar (Sec. III).

The feature in the CF₃Br photoabsorption spectrum at 15.96 eV has been assigned as a transition to a 4d Rydberg orbital converging on the fourth excited valence state of CF₃Br⁺ (\tilde{D}^2E).¹⁸ It is close in energy to the first observable peak in the F[−] ion yield at 16.1 eV, and it is possible that these two features share the same primary excitation process. The peak at 9.8 eV in the F[−] ion yield from CF₃I is very sharp and weak and appears anomalous by comparison to the rest of the spectrum. The abrupt nature of this feature points to a predissociative mechanism, and the low cross section could indicate that the extent of overlap between states is small. It has been suggested, albeit tentatively, that Rydberg states of the *ns* series converging to the $\tilde{X}^2E_{3/2}$ ionization limit lie in this energy region. Indeed there is a strong absorption band between 9.4 and 9.9 eV showing detailed structure.¹⁶

It is generally accepted that the \tilde{X}^2E electronic states of the CF₃X⁺ (X=Cl, Br, I) cations result from ionization of X lone-pair electrons and the \tilde{A}^2A_1 from ionization of a C–X bonding electron.^{6–9} The \tilde{B} , \tilde{C} , \tilde{D} , \tilde{E} , and \tilde{F} electronic states of the cations between 15 and 22 eV are most likely from fluorine lone-pair excitations. It is expected that the bonding character of the fluorine lone-pair electrons will increase with increasing ionization energy (IE).⁸ Photoexcitation of these electrons leads to the production of F[−] anions. Only F[−] produced from CF₃I is observed following photoexcitation of an electron associated with the X substituent. Even so, the resulting single peak at 9.8 eV appears isolated, and the cross section is very small compared to the rest of the spectrum. The similarities of the photoelectron spectra for the three CF₃X molecules have been highlighted by Cvitaš *et al.*,^{6,8} and they suggested that changing substituent X affects the electronic structure of the CF₃ group very little. Despite this observation, the F[−] ion yields from these three molecules differ significantly. The extent of structure and the energy range over which F[−] is observed increase as X changes from Cl to I. In addition, the AE of F[−] ions decreases. These trends appear more significant when substituting Br for I than when substituting Cl for Br. This trend possibly reflects the differing polarizabilities of the halogen atoms; the values are 2.18, 3.05, and 5.35×10^{-24} cm³ for neutral atomic Cl, Br, and I, respectively.³⁷




FIG. 2. (Color online) Cross section for Cl^- production following photoexcitation of CF_3Cl in the energy range of 12–34 eV. The total photoabsorption spectrum (Ref. 18), TPES (Ref. 10), and total fluorescence yield (Ref. 18) for CF_3Cl are included for comparative purposes. The F^- ion yield was recorded with a step size of 0.1 eV and a wavelength resolution of 6 Å. This resolution is equivalent to 0.2 eV at 20.0 eV.

B. X^- from CF_3X ($\text{X}=\text{Cl}, \text{Br}, \text{I}$)

1. Cl^- from CF_3Cl

The Cl^- ion yield from CF_3Cl is shown in Fig. 2 from 12–34 eV. For comparative purposes Fig. 2 also includes the total photoabsorption spectrum,¹⁸ TPES (Ref. 10), and total fluorescence yield¹⁸ for CF_3Cl . The numerical information is summarized in Table I. The signal in the Cl^- ion yield observed between 12 and 14 eV is considered to result from second-order effects, which are exaggerated when flux normalizing the spectrum. The Cl^- signal was shown to change linearly with CF_3Cl gas pressure, indicating that the mechanism for Cl^- formation is unimolecular ion-pair dissociation. The lowest energy ion-pair fragmentation leading to Cl^- production must also produce the cation CF_3^+ ,



The calculated enthalpy for reaction (11) is 9.2 eV. However, the experimentally observed onset to Cl^- production from CF_3Cl is 16.1 ± 0.2 eV. In the earlier work of Schenk *et al.*²² a value of 16.0 ± 0.1 eV is reported, in excellent agreement with the present work. The observed Cl^- signal at onset may be assigned to the following dissociation reaction:



The calculated enthalpy change for reaction (12) is 15.4 eV. Other onsets to features in the Cl^- ion yield, observed at 18.4, 21.3, and 23.4 eV (Fig. 2), occur where a different fragmentation reaction becomes energetically accessible,



The calculated enthalpy changes for reactions (13)–(15) are 18.4, 21.4, and 23.3 eV, respectively. We note that an experi-




FIG. 3. (Color online) (a) Br^- ion yield recorded following photoexcitation of CF_3Br between 12 and 28 eV. The TPES (Ref. 10) is superimposed in red on top of the Br^- ion yield for comparative purposes. (b) I^- ion yield recorded following photoexcitation of CF_3I between 8 and 28 eV. The 8–12 eV range of this spectrum has been blown up by a factor of 30. The TPES (Ref. 11) is superimposed in red on top of the I^- ion yield for comparative purposes. The anion spectra are not put onto an absolute scale because the signals are shown to change nonlinearly with pressure. The peak at 9.0 eV in the I^- spectrum, however, results from ion-pair formation, and the cross section at this energy is 3.8×10^{-21} cm².

mental onset occurring at a calculated thermochemical threshold suggests there is good overlap between an excited intermediate state and the new exit channel, which has energetically become open.

The production of Cl^- has similarities to that of F^- from CF_3Cl ; the fragmentation reaction assumed to occur at onset [reaction (12)] is almost identical to that assigned to F^- anions from CF_3Cl [reaction (1)]. Both ion yields show a very similar AE (Table I), and in both cases this value is much higher than the lowest energy dissociation reaction to form the respective anion as an ion pair [reactions (10) and (11)]. In addition, the cross sections for F^- and Cl^- production peak at almost identical energies (Table I) and in the range of 16–18 eV the cross sections are comparable. For example, at 17.5 eV, $\sigma_{\text{F}^-} = 5.4 \times 10^{-22}$ cm² and $\sigma_{\text{Cl}^-} = 9.2 \times 10^{-22}$ cm². Above 18 eV F^- formation increases with respect to Cl^- anions; at 21.0 eV, $\sigma_{\text{F}^-} = 1.5 \times 10^{-20}$ cm² and $\sigma_{\text{Cl}^-} = 2.2 \times 10^{-21}$ cm².

2. Br^- from CF_3Br and I^- from CF_3I

The Br^- and I^- ion yields from CF_3Br and CF_3I , respectively, are shown in Fig. 3 in the range of 8–28 eV. The TPES for CF_3Br (Ref. 10) and CF_3I (Ref. 11) are superimposed in red above the ion yields for comparative purposes. When recorded as a function of gas pressure, both the Br^- and I^- signals change nonlinearly; the rate of change in anion signal increases pseudoexponentially with increasing pressure. In instances where this trend has been seen before (e.g., SF_5^- from SF_6 and SF_5CF_3), the anions have been shown to arise from dissociative electron attachment, following photoionization of the parent molecule as the source of low-energy

electrons.²⁵ The same conclusion is reached in this study for the formation of Br⁻ and I⁻ ions from CF₃X (X=Br,I). The two-step mechanism is shown below,



CF₃Br (Refs. 38 and 39) and CF₃I (Refs. 4 and 40) are both known to attach electrons rapidly; the recommended values for the thermal electron attachment rate coefficients are $1.4 \times 10^{-8} \text{ cm}^3 \text{ s}^{-1}$ for CF₃Br (Ref. 38) and $1.9 \times 10^{-7} \text{ cm}^3 \text{ s}^{-1}$ for CF₃I (Ref. 4). In addition, the Br⁻ and I⁻ ion yields show similarities to the TPES for CF₃Br and CF₃I, respectively (Fig. 3). These similarities are much more obvious between the I⁻ ion yield and CF₃I TPES, which perhaps reflects the difference in magnitude between the attachment rate coefficients for CF₃Br and CF₃I. The apparent lack of agreement between the two spectra (ion yield versus TPES) at lower photon energies in both molecules is interesting. Only background signal is observed in the Br⁻ ion yield over the photon energy range of 12–15 eV, where the first two bands can be seen in the CF₃Br TPES. The first bands in the CF₃I TPES, representing the spin-orbit split ground state of CF₃I⁺, $\tilde{X}^2E_{3/2}$, and $\tilde{X}^2E_{1/2}$ are only observed very weakly in the I⁻ spectrum; in Fig. 3(b) the I⁻ signal over this energy region has been enlarged by a factor of 30. The ion yields in Fig. 3 are unlikely to result from dissociative electron attachment alone; Br⁻ or I⁻ anions produced by ion-pair dissociation are also detected. How much of either anion signal is due to dissociative electron attachment, and how much to ion-pair formation is unknown. However, given the evidence above it is clear that dissociative electron attachment is the more dominant mechanism contributing to the Br⁻ and I⁻ ion yields.

The agreement between the TPES and the Br⁻/I⁻ yield is slightly better at the higher energies scanned in Fig. 3, and the absence of the low-energy bands between 12 and 15 eV in the Br⁻ channel from CF₃Br and the relative weakness of the analogous bands in the I⁻ channel from CF₃I remain unexplained. Likewise, the reasons why the relative intensities between ion yield and TPES spectra are different, including the relative intensities of the $\tilde{X}^2E_{3/2}$ and $\tilde{X}^2E_{1/2}$ spin-orbit subbands in CF₃I⁺, are unclear. We note that the SF₆⁻ yield from SF₆ and the SF₅⁻ yield from SF₅CF₃ are both dominated by the two-step electron attachment mechanism over the *whole* of the valence region, and the anion yield and TPES show better agreement over a wider range of energies.²⁵ There is limited evidence from work on other polyatomic molecules (e.g., *c*-C₅F₈) that the agreement between the two spectra is enhanced if electron attachment is nondissociative.²⁵

For electron attachment to occur, the parent molecule must first be ionized. Therefore, at energies below the onset to ionization any anions produced can only arise from ion-pair dissociation. This is observed in the ion yield for I⁻ from CF₃I. The onset to ionization in CF₃I is 10.4 eV.⁴¹ However, the experimentally determined onset to I⁻ formation is at $8.8 \pm 0.2 \text{ eV}$, and a discrete peak in the signal results at 9.0




FIG. 4. Cross sections for F₂⁻ production following photoexcitation of CF₃Cl, CF₃Br, and CF₃I in the photon energy range of 12–34 eV. The ion yields were recorded with a step size of 0.1 eV and a wavelength resolution of 6 Å. This resolution is equivalent to 0.2 eV at 20.0 eV.

eV [Fig. 3(b)]. Thermochemical calculations suggest the only possible ion-pair dissociation reaction, which produces I⁻ at this energy is reaction (18),



The calculated enthalpy change for reaction (18) is 8.3 eV. We determine the cross section for I⁻ ion-pair formation at 9.0 eV to be $3.8 \times 10^{-21} \text{ cm}^2$. Normalizing this value to the total photoabsorption cross section at 9.0 eV (Ref. 16) gives a quantum yield of approximately 8×10^{-5} . An analysis of the photoabsorption spectrum of CF₃I has suggested that Rydberg states of the *ns* series converging to the $\tilde{X}^2E_{3/2}$ ionization limit lie in this energy region, and absorption features showing vibrational structure have been observed centered at energies 8.8 and 9.5 eV.¹⁶

C. F₂⁻ and FX⁻ (X=Cl, Br) from CF₃Cl, CF₃Br, and CF₃I

The F₂⁻ ion yields from CF₃X (X=Cl, Br, I) and the FX⁻ (X=Cl, Br) yields from CF₃Cl and CF₃Br in the range of 12–34 eV are shown in Figs. 4 and 5, respectively. All these anion signals show a linear increase when recorded as a function of gas pressure, indicating that F₂⁻ and FX⁻ result from unimolecular photodissociation. The figures report ab-





FIG. 5. Cross sections for FCl^- and FBr^- production following photoexcitation of CF_3Cl and CF_3Br , respectively, in the photon energy range of 12–34 eV. The ion yields were recorded with a step size of 0.1 eV and a wavelength resolution of 6 Å. This resolution is equivalent to 0.2 eV at 20.0 eV.

solite cross sections for these processes, and further numerical information is provided in Table I. The cross sections for production of FCl^- , FBr^- , and F_2^- from CF_3X are up to three orders of magnitude smaller compared to F^- production (Table I).

The onsets for F_2^- production, approximately 21, 19, and 17 eV for $X=\text{Cl}, \text{Br}, \text{I}$, occur at the thermochemical thresholds for the ion-pair dissociation reaction shown below,



The calculated dissociation enthalpy changes for reaction (19) are 21.1, 19.2, and 17.2 eV for $X=\text{Cl}, \text{Br}, \text{I}$, respectively. Two cautionary points should be made. First, the uncertainty in the values of the experimentally determined onsets (Table I, Fig. 4) is degraded by the poor signal/noise ratio in the ion yields. Second, an energy barrier resulting from forming a new F–F bond is likely. If so, the true thermochemical threshold will lie below the experimental onset, and other lower-energy dissociation reactions should be considered (e.g., F_2^-/CF^+ ion-pair formation). A similar discussion on the dissociation reactions leading to FCl^- and FBr^- from CF_3Cl and CF_3Br , respectively, is not possible due to the lack of data on the EAs of FCl and FBr .

The F_2^- ion yields all show one major feature, which most likely represents the presence of a Rydberg state converging to the fifth (\tilde{E}) or sixth (\tilde{F}) excited valence states of the CF_3X^+ molecules. As discussed in Sec. IV A 2, the origin of the excited electron is from a fluorine lone pair with significant C–F bonding character. In all three F_2^- ion yields a tentative correlation can be made between the peak energy and features in the corresponding F^- ion yields. This is unsurprising considering two F atoms must be cleaved preceding the formation of F_2^- .




FIG. 6. (Color) Cross sections for CF^- , CF_2^- , and CF_3^- production following photoexcitation of CF_3Cl , CF_3Br , and CF_3I in the photon energy range of 10–35 eV. The ion yields were recorded with a step size of 0.1 eV and a wavelength resolution of 6 Å. This resolution is equivalent to 0.2 eV at 20.0 eV.

D. CF^- , CF_2^- , and CF_3^- from CF_3Cl , CF_3Br , and CF_3I

The CF^- , CF_2^- , and CF_3^- ion yields from CF_3X ($X=\text{Cl}, \text{Br}, \text{I}$) are shown in Fig. 6. Numerical information is given in Table I. All these anion signals all show a linear rise when recorded as a function of increasing gas pressure, indicating they result from unimolecular photodissociation. The cross sections for CF_n^- ($n=1-3$) production are approximately two orders of magnitude smaller than those determined for F^- production (Table I).


Each CF_n^- ($n=1-3$) anion from each parent CF_3X molecule shows only one feature in the ion yield, with the exception of CF^- from CF_3Cl , which shows more features. We propose that the true onset for CF^- from CF_3Cl is 25.5 eV (Table I, Fig. 6) and that the observed signal in the energy range of 16–25 eV results from detecting Cl^- anions. We suggest two reasons for this. First, the mass-to-charge ratios (m/z) used when recording ion yields are close in value, 31 for CF^- and 35 for Cl^- . Although the Cl^- signal peaks at m/z 35, weak contributions can be detected at m/z values as low as 30. Combined with the fact that the CF^- signal relative to that of Cl^- is very weak, the Cl^- contribution at m/z of 31 becomes significant. Second, the ion yield of Cl^- (Sec. IV B, Fig. 2) and that of CF^- (Fig. 6) from CF_3Cl appear similar in the 16–25 eV energy range; both ion yields show an onset around 16 eV, with features at approximately 17.5 and 21 eV.

1. Onsets and thermochemistry

Unimolecular dissociation of CF₃X (X=Cl, Br, I) leading to CF₃⁻ formation must also produce the cation X⁺,



The calculated thermochemical thresholds for reaction (20) are 14.9, 13.1, and 11.0 eV when X=Cl, Br, and I, respectively; the experimentally determined onsets for CF₃⁻ anions are 15.5, 13.6, and 11.0 eV, respectively (Table I, Fig. 6). A similar dissociation process most likely produces the CF₂⁻ anions,



The calculated thermochemical thresholds for reaction (21) are 20.3, 18.5, and 16.4 eV when X=Cl, Br, and I, respectively; the experimentally determined onsets for CF₂⁻ anions are 20.2, 18.2, and 16.0 eV, respectively (Table I, Fig. 6). Dissociation of CF₃X (X=Cl, Br, I) to produce the CF₂⁻/F⁺ ion pair will only occur at excitation energies several eV above the experimental onset and is therefore not possible. Dissociation to produce the CF₂⁻/FX⁺ ion pair, however, may occur *below* the experimental onset,



The calculated thermochemical thresholds for reaction (22) are 17.4, 15.9, and 13.6 eV when X=Cl, Br, and I, respectively. If reaction (22) occurs, 2–3 eV excess energy must be accounted for. An experimental onset is always considered an upper limit, and small amounts of energy will undoubtedly be converted into translational energy of the fragment species. It should also be considered that an energy barrier to FX⁺ formation may exist, given that bonds are both broken and formed. Similar arguments are made in Sec. IV C with respect to the anions F₂⁻ and FX⁻ (X=Cl, Br, I). We consider the more likely process producing CF₂⁻ from CF₃X is reaction (21) rather than reaction (22). Low excess energies favor the production of ion pairs¹ and a bond-breaking-only dissociative reaction are favored over one where bonds are additionally formed.

The considerations discussed above are also relevant in the discussion of the CF⁻ fragment anion. The possibilities for the associated fragment cation and neutral species are greater. Several diatomic fragments, F₂, F₂⁺, FX, or FX⁺, could realistically be associated with CF⁻ ion-pair formation. The thermochemistry suggests all processes pairing CF⁻ formation with X⁺, F⁺, or F₂⁺ could be contributing to the observed CF⁻ signal from CF₃X photodissociation as observed in Fig. 6. This is perhaps reflected by the broad band, which features in all three CF⁻ ion yields.

2. Discussion of the CF_n⁻ (n=1–3) spectra

From observation in Fig. 6 it is clear that interchanging the X substituent in CF₃X with Cl, Br, or I has little effect on the structure of the ion yields of CF⁻, CF₂⁻, or CF₃⁻. However, there are consistent shifts in the AE of CF_n⁻ to lower energy as X increases in size. For example, the shift in AE for each anion is almost exactly the same when substituting Cl for Br as when substituting Br for I (Table I); the

TABLE II. Upper limits to bond dissociation energies and comparisons with literature values.

Bond	D_{298}^o (eV)	
	This work	Ref. 42
CF ₃ -F	≤(7.4 ± 0.2) ^a	5.67
CF ₃ -Cl	≤(4.4 ± 0.2) ^b	3.79
CF ₃ -Br	≤(3.6 ± 0.2) ^b	3.07
CF ₃ -I	≤(2.4 ± 0.2) ^b	2.36
CF ₂ I ⁺ -F	≤(2.7 ± 0.2) ^c	...

^aCalculated from the AE of F⁻ from CF₄ (Ref. 25).

^bCalculated from the AE of CF₃⁻ from CF₃Cl, CF₃Br, and CF₃I, respectively.

^cCalculated from the AE of F⁻ from CF₃I.

AE(CF⁻) from CF₃I is 2.0 eV lower in energy than AE(CF⁻) from CF₃Br, which is 1.9 eV lower than AE(CF⁻) from CF₃Cl. This trend is expected because all anions are observed at their thermochemical threshold, whose values decrease as the size of X increases.

The broad nature of the features in the CF⁻ ion yields does not allow any direct comparisons to be made with other spectra. In addition, the energy required to yield CF⁻ from photoexciting CF₃X (X=Cl, Br, I) is comparatively large with respect to other negative ions. Intermediate excited Rydberg states at these energies probably converge on the first inner-valence excited state of CF₃X⁺. Alternatively, these features may represent direct ion-pair formation with no involvement of an intermediate excited state. The energies of peak maxima in all the CF₂⁻ and CF₃⁻ ion yields, however, are similar to energies of features observed in other anion spectra and likely represent common excited intermediate states and hence competing ion-pair dissociation channels.

V. BOND DISSOCIATION ENERGIES

The experimental AEs for anions determined by this work may be used to calculate upper limits to bond dissociation energies, D_{298}^o .¹ For example, using the AE of CF₃⁻ can provide an upper limit to $D^o(\text{CF}_3-\text{X})$ if the IE of X and the EA of CF₃ are known, where X=Cl, Br, I,

$$\text{AE}(\text{CF}_3^-) \geq D^o(\text{CF}_3-\text{X}) + \text{IE}(\text{X}) - \text{EA}(\text{CF}_3). \quad (23)$$

Note that the AE(CF₃⁻) correlates to dissociation reaction (20). When the unimolecular dissociation involves multiple bond-breaking or the formation of a new bond, calculations performed in this way become overcomplicated and too many assumptions are made. Therefore, only AE values for anions resulting from *single* bond-breaking ion-pair dissociation are considered here. The resulting upper limits to bond dissociation energies are presented in Table II and compared to literature values. In addition $D^o(\text{CF}_3-\text{F})$ is calculated from the AE (F⁻ from CF₄)²⁵ and is also included in Table II. The uncertainty in the D^o upper limits calculated from these data is ±0.2 eV, which is taken directly from the estimated error in the AE values (Table I). The calculations for these values are explained in more detail below. We note the consistency between upper-limit values for $D^o(\text{CF}_3-\text{X})$


obtained indirectly from this ion-pair work and the accepted literature values.⁴² Furthermore, the upper-limit value for D^o tends toward the accurate value as the size of X increases from F to I. This can possibly be explained by the density of Rydberg states of CF_3X increasing as the size of X increases, and therefore the increasing likelihood that the equality of Eq. (23) holds.

As shown in Eq. (23) the AE values for CF_3^- from CF_3X (Table I, Fig. 6) are used to calculate $D^o(\text{CF}_3-\text{X})$. The EA of the CF_3 radical is 1.82 ± 0.05 eV,³³ and the IEs for Cl (12.970 eV), Br (11.816 eV), and I (10.453 eV) are taken from the JANAF thermochemical tables.²⁷ The calculation is slightly different for $D^o(\text{CF}_3-\text{F})$ because CF_3^- was not observed from CF_4 (Ref. 25), but the AE (F^- from CF_4) can be used to yield the same information if we now use the EA(F) (3.401 eV) (Ref. 29) and IE(CF_3) (9.04 ± 0.04 eV) (Ref. 34) values instead.

The formation of F^- from CF_3I at onset arises from dissociation reaction (10). Unfortunately, because the IE(CF_2I) is currently not known, an upper limit to $D^o(\text{CF}_2\text{I}-\text{F})$ cannot be calculated from the AE(F^-) value as described above. However, the relevant information is known in order to calculate an upper limit to $D^o(\text{CF}_2\text{I}^+-\text{F})$ if Eq. (24) is considered,

$$\text{AE}(\text{F}^-) \geq \text{IE}(\text{CF}_3\text{I}) + D^o(\text{CF}_2\text{I}^+-\text{F}) - \text{EA}(\text{F}). \quad (24)$$

The AE(F^-) is 9.7 ± 0.2 eV (Table I, Fig. 1), the IE(CF_3I) is 10.37 eV,⁴¹ and the EA(F) is 3.401 eV,²⁹ giving $D^o(\text{CF}_2\text{I}^+-\text{F}) \leq 2.7 \pm 0.2$ eV or 263 ± 20 kJ mol⁻¹. If $D^o(\text{CF}_2\text{I}^+-\text{F})$ is defined as the enthalpy change for reaction (25), which is valid if the Traeger and McLoughlin correction terms are ignored (Sec. III),²⁶ then an upper limit to $\Delta_f H_{298}^o(\text{CF}_2\text{I}^+)$ can be determined,



Using thermochemistry already provided (Secs. III and V) we calculate $\Delta_f H_{298}^o(\text{CF}_2\text{I}^+) \leq (598 \pm 22)$ kJ mol⁻¹.

VI. CONCLUSIONS

Negative ions have been detected following the photoexcitation of CF_3Cl , CF_3Br , and CF_3I in the photon energy range of 8–35 eV. For the fast electron-attaching gases CF_3Br and CF_3I , the Br^- and I^- signals are heavily influenced by dissociative electron attachment. All other anions detected from these three molecules result from ion-pair formation. A collection of the numerical data from this study is compiled in Tables I and II. We have shown that experimental AE values from ion-pair formation can be used to calculate upper limits for bond dissociation energies (Table II). This same point was made by Berkowitz¹ in 1996 but has rarely been implemented since. We report new data for $D^o(\text{CF}_2\text{I}^+-\text{F}) \leq 2.7 \pm 0.2$ eV and $\Delta_f H_{298}^o(\text{CF}_2\text{I}^+) \leq (598 \pm 22)$ kJ mol⁻¹.

The most surprising observation from this work is the lack of ion-pair formation detected at lower photon energies, particularly at energies below the IE of the parent molecule. This anomaly is surprising because ion-pair fragmentation is

energetically allowed and because significant structure is observed in the photoabsorption spectra below the IE. The best example of this is seen in X^- ion-pair formation from CF_3X ($\text{X}=\text{Cl}, \text{Br}, \text{I}$); a comparatively large cross section for X^- produced by reaction (26) would be predicted, but the spectra show no contribution from Cl^- or Br^- anions produced in this way. I^- anions, however, are observed below the IE of CF_3I , but the signal is surprisingly weak,



The total fluorescence yields and photoabsorption spectra correlate very little, and although there will be some contribution from fluorescence, it is not expected to be significant. Therefore, the structure observed in the photoabsorption spectra for CF_3Cl , CF_3Br , and CF_3I below the IE must almost exclusively result from neutral photodissociation. Finally, we note that ion-pair formation from CF_4 (Refs. 20 and 25) shows completely different properties from those of the CF_3X molecules studied in this paper. This should not be surprising for two reasons. First, the symmetry of the molecule changes from T_d to C_{3v} . Second, the substitution of one F by a much heavier halogen atom increases the polarizability of the molecule and therefore enhances its propensity to attach low-energy electrons.

ACKNOWLEDGMENTS

We thank Dr. David Shaw for help in running experiments on Beamline 3.1 at the Daresbury SRS, Dr. Michael Parkes for help with data collection, and Professor Ivan Powis for providing the TPE data. We also thank Dr. Sahangir Ali, whose Ph.D. thesis provided photoabsorption and fluorescence spectra for this series of molecules. Finally, we thank Dr. Michael Parkes for a critical reading of the manuscript. This collaboration between the groups in Birmingham and Belfast was partially funded by EPSRC Network Grant No. GR/N26234/01. CCLRC is thanked for the provision of SRS beamtime.

¹J. Berkowitz, in *VUV and Soft X-Ray Photoionization*, edited by U. Becker and D. A. Shirley (Plenum, New York, 1996), p. 263.

²UNEP website (<http://ozone.unep.org/>).

³S. Solomon, J. B. Burkholder, A. R. Ravishankara, and R. R. Garcia, *J. Geophys. Res. D* **99**, 20929 (1994).

⁴L. G. Christophorou and J. K. Olthoff, *J. Phys. Chem. Ref. Data* **29**, 553 (2000).

⁵Y. Li, K. O. Patten, D. Youn, and D. J. Wuebbles, *Atmos. Chem. Phys.* **6**, 4559 (2006).

⁶T. Cvitaš, H. Güsten, and L. Klasinc, *J. Chem. Phys.* **67**, 2687 (1977).

⁷R. Jadrny, L. Karlsson, L. Mattsson, and K. Siegbahn, *Phys. Scr.* **16**, 235 (1977).

⁸T. Cvitaš, H. Güsten, L. Klasinc, I. Novadj, and H. Vančik, *Z. Naturforsch [C]* **33A**, 1528 (1978).

⁹J. Doucet, P. Sauvageau, and C. Sandorfy, *J. Chem. Phys.* **58**, 3708 (1973).

¹⁰J. C. Creasey, D. M. Smith, R. P. Tuckett, K. R. Yoxall, K. Codling, and P. Hatherly, *J. Phys. Chem.* **100**, 4350 (1996).

¹¹I. Powis, O. Dutult, M. Richard-Viard, and P. Guyon, *J. Chem. Phys.* **92**, 1643 (1990).

¹²R. Gilbert, P. Sauvageau, and C. Sandorfy, *J. Chem. Phys.* **60**, 4820 (1974).

¹³H. W. Jochims, W. Lohr, and H. Baumgärtel, *Ber. Bunsenges. Phys. Chem.* **80**, 130 (1976).

¹⁴G. C. King and J. W. McConkey, *J. Phys. B* **11**, 1861 (1978).

¹⁵J. W. Au, G. R. Burton, and C. E. Brion, *Chem. Phys.* **221**, 151 (1997).

- ¹⁶ S. Eden, P. Limão-Vieira, S. V. Hoffmann, and N. J. Mason, *Chem. Phys.* **323**, 313 (2006).
- ¹⁷ M. Hoshino, K. Sunohara, C. Makohekanwa, L. Pichl, H. Cho, and H. Tanaka, *J. Chem. Phys.* **126**, 024303 (2007).
- ¹⁸ S. Ali, Ph.D. thesis, University of Birmingham, 2007.
- ¹⁹ H. Biehl, K. J. Boyle, R. P. Tuckett, H. Baumgärtel, and H. W. Jochims, *Chem. Phys.* **214**, 367 (1997).
- ²⁰ K. Mitsuke, S. Suzuki, T. Imamura, and I. Koyano, *J. Chem. Phys.* **95**, 2398 (1991).
- ²¹ K. Mitsuke, S. Suzuki, T. Imamura, and I. Koyano, *J. Chem. Phys.* **93**, 8717 (1990).
- ²² H. Schenk, H. Oertel, and H. Baumgärtel, *Ber. Bunsenges. Phys. Chem.* **83**, 683 (1979).
- ²³ C. R. Howle, S. Ali, R. P. Tuckett, D. A. Shaw, and J. B. West, *Nucl. Instrum. Methods Phys. Res. B* **237**, 656 (2005).
- ²⁴ C. A. Hunniford, S. W. J. Scully, K. F. Dunn, and C. J. Latimer, *J. Phys. B* **40**, 1225 (2007).
- ²⁵ M. J. Simpson, R. P. Tuckett, K. F. Dunn, C. A. Hunniford, C. J. Latimer, and S. W. J. Scully, *J. Chem. Phys.* **128**, 124315 (2008).
- ²⁶ J. C. Traeger and R. G. McLoughlin, *J. Am. Chem. Soc.* **103**, 3647 (1981).
- ²⁷ M. W. Chase, *J. Phys. Chem. Ref. Data Monogr.* **9**, 1 (1998).
- ²⁸ B. Ruscic, J. V. Michael, P. C. Redfern, L. A. Curtiss, and K. Raghavachari, *J. Phys. Chem. A* **102**, 10889 (1998).
- ²⁹ C. Blondel, C. Delsart, and F. Goldfarb, *J. Phys. B* **34**, L281 (2001).
- ³⁰ A. Artau, K. E. Nizzi, B. T. Hill, L. S. Sunderlin, and P. G. Wenthold, *J. Am. Chem. Soc.* **122**, 10667 (2000).
- ³¹ J. C. Rienstra-Kiracofe, G. S. Tschumper, H. F. Schaefer, S. Nandi, and B. Ellison, *Chem. Rev. (Washington, D.C.)* **102**, 231 (2002).
- ³² J. C. J. Thynne and K. A. G. MacNeil, *Int. J. Mass Spectrom. Ion Phys.* **5**, 329 (1970).
- ³³ H. J. Deyerl, L. S. Alconcel, and R. E. Continetti, *J. Phys. Chem. A* **105**, 552 (2001).
- ³⁴ G. A. Garcia, P. M. Guyon, and I. Powis, *J. Phys. Chem. A* **105**, 8296 (2001).
- ³⁵ D. P. Secombe, R. P. Tuckett, and B. O. Fisher, *J. Chem. Phys.* **114**, 4074 (2001).
- ³⁶ S. W. J. Scully, Ph.D. thesis, Queen's University Belfast, 2004.
- ³⁷ D. R. Lide, *Handbook of Chemistry and Physics*, 88th ed. (Taylor & Francis, London, 2007), Sec. 10, pp. 194–195.
- ³⁸ L. G. Christophorou, *Z. Phys. Chem.* **195**, 195 (1996).
- ³⁹ S. Marienfeld, T. Sunagawa, I. I. Fabrikant, M. Braun, M.-W. Ruf, and H. Hotop, *J. Chem. Phys.* **124**, 154316 (2006).
- ⁴⁰ S. Marienfeld, I. I. Fabrikant, M. Braun, M.-W. Ruf, and H. Hotop, *J. Phys. B* **39**, 105 (2006).
- ⁴¹ N. A. Macleod, S. Wang, J. Hennessy, T. Ridley, K. P. Lawley, and R. J. Donovan, *J. Chem. Soc., Faraday Trans.* **94**, 2689 (1998).
- ⁴² D. R. Lide, *Handbook of Chemistry and Physics*, 88th ed. (Taylor & Francis, London, 2007), Sec. 9, pp. 63–64.