# UNIVERSITY<sup>OF</sup> BIRMINGHAM University of Birmingham Research at Birmingham

## An improved lower bound for Folkman's theorem

Balogh, Jozsef; Eberhard, Sean; Narayanan, Bhargav; Treglown, Andrew; Wagner, Adam Zsolt

DOI: 10.1112/blms.12058

*License:* Other (please specify with Rights Statement)

Document Version Peer reviewed version

*Citation for published version (Harvard):* Balogh, J, Eberhard, S, Narayanan, B, Treglown, A & Wagner, AZ 2017, 'An improved lower bound for Folkman's theorem', *Bulletin of the London Mathematical Society*, vol. 49, no. 4, pp. 745-747. https://doi.org/10.1112/blms.12058

Link to publication on Research at Birmingham portal

#### Publisher Rights Statement:

This is the peer reviewed version of the following article: [FULL CITE], which has been published in final form at [Link to final article using the DOI]. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving

#### **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

•Users may freely distribute the URL that is used to identify this publication.

Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)

•Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

#### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

#### AN IMPROVED LOWER BOUND FOR FOLKMAN'S THEOREM

#### JÓZSEF BALOGH, SEAN EBERHARD, BHARGAV NARAYANAN, ANDREW TREGLOWN, AND ADAM ZSOLT WAGNER

ABSTRACT. Folkman's theorem asserts that for each  $k \in \mathbb{N}$ , there exists a natural number n = F(k) such that whenever the elements of [n] are two-coloured, there exists a set  $A \subset [n]$  of size k with the property that all the sums of the form  $\sum_{x \in B} x$ , where B is a nonempty subset of A, are contained in [n] and have the same colour. In 1989, Erdős and Spencer showed that  $F(k) \geq 2^{ck^2/\log k}$ , where c > 0 is an absolute constant; here, we improve this bound significantly by showing that  $F(k) \geq 2^{2^{k-1}/k}$  for all  $k \in \mathbb{N}$ .

#### 1. INTRODUCTION

Schur's theorem, proved in 1916, is one of the central results of Ramsey theory and asserts that whenever the elements of  $\mathbb{N}$  are finitely coloured, there exists a monochromatic set of the form  $\{x, y, x + y\}$  for some  $x, y \in \mathbb{N}$ . About fifty years ago, a wide generalisation of Schur's theorem was obtained independently by Folkman, Rado and Sanders, and this generalisation is now commonly referred to as Folkman's theorem (see [2], for example). To state Folkman's theorem, it will be convenient to have some notation. For  $n \in \mathbb{N}$ , we write [n] for the set  $\{1, 2, \ldots, n\}$ , and for a finite set  $A \subset \mathbb{N}$ , let

$$S(A) = \left\{ \sum_{x \in B} x : B \subset A \text{ and } B \neq \emptyset \right\}$$

denote the set of all *finite sums* of A. In this language, Folkman's theorem states that for all  $k, r \in \mathbb{N}$ , there exists a natural number n = F(k, r) such that whenever the elements of [n] are r-coloured, there exists a set  $A \subset [n]$  of size k such that S(A) is a monochromatic subset of [n]; of course, it is easy to see that Folkman's theorem, in the case where k = 2, implies Schur's theorem.

Date: 27 February 2017.

<sup>2010</sup> Mathematics Subject Classification. Primary 05D10; Secondary 05D40.

In this note, we shall be concerned with lower bounds for the two-colour Folkman numbers, i.e., for the quantity F(k) = F(k, 2). In 1989, Erdős and Spencer [1] proved that

$$F(k) \ge 2^{ck^2/\log k} \tag{1}$$

for all  $k \in \mathbb{N}$ , where c > 0 is an absolute constant; here, and in what follows, all logarithms are to the base 2. Our primary aim in this note is to improve (1).

Before we state and prove our main result, let us say a few words about the proof of (1). Erdős and Spencer establish (1) by considering uniformly random two-colourings. In particular, they show that if [n] is two-coloured uniformly at random and additionally  $n \leq 2^{ck^2/\log k}$  for some suitably small absolute constant c > 0, then with high probability, there is no k-set  $A \subset [n]$  for which S(A) is monochromatic. On the other hand, it is not hard to check that if  $n \geq 2^{Ck^2}$  for some suitably large absolute constant C > 0, then a two-colouring of [n] chosen uniformly at random is such that, with high probability, there exists a set  $A \subset [n]$  of size k for which S(A) is monochromatic; indeed, to see this, it is sufficient to restrict our attention to sets of the form  $\{p, 2p, \ldots, kp\}$ , where p is a prime in the interval  $[n/\log^2 n, 2n/\log^2 n]$ , and notice that the sets of finite sums of such sets all have size k(k + 1)/2 and are pairwise disjoint. With perhaps this fact in mind, in their paper, Erdős and Spencer also describe some of their attempts at removing the factor of log k in the exponent in (1); nevertheless, their bound has not been improved upon since.

Our main contribution is a new, doubly exponential, lower bound for F(k), significantly strengthening the bound due to Erdős and Spencer.

**Theorem 1.1.** For all  $k \in \mathbb{N}$ , we have

$$F(k) \ge 2^{2^{k-1}/k}.$$
(2)

This short note is organised as follows. We give the proof of Theorem 1.1 in Section 2 and conclude with some remarks in Section 3.

#### 2. Proof of the main result

In this section, we give the proof of Theorem 1.1.

Proof of Theorem 1.1. The result is easily verified when  $k \leq 3$ , so suppose that  $k \geq 4$  and let  $n = \lfloor 2^{2^{k-1}/k} \rfloor$ . In the light of our earlier remarks, a uniformly random

colouring of [n] is a poor candidate for establishing (2). Instead, we generate a (random) red-blue colouring of [n] as follows: we first red-blue colour the odd elements of [n] uniformly at random, and then extend this colouring uniquely to all of [n] by insisting that the colour of 2x be different from the colour of x for each  $x \in [n]$ ; hence, for example, if 5 is initially coloured blue, then 10 gets coloured red, 20 gets coloured blue, and so on.

Fix a set  $A \subset [n]$  of size k with  $S(A) \subset [n]$ . We have the following estimate for the probability that S(A) is monochromatic in our colouring.

### Claim 2.1. $\mathbb{P}(S(A) \text{ is monochromatic}) \leq 2^{1-2^{k-1}}$ .

Proof. First, if  $|S(A)| \leq 2^k - 2$ , then it is easy to see from the pigeonhole principle that there exist two subsets  $B_1, B_2 \subset A$  such that  $\sum_{x \in B_1} x = \sum_{x \in B_2} x$ , and by removing  $B_1 \cap B_2$  from both  $B_1$  and  $B_2$  if necessary, these sets may further be assumed to be disjoint; in particular, this implies that S(A) contains two elements one of which is twice the other. It therefore follows from the definition of our colouring that S(A) cannot be monochromatic unless  $|S(A)| = 2^k - 1$ .

Next, suppose that  $|S(A)| = 2^k - 1$ . For each odd integer  $m \in \mathbb{N}$ , we define  $G_m = \{m, 2m, 4m, \dots\} \cap [n]$ , and note that these geometric progressions partition [n]. Observe that S(A) intersects at least  $2^{k-1}$  of these progressions. Indeed, if there is an odd integer  $r \in A$  for example, then S(A) contains exactly  $2^{k-1}$  distinct odd elements and these elements must lie in different progressions. More generally, if each element of A is divisible by  $2^s$  and s is maximal, then there exists an element r of A with  $r = 2^{st}t$ , where t is odd; it is then clear that precisely  $2^{k-1}$  elements of S(A) are divisible by  $2^s$  but not by  $2^{s+1}$  and these elements must necessarily lie in different progressions. With this in mind, we define  $B_A$  to be a maximal subset of S(A) with the property  $|B_A \cap G_m| \leq 1$  for each m; for example, we may take  $B_A$  to consist of the least elements (where they exist) of the sets  $S(A) \cap G_m$ . Clearly, our red-blue colouring restricted to  $B_A$  is a uniformly random colouring, so the probability that  $B_A$  is monochromatic is  $2^{1-|B_A|} \leq 2^{1-2^{k-1}}$ .

It is now easy to see that if X is the number of sets  $A \subset [n]$  of size k for which S(A) is a monochromatic subset of [n] in our colouring, then

$$\mathbb{E}[X] \le \binom{n}{k} 2^{1-2^{k-1}} \le \left(\frac{en}{k}\right)^k 2^{1-2^{k-1}} \le \left(\frac{e2^{2^{k-1}/k}}{k}\right)^k \left(2^{1-2^{k-1}}\right) = 2\left(\frac{e}{k}\right)^k < 1,$$

where the last inequality holds for all  $k \ge 4$ . Hence, there exists a red-blue colouring of [n] without any set A of size k for which S(A) is a monochromatic subset of [n], proving the result.

#### 3. CONCLUSION

We conclude this note with two remarks. First, using the original arguments of Erdős and Spencer [1] in conjunction with an inverse Littlewood–Offord theorem of Nguyen and Vu [3], it is possible to improve (1) (up to removing the factor of log k in the exponent) by just considering uniformly random two-colourings. Second, we note that while (2) improves significantly on (1), this lower bound is still considerably far from the best upper bound for F(k), which is of tower type; see [4], for instance.

#### Acknowledgements

The first author was partially supported by NSF Grant DMS-1500121 and an Arnold O. Beckman Research Award (UIUC Campus Research Board 15006). The fourth author would like to acknowledge support from EPSRC grant EP/M016641/1.

Some of the research in this paper was carried out while the first author was a Visiting Fellow Commoner at Trinity College, Cambridge and the fourth and fifth authors were visiting the University of Cambridge; we are grateful for the hospitality of both the College and the University.

#### References

- P. Erdős and J. Spencer, *Monochromatic sumsets*, J. Combin. Theory Ser. A 50 (1989), 162–163.
- R. L. Graham, B. L. Rothschild, and J. H. Spencer, *Ramsey theory*, 2<sup>nd</sup> ed., Wiley-Interscience Series in Discrete Mathematics and Optimization, John Wiley & Sons, Inc., New York, 1990.
- H. Nguyen and V. Vu, Optimal inverse Littlewood-Offord theorems, Adv. Math. 226 (2011), 5298–5319.
- A. D. Taylor, Bounds for the disjoint unions theorem, J. Combin. Theory Ser. A 30 (1981), 339–344.

Department of Mathematics, University of Illinois, 1409 W. Green Street, Urbana IL 61801, USA

E-mail address: jobal@math.uiuc.edu

London NW53LT, UK

*E-mail address*: eberhard.math@gmail.com

DEPARTMENT OF PURE MATHEMATICS AND MATHEMATICAL STATISTICS, UNIVERSITY OF CAMBRIDGE, WILBERFORCE ROAD, CAMBRIDGE CB30WB, UK

E-mail address: b.p.narayanan@dpmms.cam.ac.uk

School of Mathematics, University of Birmingham, Edgbaston, Birmingham, B15 $2\mathrm{TT},\,\mathrm{UK}$ 

*E-mail address*: a.c.treglown@bham.ac.uk

Department of Mathematics, University of Illinois, 1409 W. Green Street, Urbana IL 61801, USA

*E-mail address*: zawagne2@illinois.edu