

UNIVERSITY OF BIRMINGHAM

University of Birmingham Research at Birmingham

ϕ -Meson production at forward rapidity in p–Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV and in pp collisions at $\sqrt{s} = 2.76$ TeV

Adam, J.; Alexandre, D.; Barnby, L.S.; Graham, K.L.; Jones, P.G.; Jusko, A.; Krivda, M.; Lee, G.R.; Lietava, R.; Villalobos Baillie, O.; Zardoshti, N.; Evans, David

DOI:

[10.1016/j.physletb.2017.01.074](https://doi.org/10.1016/j.physletb.2017.01.074)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Adam, J, Alexandre, D, Barnby, LS, Graham, KL, Jones, PG, Jusko, A, Krivda, M, Lee, GR, Lietava, R, Villalobos Baillie, O, Zardoshti, N & Evans, D 2017, ' ϕ -Meson production at forward rapidity in p–Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV and in pp collisions at $\sqrt{s} = 2.76$ TeV', *Physics Letters B*, vol. 768, pp. 203-217.
<https://doi.org/10.1016/j.physletb.2017.01.074>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.



ϕ -Meson production at forward rapidity in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV and in pp collisions at $\sqrt{s} = 2.76$ TeV

ALICE Collaboration*



ARTICLE INFO

Article history:

Received 16 November 2015

Received in revised form 8 December 2016

Accepted 9 January 2017

Available online 16 February 2017

Editor: L. Rolandi

ABSTRACT

The first study of ϕ -meson production in p-Pb collisions at forward and backward rapidity, at a nucleon-nucleon centre-of-mass energy $\sqrt{s_{NN}} = 5.02$ TeV, has been performed with the ALICE apparatus at the LHC. The ϕ -mesons have been identified in the dimuon decay channel in the transverse momentum (p_T) range $1 < p_T < 7$ GeV/c, both in the p-going ($2.03 < y < 3.53$) and the Pb-going ($-4.46 < y < -2.96$) directions – where y stands for the rapidity in the nucleon-nucleon centre-of-mass – the integrated luminosity amounting to 5.01 ± 0.19 nb $^{-1}$ and 5.81 ± 0.20 nb $^{-1}$, respectively, for the two data samples. Differential cross sections as a function of transverse momentum and rapidity are presented. The forward-backward ratio for ϕ -meson production is measured for $2.96 < |y| < 3.53$, resulting in a ratio ~ 0.5 with no significant p_T dependence within the uncertainties. The p_T dependence of the ϕ nuclear modification factor R_{Pb} exhibits an enhancement up to a factor 1.6 at $p_T = 3\text{--}4$ GeV/c in the Pb-going direction. The p_T dependence of the ϕ -meson cross section in pp collisions at $\sqrt{s} = 2.76$ TeV, which is used to determine a reference for the p-Pb results, is also presented here for $1 < p_T < 5$ GeV/c and $2.5 < y < 4$, for a 78 ± 3 nb $^{-1}$ integrated luminosity sample.

© 2017 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. Introduction

Proton-nucleus (p-A) collisions are of special interest in the context of high-energy nuclear physics for two reasons. On one hand, a precise characterisation of particle production processes in p-A collisions is needed as a reference for nucleus-nucleus data. This allows in-medium effects – linked to the formation of a deconfined phase of the QCD matter, the quark-gluon plasma (QGP) [1–3] – to be disentangled from the effects already present in cold nuclear matter. Among them, a sizeable role is played by the transverse momentum broadening of initial-state partons due to multiple scattering inside the nucleus, responsible for the Cronin effect [4], which may lead to an enhancement of intermediate- p_T hadron spectra. In addition, p-A collisions at LHC energies provide a way to probe the parton distributions of the colliding nucleus at small values of Bjorken-x, in a regime where parton densities can reach saturation [5,6]. In particular, the smallest x values contributing to the wave function of the colliding nucleus can be probed by looking at particle production at large rapidities, in the p-going direction. Such a measurement can thus extend towards lower x-values the results of the lower-energy measurements by the PHOBOS and BRAHMS experiments at RHIC [7,8]. Measurements of identified particle production may,

in particular, provide useful constraints for forthcoming theoretical studies of the saturation mechanism at small x.

We have already reported results on charged particle production in p-Pb collisions at mid-rapidity. These results focused on the pseudorapidity density [9] and the p_T dependence of the nuclear modification factor [10–12]; the latter was found to be consistent with unity for $p_T \gtrsim 2$ GeV/c. The nuclear modification factor of charged hadrons was also studied by the BRAHMS and PHOBOS Collaborations in d-Au collisions at the nucleon-nucleon centre-of-mass energy $\sqrt{s_{NN}} = 200$ GeV at RHIC [13,14], as a function of pseudorapidity, where values smaller than unity were found for $\eta \gtrsim 1$, corresponding to the d-going direction.

In this Letter we report the measurement of ϕ -meson production at forward rapidity in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV in the transverse momentum (p_T) range $1 < p_T < 7$ GeV/c, for the centre-of-mass rapidity (y) ranges $2.03 < y < 3.53$ (p-going direction) and $-4.46 < y < -2.96$ (Pb-going direction), in the dimuon decay channel with the ALICE detector. This measurement extends the investigation of light-flavour particle production to forward rapidity. At the same time, it represents an essential baseline for the understanding of ϕ production in heavy-ion collisions, where an enhancement of strange particle yields relative to the ones measured in pp collisions has been proposed long ago as a signature of the formation of a QGP phase [15–17] triggering an intense experimental effort already at SPS and RHIC energies [18–24]. It should

* E-mail address: alice-publications@cern.ch.

be noted that, despite its hidden strangeness, producing a ϕ -meson in a hadronic collision still implies the creation of a $s\bar{s}$ pair as it is the case for other strange hadrons, even if the hadronisation mechanisms can differ in reason of the different quark composition. In this context, the p–Pb data presented here will provide an important reference for future measurements in Pb–Pb collisions in the LHC Run 2, which will be performed at a comparable energy.

The differential ϕ -meson cross section as a function of transverse momentum is also presented for pp collisions at $\sqrt{s} = 2.76$ TeV. This measurement complements the ALICE results on ϕ -meson production in pp collisions at $\sqrt{s} = 7$ TeV, already reported in [25] and, combined with the latter, is used to build the pp reference for the p–Pb measurements presented here.

2. Experimental setup

A full description of the ALICE detector can be found in [26,27]. The results presented in this Letter have been obtained detecting muon pairs with the muon spectrometer, covering the pseudorapidity region $-4 < \eta_{\text{lab}} < -2.5$. Here and in the following, the sign of η_{lab} is determined by the choice of the LHC reference system. The other detectors relevant for the analysis are the Silicon Pixel Detector (SPD) of the Inner Tracking System (ITS), the V0 detector and the Zero Degree Calorimeters (ZDC).

The elements of the muon spectrometer are a hadron absorber, followed by a set of tracking stations, a dipole magnet, an iron wall acting as muon filter and a set of trigger stations. The hadron absorber is made of carbon, concrete and steel and is placed 0.9 m away from the interaction point. Its total material budget corresponds to 10 hadronic interaction lengths. The dipole magnet provides an integrated magnetic field of 3 T · m in the vertical direction. The muon tracking is provided by five tracking stations, each one composed of two cathode pad chambers. The first two stations are located upstream of the dipole magnet, the third one in the middle of its gap and the last two downstream of it. A 1.2 m thick iron wall, corresponding to 7.2 hadronic interaction lengths, is placed between the tracking and trigger detectors and absorbs the residual secondary hadrons emerging from the hadron absorber. The hadron absorber together with the iron wall stops muons with total momentum lower than ~ 4 GeV/c. The muon trigger detector consists of two stations, each one composed of two planes of resistive plate chambers, installed downstream of the muon filter.

The SPD consists of two silicon pixel layers, covering the pseudorapidity regions $|\eta_{\text{lab}}| < 2.0$ and $|\eta_{\text{lab}}| < 1.4$ for the inner and outer layer, respectively. It is used for the determination of the primary interaction vertex position. The V0 is composed of two scintillator hodoscopes covering the pseudorapidity regions $2.8 < \eta_{\text{lab}} < 5.1$ and $-3.7 < \eta_{\text{lab}} < -1.7$. It is used in the definition of the minimum bias trigger signal, and allows the offline rejection of beam-halo and beam-gas interactions to be performed. The ZDC detectors, positioned symmetrically at 112.5 m from the interaction point, are used to clean the event sample by removing beam-beam collisions not originating from nominal LHC bunches.

3. Data selection and signal extraction

The analysis presented in this Letter is based on two data samples, collected by ALICE during the 2013 p–Pb and pp runs at $\sqrt{s_{\text{NN}}} = 5.02$ TeV and $\sqrt{s} = 2.76$ TeV, respectively. In this section we present the details of the data selection, as well as the procedure followed for the extraction of the ϕ -meson signal.

3.1. Data selection

The Minimum-Bias (MB) trigger for the considered data sample is given by the logical AND of the signals in the two V0 detec-

tors [28]. Events containing a muon pair are selected by means of a specific dimuon trigger, based on the detection of two muon candidate tracks in the trigger system of the muon spectrometer, in coincidence with the MB condition. Due to the intrinsic momentum cut imposed by the detector, only muons with $p_T \gtrsim 0.5$ GeV/c manage to leave a signal in the trigger chambers.

Because of the different energy of the LHC proton and Pb beams ($E_p = 4$ TeV, $E_{\text{Pb}} = 1.58$ A · TeV), in p–Pb collisions the nucleon-nucleon centre-of-mass moves in the laboratory with a rapidity $y_0 = 0.465$ in the direction of the proton beam. The directions of the proton and Pb beam orbits were inverted during the p–Pb data taking period. This allowed the ALICE muon spectrometer to access two different rapidity regions¹: the region $2.03 < y < 3.53$ where the proton beam is directed towards the muon spectrometer (p-going direction) and the region $-4.46 < y < -2.96$ where the Pb beam is directed towards the muon spectrometer (Pb-going direction). In the following, these two rapidity ranges are also referred to as “forward” and “backward”, respectively. For pp collisions at $\sqrt{s} = 2.76$ TeV the muon spectrometer covers the rapidity region $2.5 < y < 4$.²

Background events not coming from beam-beam interactions are rejected by performing an offline selection, based on the requirement that the timing signals from the V0 and ZDC detectors are compatible with a collision occurring in the fiducial interaction region $|z_{\text{vtx}}| \lesssim 10$ cm.

The integrated luminosity for the p–Pb data samples was evaluated as $L_{\text{int}} = N_{\text{MB}}/\sigma_{\text{MB}}$, where N_{MB} is the number of MB events corresponding to the analysed triggered events, and σ_{MB} the MB trigger cross section. The value of N_{MB} was obtained by averaging the results of two different methods – one based on the ratio of trigger rates and the other based on the offline selection of dimuon events in the MB data sample [29] – while the MB trigger cross sections σ_{MB} were measured with a van der Meer scan and found to be 2.09 ± 0.07 b and 2.12 ± 0.07 b, respectively, for the beam configurations corresponding to the forward and backward rapidity coverage of the muon spectrometer [30]. For the pp data sample, the integrated luminosity is calculated with the method described in [31], using as reference the MB trigger cross section $\sigma_{\text{MB}} = 47.7 \pm 0.9$ mb, measured in a van der Meer scan [32].

The resulting values of L_{int} for the analysed p–Pb data samples are 5.01 ± 0.19 nb⁻¹ and 5.81 ± 0.20 nb⁻¹ [29,30] – corresponding to $\sim 24\,000$ and $\sim 26\,000$ reconstructed $\phi \rightarrow \mu\mu$ decays (see next section) – respectively for the forward and backward rapidity regions. For the pp data sample, the integrated luminosity amounts to 78 ± 3 nb⁻¹ for a total number of $\sim 1\,400$ reconstructed $\phi \rightarrow \mu\mu$ decays.

Track reconstruction in the muon spectrometer is based on a Kalman filter algorithm [25,33,34]. Muon identification is performed by requiring the candidate track to match a track segment in the trigger chambers (trigger tracklet). This request selects muons with $p_{T,\mu} \gtrsim 0.5$ GeV/c and, as a consequence, significantly affects the collected statistics for dimuons with invariant mass $\lesssim 1$ GeV/c² and $p_T \lesssim 1$ GeV/c. It is also required that muon tracks lie in the pseudorapidity interval $-4 < \eta_\mu < -2.5$, where η_μ is defined in the laboratory frame, in order to remove the tracks close to the acceptance borders of the spectrometer, where the acceptance drops abruptly. Selected tracks are finally required to exit the hadron absorber at a radial distance from the beam axis, R_{abs} , in the range $17.6 < R_{\text{abs}} < 89.5$ cm: this cut, for all practical purposes equivalent to the one on η_μ , explicitly ensures the rejection

¹ The sign of y is defined by assuming the proton beam to have positive rapidity.

² In this case the sign of y is defined by assuming the proton beam entering the muon spectrometer to have positive rapidity.




Fig. 1. Dimuon mass spectrum after combinatorial background subtraction: p_T -integrated pp sample (top panels) and p_T -integrated p-Pb sample in the backward (centre panels) and forward (bottom panels) rapidity regions, compared to the result of the hadronic-cocktail and the empirical-function fits (left- and right-column panels, respectively). Error boxes on data points (well visible only in some regions on the plots) represent the systematic uncertainty due to the combinatorial background subtraction, while error bars account for the statistical uncertainty. The width of the hadronic-cocktail fit result (red band) combines the statistical uncertainties of the free fit parameters with the systematic uncertainties on the fixed parameters (see text). (For interpretation of the references to colour in this figure, the reader is referred to the web version of this article.)

of tracks crossing the region of the absorber with the highest density material, where multiple scattering and energy loss effects are large and can affect the mass resolution. Muon pairs are built combining two muon tracks that satisfy the above cuts.

3.2. Signal extraction

The Opposite-Sign (OS) muon pairs are composed of correlated and uncorrelated pairs. The former contain the signal of interest for the present analysis, while the latter – mainly coming from semi-muonic decays of pions and kaons – form a combinatorial background. The contribution of the combinatorial background to the OS mass spectrum was evaluated using an event mixing tech-

nique in which uncorrelated pairs are formed with muons taken from different events. A detailed description of the technique can be found in [25]. The ratio between correlated and uncorrelated OS dimuons at the ϕ -meson mass is ~ 0.65 (~ 0.40) in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV at forward (backward) rapidity, and ~ 1.30 in pp collisions at $\sqrt{s} = 2.76$ TeV. A direct comparison of the raw OS mass spectrum and the associated combinatorial background is presented in [35], for each of the p_T intervals considered in the analysis.

The invariant mass spectra in pp and p-Pb collisions, obtained after combinatorial background subtraction, are shown in Fig. 1 for the p_T -integrated samples. In the left-column panels of Fig. 1, the signal is described in the low-mass region (from the threshold

up to $\sim 1.5 \text{ GeV}/c^2$) by the superposition of a so-called hadronic cocktail and the open charm and open beauty processes. The processes included in the hadronic cocktail are the two-body and Dalitz decays of the light neutral mesons η , ρ , ω , η' and ϕ , which dominate dimuon production for invariant masses below $\sim 1 \text{ GeV}/c^2$. The open charm and open beauty contributions arise from correlated semi-muonic decays of charm and beauty mesons and baryons.

The hadronic cocktail was simulated with a dedicated generator described in [25], tuned to the existing measurements whenever possible, otherwise based on the kinematic distributions extracted from PYTHIA [36]. In particular, the kinematic distributions of the ϕ -meson have been tuned by means of an iterative procedure to the results presented in this Letter to ensure self-consistency for this analysis. The open charm and beauty generation is based on a parameterisation of the spectra generated with PYTHIA [33]. The detector response for all these processes is obtained with a simulation based on the GEANT3 [37] transport code. Simulated events are then subjected to the same reconstruction and selection procedure as the data.

When describing the signal with the superposition of the aforementioned contributions, four parameters are adjusted in the fit procedure in each of the p_T or rapidity intervals considered in the analysis: the yield of the η , ω and ϕ -mesons, and the one of the open charm and beauty processes, with the relative beauty/charm contribution fixed (see later in this paragraph). In this way, each parameter is linked to a process dominating in at least one region of the considered mass spectrum. The remaining degrees of freedom are fixed either according to the relative branching ratios known from literature [38], or assuming specific hypotheses on the cross section ratios. In particular, the production cross section of the ρ -meson is assumed to be the same as for the ω as suggested from both models and pp data [25], while the η' contribution was derived from the η cross section by applying the ratio of the corresponding cross sections $\sigma_{\eta'}/\sigma_\eta = 0.3$ taken from the PYTHIA tunes ATLAS-CSC and D6T which best describe the available low-mass dimuon measurements at the LHC energies [25]. The open beauty normalisation is fixed to the open charm one via a fit of the p_T - and rapidity-integrated mass spectra in which the yields from both processes are free parameters; when performing differential studies, the beauty/charm ratio is scaled according to the differential distributions for the two processes, given by the Monte Carlo (MC) simulations.

For each p_T and rapidity interval, the raw number of ϕ -mesons is determined via a fit procedure based on a χ^2 minimisation, performed on the signal obtained after the subtraction of the combinatorial background, shown in Fig. 1 for the p_T -integrated samples. Several tests have been performed to evaluate the robustness of the signal extraction and estimate an appropriate systematic uncertainty for it. They include in particular:

- Replacing the fit based on the full MC hadronic cocktail with a fit based on the superposition of various empirical functions. In this case, illustrated in the right-column panels of Fig. 1, the continuum is modelled either with exponential functions or variable-width Gaussians, while the $\rho+\omega$ and ϕ peaks are described by Crystal Ball functions [39] tuned on the MC.
- Varying the ratio between the yields of open beauty and open charm processes. It was verified that for perturbations as large as $\pm 50\%$ (resulting in a reasonably wide range of variation for the shape of the total continuum) no significant systematic effect is visible.
- Varying the ratios between the two-body and Dalitz branching ratios of the η and ω -mesons, as well as the cross section ratios $\sigma_\rho/\sigma_\omega$ and $\sigma_{\eta'}/\sigma_\eta$, within the uncertainties coming either

from the available measurements or from the differences between the PYTHIA tunes considered in the analysis of the pp data. The branching ratio $BR_{\omega \rightarrow \mu\mu}$ was taken as the average (weighted by the corresponding uncertainties) of the available measurements of $BR_{\omega \rightarrow \mu\mu}$ and $BR_{\omega \rightarrow ee}$ [38], assuming lepton universality.

- Varying the considered fit range: in particular, the fit was performed both including and excluding the mass region from 0.4 to $0.65 \text{ GeV}/c^2$ where the quality of the comparison between the data and the sum of the MC sources turns out to be lower.

The total systematic uncertainty on the signal extraction was taken as the quadratic sum of the above sources. The systematic uncertainty on the combinatorial background is estimated by comparing the shape of the Like-Sign dimuon contributions coming from the event mixing procedure and from the raw data [25]. This uncertainty depends on the mass, its relative contribution being maximal in the mass window $0.5\text{--}0.8 \text{ GeV}/c^2$ and minimal around the ϕ -meson peak, and it is added in quadrature, for each point of the mass spectrum, to the statistical uncertainty of the signal: in this way, this source of systematics is accounted for by the χ^2 minimisation procedure, and automatically propagated when evaluating the ϕ -meson raw signal from the fit parameters. The uncertainty associated to the sum of the MC sources (red band in the left-column plots of Fig. 1) is evaluated by combining the uncertainties on the normalisation of each considered process. For the processes whose normalisation is left free in the fit, this uncertainty is the statistical one resulting from the fit procedure itself; for the rest of the processes, we also propagate the systematic uncertainty on the parameters (branching ratios or cross section ratios) which fix their normalisations to those of the free processes.

4. Results

The results of the ϕ -meson analysis are presented as follows. We first present the measurement of the production cross sections, starting with its p_T -dependence in pp collisions at $\sqrt{s} = 2.76 \text{ TeV}$, followed by p-Pb collision results as a function of p_T and rapidity. Then, we show the ratio of the cross sections measured in the forward and backward regions, obtained in the common rapidity interval $2.96 < |y| < 3.53$. Finally, the measurement of the nuclear modification factor R_{pPb} as a function of p_T is presented, separately for the p-going and the Pb-going directions.

4.1. Production cross section in pp and p–Pb collisions

The cross section σ_ϕ was evaluated for each p_T and rapidity interval as:

$$\sigma_\phi(x) = \frac{N_{\phi \rightarrow \mu\mu}^{\text{raw}}(x)}{[A \cdot \varepsilon](x) \cdot BR_{\phi \rightarrow \mu\mu} \cdot L_{\text{int}}} ,$$

where x stands for any specific p_T or rapidity interval considered. The total systematic uncertainty on $N_{\phi \rightarrow \mu\mu}^{\text{raw}}(x)$, after combining the different sources described above, ranges between 3% and 8% depending on the collision system and kinematic range. The branching ratio $BR_{\phi \rightarrow \mu\mu}$ was taken from [38] as the average (weighted by the corresponding uncertainties) of the available measurements of $BR_{\phi \rightarrow \mu\mu}$ and $BR_{\phi \rightarrow ee}$, assuming lepton universality, resulting in a final uncertainty of approximately 1%. The product of the geometrical acceptance A and the reconstruction efficiency ε has been evaluated by means of MC simulations, using the cocktail predictions for the differential input spectra. The values are obtained as the ratio between the number of dimuons at the output of the reconstruction chain – including the effect of the event selection




Fig. 2. ϕ -meson cross section as a function of p_T in pp collisions at $\sqrt{s} = 2.76$ TeV. Error bars and boxes represent statistical and systematic uncertainties, respectively. Predictions from PHOJET [42] and the PYTHIA tunes ATLAS-CSC [44], D6T [45], Perugia0 and Perugia11 [43] are also shown for comparison, as well as the result of a fit with the Levy-Tsallis function defined by Eq. (1).

criteria imposed on the data – and the number of dimuons injected as input.

The uncertainty on $[A \cdot \varepsilon]$ mainly originates from the systematic uncertainty on the dimuon tracking and trigger efficiencies. The systematic uncertainty on the tracking efficiency, amounting to 6% and 4% for the backward and forward rapidity regions, respectively, comes from the residual differences between the results of the efficiency-determination method based on reconstructed tracks [29,40], applied to both data and MC. For the systematic uncertainty on the trigger efficiency, we also refer to the procedure discussed in [29], resulting in an uncertainty of 3.2% and 2.8%, respectively, for the backward and forward rapidity regions considered in the analysis. In order to test possible additional systematic effects related to the hardware trigger p_T cut, imposing a non-sharp threshold around 0.5 GeV/c, the analysis was repeated with the additional offline sharp cuts $p_{T,\mu} > 0.5$ GeV/c and $p_{T,\mu} > 1$ GeV/c on single muons. For each of the two alternative scenarios, the corresponding measurement of the ϕ -meson cross section was compared to the one coming from the reference analysis: the difference between the results was found to be smaller than the quadratic difference of the statistical uncertainties, showing that no significant bias related to the trigger threshold affects the results [41].

The reported values correspond to a zero-polarisation scenario for the 2-body decay of the ϕ -meson, in the absence of evidence supporting less trivial assumptions (in particular, no measurement of ϕ -meson polarisation is currently available at the LHC energies).

4.1.1. Production cross section in pp collisions

The inclusive, p_T -differential ϕ -meson cross section in pp collisions at $\sqrt{s} = 2.76$ TeV is shown in Fig. 2. The data points, also summarised in Table 1, are compared with the predictions from PHOJET [42] and PYTHIA [36], where for the latter the Perugia0, Perugia11 [43], ATLAS-CSC [44], and D6T [45] tunes are considered. An overall good agreement is found between predictions and data, with the exception of the Perugia0 and Perugia11 tunes of PYTHIA which underestimate the measured cross section by a factor of two, as already observed for the ϕ -meson measurements at $\sqrt{s} = 7$ TeV [25,46]. It is worth to note that the D6T tune is not successful in describing the p_T evolution of the K/π ratio at mid-rapidity in pp collisions at $\sqrt{s} = 2.76$ TeV, as measured by the CMS Collaboration [47]: this suggests that hidden strangeness is better reproduced than open strangeness in this

Table 1

p_T -differential production cross section for the ϕ -meson in pp collisions at $\sqrt{s} = 2.76$ TeV, for $2.5 < y < 4$. The first uncertainty is statistical and the second is the bin-to-bin uncorrelated systematic. The bin-to-bin correlated relative systematic uncertainty is 3.9%. The χ^2/ndf values are relative to the hadronic-cocktail fit and the $[0.8, 1.2 \text{ GeV}/c^2]$ mass region, where $\text{ndf} = 10$.

p_T (GeV/c)	χ^2/ndf	$d^2\sigma_\phi/(dydp_T)$ (mb/(GeV/c))
[1.0, 1.5]	1.1	$0.423 \pm 0.067 \pm 0.043$
[1.5, 2.0]	1.7	$0.182 \pm 0.025 \pm 0.018$
[2.0, 2.5]	1.1	$0.089 \pm 0.011 \pm 0.007$
[2.5, 3.0]	1.1	$0.0340 \pm 0.0056 \pm 0.0020$
[3.0, 3.5]	0.9	$0.0139 \pm 0.0032 \pm 0.0011$
[3.5, 4.0]	1.1	$0.0087 \pm 0.0022 \pm 0.0006$
[4.0, 5.0]	1.1	$0.0028 \pm 0.0012 \pm 0.0002$

Table 2

Systematic uncertainties (in percent) contributing to the measurement of the ϕ cross section in pp collisions at $\sqrt{s} = 2.76$ TeV. When the uncertainty values depend on the p_T interval, their minimum and maximum values are quoted.

Source	Syst. uncertainty on σ_ϕ^{pp}
<i>Uncorrelated</i>	
Signal extraction	3–8%
Tracking efficiency	4%
Trigger efficiency	3%
<i>Correlated</i>	
L_{int}	3.8%
$BR(\phi \rightarrow \ell\ell)$	1%

specific PYTHIA tune. Data points were fitted with a Levy-Tsallis function [48]

$$\frac{1}{p_T} \frac{dN}{dp_T} \propto \left(1 + \frac{m_T - m_\phi}{nT}\right)^{-n}, \quad (1)$$

where $m_T = \sqrt{p_T^2 + m_\phi^2}$ stands for the transverse mass, obtaining the values $n = 10.2 \pm 4.8$ and $T = 284 \pm 72$ MeV for the fit parameters, where the errors reflect the statistical uncertainties only. The cross section integrated over the accessible p_T range $1 < p_T < 5$ GeV/c is $\sigma_\phi = 0.566 \pm 0.055$ (stat.) ± 0.044 (syst.) mb. The systematic uncertainties for this measurement are summarised in Table 2.

4.1.2. Production cross section in p-Pb collisions

The ϕ cross section as a function of p_T in p-Pb collisions is shown in Fig. 3 for the forward and backward rapidity regions considered in the analysis. The results, also reported in Table 3, are fitted with the Levy-Tsallis distribution defined in Eq. (1), the resulting fit parameters being $\beta = 9.6 \pm 1.3$ and $T = 366 \pm 30$ MeV for the forward rapidity region and $\beta = 11.4 \pm 1.4$ and $T = 384 \pm 24$ MeV for the backward one, where the errors reflect the statistical uncertainties only. The predictions from HIJING (with gluon shadowing) [49] and DPMJET [50] are also shown: these generators provided a good description of the ALICE $dN_{\text{ch}}/d\eta_{\text{lab}}$ results at mid-rapidity [9]. Averaging over the available p_T range, the discrepancy between the data and the predictions from HIJING and DPMJET amounts to $\sim 18\%$ and $\sim 57\%$, respectively, at backward rapidity (the Pb-going direction) and $\sim 5\%$ and $\sim 9.5\%$, respectively, at forward rapidity (the p-going direction). In all the cases, the generators underestimate the data points.

The ϕ cross section in p-Pb collisions, integrated over the accessible p_T range, $1 < p_T < 7$ GeV/c, is shown as a function of rapidity in Fig. 4. The data points, also summarised in Table 4, exhibit a significant asymmetry between the forward and backward rapidity regions. The data point from the ϕ -meson analysis at mid-rapidity in the K^+K^- channel [51], also shown for the




Fig. 3. ϕ -meson cross section in p-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV as a function of p_{T} in the backward (left) and forward (right) rapidity regions. Error bars (smaller than the markers) and boxes represent statistical and systematic uncertainties, respectively. Predictions by HIJING [49] and DPMJET [50] are also shown, together with the result of a fit with the Levy-Tsallis function (Eq. (1)).

Table 3
Production cross section for the ϕ -meson in p-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV, as a function of p_{T} , in the backward and forward rapidity regions. The first uncertainty is statistical and the second is the bin-to-bin uncorrelated systematic. The bin-to-bin correlated relative systematic uncertainty is 3.6% and 3.9%, respectively, for the backward and the forward regions. The χ^2/ndf values are relative to the hadronic-cocktail fit and the $[0.8, 1.2 \text{ GeV}/c^2]$ mass region.

p_{T} (GeV/c)	$-4.46 < y < -2.96$		$2.03 < y < 3.53$	
	χ^2/ndf	$d^2\sigma_{\phi}^{\text{ppb}}/(dydp_{\text{T}})$ (mb/(GeV/c))	χ^2/ndf	$d^2\sigma_{\phi}^{\text{ppb}}/(dydp_{\text{T}})$ (mb/(GeV/c))
[1.0, 1.5]	0.7	$102 \pm 8 \pm 12$	1.5	$73.3 \pm 5.6 \pm 8.0$
[1.5, 2.0]	1.2	$58.6 \pm 3.3 \pm 5.5$	1.9	$42.1 \pm 2.5 \pm 4.3$
[2.0, 2.5]	2.5	$28.3 \pm 1.4 \pm 2.9$	1.7	$21.0 \pm 1.2 \pm 2.0$
[2.5, 3.0]	4.2	$15.0 \pm 0.7 \pm 1.2$	3.1	$10.07 \pm 0.77 \pm 0.97$
[3.0, 3.5]	2.6	$7.66 \pm 0.40 \pm 0.70$	2.0	$6.38 \pm 0.41 \pm 0.61$
[3.5, 4.0]	1.9	$4.20 \pm 0.24 \pm 0.34$	1.2	$3.96 \pm 0.30 \pm 0.36$
[4.0, 4.5]	0.7	$2.15 \pm 0.17 \pm 0.16$	1.0	$1.99 \pm 0.20 \pm 0.15$
[4.5, 5.0]	0.9	$1.20 \pm 0.11 \pm 0.10$	0.9	$1.06 \pm 0.13 \pm 0.08$
[5.0, 6.0]	1.0	$0.560 \pm 0.052 \pm 0.054$	1.0	$0.570 \pm 0.088 \pm 0.043$
[6.0, 7.0]	1.2	$0.201 \pm 0.030 \pm 0.028$	0.9	$0.199 \pm 0.045 \pm 0.016$




Fig. 4. ϕ cross section in p-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV as a function of rapidity, integrated over the range $1 < p_{\text{T}} < 7 \text{ GeV}/c$. Error bars and boxes represent statistical and systematic uncertainties, respectively. Predictions by HIJING and DPMJET are also shown, together with the mid-rapidity data point from the ϕ -meson measurement in the K^+K^- channel [51], also evaluated in the range $1 < p_{\text{T}} < 7 \text{ GeV}/c$.

$1 < p_{\text{T}} < 7 \text{ GeV}/c$ p_{T} range, fits well into the trend defined by the two series of points in the backward and forward rapidity regions. This observation complements the previous measurements of light-flavour particle production (charged unidentified particles) reported in p-Pb by ALICE at the LHC at mid-rapidity [9], and in d-Au by PHOBOS at RHIC ranging from mid to forward rapidity [14]. The comparison between the data and the predictions by

Table 4
Production cross section for the ϕ -meson in p-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV, as a function of rapidity, integrated over the range $1 < p_{\text{T}} < 7 \text{ GeV}/c$. The first uncertainty is statistical and the second is the bin-to-bin uncorrelated systematic. The bin-to-bin correlated relative systematic uncertainty is 3.6% and 3.9%, respectively, for the backward and the forward regions. The χ^2/ndf values are relative to the hadronic-cocktail fit and the $[0.8, 1.2 \text{ GeV}/c^2]$ mass region.

y	χ^2/ndf	$d\sigma_{\phi}^{\text{ppb}}/dy$ (mb)	y	χ^2/ndf	$d\sigma_{\phi}^{\text{ppb}}/dy$ (mb)
[-4.46, -4.25]	0.9	$89 \pm 10 \pm 9$	[2.03, 2.35]	2.6	$104 \pm 11 \pm 6$
[-4.25, -4.05]	1.8	$89 \pm 6 \pm 7$	[2.35, 2.55]	1.5	$102 \pm 7 \pm 5$
[-4.05, -3.85]	0.9	$103 \pm 5 \pm 8$	[2.55, 2.75]	2.0	$96 \pm 5 \pm 6$
[-3.85, -3.65]	2.9	$117 \pm 6 \pm 9$	[2.75, 2.95]	1.6	$86 \pm 4 \pm 5$
[-3.65, -3.45]	1.2	$128 \pm 7 \pm 9$	[2.95, 3.15]	2.3	$68 \pm 4 \pm 4$
[-3.45, -3.25]	3.6	$133 \pm 9 \pm 9$	[3.15, 3.35]	1.0	$66 \pm 5 \pm 5$
[-3.25, -2.96]	1.2	$136 \pm 14 \pm 11$	[3.35, 3.53]	1.2	$45 \pm 8 \pm 6$

HIJING and DPMJET, illustrated in Fig. 4, clearly shows how the models – which successfully described charged particle production at mid-rapidity in the same collision system [9] – fail to properly reproduce the shape and the normalisation of the observed rapidity dependence of the ϕ cross section. Still, the HIJING prediction qualitatively reproduces the forward-backward asymmetry observed in the data, as well as – ignoring the normalisation – the shape of the y -dependence in the backward region. DPMJET, on the contrary, fails to reproduce even qualitatively the observed forward-backward asymmetry.

4.2. Forward-backward ratio in p-Pb collisions

To establish a more direct comparison of the cross section in the p-going and Pb-going directions, $\sigma_{\phi}^{\text{ppb}}$ was extracted as a func-




Fig. 5. Forward-backward ratio for the ϕ -meson in p-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02 \text{ TeV}$ as a function of p_T , in the rapidity range $2.96 < |y| < 3.53$ common to the two rapidity regions considered in the analysis. Error bars and boxes represent statistical and systematic uncertainties, respectively. The blue box on the left represents the bin-to-bin correlated systematic uncertainty, see Table 7. Predictions from HIJING and DPMJET are also shown for comparison. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

tion of p_T in the common $|y|$ range $2.96 < |y| < 3.53$. The p_T interval $1.0 < p_T < 1.5 \text{ GeV}/c$ was discarded in this measurement because of the poor statistics available in this limited rapidity range, resulting in an uncertainty larger than 50%.

The ratio between the forward and backward cross section, R_{FB} , is shown as a function of p_T in Fig. 5. The data points exhibit no significant p_T dependence within the experimental uncertainties. Predictions by HIJING and DPMJET are also shown, with HIJING slightly overestimating the data points and DPMJET clearly failing to reproduce the observed values, staying above $R_{\text{FB}} = 1$ in the whole p_T range considered here. This observation is consistent with the observations in Fig. 4, where the forward-backward asymmetry of the ϕ -meson yield was better reproduced by HIJING than by DPMJET.

4.3. Nuclear modification factor in p-Pb collisions

The ϕ -meson nuclear modification factor R_{pPb} is defined as the ratio between the production cross section $\sigma_{\phi}^{\text{pPb}}(p_T)$ in p-Pb collisions and the cross section $\sigma_{\phi}^{\text{pp}}(p_T)$ in pp collisions – evaluated at $\sqrt{s} = 5.02 \text{ TeV}$ as described in the following – scaled by A_{Pb} :

$$R_{\text{pPb}}(p_T) = \frac{\sigma_{\phi}^{\text{pPb}}(p_T)}{\sigma_{\phi}^{\text{pp}}(p_T) \cdot A_{\text{Pb}}}, \quad (2)$$

where A_{Pb} is the nuclear mass number for the Pb nucleus. Since for the pp cross section $\sigma_{\phi}^{\text{pp}}$ at $\sqrt{s} = 5.02 \text{ TeV}$ no direct measurement is currently available, it was evaluated by interpolating the measurements in the rapidity interval $2.5 < y < 4$ at $\sqrt{s} = 2.76$ (see Section 4.1.1) and 7 TeV [25]. For each p_T interval, the \sqrt{s} dependence of the differential cross section $d^2\sigma_{\phi}^{\text{pp}}/(dy dp_T)$ was described with a power law $\sigma^{\text{pp}}(\sqrt{s}) = C \cdot (\sqrt{s})^\alpha$, where C and α are determined using the data at 2.76 and 7 TeV. Alternative parameterisations were also considered [52], namely a linear and an exponential function, and the mean of the results obtained with the three functions was taken. Since the pp measurements are limited to $1 < p_T < 5 \text{ GeV}/c$, the cross section at $\sqrt{s_{\text{NN}}} = 5.02 \text{ TeV}$ was extrapolated towards higher p_T by means of a Levy-Tsallis function, which describes the calculated differential cross section in the p_T range covered by the measurements. The uncertainty

Table 5

Differential cross section for the ϕ -meson in pp collisions at $\sqrt{s} = 5.02 \text{ TeV}$ in the backward and forward rapidity regions of interest for the analysis of the p-Pb data, as obtained interpolating the available measurements at $\sqrt{s} = 2.76$ and 7 TeV. Total uncertainties, combining statistical and systematic sources, are reported.

p_T (GeV/c)	$d^2\sigma_{\phi}/dy dp_T$ (mb/(GeV/c))	$-4.46 < y < -2.96$	$2.03 < y < 3.53$
	$-4.46 < y < -2.96$		
[1.0, 1.5]	0.491 ± 0.067	0.656 ± 0.090	
[1.5, 2.0]	0.223 ± 0.015	0.297 ± 0.020	
[2.0, 2.5]	0.0995 ± 0.0071	0.1328 ± 0.0095	
[2.5, 3.0]	0.0467 ± 0.0032	0.0623 ± 0.0043	
[3.0, 3.5]	0.0234 ± 0.0015	0.0312 ± 0.0020	
[3.5, 4.0]	0.0125 ± 0.0011	0.0167 ± 0.0015	
[4.0, 4.5]	0.00706 ± 0.00094	0.0094 ± 0.0012	
[4.5, 5.0]	0.00419 ± 0.00082	0.0056 ± 0.0011	
[5.0, 6.0]	0.00213 ± 0.00060	0.00284 ± 0.00081	
[6.0, 7.0]	0.00093 ± 0.00039	0.00124 ± 0.00052	

on the interpolated cross sections arises from the choice of the function used for the interpolation, from the uncertainties in the measurements at 2.76 and 7 TeV, and – for $p_T > 5 \text{ GeV}/c$ – from the extrapolation based on the Levy-Tsallis fit. They range from about 7% for $p_T = 1 \text{ GeV}/c$ to 20% for $p_T = 5 \text{ GeV}/c$, and exceed 30% for $p_T > 5 \text{ GeV}/c$, representing the major source of systematic uncertainty for the measurement of the nuclear modification factor. The interpolated cross section, which refers to the rapidity range $2.5 < y < 4$, was finally scaled to the forward and backward rapidity windows $2.03 < y < 3.53$ and $-4.46 < y < -2.96$, considered for the analysis of the p-Pb data. The relative scaling factors $f_{\text{fwd}} = 1.135 \pm 0.031$ and $f_{\text{bkw}} = 0.850 \pm 0.028$ were evaluated as an average from simulations with PHOJET and the Perugia0, Perugia11, ATLAS-CSC and D6T PYTHIA tunes. In doing so, we also retained the PYTHIA tunes which were observed to fail in describing the pp data (see Section 4.1.1): the reason is that the disagreement between models and data concerns in this case the absolute normalisation more than the shape of the kinematic distributions, which is the only relevant feature in the evaluation of the f_{fwd} and f_{bkw} factors. The uncertainties (amounting to about 3%) correspond to the differences between the considered MC predictions. The numerical values are reported in Table 5.

The nuclear modification factor R_{pPb} as a function of p_T is shown in the two panels of Fig. 6 for the backward and forward rapidity regions considered in the analysis. The numerical values are also quoted in Table 6. For each p_T interval, the systematic uncertainty detailed in Table 7 results from the quadratic sum of the uncertainty on the ϕ cross section in p-Pb and the one of the pp reference. A rising trend of R_{pPb} when going from $p_T = 1 \text{ GeV}/c$ to $p_T \approx 3-4 \text{ GeV}/c$ can be observed both at backward and forward rapidity. The values of R_{pPb} in the two rapidity ranges, however, are significantly different. In particular, at backward rapidity we observe an enhancement of the ϕ cross section with respect to the scaled pp reference peaked around $p_T = 3-4 \text{ GeV}/c$. This enhancement, absent in the forward rapidity region, reaches a factor of up to ~ 1.6 and could be associated either to an initial-state effect (including a possible Cronin-like enhancement [4,53]) or to a final state effect related to radial flow in p-Pb as proposed for recent ALICE measurements at mid-rapidity [12]. Discriminating between these two effects requires more detailed investigations, including differential analyses as a function of global event properties like collision centrality.

Concerning the behaviour at high p_T , we observe that the ϕ -meson R_{pPb} is compatible with unity for $p_T \gtrsim 4 \text{ GeV}/c$ in the p-going direction, similar to what was observed for the R_{pPb} of charged particle production at mid-rapidity [10,12]. The observations in the Pb-going direction do not allow a clear trend of the R_{pPb} factor at high p_T to be established. A possible saturation at




Fig. 6. Nuclear modification factor $R_{p\text{Pb}}$ in p–Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV for the ϕ -meson as a function of p_{T} , in the backward (left) and forward (right) rapidity regions considered in the analysis. Error bars and boxes represent statistical and bin-to-bin uncorrelated systematic uncertainties, respectively. The blue box on the left represents the bin-to-bin correlated systematic uncertainty, see Table 7. (For interpretation of colour in this figure legend, the reader is referred to the web version of this article.)

Table 6

Nuclear modification factor $R_{p\text{Pb}}$ in p–Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV for the ϕ -meson as a function of p_{T} in the backward and forward rapidity regions. The first uncertainty is statistical and the second is the bin-to-bin uncorrelated systematic. The bin-to-bin correlated relative systematic uncertainty is 8%.

p_{T} (GeV/c)	$R_{p\text{Pb}}$	
	$-4.46 < y < -2.96$	$2.03 < y < 3.53$
[1.0, 1.5]	$1.00 \pm 0.08 \pm 0.18$	$0.537 \pm 0.041 \pm 0.094$
[1.5, 2.0]	$1.26 \pm 0.07 \pm 0.15$	$0.681 \pm 0.040 \pm 0.083$
[2.0, 2.5]	$1.37 \pm 0.07 \pm 0.17$	$0.760 \pm 0.043 \pm 0.091$
[2.5, 3.0]	$1.54 \pm 0.07 \pm 0.16$	$0.777 \pm 0.059 \pm 0.092$
[3.0, 3.5]	$1.57 \pm 0.08 \pm 0.18$	$0.98 \pm 0.06 \pm 0.11$
[3.5, 4.0]	$1.62 \pm 0.09 \pm 0.19$	$1.14 \pm 0.09 \pm 0.15$
[4.0, 4.5]	$1.46 \pm 0.12 \pm 0.22$	$1.02 \pm 0.10 \pm 0.15$
[4.5, 5.0]	$1.38 \pm 0.13 \pm 0.29$	$0.91 \pm 0.11 \pm 0.19$
[5.0, 6.0]	$1.26 \pm 0.12 \pm 0.38$	$0.97 \pm 0.15 \pm 0.29$
[6.0, 7.0]	$1.04 \pm 0.16 \pm 0.46$	$0.77 \pm 0.17 \pm 0.33$

Table 7

Systematic uncertainties (in percent) contributing to the measurement of the ϕ cross section and nuclear modification factor in the backward and forward rapidity regions in p–Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV. When the uncertainty values depend on the p_{T} interval, their minimum and maximum values are quoted.

Source	Syst. uncertainty on $\sigma_{\phi}^{\text{pp}}$ and $R_{p\text{Pb}}$	
	$-4.46 < y < -2.96$	$2.03 < y < 3.53$
<i>Uncorrelated</i>		
Signal extraction	3–5%	4–8%
Tracking efficiency	6%	4%
Trigger efficiency	3.2%	2.8%
$\sigma_{\phi}^{\text{pp}}$	7–30%	7–30%
<i>Correlated</i>		
f_{int}	3.5%	3.8%
$BR(\phi \rightarrow \ell\ell)$	1%	1%
f_{bkw}	3.3%	–
f_{fwd}	–	2.7%

$R_{p\text{Pb}} \approx 1$ for $p_{\text{T}} \gtrsim 5$ GeV/c is, however, still compatible with the measurements.

Only few other existing measurements can be compared to our data. In particular, results on ϕ -meson production in d–Au collisions at $\sqrt{s_{\text{NN}}} = 200$ GeV have been recently released by the PHENIX Collaboration [54]. The p_{T} -dependence of the $R_{d\text{Au}}$ measured by PHENIX, as well as its evolution from backward to forward rapidity, is found to be similar to what is observed in our results for $R_{p\text{Pb}}$. Mid-rapidity data on $R_{d\text{Au}}$, also presented by the PHENIX Collaboration for the ϕ -meson, seem to sit between the

forward- and backward-rapidity results. Forward-rapidity measurements in d–Au collisions at $\sqrt{s_{\text{NN}}} = 200$ GeV at RHIC [13,14] are also available for unidentified charged particles, although for the d-going direction only. These data exhibit, similar to our ϕ -meson results in the p-going direction, a rise of $R_{d\text{Au}}$ from ~ 0.5 to ~ 1 between $p_{\text{T}} \sim 1$ GeV/c and $p_{\text{T}} \sim 4$ GeV/c. A similar rise of $R_{p\text{Pb}}$ in p–Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV is also observed in the already cited measurement of unidentified charged particle and identified charged pion and kaon production at mid-rapidity performed by ALICE [10,12]. A recent study of ϕ -meson production in p–Pb collisions at mid-rapidity by ALICE [51] does not currently include results on $R_{p\text{Pb}}$.

5. Conclusions

We have presented results on ϕ -meson production in the dimuon channel in p–Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV obtained by the ALICE experiment at the LHC. Cross section and nuclear modification factor measurements were performed for $1 < p_{\text{T}} < 7$ GeV/c in the rapidity windows $2.03 < y < 3.53$ (p-going direction) and $-4.46 < y < -2.96$ (Pb-going direction). A corresponding cross section measurement in pp collisions at $\sqrt{s} = 2.76$ TeV has also been reported, for $1 < p_{\text{T}} < 5$ GeV/c in the region $2.5 < y < 4$. Predictions from HIJING and DPMJET are compared to the p–Pb cross sections and are found to underestimate the data both at backward (by about 18% and 57% on average, respectively) and at forward rapidity (by about 5% and 9.5% on average, respectively). The forward–backward ratio in the ϕ -meson cross section in p–Pb collisions was measured in the rapidity range $2.96 < |y| < 3.53$, and no significant p_{T} dependence was found within uncertainties. In this case, the data points are significantly overestimated by the DPMJET model, while only a slight disagreement is observed with respect to the HIJING prediction.

In the p-going direction a rising trend of the nuclear modification factor $R_{p\text{Pb}}$ is observed from ~ 0.5 to ~ 1 , when going from $p_{\text{T}} = 1$ GeV/c to $p_{\text{T}} = 4$ GeV/c. This observation is compatible with the behaviour of charged particles at forward rapidity at RHIC energies, and at mid-rapidity at LHC energies. In the Pb-going direction, on the other hand, an enhancement is observed for $R_{p\text{Pb}}$, reaching values as large as ~ 1.6 around $p_{\text{T}} = 3$ –4 GeV/c. An interpretation of these results, either in terms of an initial-state (Cronin-like) effect or a final-state effect related to radial flow in p–Pb, is not possible yet, due to a general lack of theoretical predictions for particle production in the light-flavour sector at forward rapidity in p–A collisions at the LHC energies.

Acknowledgements

The ALICE Collaboration would like to thank all its engineers and technicians for their invaluable contributions to the construction of the experiment and the CERN accelerator teams for the outstanding performance of the LHC complex. The ALICE Collaboration gratefully acknowledges the resources and support provided by all Grid centres and the Worldwide LHC Computing Grid (WLCG) Collaboration. The ALICE Collaboration acknowledges the following funding agencies for their support in building and running the ALICE detector: A. I. Alikhanyan National Science Laboratory (Yerevan Physics Institute) Foundation (ANSL), State Committee of Science and World Federation of Scientists (WFS), Armenia; Austrian Academy of Sciences and Nationalstiftung für Forschung, Technologie und Entwicklung, Austria; Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Universidade Federal do Rio Grande do Sul (UFRGS), Financiadora de Estudos e Projetos (Finep) and Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP), Brazil; Ministry of Science and Technology of the People's Republic of China (MOST), National Natural Science Foundation of China (NSFC) and Ministry of Education of China (MOE), China; Ministry of Science, Education and Sports and Croatian Science Foundation, Croatia; Ministry of Education, Youth and Sports of the Czech Republic, Czech Republic; The Danish Council for Independent Research – Natural Sciences, the Carlsberg Foundation and Danish National Research Foundation (DNRF), Denmark; Helsinki Institute of Physics (HIP), Finland; Commissariat à l'Energie Atomique (CEA) and Institut National de Physique Nucléaire et de Physique des Particules (IN2P3) and Centre National de la Recherche Scientifique (CNRS), France; Bundesministerium für Bildung, Wissenschaft, Forschung und Technologie (BMBF) and GSI Helmholtzzentrum für Schwerionenforschung GmbH, Germany; Ministry of Education, Research and Religious Affairs, Greece; National Research, Development and Innovation Office, Hungary; Department of Atomic Energy, Government of India (DAE) and Council of Scientific and Industrial Research (CSIR), New Delhi, India; Indonesian Institute of Science, Indonesia; Centro Fermi – Museo Storico della Fisica e Centro Studi e Ricerche Enrico Fermi and Instituto Nazionale di Fisica Nucleare (INFN), Italy; Institute for Innovative Science and Technology, Nagasaki Institute of Applied Science (IIIST), Japan Society for the Promotion of Science (JSPS) KAKENHI and Japanese Ministry of Education, Culture, Sports, Science, and Technology (MEXT), Japan; Consejo Nacional de Ciencia (CONACYT) y Tecnología, through Fondo de Cooperación Internacional en Ciencia y Tecnología (FONCICYT) and Dirección General de Asuntos del Personal Académico (DGAPA), Mexico; Nationaal instituut voor subatomaire fysica (Nikhef), Netherlands; The Research Council of Norway, Norway; Commission on Science and Technology for Sustainable Development in the South (COMSATS), Pakistan; Pontificia Universidad Católica del Perú, Peru; Ministry of Science and Higher Education and National Science Centre, Poland; Korea Institute of Science and Technology Information and National Research Foundation of Korea (NRF), Republic of Korea; Ministry of Education and Scientific Research, Institute of Atomic Physics and Romanian National Agency for Science, Technology and Innovation, Romania; Joint Institute for Nuclear Research (JINR), Ministry of Education and Science of the Russian Federation and National Research Centre Kurchatov Institute, Russia; Ministry of Education, Science, Research and Sport of the Slovak Republic, Slovakia; National Research Foundation of South Africa, South Africa; Centro de Aplicaciones Tecnológicas y Desarrollo Nuclear (CEADEN), Cubaenergía, Cuba, Ministerio de Ciencia e Innovación and Centro de Investigaciones Energéticas, Medioambientales y Tecnológicas (CIEMAT), Spain; Swedish Research Council (VR) and Knut and Alice Wal-

lenberg Foundation (KAW), Sweden; European Organization for Nuclear Research, Switzerland; National Science and Technology Development Agency (NSDTA), Suranaree University of Technology (SUT) and Office of the Higher Education Commission under NRU project of Thailand, Thailand; Turkish Atomic Energy Agency (TAEK), Turkey; National Academy of Sciences of Ukraine, Ukraine; Science and Technology Facilities Council (STFC), United Kingdom; National Science Foundation of the United States of America (NSF) and United States Department of Energy, Office of Nuclear Physics (DOE NP), United States of America.

References

- [1] A. Bazavov, T. Bhattacharya, M. Cheng, C. DeTar, H. Ding, et al., The chiral and deconfinement aspects of the QCD transition, *Phys. Rev. D* 85 (2012) 054503, arXiv:1111.1710 [hep-lat].
- [2] Wuppertal-Budapest Collaboration, S. Borsanyi, et al., Is there still any T_c mystery in lattice QCD? Results with physical masses in the continuum limit III, *J. High Energy Phys.* 1009 (2010) 073, arXiv:1005.3508 [hep-lat].
- [3] S. Borsanyi, G. Endrodi, Z. Fodor, A. Jakovac, S.D. Katz, et al., The QCD equation of state with dynamical quarks, *J. High Energy Phys.* 1011 (2010) 077, arXiv: 1007.2580 [hep-lat].
- [4] A. Accardi, Cronin effect in proton nucleus collisions: a survey of theoretical models, arXiv:hep-ph/0212148.
- [5] C. Salgado, J. Alvarez-Muniz, F. Arleo, N. Armesto, M. Botje, et al., Proton-nucleus collisions at the LHC: scientific opportunities and requirements, *J. Phys. G* 39 (2012) 015010, arXiv:1105.3919 [hep-ph].
- [6] M. Brandt, M. Klasen, F. König, Nuclear parton density modifications from low-mass lepton pair production at the LHC, *Nucl. Phys. A* 927 (2014) 78–90, arXiv: 1401.6817 [hep-ph].
- [7] BRAHMS Collaboration, I. Arsene, et al., Quark gluon plasma and color glass condensate at RHIC? The perspective from the BRAHMS experiment, *Nucl. Phys. A* 757 (2005) 1–27, arXiv:nucl-ex/0410020.
- [8] B. Back, M. Baker, M. Ballintijn, D. Barton, B. Becker, et al., The PHOBOS perspective on discoveries at RHIC, *Nucl. Phys. A* 757 (2005) 28–101, arXiv:nucl-ex/0410022.
- [9] ALICE Collaboration, B. Abelev, et al., Pseudorapidity density of charged particles p -Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, *Phys. Rev. Lett.* 110 (2013) 032301, arXiv:1210.3615 [nucl-ex].
- [10] ALICE Collaboration, B. Abelev, et al., Transverse momentum distribution and nuclear modification factor of charged particles in p -Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, *Phys. Rev. Lett.* 110 (2013) 082302, arXiv:1210.4520 [nucl-ex].
- [11] ALICE Collaboration, B. Abelev, et al., Transverse momentum dependence of inclusive primary charged-particle production in p -Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, *Eur. Phys. J. C* 74 (9) (2014) 3054, arXiv:1405.2737 [nucl-ex].
- [12] ALICE Collaboration, J. Adam, et al., Multiplicity dependence of charged pion, kaon, and (anti)proton production at large transverse momentum in p -Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, *Phys. Lett. B* 760 (2016) 720–735, arXiv:1601.03658 [nucl-ex].
- [13] BRAHMS Collaboration, I. Arsene, et al., On the evolution of the nuclear modification factors with rapidity and centrality in d +Au collisions at $\sqrt{s_{NN}} = 200$ GeV, *Phys. Rev. Lett.* 93 (2004) 242303, arXiv:nucl-ex/0403005.
- [14] PHOBOS Collaboration, B. Back, et al., Pseudorapidity distribution of charged particles in d +Au collisions at $\sqrt{s_{NN}} = 200$ GeV, *Phys. Rev. Lett.* 93 (2004) 082301, arXiv:nucl-ex/0311009.
- [15] J. Rafelski, B. Müller, Strangeness production in the quark-gluon plasma, *Phys. Rev. Lett.* 48 (1982) 1066.
- [16] A. Shor, ϕ -Meson production as a probe of the quark gluon plasma, *Phys. Rev. Lett.* 54 (1985) 1122–1125.
- [17] P. Koch, B. Müller, J. Rafelski, Strangeness in relativistic heavy ion collisions, *Phys. Rep.* 142 (1986) 167–262.
- [18] NA49 Collaboration, C. Alt, et al., System-size dependence of strangeness production in nucleus-nucleus collisions at $\sqrt{s_{NN}} = 17.3$ GeV, *Phys. Rev. Lett.* 94 (2005) 052301, arXiv:nucl-ex/0406031.
- [19] NA49 Collaboration, C. Alt, et al., Energy dependence of ϕ meson production in central Pb+Pb collisions at $\sqrt{s_{NN}} = 6$ to 17 GeV, *Phys. Rev. C* 78 (2008) 044907, arXiv:0806.1937 [nucl-ex].
- [20] NA50 Collaboration, B. Alessandro, et al., ϕ production in Pb-Pb collisions at 158-GeV/c per nucleon incident momentum, *Phys. Lett. B* 555 (2003) 147–155; NA50 Collaboration, B. Alessandro, et al., *Phys. Lett. B* 561 (2003) 294 (Erratum).
- [21] NA60 Collaboration, K. Banicz, et al., ϕ production in In-In collisions at 158-A-GeV, *Eur. Phys. J. C* 64 (2009) 1–18, arXiv:0906.1102 [hep-ex].
- [22] NA60 Collaboration, R. Arnaldi, et al., A comparative measurement of $\phi \rightarrow K^+ K^-$ and $\phi \rightarrow \mu^+ \mu^-$ in In-In collisions at the CERN SPS, *Phys. Lett. B* 699 (2011) 325–329, arXiv:1104.4060 [nucl-ex].

- [23] STAR Collaboration, B. Abelev, et al., Measurements of ϕ meson production in relativistic heavy-ion collisions at RHIC, Phys. Rev. C 79 (2009) 064903, arXiv: 0809.4737 [nucl-ex].
- [24] PHENIX Collaboration, S.S. Adler, et al., Production of ϕ mesons at mid-rapidity in $\sqrt{s_{NN}} = 200$ GeV Au+Au collisions at RHIC, Phys. Rev. C 72 (2005) 014903, arXiv:nucl-ex/0410012.
- [25] ALICE Collaboration, B. Abelev, et al., Light vector meson production in pp collisions at $\sqrt{s} = 7$ TeV, Phys. Lett. B 710 (2012) 557–568, arXiv:1112.2222 [nucl-ex].
- [26] ALICE Collaboration, K. Aamodt, et al., The ALICE experiment at the CERN LHC, J. Instrum. 3 (2008) S08002.
- [27] ALICE Collaboration, B. Abelev, et al., Performance of the ALICE experiment at the CERN LHC, Int. J. Mod. Phys. A 29 (2014) 1430044, arXiv:1402.4476 [nucl-ex].
- [28] ALICE Collaboration, M. Gagliardi, Measurement of reference cross sections in pp and Pb-Pb collisions at the LHC in van der Meer scans with the ALICE detector, AIP Conf. Proc. 1422 (2012) 110–116, arXiv:1109.5369 [hep-ex].
- [29] ALICE Collaboration, B. Abelev, et al., J/ψ production and nuclear effects in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, J. High Energy Phys. 1402 (2014) 073, arXiv:1308.6726 [nucl-ex].
- [30] ALICE Collaboration, B. Abelev, et al., Measurement of visible cross sections in proton-lead collisions at $\sqrt{s_{NN}} = 5.02$ TeV in van der Meer scans with the ALICE detector, J. Instrum. 9 (11) (2014) P11003, arXiv:1405.1849 [nucl-ex].
- [31] ALICE Collaboration, B. Abelev, et al., Measurement of quarkonium production at forward rapidity in pp collisions at $\sqrt{s} = 7$ TeV, Eur. Phys. J. C 74 (8) (2014) 2974, arXiv:1403.3648 [nucl-ex].
- [32] ALICE Collaboration, B. Abelev, et al., Measurement of inelastic, single- and double-diffractive cross sections in proton-proton collisions at the LHC with ALICE, Eur. Phys. J. C 73 (6) (2013) 2456, arXiv:1208.4968 [hep-ex].
- [33] ALICE Collaboration, L. Aphecetche, et al., Numerical Simulations and Offline Reconstruction of the Muon Spectrometer of ALICE, 2009, ALICE-INT-2009-044.
- [34] ALICE Collaboration, G. Chabratova, et al., Development of the Kalman Filter for Tracking in the Forward Muon Spectrometer of ALICE, 2003, ALICE-INT-2003-002.
- [35] ALICE Collaboration, Supplemental figures for “ ϕ -Meson production at forward rapidity in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV and in pp collisions at $\sqrt{s} = 2.76$ TeV”, <https://cds.cern.ch/record/2195308>.
- [36] T. Sjostrand, S. Mrenna, P.Z. Skands, PYTHIA 6.4 physics and manual, J. High Energy Phys. 0605 (2006) 026, arXiv:hep-ph/0603175.
- [37] R. Brun, F. Carminati, S. Giani, GEANT Detector Description and Simulation Tool, CERN-W-5013, 1994.
- [38] Particle Data Group Collaboration, J. Beringer, et al., Review of Particle Physics (RPP), Phys. Rev. D 86 (2012) 010001.
- [39] M. Oreiglia, A Study of the Reactions $\psi' \rightarrow \gamma\gamma\psi$, PhD thesis, SLAC, 1980, <http://search.proquest.com/docview/303036283>.
- [40] ALICE Collaboration, K. Aamodt, et al., Rapidity and transverse momentum dependence of inclusive J/ψ production in pp collisions at $\sqrt{s} = 7$ TeV, Phys. Lett. B 704 (2011) 442–455, arXiv:1105.0380 [hep-ex]; ALICE Collaboration, K. Aamodt, et al., Phys. Lett. B 718 (2012) 692 (Erratum).
- [41] R. Barlow, Systematic errors: facts and fictions, arXiv:hep-ex/0207026.
- [42] R. Engel, J. Ranft, Hadronic photon-photon interactions at high-energies, Phys. Rev. D 54 (1996) 4244–4262, arXiv:hep-ph/9509373.
- [43] P.Z. Skands, Tuning Monte Carlo generators: the Perugia tunes, Phys. Rev. D 82 (2010) 074018, arXiv:1005.3457 [hep-ph].
- [44] C. Buttar, D. Clements, I. Dawson, A. Moraes, Simulations of minimum bias events and the underlying event, MC tuning and predictions for the LHC, Acta Phys. Pol. B 35 (2004) 433–441.
- [45] R. Field, Physics at the Tevatron, Acta Phys. Pol. B 39 (2008) 2611–2672.
- [46] ALICE Collaboration, B. Abelev, et al., Production of $K^*(892)^0$ and $\phi(1020)$ in pp collisions at $\sqrt{s} = 7$ TeV, Eur. Phys. J. C 72 (2012) 2183, arXiv:1208.5717 [hep-ex].
- [47] CMS Collaboration, S. Chatrchyan, et al., Study of the inclusive production of charged pions, kaons, and protons in pp collisions at $\sqrt{s} = 0.9$, 2.76, and 7 TeV, Eur. Phys. J. C 72 (2012) 2164, arXiv:1207.4724 [hep-ex].
- [48] C. Tsallis, Possible generalization of Boltzmann-Gibbs statistics, J. Stat. Phys. 52 (1988) 479–487.
- [49] X.-N. Wang, M. Gyulassy, HIJING: a Monte Carlo model for multiple jet production in p , pA and AA collisions, Phys. Rev. D 44 (1991) 3501–3516.
- [50] S. Roesler, R. Engel, J. Ranft, The Monte Carlo event generator DPMJET-III, in: Advanced Monte Carlo for Radiation Physics, Particle Transport Simulation and Applications. Proceedings, Conference, MC2000, Lisbon, Portugal, October 23–26, 2000, 2000, pp. 1033–1038, arXiv:hep-ph/0012252, <http://www-public.slac.stanford.edu/sciDoc/docMeta.aspx?slacPubNumber=SLAC-PUB-8740>.
- [51] ALICE Collaboration, J. Adam, et al., Production of $K^*(892)^0$ and $\phi(1020)$ in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Eur. Phys. J. C 76 (5) (2016) 245, arXiv: 1601.07868 [nucl-ex].
- [52] ALICE, LHCb Collaboration, Reference pp cross-sections for J/ψ studies in proton-lead collisions at $\sqrt{s_{NN}} = 5.02$ TeV and comparisons between ALICE and LHCb results, <http://cds.cern.ch/record/1639617>, CONF-2013-013.
- [53] HERA-B Collaboration, I. Abt, et al., K^0 and ϕ meson production in proton-nucleus interactions at $\sqrt{s} = 41.6$ GeV, Eur. Phys. J. C 50 (2007) 315–328, arXiv:hep-ex/0606049.
- [54] PHENIX Collaboration, A. Adare, et al., ϕ meson production in d +Au collisions at $\sqrt{s_{NN}} = 200$ GeV, Phys. Rev. C 92 (4) (2015) 044909, arXiv:1506.08181 [nucl-ex].

The ALICE Collaboration

J. Adam ⁴⁰, D. Adamová ⁸³, M.M. Aggarwal ⁸⁷, G. Aglieri Rinella ³⁶, M. Agnello ¹¹¹, N. Agrawal ⁴⁸, Z. Ahammed ¹³², S.U. Ahn ⁶⁸, I. Aimo ^{94,111}, S. Aiola ¹³⁶, M. Ajaz ¹⁶, A. Akindinov ⁵⁸, S.N. Alam ¹³², D. Aleksandrov ¹⁰⁰, B. Alessandro ¹¹¹, D. Alexandre ¹⁰², R. Alfaro Molina ⁶⁴, A. Alici ^{105,12}, A. Alkin ³, J.R.M. Almaraz ¹¹⁹, J. Alme ³⁸, T. Alt ⁴³, S. Altinpinar ¹⁸, I. Altsybeev ¹³¹, C. Alves Garcia Prado ¹²⁰, C. Andrei ⁷⁸, A. Andronic ⁹⁷, V. Anguelov ⁹³, J. Anielski ⁵⁴, T. Antićić ⁹⁸, F. Antinori ¹⁰⁸, P. Antonioli ¹⁰⁵, L. Aphecetche ¹¹³, H. Appelshäuser ⁵³, S. Arcelli ²⁸, N. Armesto ¹⁷, R. Arnaldi ¹¹¹, I.C. Arsene ²², M. Arslanbek ⁵³, B. Audurier ¹¹³, A. Augustinus ³⁶, R. Averbeck ⁹⁷, M.D. Azmi ¹⁹, M. Bach ⁴³, A. Badalà ¹⁰⁷, Y.W. Baek ⁴⁴, S. Bagnasco ¹¹¹, R. Bailhache ⁵³, R. Bala ⁹⁰, A. Baldissari ¹⁵, F. Baltasar Dos Santos Pedrosa ³⁶, R.C. Baral ⁶¹, A.M. Barbano ¹¹¹, R. Barbera ²⁹, F. Barile ³³, G.G. Barnaföldi ¹³⁵, L.S. Barnby ¹⁰², V. Barret ⁷⁰, P. Bartalini ⁷, K. Barth ³⁶, J. Bartke ¹¹⁷, E. Bartsch ⁵³, M. Basile ²⁸, N. Bastid ⁷⁰, S. Basu ¹³², B. Bathen ⁵⁴, G. Batigne ¹¹³, A. Batista Camejo ⁷⁰, B. Batyunya ⁶⁶, P.C. Batzing ²², I.G. Bearden ⁸⁰, H. Beck ⁵³, C. Bedda ¹¹¹, N.K. Behera ^{48,49}, I. Belikov ⁵⁵, F. Bellini ²⁸, H. Bello Martinez ², R. Bellwied ¹²², R. Belmont ¹³⁴, E. Belmont-Moreno ⁶⁴, V. Belyaev ⁷⁶, G. Bencedi ¹³⁵, S. Beole ²⁷, I. Berceanu ⁷⁸, A. Bercuci ⁷⁸, Y. Berdnikov ⁸⁵, D. Berenyi ¹³⁵, R.A. Bertens ⁵⁷, D. Berzana ^{36,27}, L. Betev ³⁶, A. Bhasin ⁹⁰, I.R. Bhat ⁹⁰, A.K. Bhati ⁸⁷, B. Bhattacharjee ⁴⁵, J. Bhom ¹²⁸, L. Bianchi ¹²², N. Bianchi ⁷², C. Bianchin ^{134,57}, J. Bielčík ⁴⁰, J. Bielčíková ⁸³, A. Bilandžić ⁸⁰, R. Biswas ⁴, S. Biswas ⁷⁹, S. Bjelogrlic ⁵⁷, J.T. Blair ¹¹⁸, F. Blanco ¹⁰, D. Blau ¹⁰⁰, C. Blume ⁵³, F. Bock ^{93,74}, A. Bogdanov ⁷⁶, H. Bøggild ⁸⁰, L. Boldizsár ¹³⁵, M. Bombara ⁴¹, J. Book ⁵³, H. Borel ¹⁵, A. Borissov ⁹⁶, M. Borri ⁸², F. Bossú ⁶⁵, E. Botta ²⁷, S. Böttger ⁵², P. Braun-Munzinger ⁹⁷, M. Bregant ¹²⁰, T. Breitner ⁵², T.A. Broker ⁵³, T.A. Browning ⁹⁵, M. Broz ⁴⁰, E.J. Brucken ⁴⁶, E. Bruna ¹¹¹, G.E. Bruno ³³, D. Budnikov ⁹⁹, H. Buesching ⁵³, S. Bufalino ^{27,111}, P. Buncic ³⁶, O. Busch ^{128,93}, Z. Buthelezi ⁶⁵, J.B. Butt ¹⁶, J.T. Buxton ²⁰, D. Caffarri ³⁶, X. Cai ⁷, H. Caines ¹³⁶,

- L. Calero Diaz ⁷², A. Caliva ⁵⁷, E. Calvo Villar ¹⁰³, P. Camerini ²⁶, F. Carena ³⁶, W. Carena ³⁶,
 F. Carnesecchi ²⁸, J. Castillo Castellanos ¹⁵, A.J. Castro ¹²⁵, E.A.R. Casula ²⁵, C. Cavicchioli ³⁶,
 C. Ceballos Sanchez ⁹, J. Cepila ⁴⁰, P. Cerello ¹¹¹, J. Cerkala ¹¹⁵, B. Chang ¹²³, S. Chapeland ³⁶,
 M. Chartier ¹²⁴, J.L. Charvet ¹⁵, S. Chattopadhyay ¹³², S. Chattopadhyay ¹⁰¹, V. Chelnokov ³, M. Cherney ⁸⁶,
 C. Cheshkov ¹³⁰, B. Cheynis ¹³⁰, V. Chibante Barroso ³⁶, D.D. Chinellato ¹²¹, P. Chochula ³⁶, K. Choi ⁹⁶,
 M. Chojnacki ⁸⁰, S. Choudhury ¹³², P. Christakoglou ⁸¹, C.H. Christensen ⁸⁰, P. Christiansen ³⁴, T. Chujo ¹²⁸,
 S.U. Chung ⁹⁶, Z. Chunhui ⁵⁷, C. Cicalo ¹⁰⁶, L. Cifarelli ^{12,28}, F. Cindolo ¹⁰⁵, J. Cleymans ⁸⁹, F. Colamaria ³³,
 D. Colella ^{36,33,59}, A. Collu ²⁵, M. Colocci ²⁸, G. Conesa Balbastre ⁷¹, Z. Conesa del Valle ⁵¹,
 M.E. Connors ¹³⁶, J.G. Contreras ^{11,40}, T.M. Cormier ⁸⁴, Y. Corrales Morales ²⁷, I. Cortés Maldonado ²,
 P. Cortese ³², M.R. Cosentino ¹²⁰, F. Costa ³⁶, P. Crochet ⁷⁰, R. Cruz Albino ¹¹, E. Cuautle ⁶³, L. Cunqueiro ³⁶,
 T. Dahms ^{92,37}, A. Dainese ¹⁰⁸, A. Danu ⁶², D. Das ¹⁰¹, I. Das ^{101,51}, S. Das ⁴, A. Dash ¹²¹, S. Dash ⁴⁸,
 S. De ¹²⁰, A. De Caro ^{31,12}, G. de Cataldo ¹⁰⁴, J. de Cuveland ⁴³, A. De Falco ²⁵, D. De Gruttola ^{12,31},
 N. De Marco ¹¹¹, S. De Pasquale ³¹, A. Deisting ^{97,93}, A. Deloff ⁷⁷, E. Dénes ^{135,i}, G. D'Erasmo ³³,
 D. Di Bari ³³, A. Di Mauro ³⁶, P. Di Nezza ⁷², M.A. Diaz Corchero ¹⁰, T. Dietel ⁸⁹, P. Dillenseger ⁵³,
 R. Divià ³⁶, Ø. Djupsland ¹⁸, A. Dobrin ^{57,81}, T. Dobrowolski ^{77,i}, D. Domenicis Gimenez ¹²⁰, B. Dönigus ⁵³,
 O. Dordic ²², T. Drozhzhova ⁵³, A.K. Dubey ¹³², A. Dubla ⁵⁷, L. Ducroux ¹³⁰, P. Dupieux ⁷⁰, R.J. Ehlers ¹³⁶,
 D. Elia ¹⁰⁴, H. Engel ⁵², B. Erazmus ^{36,113}, I. Erdemir ⁵³, F. Erhardt ¹²⁹, D. Eschweiler ⁴³, B. Espagnon ⁵¹,
 M. Estienne ¹¹³, S. Esumi ¹²⁸, J. Eum ⁹⁶, D. Evans ¹⁰², S. Evdokimov ¹¹², G. Eyyubova ⁴⁰, L. Fabbietti ^{37,92},
 D. Fabris ¹⁰⁸, J. Faivre ⁷¹, A. Fantoni ⁷², M. Fasel ⁷⁴, L. Feldkamp ⁵⁴, D. Felea ⁶², A. Feliciello ¹¹¹,
 G. Feofilov ¹³¹, J. Ferencei ⁸³, A. Fernández Téllez ², E.G. Ferreiro ¹⁷, A. Ferretti ²⁷, A. Festanti ³⁰,
 V.J.G. Feuillard ^{15,70}, J. Figiel ¹¹⁷, M.A.S. Figueiredo ^{124,120}, S. Filchagin ⁹⁹, D. Finogeev ⁵⁶, F.M. Fionda ²⁵,
 E.M. Fiore ³³, M.G. Fleck ⁹³, M. Floris ³⁶, S. Foertsch ⁶⁵, P. Foka ⁹⁷, S. Fokin ¹⁰⁰, E. Fragiacomo ¹¹⁰,
 A. Francescon ^{36,30}, U. Frankenfeld ⁹⁷, U. Fuchs ³⁶, C. Furget ⁷¹, A. Furs ⁵⁶, M. Fusco Girard ³¹,
 J.J. Gaardhøje ⁸⁰, M. Gagliardi ²⁷, A.M. Gago ¹⁰³, M. Gallio ²⁷, D.R. Gangadharan ⁷⁴, P. Ganoti ⁸⁸, C. Gao ⁷,
 C. Garabatos ⁹⁷, E. Garcia-Solis ¹³, C. Gargiulo ³⁶, P. Gasik ^{92,37}, M. Germain ¹¹³, A. Gheata ³⁶,
 M. Gheata ^{62,36}, P. Ghosh ¹³², S.K. Ghosh ⁴, P. Gianotti ⁷², P. Giubellino ^{36,111}, P. Giubilato ³⁰,
 E. Gladysz-Dziadus ¹¹⁷, P. Glässel ⁹³, D.M. Goméz Coral ⁶⁴, A. Gomez Ramirez ⁵², P. González-Zamora ¹⁰,
 S. Gorbunov ⁴³, L. Görlich ¹¹⁷, S. Gotovac ¹¹⁶, V. Grabski ⁶⁴, L.K. Graczykowski ¹³³, K.L. Graham ¹⁰²,
 A. Grelli ⁵⁷, A. Grigoras ³⁶, C. Grigoras ³⁶, V. Grigoriev ⁷⁶, A. Grigoryan ¹, S. Grigoryan ⁶⁶, B. Grinyov ³,
 N. Grion ¹¹⁰, J.F. Grosse-Oetringhaus ³⁶, J-Y. Grossiord ¹³⁰, R. Grossi ³⁶, F. Guber ⁵⁶, R. Guernane ⁷¹,
 B. Guerzoni ²⁸, K. Gulbrandsen ⁸⁰, H. Gulkanyan ¹, T. Gunji ¹²⁷, A. Gupta ⁹⁰, R. Gupta ⁹⁰, R. Haake ⁵⁴,
 Ø. Haaland ¹⁸, C. Hadjidakis ⁵¹, M. Haiduc ⁶², H. Hamagaki ¹²⁷, G. Hamar ¹³⁵, A. Hansen ⁸⁰, J.W. Harris ¹³⁶,
 H. Hartmann ⁴³, A. Harton ¹³, D. Hatzifotiadou ¹⁰⁵, S. Hayashi ¹²⁷, S.T. Heckel ⁵³, M. Heide ⁵⁴,
 H. Helstrup ³⁸, A. Hergheliegiu ⁷⁸, G. Herrera Corral ¹¹, B.A. Hess ³⁵, K.F. Hetland ³⁸, T.E. Hilden ⁴⁶,
 H. Hillemanns ³⁶, B. Hippolyte ⁵⁵, R. Hosokawa ¹²⁸, P. Hristov ³⁶, M. Huang ¹⁸, T.J. Humanic ²⁰,
 N. Hussain ⁴⁵, T. Hussain ¹⁹, D. Hutter ⁴³, D.S. Hwang ²¹, R. Ilkaev ⁹⁹, I. Ilkiv ⁷⁷, M. Inaba ¹²⁸,
 M. Ippolitov ^{76,100}, M. Irfan ¹⁹, M. Ivanov ⁹⁷, V. Ivanov ⁸⁵, V. Izucheev ¹¹², P.M. Jacobs ⁷⁴, S. Jadlovska ¹¹⁵,
 C. Jahnke ¹²⁰, H.J. Jang ⁶⁸, M.A. Janik ¹³³, P.H.S.Y. Jayaratna ¹²², C. Jena ³⁰, S. Jena ¹²²,
 R.T. Jimenez Bustamante ⁹⁷, P.G. Jones ¹⁰², H. Jung ⁴⁴, A. Jusko ¹⁰², P. Kalinak ⁵⁹, A. Kalweit ³⁶, J. Kamin ⁵³,
 J.H. Kang ¹³⁷, V. Kaplin ⁷⁶, S. Kar ¹³², A. Karasu Uysal ⁶⁹, O. Karavichev ⁵⁶, T. Karavicheva ⁵⁶,
 L. Karayan ^{93,97}, E. Karpechev ⁵⁶, U. Kebschull ⁵², R. Keidel ¹³⁸, D.L.D. Keijdener ⁵⁷, M. Keil ³⁶, K.H. Khan ¹⁶,
 M.M. Khan ¹⁹, P. Khan ¹⁰¹, S.A. Khan ¹³², A. Khanzadeev ⁸⁵, Y. Kharlov ¹¹², B. Kileng ³⁸, B. Kim ¹³⁷,
 D.W. Kim ^{44,68}, D.J. Kim ¹²³, H. Kim ¹³⁷, J.S. Kim ⁴⁴, M. Kim ⁴⁴, M. Kim ¹³⁷, S. Kim ²¹, T. Kim ¹³⁷,
 S. Kirsch ⁴³, I. Kisel ⁴³, S. Kiselev ⁵⁸, A. Kisiel ¹³³, G. Kiss ¹³⁵, J.L. Klay ⁶, C. Klein ⁵³, J. Klein ^{36,93},
 C. Klein-Bösing ⁵⁴, A. Kluge ³⁶, M.L. Knichel ⁹³, A.G. Knospe ¹¹⁸, T. Kobayashi ¹²⁸, C. Kobdaj ¹¹⁴,
 M. Kofarago ³⁶, T. Kollegger ^{97,43}, A. Kolojvari ¹³¹, V. Kondratiev ¹³¹, N. Kondratyeva ⁷⁶, E. Kondratyuk ¹¹²,
 A. Konevskikh ⁵⁶, M. Kopcik ¹¹⁵, M. Kour ⁹⁰, C. Kouzinopoulos ³⁶, O. Kovalenko ⁷⁷, V. Kovalenko ¹³¹,
 M. Kowalski ¹¹⁷, G. Koyithatta Meethaleveedu ⁴⁸, J. Kral ¹²³, I. Králik ⁵⁹, A. Kravčáková ⁴¹, M. Kretz ⁴³,
 M. Krivda ^{59,102}, F. Krizek ⁸³, E. Kryshen ³⁶, M. Krzewicki ⁴³, A.M. Kubera ²⁰, V. Kučera ⁸³, T. Kugathasan ³⁶,
 C. Kuhn ⁵⁵, P.G. Kuijer ⁸¹, A. Kumar ⁹⁰, J. Kumar ⁴⁸, L. Kumar ^{79,87}, P. Kurashvili ⁷⁷, A. Kurepin ⁵⁶,
 A.B. Kurepin ⁵⁶, A. Kuryakin ⁹⁹, S. Kushpil ⁸³, M.J. Kweon ⁵⁰, Y. Kwon ¹³⁷, S.L. La Pointe ¹¹¹, P. La Rocca ²⁹,
 C. Lagana Fernandes ¹²⁰, I. Lakomov ³⁶, R. Langoy ⁴², C. Lara ⁵², A. Lardeux ¹⁵, A. Lattuca ²⁷, E. Laudi ³⁶

- R. Lea ²⁶, L. Leardini ⁹³, G.R. Lee ¹⁰², S. Lee ¹³⁷, I. Legrand ³⁶, F. Lehas ⁸¹, R.C. Lemmon ⁸², V. Lenti ¹⁰⁴, E. Leogrande ⁵⁷, I. León Monzón ¹¹⁹, M. Leoncino ²⁷, P. Lévai ¹³⁵, S. Li ^{7,70}, X. Li ¹⁴, J. Lien ⁴², R. Lietava ¹⁰², S. Lindal ²², V. Lindenstruth ⁴³, C. Lippmann ⁹⁷, M.A. Lisa ²⁰, H.M. Ljunggren ³⁴, D.F. Lodato ⁵⁷, P.I. Loenne ¹⁸, V. Loginov ⁷⁶, C. Loizides ⁷⁴, X. Lopez ⁷⁰, E. López Torres ⁹, A. Lowe ¹³⁵, P. Luettig ⁵³, M. Lunardon ³⁰, G. Luparello ²⁶, P.H.F.N.D. Luz ¹²⁰, A. Maevskaya ⁵⁶, M. Mager ³⁶, S. Mahajan ⁹⁰, S.M. Mahmood ²², A. Maire ⁵⁵, R.D. Majka ¹³⁶, M. Malaev ⁸⁵, I. Maldonado Cervantes ⁶³, L. Malinina ^{66,ii}, D. Mal'Kevich ⁵⁸, P. Malzacher ⁹⁷, A. Mamonov ⁹⁹, V. Manko ¹⁰⁰, F. Manso ⁷⁰, V. Manzari ^{36,104}, M. Marchisone ²⁷, J. Mareš ⁶⁰, G.V. Margagliotti ²⁶, A. Margotti ¹⁰⁵, J. Margutti ⁵⁷, A. Marín ⁹⁷, C. Markert ¹¹⁸, M. Marquard ⁵³, N.A. Martin ⁹⁷, J. Martin Blanco ¹¹³, P. Martinengo ³⁶, M.I. Martínez ², G. Martínez García ¹¹³, M. Martinez Pedreira ³⁶, Y. Martynov ³, A. Mas ¹²⁰, S. Masciocchi ⁹⁷, M. Masera ²⁷, A. Masoni ¹⁰⁶, L. Massacrier ¹¹³, A. Mastroserio ³³, H. Masui ¹²⁸, A. Matyja ¹¹⁷, C. Mayer ¹¹⁷, J. Mazer ¹²⁵, M.A. Mazzoni ¹⁰⁹, D. McDonald ¹²², F. Meddi ²⁴, Y. Melikyan ⁷⁶, A. Menchaca-Rocha ⁶⁴, E. Meninno ³¹, J. Mercado Pérez ⁹³, M. Meres ³⁹, Y. Miake ¹²⁸, M.M. Mieskolainen ⁴⁶, K. Mikhaylov ^{66,58}, L. Milano ³⁶, J. Milosevic ²², L.M. Minervini ^{104,23}, A. Mischke ⁵⁷, A.N. Mishra ⁴⁹, D. Miśkowiec ⁹⁷, J. Mitra ¹³², C.M. Mitu ⁶², N. Mohammadi ⁵⁷, B. Mohanty ^{132,79}, L. Molnar ⁵⁵, L. Montaño Zetina ¹¹, E. Montes ¹⁰, M. Morando ³⁰, D.A. Moreira De Godoy ^{113,54}, S. Moretto ³⁰, A. Morreale ¹¹³, A. Morsch ³⁶, V. Muccifora ⁷², E. Mudnic ¹¹⁶, D. Mühlheim ⁵⁴, S. Muhuri ¹³², M. Mukherjee ¹³², J.D. Mulligan ¹³⁶, M.G. Munhoz ¹²⁰, S. Murray ⁶⁵, L. Musa ³⁶, J. Musinsky ⁵⁹, B.K. Nandi ⁴⁸, R. Nania ¹⁰⁵, E. Nappi ¹⁰⁴, M.U. Naru ¹⁶, C. Natrass ¹²⁵, K. Nayak ⁷⁹, T.K. Nayak ¹³², S. Nazarenko ⁹⁹, A. Nedosekin ⁵⁸, L. Nellen ⁶³, F. Ng ¹²², M. Nicassio ⁹⁷, M. Niculescu ^{62,36}, J. Niedziela ³⁶, B.S. Nielsen ⁸⁰, S. Nikolaev ¹⁰⁰, S. Nikulin ¹⁰⁰, V. Nikulin ⁸⁵, F. Noferini ^{105,12}, P. Nomokonov ⁶⁶, G. Nooren ⁵⁷, J.C.C. Noris ², J. Norman ¹²⁴, A. Nyanin ¹⁰⁰, J. Nystrand ¹⁸, H. Oeschler ⁹³, S. Oh ¹³⁶, S.K. Oh ⁶⁷, A. Ohlson ³⁶, A. Okatan ⁶⁹, T. Okubo ⁴⁷, L. Olah ¹³⁵, J. Oleniacz ¹³³, A.C. Oliveira Da Silva ¹²⁰, M.H. Oliver ¹³⁶, J. Onderwaater ⁹⁷, C. Oppedisano ¹¹¹, R. Orava ⁴⁶, A. Ortiz Velasquez ⁶³, A. Oskarsson ³⁴, J. Otwinowski ¹¹⁷, K. Oyama ⁹³, M. Ozdemir ⁵³, Y. Pachmayer ⁹³, P. Pagano ³¹, G. Paić ⁶³, C. Pajares ¹⁷, S.K. Pal ¹³², J. Pan ¹³⁴, A.K. Pandey ⁴⁸, D. Pant ⁴⁸, P. Papcun ¹¹⁵, V. Papikyan ¹, G.S. Pappalardo ¹⁰⁷, P. Pareek ⁴⁹, W.J. Park ⁹⁷, S. Parmar ⁸⁷, A. Passfeld ⁵⁴, V. Paticchio ¹⁰⁴, R.N. Patra ¹³², B. Paul ¹⁰¹, T. Peitzmann ⁵⁷, H. Pereira Da Costa ¹⁵, E. Pereira De Oliveira Filho ¹²⁰, D. Peresunko ^{100,76}, C.E. Pérez Lara ⁸¹, E. Perez Lezama ⁵³, V. Peskov ⁵³, Y. Pestov ⁵, V. Petráček ⁴⁰, V. Petrov ¹¹², M. Petrovici ⁷⁸, C. Petta ²⁹, S. Piano ¹¹⁰, M. Pikna ³⁹, P. Pillot ¹¹³, O. Pinazza ^{105,36}, L. Pinsky ¹²², D.B. Piyarathna ¹²², M. Płoskoń ⁷⁴, M. Planinic ¹²⁹, J. Pluta ¹³³, S. Pochybova ¹³⁵, P.L.M. Podesta-Lerma ¹¹⁹, M.G. Poghosyan ^{86,84}, B. Polichtchouk ¹¹², N. Poljak ¹²⁹, W. Poonsawat ¹¹⁴, A. Pop ⁷⁸, S. Porteboeuf-Houssais ⁷⁰, J. Porter ⁷⁴, J. Pospisil ⁸³, S.K. Prasad ⁴, R. Preghenella ^{36,105}, F. Prino ¹¹¹, C.A. Pruneau ¹³⁴, I. Pshenichnov ⁵⁶, M. Puccio ¹¹¹, G. Puddu ²⁵, P. Pujahari ¹³⁴, V. Punin ⁹⁹, J. Putschke ¹³⁴, H. Qvigstad ²², A. Rachevski ¹¹⁰, S. Raha ⁴, S. Rajput ⁹⁰, J. Rak ¹²³, A. Rakotozafindrabe ¹⁵, L. Ramello ³², F. Rami ⁵⁵, R. Raniwala ⁹¹, S. Raniwala ⁹¹, S.S. Räsänen ⁴⁶, B.T. Rascanu ⁵³, D. Rathee ⁸⁷, K.F. Read ¹²⁵, J.S. Real ⁷¹, K. Redlich ⁷⁷, R.J. Reed ¹³⁴, A. Rehman ¹⁸, P. Reichelt ⁵³, F. Reidt ^{93,36}, X. Ren ⁷, R. Renfordt ⁵³, A.R. Reolon ⁷², A. Reshetin ⁵⁶, F. Rettig ⁴³, J.-P. Revol ¹², K. Reygers ⁹³, V. Riabov ⁸⁵, R.A. Ricci ⁷³, T. Richert ³⁴, M. Richter ²², P. Riedler ³⁶, W. Riegler ³⁶, F. Riggi ²⁹, C. Ristea ⁶², A. Rivetti ¹¹¹, E. Rocco ⁵⁷, M. Rodríguez Cahuantzi ², A. Rodriguez Manso ⁸¹, K. Røed ²², E. Rogochaya ⁶⁶, D. Rohr ⁴³, D. Röhrlrich ¹⁸, R. Romita ¹²⁴, F. Ronchetti ⁷², L. Ronflette ¹¹³, P. Rosnet ⁷⁰, A. Rossi ^{30,36}, F. Roukoutakis ⁸⁸, A. Roy ⁴⁹, C. Roy ⁵⁵, P. Roy ¹⁰¹, A.J. Rubio Montero ¹⁰, R. Rui ²⁶, R. Russo ²⁷, E. Ryabinkin ¹⁰⁰, Y. Ryabov ⁸⁵, A. Rybicki ¹¹⁷, S. Sadovsky ¹¹², K. Šafařík ³⁶, B. Sahlmuller ⁵³, P. Sahoo ⁴⁹, R. Sahoo ⁴⁹, S. Sahoo ⁶¹, P.K. Sahu ⁶¹, J. Saini ¹³², S. Sakai ⁷², M.A. Saleh ¹³⁴, C.A. Salgado ¹⁷, J. Salzwedel ²⁰, S. Sambyal ⁹⁰, V. Samsonov ⁸⁵, X. Sanchez Castro ⁵⁵, L. Šándor ⁵⁹, A. Sandoval ⁶⁴, M. Sano ¹²⁸, D. Sarkar ¹³², E. Scapparone ¹⁰⁵, F. Scarlassara ³⁰, R.P. Scharenberg ⁹⁵, C. Schiaua ⁷⁸, R. Schicker ⁹³, C. Schmidt ⁹⁷, H.R. Schmidt ³⁵, S. Schuchmann ⁵³, J. Schukraft ³⁶, M. Schulc ⁴⁰, T. Schuster ¹³⁶, Y. Schutz ^{113,36}, K. Schwarz ⁹⁷, K. Schweda ⁹⁷, G. Scioli ²⁸, E. Scomparin ¹¹¹, R. Scott ¹²⁵, J.E. Seger ⁸⁶, Y. Sekiguchi ¹²⁷, D. Sekihata ⁴⁷, I. Selyuzhenkov ⁹⁷, K. Senosi ⁶⁵, J. Seo ^{96,67}, E. Serradilla ^{64,10}, A. Sevcenco ⁶², A. Shabanov ⁵⁶, A. Shabetai ¹¹³, O. Shadura ³, R. Shahoyan ³⁶, A. Shangaraev ¹¹², A. Sharma ⁹⁰, M. Sharma ⁹⁰, M. Sharma ⁹⁰, N. Sharma ^{125,61}, K. Shigaki ⁴⁷, K. Shtejer ^{9,27}, Y. Sibiriak ¹⁰⁰, S. Siddhanta ¹⁰⁶, K.M. Sielewicz ³⁶, T. Siemiarczuk ⁷⁷, D. Silvermyr ^{84,34}, C. Silvestre ⁷¹, G. Simatovic ¹²⁹, G. Simonetti ³⁶, R. Singaraju ¹³², R. Singh ⁷⁹, S. Singha ^{132,79}, V. Singhal ¹³², B.C. Sinha ¹³², T. Sinha ¹⁰¹, B. Sitar ³⁹,

M. Sitta ³², T.B. Skaali ²², M. Slupecki ¹²³, N. Smirnov ¹³⁶, R.J.M. Snellings ⁵⁷, T.W. Snellman ¹²³,
 C. Søgaard ³⁴, R. Soltz ⁷⁵, J. Song ⁹⁶, M. Song ¹³⁷, Z. Song ⁷, F. Soramel ³⁰, S. Sorensen ¹²⁵, M. Spacek ⁴⁰,
 E. Spiriti ⁷², I. Sputowska ¹¹⁷, M. Spyropoulou-Stassinaki ⁸⁸, B.K. Srivastava ⁹⁵, J. Stachel ⁹³, I. Stan ⁶²,
 G. Stefanek ⁷⁷, M. Steinpreis ²⁰, E. Stenlund ³⁴, G. Steyn ⁶⁵, J.H. Stiller ⁹³, D. Stocco ¹¹³, P. Strmen ³⁹,
 A.A.P. Suaide ¹²⁰, T. Sugitate ⁴⁷, C. Suire ⁵¹, M. Suleymanov ¹⁶, R. Sultanov ⁵⁸, M. Šumbera ⁸³,
 T.J.M. Symons ⁷⁴, A. Szabo ³⁹, A. Szanto de Toledo ^{120,i}, I. Szarka ³⁹, A. Szczepankiewicz ³⁶,
 M. Szymanski ¹³³, U. Tabassam ¹⁶, J. Takahashi ¹²¹, G.J. Tambave ¹⁸, N. Tanaka ¹²⁸, M.A. Tangaro ³³,
 J.D. Tapia Takaki ^{51,iii}, A. Tarantola Peloni ⁵³, M. Tarhini ⁵¹, M. Tariq ¹⁹, M.G. Tarzila ⁷⁸, A. Tauro ³⁶,
 G. Tejeda Muñoz ², A. Telesca ³⁶, K. Terasaki ¹²⁷, C. Terrevoli ^{30,25}, B. Teyssier ¹³⁰, J. Thäder ^{74,97},
 D. Thomas ¹¹⁸, R. Tieulent ¹³⁰, A.R. Timmins ¹²², A. Toia ⁵³, S. Trogolo ¹¹¹, V. Trubnikov ³, W.H. Trzaska ¹²³,
 T. Tsuji ¹²⁷, A. Tumkin ⁹⁹, R. Turrisi ¹⁰⁸, T.S. Tveter ²², K. Ullaland ¹⁸, A. Uras ¹³⁰, G.L. Usai ²⁵,
 A. Utrobitic ¹²⁹, M. Vajzer ⁸³, M. Vala ⁵⁹, L. Valencia Palomo ⁷⁰, S. Vallero ²⁷, J. Van Der Maarel ⁵⁷,
 J.W. Van Hoorn ³⁶, M. van Leeuwen ⁵⁷, T. Vanat ⁸³, P. Vande Vyvre ³⁶, D. Varga ¹³⁵, A. Vargas ²,
 M. Vargyas ¹²³, R. Varma ⁴⁸, M. Vasileiou ⁸⁸, A. Vasiliev ¹⁰⁰, A. Vauthier ⁷¹, V. Vechernin ¹³¹, A.M. Veen ⁵⁷,
 M. Veldhoen ⁵⁷, A. Velure ¹⁸, M. Venaruzzo ⁷³, E. Vercellin ²⁷, S. Vergara Limón ², R. Vernet ⁸,
 M. Verweij ^{134,36}, L. Vickovic ¹¹⁶, G. Viesti ^{30,i}, J. Viinikainen ¹²³, Z. Vilakazi ¹²⁶, O. Villalobos Baillie ¹⁰²,
 A. Vinogradov ¹⁰⁰, L. Vinogradov ¹³¹, Y. Vinogradov ^{99,i}, T. Virgili ³¹, V. Vislavicius ³⁴, Y.P. Viyogi ¹³²,
 A. Vodopyanov ⁶⁶, M.A. Völkl ⁹³, K. Voloshin ⁵⁸, S.A. Voloshin ¹³⁴, G. Volpe ^{135,36}, B. von Haller ³⁶,
 I. Vorobyev ^{37,92}, D. Vranic ^{36,97}, J. Vrláková ⁴¹, B. Vulpescu ⁷⁰, A. Vyushin ⁹⁹, B. Wagner ¹⁸, J. Wagner ⁹⁷,
 H. Wang ⁵⁷, M. Wang ^{7,113}, Y. Wang ⁹³, D. Watanabe ¹²⁸, Y. Watanabe ¹²⁷, M. Weber ³⁶, S.G. Weber ⁹⁷,
 J.P. Wessels ⁵⁴, U. Westerhoff ⁵⁴, J. Wiechula ³⁵, J. Wikne ²², M. Wilde ⁵⁴, G. Wilk ⁷⁷, J. Wilkinson ⁹³,
 M.C.S. Williams ¹⁰⁵, B. Windelband ⁹³, M. Winn ⁹³, C.G. Yaldo ¹³⁴, H. Yang ⁵⁷, P. Yang ⁷, S. Yano ⁴⁷, Z. Yin ⁷,
 H. Yokoyama ¹²⁸, I-K. Yoo ⁹⁶, V. Yurchenko ³, I. Yushmanov ¹⁰⁰, A. Zaborowska ¹³³, V. Zaccolo ⁸⁰,
 A. Zaman ¹⁶, C. Zampolli ¹⁰⁵, H.J.C. Zanolli ¹²⁰, S. Zaporozhets ⁶⁶, N. Zardoshti ¹⁰², A. Zarochentsev ¹³¹,
 P. Závada ⁶⁰, N. Zaviyalov ⁹⁹, H. Zbroszczyk ¹³³, I.S. Zgura ⁶², M. Zhalov ⁸⁵, H. Zhang ^{18,7}, X. Zhang ⁷⁴,
 Y. Zhang ⁷, C. Zhao ²², N. Zhigareva ⁵⁸, D. Zhou ⁷, Y. Zhou ^{80,57}, Z. Zhou ¹⁸, H. Zhu ^{18,7}, J. Zhu ^{7,113},
 X. Zhu ⁷, A. Zichichi ^{28,12}, A. Zimmermann ⁹³, M.B. Zimmermann ^{36,54}, G. Zinovjev ³, M. Zyzak ⁴³

¹ A.I. Alikhanyan National Science Laboratory (Yerevan Physics Institute) Foundation, Yerevan, Armenia² Benemérita Universidad Autónoma de Puebla, Puebla, Mexico³ Bogolyubov Institute for Theoretical Physics, Kiev, Ukraine⁴ Bose Institute, Department of Physics and Centre for Astroparticle Physics and Space Science (CAPSS), Kolkata, India⁵ Budker Institute for Nuclear Physics, Novosibirsk, Russia⁶ California Polytechnic State University, San Luis Obispo, CA, United States⁷ Central China Normal University, Wuhan, China⁸ Centre de Calcul de l'IN2P3, Villeurbanne, France⁹ Centro de Aplicaciones Tecnológicas y Desarrollo Nuclear (CEADEN), Havana, Cuba¹⁰ Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain¹¹ Centro de Investigación y de Estudios Avanzados (CINVESTAV), Mexico City and Mérida, Mexico¹² Centro Fermi – Museo Storico della Fisica e Centro Studi e Ricerche “Enrico Fermi”, Rome, Italy¹³ Chicago State University, Chicago, IL, United States¹⁴ China Institute of Atomic Energy, Beijing, China¹⁵ Commissariat à l’Energie Atomique, IRFU, Saclay, France¹⁶ COMSATS Institute of Information Technology (CIIT), Islamabad, Pakistan¹⁷ Departamento de Física de Partículas and IGFAE, Universidad de Santiago de Compostela, Santiago de Compostela, Spain¹⁸ Department of Physics and Technology, University of Bergen, Bergen, Norway¹⁹ Department of Physics, Aligarh Muslim University, Aligarh, India²⁰ Department of Physics, Ohio State University, Columbus, OH, United States²¹ Department of Physics, Sejong University, Seoul, South Korea²² Department of Physics, University of Oslo, Oslo, Norway²³ Dipartimento di Elettrotecnica ed Elettronica del Politecnico, Bari, Italy²⁴ Dipartimento di Fisica dell’Università ‘La Sapienza’ and Sezione INFN Rome, Italy²⁵ Dipartimento di Fisica dell’Università and Sezione INFN, Cagliari, Italy²⁶ Dipartimento di Fisica dell’Università and Sezione INFN, Trieste, Italy²⁷ Dipartimento di Fisica dell’Università and Sezione INFN, Turin, Italy²⁸ Dipartimento di Fisica e Astronomia dell’Università and Sezione INFN, Bologna, Italy²⁹ Dipartimento di Fisica e Astronomia dell’Università and Sezione INFN, Catania, Italy³⁰ Dipartimento di Fisica e Astronomia dell’Università and Sezione INFN, Padova, Italy³¹ Dipartimento di Fisica ‘E.R. Caianiello’ dell’Università and Gruppo Collegato INFN, Salerno, Italy³² Dipartimento di Scienze e Innovazione Tecnologica dell’Università del Piemonte Orientale and Gruppo Collegato INFN, Alessandria, Italy³³ Dipartimento Interateneo di Fisica ‘M. Merlin’ and Sezione INFN, Bari, Italy³⁴ Division of Experimental High Energy Physics, University of Lund, Lund, Sweden³⁵ Eberhard Karls Universität Tübingen, Tübingen, Germany³⁶ European Organization for Nuclear Research (CERN), Geneva, Switzerland³⁷ Excellence Cluster Universe, Technische Universität München, Munich, Germany

- ³⁸ Faculty of Engineering, Bergen University College, Bergen, Norway
³⁹ Faculty of Mathematics, Physics and Informatics, Comenius University, Bratislava, Slovakia
⁴⁰ Faculty of Nuclear Sciences and Physical Engineering, Czech Technical University in Prague, Prague, Czechia
⁴¹ Faculty of Science, P.J. Šafárik University, Košice, Slovakia
⁴² Faculty of Technology, Buskerud and Vestfold University College, Vestfold, Norway
⁴³ Frankfurt Institute for Advanced Studies, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany
⁴⁴ Gangneung-Wonju National University, Gangneung, South Korea
⁴⁵ Gauhati University, Department of Physics, Guwahati, India
⁴⁶ Helsinki Institute of Physics (HIP), Helsinki, Finland
⁴⁷ Hiroshima University, Hiroshima, Japan
⁴⁸ Indian Institute of Technology Bombay (IIT), Mumbai, India
⁴⁹ Indian Institute of Technology Indore, Indore (IITI), India
⁵⁰ Inha University, Incheon, South Korea
⁵¹ Institut de Physique Nucléaire d'Orsay (IPNO), Université Paris-Sud, CNRS-IN2P3, Orsay, France
⁵² Institut für Informatik, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany
⁵³ Institut für Kernphysik, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany
⁵⁴ Institut für Kernphysik, Westfälische Wilhelms-Universität Münster, Münster, Germany
⁵⁵ Institut Pluridisciplinaire Hubert Curien (IPHC), Université de Strasbourg, CNRS-IN2P3, Strasbourg, France
⁵⁶ Institute for Nuclear Research, Academy of Sciences, Moscow, Russia
⁵⁷ Institute for Subatomic Physics of Utrecht University, Utrecht, Netherlands
⁵⁸ Institute for Theoretical and Experimental Physics, Moscow, Russia
⁵⁹ Institute of Experimental Physics, Slovak Academy of Sciences, Košice, Slovakia
⁶⁰ Institute of Physics, Academy of Sciences of the Czech Republic, Prague, Czechia
⁶¹ Institute of Physics, Bhubaneswar, India
⁶² Institute of Space Science (ISS), Bucharest, Romania
⁶³ Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, Mexico City, Mexico
⁶⁴ Instituto de Física, Universidad Nacional Autónoma de México, Mexico City, Mexico
⁶⁵ iThemba LABS, National Research Foundation, Somerset West, South Africa
⁶⁶ Joint Institute for Nuclear Research (JINR), Dubna, Russia
⁶⁷ Konkuk University, Seoul, South Korea
⁶⁸ Korea Institute of Science and Technology Information, Daejeon, South Korea
⁶⁹ KTO Karatay University, Konya, Turkey
⁷⁰ Laboratoire de Physique Corpusculaire (LPC), Clermont Université, Université Blaise Pascal, CNRS-IN2P3, Clermont-Ferrand, France
⁷¹ Laboratoire de Physique Subatomique et de Cosmologie, Université Grenoble-Alpes, CNRS-IN2P3, Grenoble, France
⁷² Laboratori Nazionali di Frascati, INFN, Frascati, Italy
⁷³ Laboratori Nazionali di Legnaro, INFN, Legnaro, Italy
⁷⁴ Lawrence Berkeley National Laboratory, Berkeley, CA, United States
⁷⁵ Lawrence Livermore National Laboratory, Livermore, CA, United States
⁷⁶ Moscow Engineering Physics Institute, Moscow, Russia
⁷⁷ National Centre for Nuclear Studies, Warsaw, Poland
⁷⁸ National Institute for Physics and Nuclear Engineering, Bucharest, Romania
⁷⁹ National Institute of Science Education and Research, Bhubaneswar, India
⁸⁰ Niels Bohr Institute, University of Copenhagen, Copenhagen, Denmark
⁸¹ Nikhef, Nationaal instituut voor subatomaire fysica, Amsterdam, Netherlands
⁸² Nuclear Physics Group, STFC Daresbury Laboratory, Daresbury, United Kingdom
⁸³ Nuclear Physics Institute, Academy of Sciences of the Czech Republic, Řež u Prahy, Czechia
⁸⁴ Oak Ridge National Laboratory, Oak Ridge, TN, United States
⁸⁵ Petersburg Nuclear Physics Institute, Gatchina, Russia
⁸⁶ Physics Department, Creighton University, Omaha, NE, United States
⁸⁷ Physics Department, Panjab University, Chandigarh, India
⁸⁸ Physics Department, University of Athens, Athens, Greece
⁸⁹ Physics Department, University of Cape Town, Cape Town, South Africa
⁹⁰ Physics Department, University of Jammu, Jammu, India
⁹¹ Physics Department, University of Rajasthan, Jaipur, India
⁹² Physik Department, Technische Universität München, Munich, Germany
⁹³ Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany
⁹⁴ Politecnico di Torino, Turin, Italy
⁹⁵ Purdue University, West Lafayette, IN, United States
⁹⁶ Pusan National University, Pusan, South Korea
⁹⁷ Research Division and ExtReMe Matter Institute EMMI, GSI Helmholtzzentrum für Schwerionenforschung, Darmstadt, Germany
⁹⁸ Rudjer Bošković Institute, Zagreb, Croatia
⁹⁹ Russian Federal Nuclear Center (VNIIEF), Sarov, Russia
¹⁰⁰ Russian Research Centre Kurchatov Institute, Moscow, Russia
¹⁰¹ Saha Institute of Nuclear Physics, Kolkata, India
¹⁰² School of Physics and Astronomy, University of Birmingham, Birmingham, United Kingdom
¹⁰³ Sección Física, Departamento de Ciencias, Pontificia Universidad Católica del Perú, Lima, Peru
¹⁰⁴ Sezione INFN, Bari, Italy
¹⁰⁵ Sezione INFN, Bologna, Italy
¹⁰⁶ Sezione INFN, Cagliari, Italy
¹⁰⁷ Sezione INFN, Catania, Italy
¹⁰⁸ Sezione INFN, Padova, Italy
¹⁰⁹ Sezione INFN, Rome, Italy
¹¹⁰ Sezione INFN, Trieste, Italy
¹¹¹ Sezione INFN, Turin, Italy
¹¹² SSC IHEP of NRC Kurchatov institute, Protvino, Russia
¹¹³ SUBATECH, Ecole des Mines de Nantes, Université de Nantes, CNRS-IN2P3, Nantes, France
¹¹⁴ Suranaree University of Technology, Nakhon Ratchasima, Thailand
¹¹⁵ Technical University of Košice, Košice, Slovakia
¹¹⁶ Technical University of Split FESB, Split, Croatia

- ¹¹⁷ The Henryk Niewodniczanski Institute of Nuclear Physics, Polish Academy of Sciences, Cracow, Poland
¹¹⁸ The University of Texas at Austin, Physics Department, Austin, TX, United States
¹¹⁹ Universidad Autónoma de Sinaloa, Culiacán, Mexico
¹²⁰ Universidade de São Paulo (USP), São Paulo, Brazil
¹²¹ Universidade Estadual de Campinas (UNICAMP), Campinas, Brazil
¹²² University of Houston, Houston, TX, United States
¹²³ University of Jyväskylä, Jyväskylä, Finland
¹²⁴ University of Liverpool, Liverpool, United Kingdom
¹²⁵ University of Tennessee, Knoxville, TN, United States
¹²⁶ University of the Witwatersrand, Johannesburg, South Africa
¹²⁷ University of Tokyo, Tokyo, Japan
¹²⁸ University of Tsukuba, Tsukuba, Japan
¹²⁹ University of Zagreb, Zagreb, Croatia
¹³⁰ Université de Lyon, Université Lyon 1, CNRS/IN2P3, IPN-Lyon, Villeurbanne, France
¹³¹ V. Fock Institute for Physics, St. Petersburg State University, St. Petersburg, Russia
¹³² Variable Energy Cyclotron Centre, Kolkata, India
¹³³ Warsaw University of Technology, Warsaw, Poland
¹³⁴ Wayne State University, Detroit, MI, United States
¹³⁵ Wigner Research Centre for Physics, Hungarian Academy of Sciences, Budapest, Hungary
¹³⁶ Yale University, New Haven, CT, United States
¹³⁷ Yonsei University, Seoul, South Korea
¹³⁸ Zentrum für Technologietransfer und Telekommunikation (ZTT), Fachhochschule Worms, Worms, Germany

ⁱ Deceased.ⁱⁱ Also at: M.V. Lomonosov Moscow State University, D.V. Skobeltsyn Institute of Nuclear Physics, Moscow, Russia.ⁱⁱⁱ Also at: University of Kansas, Lawrence, KS, United States.