

# Predicting inpatient hypoglycaemia in hospitalized patients with diabetes: a retrospective analysis of 9584 admissions with diabetes

Stuart, Kevin; Adderley, Nicola; Marshall, Tom; Rayman, Gerry; Sitch, Alice; Manley, Susan E.; Ghosh, Sandip; Toulis, Konstantinos; Nirantharakumar, Krishnarajah

DOI:

[10.1111/dme.13409](https://doi.org/10.1111/dme.13409)

## Document Version

Peer reviewed version

## Citation for published version (Harvard):

Stuart, K, Adderley, N, Marshall, T, Rayman, G, Sitch, A, Manley, SE, Ghosh, S, Toulis, K & Nirantharakumar, K 2017, 'Predicting inpatient hypoglycaemia in hospitalized patients with diabetes: a retrospective analysis of 9584 admissions with diabetes', *Diabetic Medicine*, vol. 34, no. 10, pp. 1385-1391. <https://doi.org/10.1111/dme.13409>

[Link to publication on Research at Birmingham portal](#)

## Publisher Rights Statement:

Checked for eligibility: 06/06/2017

This is the peer reviewed version of the following article: Stuart et. al., Predicting inpatient hypoglycaemia in hospitalized patients with diabetes: a retrospective analysis of 9584 admissions with diabetes, *Diabetic Medicine*, [2017], which has been published in final form at 10.1111/dme.13409. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.

## General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

**Title: Predicting inpatient hypoglycaemia in hospitalized patients with diabetes: a retrospective analysis of 9,584 admissions with diabetes**

**Running head:** Predicting inpatient hypoglycaemia in hospitalized patients with diabetes

**Authors:** K. Stuart<sup>1,2\*</sup>, N. J. Adderley<sup>1\*</sup>, T. Marshall<sup>1</sup>, G. Rayman<sup>3</sup>, A. Sitch<sup>1</sup>, S. Manley<sup>4</sup>, S. Ghosh<sup>4</sup>, K. A. Toulis<sup>1,5\*</sup>, K. Nirantharakumar<sup>1,4\*</sup>

\*Contributed equally

**Affiliations:**

1. Institute of Applied Health Research, University of Birmingham, UK
2. West Hertfordshire NHS Trust, UK
3. Ipswich Hospital NHS Trust, UK
4. Diabetes Department, University Hospitals Birmingham NHS Foundation Trust, UK
5. 424 General Military Hospital, Thessaloniki, Greece

**Corresponding Authors:**

Dr. Nicola Adderley  
Research Fellow  
Institute of Applied Health Research  
University of Birmingham  
Birmingham  
B15 2TT  
[n.j.adderley@bham.ac.uk](mailto:n.j.adderley@bham.ac.uk)

Dr. Krish Nirantharakumar  
Senior Clinical Lecturer  
Institute of Applied Health Research  
University of Birmingham  
Birmingham  
B15 2TT  
[k.nirantharan@bham.ac.uk](mailto:k.nirantharan@bham.ac.uk)

**Word count:**

Abstract	250
Main Text	1573
References	31
Tables	4 + 3 supplementary
Figures	4

**Funding**

Tom Marshall and Nicola Adderley were partly funded by the National Institute for Health Research (NIHR) through the Collaborations for Leadership in Applied Health Research and Care for West Midlands (CLAHRC-WM). This article presents independent research funded by the National Institute for Health Research (NIHR). The views expressed in this publication are not necessarily those of the NIHR, the Department of Health, NHS Partner Trusts, University of Birmingham or the CLAHRC WM Management Group.

**Conflicts of interest**

None.

**What's new?**

- Hypoglycaemia is associated with worse outcomes in inpatients with diabetes. At present there is no targeted and validated prediction model for identifying patients with an increased risk of hypoglycaemia.
- We performed a retrospective analysis of inpatient admissions with diabetes to develop a prediction model for hypoglycaemia incorporating routinely collected biochemical data.
- We found that the occurrence of hypoglycaemia could be predicted by a model incorporating background medication, ethnicity, age, admission type and laboratory measurements.
- Model performance indicates potential clinical utility to identify patients at risk of hypoglycaemia during their inpatient stay, which could lead to improved patient management and outcomes.

## **Abstract**

### **Aims**

Inpatient hypoglycaemia in patients with diabetes mellitus is associated with excess mortality, increased length of stay and increased complication rate. The objective of this study was to explore whether a quantitative approach to identify hospitalized patients with diabetes at risk of hypoglycaemia could be feasible by incorporating routine biochemical, haematological and prescription data.

### **Methods**

A retrospective cross-sectional analysis of all diabetic admissions (n=9,584) from 1st January 2014 to 31st December 2014 was performed. Hypoglycaemia was defined as a blood glucose level of  $< 4$  mmol/L. The prediction model was constructed using multivariable logistic regression, populated by clinically important variables and routine laboratory data.

### **Results**

Using a pre-specified variable selection strategy, it was shown that the occurrence of inpatient hypoglycaemia could be predicted by a combined model taking into account background medication (type of insulin, use of sulphonylurea), ethnicity (Black and Asian), age (75+ years old), type of admission (emergency) and laboratory measurements (eGFR, CRP, sodium and albumin). ROC curve analysis revealed that the Area Under the Curve (AUC) was 0.733 (95% CI 0.719 to 0.747). The cut-off point chosen to maximize both sensitivity and specificity was 0.15. AUC obtained from internal validation did not differ from the primary model (0.731 (95% CI 0.717 to 0.746)).

## **Conclusions**

The inclusion of routine biochemical data, available at the time of admission, can add prognostic value to demographic and medication history. The predictive performance of the constructed model indicates potential clinical utility to identify patients at risk of hypoglycaemia during their inpatient stay.

## **Introduction**

Hospitalised patients with diabetes have high infection rates [1-4], longer length of stay [5-7] and increased mortality (10% higher) [8]. Hypoglycaemia is one of the important determining events for this worse prognosis, which can be reflected in excess mortality, increased length of stay and increased complications amongst patients with diabetes [9-11]. Hence, predicting the risk of hypoglycaemia in hospitalized patients with diabetes mellitus and preventing its occurrence in a selected, high-risk subset of them through active monitoring or therapeutic modifications may be an efficient way to improve outcomes.

However, a targeted and validated prediction model for identifying inpatients with diabetes at an increased risk of hypoglycaemia is still missing. The main reason is the heterogeneity in populations, thresholds for hypoglycaemia, underlying diseases and co-morbidities across relevant studies [11-19]. This is why the Joint British Diabetes Societies' (JBDS) position currently advocates a non-quantitative approach with regard to risk of inpatient hypoglycaemia in patients with diabetes [20]. Specifically, the recommended approach would be to consider medical or lifestyle risk factors, such as strict glucose management, history of severe hypoglycemic events, duration of insulin use, severe liver failure, renal failure, terminal illness or increasing age and alcohol use respectively.

To explore whether a quantitative approach to identify hospitalized patients with diabetes at risk of hypoglycaemia could be feasible and construct a relevant prediction model incorporating routine biochemical data collected at admission, we performed a retrospective analysis of admissions with diabetes.

## **Methods**

All hospitalized adult (>16 years old) patients with diabetes mellitus in the general ward of our Institution were considered as potentially eligible, irrespective of the primary diagnosis. The observation period was from 1st January 2014 to 31st December 2014. The diagnosis of diabetes mellitus was consolidated by the presence of relevant Patient Administration System (PAS) discharge codes for diabetes or the presence of a continuous anti-hyperglycemic medication in the electronic medical record system (Patient Information and Communication System (PICS)). The diagnosis of diabetes was also ensured by cross-checking the PAS database for the International Classification of Diseases version 10 (ICD-10) diabetes codes (E10 – E14) retrospectively for the previous 10 years, up to 2004. Prescription of metformin or short acting insulin in the absence of a diabetes code was considered an exclusion criterion to ensure case (diabetes mellitus) definition. Patients admitted to ITU were excluded from the analysis.

The outcome of interest was the occurrence of hypoglycaemia, which was defined as a blood glucose level of < 4 mmol/L (point of care or laboratory blood glucose measurement).

### *Statistical Analysis*

Baseline differences in continuous data between hypoglycaemic and non hypoglycaemic groups were explored using the Student's T-test or Mann-Whitney U test according to distribution. Differences in dichotomous data were explored using a Chi-squared test.

The prediction model was constructed on the basis of a multivariable logistic regression model with hypoglycaemia as the dependent outcome. This model was populated by clinically important variables, as identified by previous literature, including laboratory results as recorded in PICS. All variables with a  $p$ -value at significance level of 20% ( $p = 0.20$ ) in the univariate analysis were included in the multivariable logistic regression model (Supplementary Table 1). In cases where two variables were statistically significant but highly correlated, one was chosen based on clinical familiarity. Continuous data (for example age) were also analyzed as categories to inspect potential non-linear effects. Area Under the Curve analysis was performed to assess the predictive performance of the final model. A decision threshold was selected with the intent of maximizing both sensitivity and specificity. Positive and negative predictive values were calculated using the prevalence of hypoglycaemia in this dataset.

Internal validation was assessed using bootstrapping, in which sample datasets the same size as the original dataset are repeatedly resampled, with replacement, and model parameters are re-estimated and averaged over the samples [21]; this identifies any shrinkage necessary to reduce model over-optimism due to over-fitting. Missing data were handled by a chained multiple imputation technique with predictive mean matching. Bootstrapped results from each of the imputed datasets were combined using Rubin's rule. In sensitivity analysis, missing values were replaced by values in

the normal range for the given clinical pathology test. Model calibration was assessed by plotting predicted probability of hypoglycaemia against observed probability. Significance level was set at 5% and all analyses were implemented in Stata 13.0.

## **Results**

### *Flow chart*

From a total of 106,580 patients admitted to our Institution during 2014, 57,922 admissions were eligible (emergency or elective admissions) and of those, 9,584 admissions were in patients with diabetes. Hypoglycaemia occurred in 1,327 (13.8%) admissions with diabetes while the remaining 8,257 (86.2%) did not have a hypoglycaemic event. A flow chart summarizing the selection process along with reasons for exclusion is presented in Figure 1.

6,187 unique patients made up the 9,584 admissions included in the analysis. Of the 1,792 patients who contributed two or more admissions, 664 experienced a hypoglycaemic episode; none of these 664 patients had a hypoglycaemic episode in more than one admission.

### *Baseline characteristics*

A table summarizing differences in the baseline demographics and medication history is presented in Table 1 and 2. Patients in the hypoglycaemia group were older, more deprived, with higher comorbidities and more likely to belong to an ethnic minority group compared to the group without hypoglycaemia. Insulin use and sulphonylureas were also proportionally more common in the hypoglycaemia group (Table 2). A table summarizing differences in the baseline laboratory measurements is presented in Table 3. Patients in the hypoglycaemia group were more likely to have elevated



inflammation markers (C-reactive protein, CRP levels), electrolyte disturbances or have anaemia.

#### *Final model*

Using a pre-specified variable selection strategy, younger age categories, female gender, hyperkalaemia, an elevated neutrophil count and anaemia were not found to be significant predictors of hypoglycaemia in the multivariable logistic regression. Elderly greater than 75 years, Insulin and sulphonylureas therapy, Black and Asian ethnicity, emergency admissions, lower eGFR, higher CRP, hyponatraemia (<125mmol/L) and hypo-albuminaemia were the strongest predictors of hypoglycaemia in patients with diabetes mellitus admitted in the general ward. The final model (prior to bootstrapping) is presented in the Table 4.

In sensitivity analysis, we used multilevel modelling (xtlogit) to allow for any possible correlation between multiple admissions for an individual patient; this made very little difference to the model parameters.

#### *Model performance and internal validity*

ROC curve analysis revealed that the Area Under the Curve (AUC) was 0.733 (95% CI 0.719 to 0.747) (Figure 2). The cut-off point chosen to maximize both sensitivity and specificity was 0.15 predicted probability of hypoglycaemia, with these corresponding values being 59.3% and 73.7% respectively. At this cut-off the positive predictive value was 26.6% and the negative predictive value was 91.8%. A confusion matrix describing model performance is shown in Figure 3. The model was internally validated by bootstrapping; model coefficients were unchanged and test statistics were similar to the primary model, indicating no evidence of over-fitting: AUC 0.731 (95% CI 0.717 to 0.746), sensitivity 59.3%, specificity 73.4%, positive predictive value 26.4%, and negative predictive value 91.8%. Visual assessment of the

calibration plot (Figure 4) suggests that model calibration is good (i.e. predicted probabilities of hypoglycaemia are similar to observed probabilities).

In sensitivity analysis, missing clinical pathology test results were replaced with values in the normal range (Supplementary Table 2). This had little effect on the model performance: AUC 0.735 (95% CI 0.721 to 0.749), sensitivity 60.4%, specificity 73.4%, positive predictive value 26.8%, and negative predictive value 92.0%.

## **Discussion**

In this retrospective analysis of all diabetic admissions, we found that the occurrence of inpatient hypoglycaemia could be predicted by a combined model taking into account background medication (type of insulin, use of sulphonylurea use), ethnicity (Black and Asian), age (75+ years old), type of admissions (emergency) and laboratory measurements (eGFR, CRP, sodium and albumin).

Considering that inpatient hypoglycaemia is significantly associated with inpatient mortality and increased length of hospital stay [10, 11] and that real-time generated alerts using a validated computerized predictive algorithm may significantly reduce its occurrence [16], the findings of the present study could provide a robustly validated basis to improve inpatient safety and outcomes. Importantly, this study also showed that the inclusion of routine biochemical data (such as CRP, albumin, eGFR and sodium), available at the time of admission, could add prognostic value to demographic and medication history, thus providing a more holistic and optimized approach in the prediction of hypoglycemic events.

On the other hand, the findings of the present study should be interpreted in the context of its limitations. First external validity, that is assessing the model in a

different environment on a different population, would be a prerequisite before recommending its adoption in clinical practice. Moreover, the extent of missing data with regard to glycated hemoglobin (HbA1c) prevented the plausible exploration of tight versus poor glycemic control as a predictor of hypoglycaemia. The study used routinely collected observational data, and therefore there may be unobserved confounding for which it was not possible make adjustments in the analysis. However, this does not vitiate the utility of the model or the observed association between the predictor variables and the outcome.

Taking the above into consideration and using the largest dataset of diabetic admissions to date, we suggest that the occurrence of inpatient hypoglycaemia may be predicted by a combined model using background medication ethnicity, age, type of admission and routine laboratory measurements as independent predictors. The predictive performance of the constructed model indicates potential clinical utility in identifying patients at high risk of inpatient hypoglycaemia.

## **Funding**

Tom Marshall and Nicola Adderley were partly funded by the National Institute for Health Research (NIHR) through the Collaborations for Leadership in Applied Health Research and Care for West Midlands (CLAHRC-WM). This article presents independent research funded by the National Institute for Health Research (NIHR). The views expressed in this publication are not necessarily those of the NIHR, the Department of Health, NHS Partner Trusts, University of Birmingham or the CLAHRC WM Management Group.

**Conflicts of interest**

None.

**Author contributions**

KS, NA and KN developed the model. AS advised on the statistical analysis. KT, TM, GR, SG, SM and KN provided clinical input. All authors wrote the paper or revised it critically for important intellectual content.

## References

1. Laird AM, Miller PR, Kilgo PD, Meredith JW, Chang MC. Relationship of early hyperglycemia to mortality in trauma patients. *Journal of Trauma-Injury Infection & Critical Care*. 2004;56(5):1058-62.
2. Marchant MH, Jr., Viens NA, Cook C, Vail TP, Bolognesi MP. The impact of glycemic control and diabetes mellitus on perioperative outcomes after total joint arthroplasty. *Journal of Bone & Joint Surgery - American Volume*. 2009;91(7):1621-9.
3. Pomposelli JJ, Baxter JK, III, Babineau TJ, Pomfret EA, Driscoll DF, Forse RA, et al. Early postoperative glucose control predicts nosocomial infection rate in diabetic patients. *Jpen: Journal of Parenteral & Enteral Nutrition*. 1998;22(2):77-81.
4. Matros E, Aranki SF, Bayer LR, McGurk S, Neuwalder J, Orgill DP. Reduction in incidence of deep sternal wound infections: Random or real? *The Journal of Thoracic and Cardiovascular Surgery*. 2010;139(3):680-5.
5. Sampson MJ, Dozio N, Ferguson B, Dhatariya K. Total and excess bed occupancy by age, specialty and insulin use for nearly one million diabetes patients discharged from all English Acute Hospitals. *Diabetes ResClinPract*. 2007;77(1):92-8.
6. Health and Social Care Information Centre. National diabetes inpatient audit 2013 2014 [updated 01/01/2015; cited 2015 01/01/2015]. Available from: [http://www.diabetes.nhs.uk/information\\_and\\_data/diabetes\\_audits/national\\_diabetes\\_inpatient\\_audit\\_2011\\_and\\_2012/](http://www.diabetes.nhs.uk/information_and_data/diabetes_audits/national_diabetes_inpatient_audit_2011_and_2012/).
7. Health and Social Care Information Centre. National diabetes inpatient audit 2011 2012 [updated 8/10/2012; cited 2012 8/21/2012]. Available from: [http://www.diabetes.nhs.uk/information\\_and\\_data/diabetes\\_audits/national\\_diabetes\\_inpatient\\_audit\\_2011\\_and\\_2012/](http://www.diabetes.nhs.uk/information_and_data/diabetes_audits/national_diabetes_inpatient_audit_2011_and_2012/).
8. YHPHO. Mortality among inpatients with diabetes Yorkshire 2012 [updated 2012; cited 2013 2/1/2013]. Available from: <http://www.yhpho.org.uk/default.aspx?RID=139172>.
9. Khunti K, Davies M, Majeed A, Thorsted BL, Wolden ML, Paul SK. Hypoglycemia and risk of cardiovascular disease and all-cause mortality in insulin-treated people with type 1 and type 2 diabetes: a cohort study. *Diabetes care*. 2015;38(2):316-22.
10. Nirantharakumar K, Marshall T, Kennedy A, Narendran P, Hemming K, Coleman JJ. Hypoglycaemia is associated with increased length of stay and mortality

in people with diabetes who are hospitalized. *Diabetic medicine : a journal of the British Diabetic Association*. 2012;29(12):e445-8.

11. Turchin A, Matheny ME, Shubina M, Scanlon JV, Greenwood B, Pendergrass ML. Hypoglycemia and clinical outcomes in patients with diabetes hospitalized in the general ward. *Diabetes Care*. 2009;32(7):1153-7.
12. Arinzon Z, Fidelman Z, Berner YN, Adunsky A. Infection-related hypoglycemia in institutionalized demented patients: a comparative study of diabetic and nondiabetic patients. *Archives of gerontology and geriatrics*. 2007;45(2):191-200.
13. Elliott MB, Schafers SJ, McGill JB, Tobin GS. Prediction and prevention of treatment-related inpatient hypoglycemia. *Journal of diabetes science and technology*. 2012;6(2):302-9.
14. Farrokhi F, Klindukhova O, Chandra P, Peng L, Smiley D, Newton C, et al. Risk factors for inpatient hypoglycemia during subcutaneous insulin therapy in non-critically ill patients with type 2 diabetes. *Journal of diabetes science and technology*. 2012;6(5):1022-9.
15. Kagansky N, Levy S, Rimón E, Cojocaru L, Fridman A, Ozer Z, et al. Hypoglycemia as a predictor of mortality in hospitalized elderly patients. *Archives of internal medicine*. 2003;163(15):1825-9.
16. Kilpatrick CR, Elliott MB, Pratt E, Schafers SJ, Blackburn MC, Heard K, et al. Prevention of inpatient hypoglycemia with a real-time informatics alert. *Journal of hospital medicine*. 2014;9(10):621-6.
17. Krinsley JS, Grover A. Severe hypoglycemia in critically ill patients: risk factors and outcomes. *Critical care medicine*. 2007;35(10):2262-7.
18. Lipska KJ, Warton EM, Huang ES, Moffet HH, Inzucchi SE, Krumholz HM, et al. HbA1c and risk of severe hypoglycemia in type 2 diabetes: the Diabetes and Aging Study. *Diabetes care*. 2013;36(11):3535-42.
19. Pfortmueller CA, Wiemann C, Funk GC, Leichtle AB, Fiedler GM, Exadaktylos AK, et al. Hypoglycemia is associated with increased mortality in patients with acute decompensated liver cirrhosis. *Journal of critical care*. 2014;29(2):316 e7-12.
20. Joint British Diabetes Societies Inpatient Care Group. The hospital management of Hypoglycaemia in adults with Diabetes Mellitus. 2013.

21. Steyerberg EW. Selection of main effects. In EW Steyerberg (ed.), *Clinical Prediction Models: A Practical Approach to Development, Validation, and Updating*. Netherlands 2010.

**Table 1. Baseline demographics**

<b>Patient Demographics</b>	<b>Normoglycaemia (&gt;3.9 mmol/L; N= 8,257)</b>	<b>Hypoglycaemia (≤3.9 mmol/L; N= 1,327)</b>	<b>P-value</b>
Age median (IQR)	68 (56-78)	70 (57-80)	0.032
Age categories N (%)			< 0.001
16-44	723 (8.8)	128 (9.7)	
45-54	1,118 (13.5)	154 (11.6)	
55-64	1,574 (19.1)	218 (16.4)	
65-74	2,095 (25.4)	294 (22.2)	
75-84	1,858 (22.5)	361 (27.2)	
≥ 85	889 (10.8)	172 (13.0)	
Sex N (%) <sup>0</sup>			0.026
Male	4,566 (55.4)	691 (52.1)	
Female	3,682 (44.6)	636 (47.9)	
Ethnicity N (%) <sup>0</sup>			< 0.001
Caucasian	5,657 (68.5)	855 (64.4)	
Black	455 (5.5)	117 (8.8)	
Asian	1,765 (21.4)	298 (22.5)	
Other	380 (4.6)	57 (4.3)	
IMD deprivation quintile N (%) <sup>0</sup>			0.025
Least deprived 1	366 (4.5)	47 (3.6)	
2	531 (6.5)	85 (6.5)	
3	1,657 (20.4)	239 (18.3)	
4	1,675 (20.6)	246 (18.8)	
Most deprived 5	3,915 (48.1)	689 (52.8)	
Type of Admission N (%)			< 0.001
Elective	1,882 (22.8)	144 (10.9)	
Emergency	6,375 (77.2)	1,183 (89.2)	
Modified <sup>†</sup> Charlson Comorbidity score N (%)			< 0.001
0	3,807 (46.1)	429 (32.3)	
1	1,524 (18.5)	214 (16.1)	
2 or more	2,926 (35.4)	684 (51.5)	

IMD: Index of Multiple Deprivation. Missing Data: <sup>0</sup> <1.5%, otherwise nil.

<sup>†</sup>Charlson score with diabetes scores removed.



**Table 2: Medication history**

<b>Patient Medications</b>	<b>Normoglycaemia (&gt;3.9 mmol/L; N= 8,257)</b>	<b>Hypoglycaemia (&lt;3.9 mmol/L; N= 1,327)</b>	<b>P-value</b>
Insulin N (%)			
Short Acting	887 (10.7)	373 (28.1)	< 0.001
Intermediate Acting	184 (2.2)	88 (6.6)	< 0.001
Long Acting	1,085 (13.1)	381 (28.7)	< 0.001
Sulphonylureas N (%)	1,370 (16.6)	334 (25.2)	< 0.001
Metformin N (%)	3,145 (38.1)	501 (37.8)	0.816
Thiazolidinediones N (%)	139 (1.7)	29 (2.2)	0.196
Incretin mimetics N (%)	107 (1.3)	13 (1.0)	0.336
DPP-4 inhibitors N (%)	691 (8.4)	124 (9.3)	0.237
Other antidiabetic medications <sup>†</sup> N (%)	36 (0.4)	14 (1.1)	0.004

Reg.: Regulator. <sup>†</sup> Alpha-glucosidase, prandial glucose regulator, SGLT2

inhibitors. Common combinations of medications are shown in Supplementary Table 3.

**Table 3. Baseline laboratory characteristics considered for the prediction model**


Patient Laboratory Results	Normoglycaemia (>3.9 mmol/L; N= 8,257)	Hypoglycaemia (≤3.9 mmol/L; N= 1,327)	P-value
<b>Electrolytes and renal function</b>			
Sodium mean (SD) <sup>◊</sup>	137.0 (4.8)	136.3 (5.7)	< 0.001
Sodium N (%) <sup>◊</sup>			< 0.001
< 125 mmol/L	110 (1.6)	43 (3.4)	
125-134 mmol/L	1,611 (23.1)	377 (29.9)	
135-144 mmol/L	5,076 (72.9)	796 (63.2)	
145-154 mmol/L	160 (2.3)	37 (2.9)	
≥ 155 mmol/L	8 (0.1)	7 (0.6)	
Potassium mean (SD) <sup>◊</sup>	4.5 (0.6)	4.6 (0.8)	< 0.001
Potassium N (%) <sup>◊</sup>			< 0.001
< 3 mmol/L	28 (0.4)	9 (0.8)	
3-6 mmol/L	6,139 (97.2)	1,046 (94.2)	
≥ 6 mmol/L	150 (2.4)	56 (5.0)	
Creatinine median (IQR) <sup>◊</sup>	88 (68, 127)	105 (75, 173)	< 0.001
eGFR median (IQR) <sup>◊</sup>	66 (43, 88)	51.5 (30, 78)	< 0.001
eGFR N (%) <sup>◊</sup>			< 0.001
≥ 90	1,686 (24.2)	230 (18.3)	
60-89	2,352 (33.8)	303 (24.1)	
30-59	1,860 (26.7)	411 (32.7)	
15-29	533 (7.7)	168 (13.4)	
< 15	524 (7.5)	146 (11.6)	
<b>Infection / Inflammation marker</b>			
CRP median (IQR)*	13 (4, 47)	24 (6, 77)	< 0.001
CRP N (%)*			< 0.001
< 10 mg/L	2,622 (44.1)	413 (34.3)	
10-49 mg/L	1,902 (32.0)	377 (31.3)	
50-99 mg/L	689 (11.6)	175 (14.5)	
≥ 100 mg/L	728 (12.3)	240 (19.9)	
<b>Haematology Tests</b>			
Haemoglobin mean (SD) <sup>◊</sup>	120.8 (21.4)	115.5 (21.4)	< 0.001
Haemoglobin <sup>◊</sup>			< 0.001
< 80 g/L	220 (3.2)	60 (4.8)	
80-109 g/L	1,732 (25.2)	413 (33.0)	
≥ 110 g/L	4,911 (71.6)	780 (62.3)	
Neutrophil median (IQR) <sup>◊</sup>	6.3 (4.5, 8.9)	7.0 (4.8, 10.3)	< 0.001
Neutrophil N (%) <sup>◊</sup>			< 0.001
< 2	157 (2.3)	32 (2.5)	
2-7.9	4,476 (65.0)	718 (57.1)	
≥ 8	2,261 (32.8)	508 (40.3)	
<b>Liver function test</b>			
Albumin mean (SD) <sup>◊</sup>	40.1 (5.4)	37.9 (6.3)	< 0.001
Albumin N (%) <sup>◊</sup>			< 0.001
< 25	77 (1.8)	34 (2.8)	
25-35	862 (13.1)	305 (24.7)	
≥ 35	5,654 (85.8)	898 (72.6)	
Bilirubin median (IQR) <sup>◊</sup>	7 (5, 11)	7 (5, 12)	0.742

Missing Data: <sup>◊</sup> 5-20%, \* 21-30%


**Table 4. Results of the multivariable logistic regression analysis**

Variables (n, % observations with imputed data)	Regression Coefficient (95% CI)	OR (95% CI)	P-value
Age categories			
16-44	0.19 (-0.08 to 0.45)	1.20 (0.92 to 1.58)	0.177
*45-54	-	1	-
55-64	0.03 (-0.20 to 0.26)	1.03 (0.82 to 1.30)	0.786
65-74	0.03 (-0.20 to 0.25)	1.03 (0.82 to 1.28)	0.823
≥ 75	0.25 (0.04 to 0.46)	1.28 (1.04 to 1.59)	0.022
Sex			
*Male	-	1	-
Female	0.06 (-0.07 to 0.18)	1.06 (0.94 to 1.20)	0.362
Ethnicity			
*Caucasian	-	1	-
Black	0.53 (0.30 to 0.77)	1.71 (1.35 to 2.16)	< 0.001
Asian	0.17 (0.02 to 0.33)	1.19 (1.02 to 1.39)	0.027
Other	0.20 (-0.10 to 0.50)	1.22 (0.90 to 1.66)	0.198
Admission Type			
*Elective	-	1	-
Emergency	0.70 (0.51 to 0.89)	2.01 (1.66 to 2.43)	< 0.001
Insulin			
Short Acting	0.86 (0.71 to 1.02)	2.37 (2.03 to 2.78)	< 0.001
Intermediate Acting	0.71 (0.42 to 0.99)	2.03 (1.53 to 2.69)	< 0.001
Long Acting	0.70 (0.55 to 0.85)	2.02 (1.73 to 2.35)	< 0.001
Sulphonylureas	0.57 (0.42 to 0.72)	1.77 (1.53 to 2.05)	< 0.001
Sodium (1,350, 14.1%)			
<125 mmol/L	0.55 (0.16 to 0.93)	1.73 (1.17 to 2.54)	0.006
125-134 mmol/L	0.08 (-0.07 to 0.23)	1.08 (0.93 to 1.25)	0.291
*135-144 mmol/L	-	1	-
145-154 mmol/L	0.09 (-0.30 to 0.48)	1.09 (0.74 to 1.61)	0.668
≥ 155 mmol/L	1.09 (-0.01 to 2.19)	2.97 (0.99 to 8.91)	0.052
Potassium (2,147, 22.4%)			
< 3 mmol/L	0.34 (-0.47 to 1.14)	1.40 (0.63 to 3.13)	0.413
*3 – 6 mmol/L	-	1	-
≥ 6 mmol/L	0.08 (-0.28 to 0.43)	1.08 (0.76 to 1.54)	0.672
eGFR (1,362, 14.2%)			
*≥ 90	-	1	-
60-89	-0.02 (-0.22 to 0.17)	0.98 (0.80 to 1.19)	0.815
30-59	0.34 (0.14 to 0.54)	1.40 (1.15 to 1.71)	0.001
15-29	0.54 (0.28 to 0.79)	1.71 (1.32 to 2.20)	< 0.001
< 15	0.41 (0.14 to 0.67)	1.50 (1.16 to 1.95)	0.002
CRP (2,429, 25.3%)			
*0 - 10	-	1	-
10 - 49	0.10 (-0.06 to 0.26)	1.10 (0.94 to 1.30)	0.230
50 - 99	0.27 (0.05 to 0.49)	1.31 (1.06 to 1.63)	0.014
≥ 100	0.37 (0.15 to 0.58)	1.44 (1.16 to 1.78)	0.001
Albumin (1,745, 18.2%)			
< 25	0.61 (0.17 to 1.06)	1.85 (1.19 to 2.88)	0.007
25-35	0.49 (0.31 to 0.66)	1.63 (1.37 to 1.94)	< 0.001
*≥ 35	-	1	-
Neutrophil count (1,423, 14.8%)			
<2	0.11 (-0.29 to 0.52)	1.12 (0.75 to 1.66)	0.584
*2-8	-	1	-
≥ 8	0.06 (-0.08 to 0.20)	1.06 (0.92 to 1.23)	0.405
Haemoglobin (1,459, 15.2%)			
< 80	-0.03 (-0.35 to 0.30)	0.98 (0.70 to 1.35)	0.880
80-110	0.02 (-0.12 to 0.17)	1.02 (0.88 to 1.19)	0.742
*≥ 110	-	1	-
Constant	-3.57 (-3.86 to -3.27)		< 0.001

**Figure 1. Flow diagram of included patients with diabetes**




**Figure 2. ROC curve for fitted model**



**Figure 3. Confusion matrix showing actual and predicted numbers of hypoglycaemic outcomes**

		Predicted		
		Yes	No	
Actual	Yes	787	540	1,327
	No	2,182	6,075	8,257
		2,969	6,615	

**Figure 4. Calibration plot showing predicted and observed probabilities of hypoglycaemia**



**Supplementary Table 1.** Findings of the univariate analyses.

<b>Variables</b>	<b>Odds Ratio</b>	<b>95% CI</b>	<b>P-value</b>
<b>Age categories</b>			
16-44	1.29	1.00 to 1.65	0.051
*45-54	1	-	-
55-64	1.01	0.81 to 1.25	0.961
65-74	1.02	0.83 to 1.25	0.861
≥ 75	1.41	1.16 to 1.71	< 0.001
<b>Gender</b>			
*Male	1	-	-
Female	1.14	1.02 to 1.28	0.026
<b>Ethnicity</b>			
*Caucasian	1	-	-
Black	1.70	1.37 to 2.11	< 0.001
Asian	1.12	0.97 to 1.29	0.127
Other	0.99	0.74 to 1.32	0.959
<b>IMD deprivation quintile</b>			
*Least deprived 1	1	-	-
2	1.25	0.85 to 1.82	0.256
3	1.12	0.81 to 1.57	0.494
4	1.14	0.82 to 1.59	0.428
Most deprived 5	1.37	1.00 to 1.88	0.049
<b>Type of Admission</b>			
*Elective			
Emergency	2.43	2.02 to 2.90	<0.001
<b>Modified<sup>†</sup> Charlson</b>			
*0	1	-	-
1	1.25	1.05 to 1.48	0.013
≥ 2	2.07	1.82 to 2.36	< 0.001

IMD: Index of Multiple Deprivation. <sup>†</sup>Modified Charlson: Charlson score minus Diabetes.

<b>Variables</b>	<b>Odds Ratio</b>	<b>95% CI</b>	<b>P-value</b>
<b>Insulin</b>			
Short Acting	3.25	2.83 to 3.73	< 0.001
Intermediate Acting	3.12	2.40 to 4.05	< 0.001
Long Acting	2.66	2.33 to 3.05	< 0.001
Any	3.36	2.98 to 3.79	< 0.001
Sulphonylureas	1.69	1.47 to 1.94	< 0.001
Metformin	0.99	0.87 to 1.11	0.816
Thiazolidinediones	1.30	0.87 to 1.96	0.197
Incretin mimetics	0.75	0.42 to 1.34	0.338
DPP-4 inhibitors	1.13	0.92 to 1.38	0.237

<b>Variables</b>	<b>Odds Ratio</b>	<b>95% CI</b>	<b>P-value</b>
<b>Sodium</b>			
<125 mmol/L	2.49	1.73 to 3.58	< 0.001
125-134 mmol/L	1.53	1.34 to 1.75	< 0.001
*135-144 mmol/L	1	-	-
145-154 mmol/L	1.46	1.01 to 2.11	0.044
≥155 mmol/L	5.78	2.06 to 16.24	0.001
<b>Potassium</b>			
<3 mmol/L	1.78	0.85 to 3.75	0.129
*3-6 mmol/L	1	-	-
≥6 mmol/L	2.26	1.66 to 3.06	< 0.001
<b>eGFR</b>			
*≥ 90	1	-	-
60-89	0.93	0.77 to 1.12	0.424
30-59	1.62	1.35 to 1.93	< 0.001
15-29	2.34	1.86 to 2.94	< 0.001
< 15	2.12	1.69 to 2.67	< 0.001
<b>CRP</b>			
*0-10	1	-	-
10-49	1.32	1.13 to 1.53	< 0.001
50-99	1.74	1.43 to 2.11	< 0.001
≥ 100	2.28	1.91 to 2.72	< 0.001
<b>Albumin</b>			
< 25	2.79	1.84 to 4.23	< 0.001
25-35	2.25	1.93 to 2.61	<0.001
*≥ 35	1	-	-
<b>Neutrophil count</b>			
< 2	1.21	0.83 to 1.77	0.330
*2-8	1	-	-
≥ 8	1.44	1.27 to 1.63	< 0.001
<b>Haemoglobin</b>			
< 80	1.76	1.31 to 2.37	< 0.001
80-109	1.49	1.31 to 1.70	< 0.001
*≥ 110	1	-	-

\*Comparator.



**Supplementary Table 2.** Results of analysis replacing missing clinical pathology test results with values in the normal range.

Variables		Regression Coefficient (95% CI)	P-value	Boostrapped Regression Coefficient (95% CI)	P-value
Age categories					
	16-44	0.20 (-0.07 to 0.47)	0.139	0.20 (-0.07 to 0.48)	0.144
	*45-54	-	-	-	-
	55-64	0.02 (-0.21 to 0.26)	0.836	0.02 (-0.21 to 0.26)	0.836
	65-74	0.01 (-0.21 to 0.23)	0.934	0.01 (-0.22 to 0.23)	0.935
	≥ 75	0.24 (0.02 to 0.45)	0.029	0.24 (0.02 to 0.45)	0.033
Sex					
	*Male	-	-	-	-
	Female	0.05 (-0.07 to 0.18)	0.395	0.05 (-0.07 to 0.18)	0.399
Ethnicity					
	* Caucasian	-	-	-	-
	Black	0.56 (0.33 to 0.80)	< 0.001	0.56 (0.32 to 0.80)	< 0.001
	Asian	0.18 (0.02 to 0.33)	0.025	0.18 (0.02 to 0.34)	0.029
	Other	0.23 (-0.07 to 0.54)	0.131	0.23 (-0.07 to 0.54)	0.132
Admission Type					
	*Elective	-	-	-	-
	Emergency	0.50 (0.30 to 0.70)	< 0.001	0.50 (0.30 to 0.70)	< 0.001
Insulin					
	Short Acting	0.82 (0.66 to 0.98)	< 0.001	0.82 (0.65 to 0.99)	< 0.001
	Intermediate Acting	0.68 (0.40 to 0.97)	< 0.001	0.68 (0.38 to 0.98)	< 0.001
	Long Acting	0.69 (0.54 to 0.84)	< 0.001	0.69 (0.53 to 0.85)	< 0.001
	Sulphonylureas	0.54 (0.40 to 0.69)	< 0.001	0.54 (0.39 to 0.70)	< 0.001
Sodium					
	<125 mmol/L	0.57 (0.18 to 0.96)	0.004	0.57 (0.17 to 0.97)	0.005
	125-134 mmol/L	0.06 (-0.08 to 0.21)	0.413	0.06 (-0.09 to 0.22)	0.440
	*135-144 mmol/L	-	-	-	-
	145-154 mmol/L	0.08 (-0.31 to 0.48)	0.672	0.08 (-0.35 to 0.51)	0.700
	≥ 155 mmol/L	1.05 (-0.05 to 2.14)	0.061	1.05 (-0.21 to 2.30)	0.102
Potassium					
	< 3 mmol/L	0.42 (-0.39 to 1.23)	0.311	0.42 (-0.37 to 1.22)	0.300
	*3 – 6 mmol/L	-	-	-	-
	≥ 6 mmol/L	0.02 (-0.33 to 0.36)	0.931	0.02 (-0.35 to 0.38)	0.934
eGFR					
	*≥ 90	-	-	-	-
	60-89	0.03 (-0.15 to 0.21)	0.727	0.03 (-0.15 to 0.21)	0.729
	30-59	0.39 (0.21 to 0.57)	< 0.001	0.39 (0.20 to 0.57)	< 0.001
	15-29	0.57 (0.33 to 0.81)	< 0.001	0.57 (0.33 to 0.81)	< 0.001
	< 15	0.40 (0.15 to 0.65)	0.002	0.40 (0.14 to 0.66)	0.003
CRP					
	*0 - 10	-	-	-	-
	10 - 49	0.23 (0.08 to 0.39)	0.003	0.23 (0.07 to 0.39)	0.004
	50 - 99	0.41 (0.20 to 0.62)	< 0.001	0.41 (0.19 to 0.62)	< 0.001
	≥ 100	0.49 (0.28 to 0.70)	< 0.001	0.49 (0.27 to 0.71)	< 0.001
Albumin					
	< 25	0.62 (0.18 to 1.07)	0.006	0.62 (0.16 to 1.09)	0.009
	25-35	0.48 (0.31 to 0.65)	< 0.001	0.48 (0.30 to 0.66)	< 0.001
	*≥ 35	-	-	-	-
Neutrophil count					
	<2	0.15 (-0.26 to 0.56)	0.475	0.15 (-0.25 to 0.55)	0.467
	*2-8	-	-	-	-
	≥ 8	0.06 (-0.08 to 0.20)	0.379	0.06 (-0.08 to 0.21)	0.383
Haemoglobin					
	< 80	-0.01 (-0.34 to 0.31)	0.935	-0.01 (-0.34 to 0.31)	0.936
	80-110	0.03 (-0.12 to 0.18)	0.663	0.03 (-0.12 to 0.19)	0.676
	*≥ 110	-	-	-	-
Constant		-3.43 (-3.69 to -3.16)	< 0.001	-3.43 (-3.70 to -3.16)	< 0.001

**Supplementary Table 3.** Common drug combinations.

<b>Drug combination</b>		<b>n % of study population (n = 9,584)</b>	
2 drugs	Sulphonylureas + metformin	1,012	10.6
	Insulin + metformin	743	7.8
	Metformin + DPP-4 inhibitors	458	4.8
	Insulin + sulphonylureas	370	3.9
	Sulphonylureas + DPP-4 inhibitors	351	3.7
	Insulin + DPP-4 inhibitors	233	2.4
3 drugs	Sulphonylureas + metformin + DPP-4	223	2.3
	Sulphonylureas + insulin + metformin	201	2.1