

# The onset of fabric development in deep marine sediments

Maffione, Marco; Morris, Antony

DOI:

[10.1016/j.epsl.2017.06.018](https://doi.org/10.1016/j.epsl.2017.06.018)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

*Document Version*

Peer reviewed version

*Citation for published version (Harvard):*

Maffione, M & Morris, A 2017, 'The onset of fabric development in deep marine sediments', *Earth and Planetary Science Letters*, vol. 474, pp. 32-39. <https://doi.org/10.1016/j.epsl.2017.06.018>

[Link to publication on Research at Birmingham portal](#)

**Publisher Rights Statement:**

Checked for eligibility: 08/06/2017

**General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

**Take down policy**

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

## Highlights

- High-resolution AMS study of deep marine clay-rich sediments from IODP Site U1438
- Compaction-related fabrics initiate sharply at 83 mbsf (5.7 Ma after deposition)
- Fabrics continue to form over a 6 Ma long, 30 m thick 'initial compaction window'
- An effective stress threshold triggers compaction-related fabric development

1 **The onset of fabric development in deep marine sediments**

2 Marco Maffione<sup>1,2</sup> and Antony Morris<sup>3</sup>

3 <sup>1</sup>School of Geography, Earth and Environmental Sciences, University of

4 Birmingham, Edgbaston, Birmingham, B15 2TT, UK

5 <sup>2</sup>Department of Earth Sciences, Utrecht University, Heidelberglaan 2, 3584 CS

6 Utrecht, Netherlands

7 <sup>3</sup>School of Geography, Earth and Environmental Sciences, Plymouth University,

8 Drake Circus, Plymouth, PL4 8AA, UK

9

10

11 Keywords

12 Sedimentary fabric; deep marine; compaction; magnetic anisotropy; AMS;

13 International Ocean Discovery Program

14

15 **Abstract**

16 Post-depositional compaction is a key stage in the formation of sedimentary

17 rocks that results in porosity reduction, grain realignment and the production of

18 sedimentary fabrics. The progressive time-depth evolution of the onset of fabric

19 development in deep marine sediments is poorly constrained due to the limited

20 quantity and resolution of existing data. Here we present high-resolution

21 anisotropy of magnetic susceptibility (AMS) results from clay-rich deep marine

22 sediments recovered at International Ocean Discovery Program Site U1438

23 (Philippine Sea). AMS is a petrofabric tool sensitive to the preferred orientation

24 of grains in rocks. Down-section variations of AMS parameters, density, porosity

25 and the inclination of magnetic remanences demonstrate that fabrics develop in  
26 response to compaction and dewatering but also that they do not develop  
27 progressively with depth below the mudline. Instead, a horizontal foliation first  
28 forms at 83 mbsf once the sediment load reaches an effective stress threshold for  
29 the onset of compaction and is then continuously enhanced down to 113 mbsf,  
30 defining a 30 m-thick 'initial compaction window'. The magnetostratigraphic age  
31 model for IODP Site U1438 indicates a delay of 5.7 Ma in initial fabric formation  
32 following sediment deposition, with strongly defined fabrics then taking an  
33 additional 6.5 Ma to develop.

34

## 35 **1. Introduction**

36 During deposition of (hemi)pelagic sediments in deep-sea environments, platy  
37 minerals (mainly phyllosilicates) tend to align with their long axes parallel to the  
38 water-sediment interface. Bottom-current disturbance and bioturbation,  
39 together with the natural predisposition of clay flakes to form edge-to-edge and  
40 edge-to-face contacts due to surface electric charge distribution (Bennett et al.,  
41 1991), eventually result in an uppermost sedimentary interval characterized by  
42 a chaotic internal structure (i.e., isotropic fabric) and high porosity and water  
43 content (Bennett et al., 1991; Reynolds and Gorsline, 1992). With increasing  
44 vertical burial load at depth, clays particles rotate to form horizontal face-to-face  
45 contacts, accompanied by simultaneous dewatering and porosity reduction  
46 (Bennett et al., 1981; Bennett and Hulbert, 1986). The effect of this process on  
47 the microstructure of the sediment is the formation of a fabric characterized by a  
48 well-defined horizontal foliation plane.

49 The majority of dewatering and compaction in pelagic sedimentary sequences is  
50 thought to occur progressively within the uppermost stratigraphic intervals (e.g.,  
51 Arason and Levi, 1990). However, only a few previous studies have investigated  
52 microstructure changes during compaction and initial fabric development in  
53 deep marine environments (Kopf and Behrmann, 1997; Hirano et al., 2001;  
54 Kawamura and Ogawa, 2002; 2004), and none have a sufficiently high spatial  
55 and temporal resolution to describe in detail the evolution of this process.  
56 Here we present results of a high-resolution anisotropy of magnetic  
57 susceptibility (AMS) analysis of a sequence of unconsolidated deep marine  
58 sediments recovered in the Philippine Sea by International Ocean Discovery  
59 Program (IODP) Expedition 351 (Arculus et al., 2015a), where  
60 magnetostratigraphy provides an accurate age model. We use AMS as a sensitive  
61 measure of fabric development in these sediments (e.g., Rochette et al., 1992;  
62 Tarling and Hrouda, 1993; Borradaile and Jackson, 2004), and compare magnetic  
63 fabric parameters with other physical properties to tightly constrain the depth  
64 and timing of the onset and evolution of fabric development in a deep marine  
65 environment.

66

## 67 **2. Geological background and sampling**

### 68 *2.1. Tectonic framework of the Philippine Sea Plate and IODP Site U1438*

69 The Philippine Sea Plate (PSP) is an oceanic plate located between the Eurasia  
70 and Pacific plates and bordered by active subduction zones (Figure 1). The PSP  
71 can be subdivided into three tectono-stratigraphic domains: (i) a western  
72 domain floored by a complex array of oceanic plateaux, ridges, and basins of  
73 Cretaceous to Oligocene age and bordered to the east by the Eo-Oligocene (~55-

74 25 Ma) Kyushu-Palau Ridge (e.g., Deschamps and Lallemand, 2003; Okino and  
75 Fuijoka, 2003; Ishizuka et al., 2011a, 2013); (ii) a central domain dominated by  
76 the Miocene (~25-12 Ma) Parece Vela-Shikoku back-arc basin (Okino et al.,  
77 1994) occurring between the Kyushu-Palau Ridge and the modern Izu-Bonin-  
78 Mariana (IBM) arc; and (iii) an eastern domain forming the forearc region  
79 between the modern IBM arc and the IBM trench, developed upon initiation of  
80 subduction of the Pacific plate below the PSP at ~52 Ma (Ishizuka et al., 2011b;  
81 Reagan et al., 2010; Arculus et al., 2015b).

82 During June and July 2014, IODP Expedition 351 (Arculus et al., 2015a)  
83 recovered a suite of sedimentary and volcanic rocks at Site U1438 (4700 m  
84 water depth) in the western domain of the PSP (Amami Sankaku Basin; 27.38°N,  
85 134.32°E; Figure 1a). Four progressively deeper holes were drilled at this site,  
86 down to 26.5, 257.3, 897.8, and 1611 metres below seafloor (mbsf), at Holes  
87 U1438A, B, D and E respectively. Recovered rocks (Figure 1b) consist of a thick  
88 sedimentary sequence (Units I-IV) deposited since the Early Eocene (~55 Ma)  
89 and recording the birth, evolution and death of the Kyushu-Palau Ridge,  
90 overlying igneous basement rocks (Unit 1).

91

## 92 *2.2. Deep marine sedimentation in the west PSP (Unit I)*

93 Unit I (0-160.3 mbsf; Hole U1438B) is an unconsolidated fine-grained pelagic  
94 and hemipelagic sedimentary sequence composed of mud, tuffaceous mud, mud  
95 with ash, and clay with discrete ash beds. Paleomagnetic and biostratigraphic  
96 constraints place the base of Unit I at the Miocene-Oligocene transition (Arculus  
97 et al., 2015a). Deposition of Unit I therefore started soon after the demise of  
98 Kyushu-Palau Ridge volcanism and initial opening of the Parece Vela-Shikoku

99 back-arc basin to the east (thought to occur at ~25 Ma; Okino et al., 1994). Since  
100 no significant tectonic events occurred since that time at the location of Site  
101 U1438, Unit I represents the product of deep marine sedimentation, with  
102 sedimentation rates ranging from ~2 to ~0.5 cm/kyr (Arculus et al., 2015a).  
103 The average grain size of Unit I changes slightly down-core, with the uppermost  
104 interval (0-45 mbsf) mainly represented by siltstones with a significant (up to  
105 60%) biogenic, mainly siliceous component, a central interval (45-93 mbsf)  
106 dominated by claystones with a lower biogenic content, and a bottom interval  
107 (93-160.3 mbsf) represented again predominantly by siltstones. Bioturbation is  
108 rare in the upper interval of Unit I (< 93 mbsf) and increases at depth, with more  
109 bioturbated intervals occurring below 121 mbsf. X-Ray diffraction analyses  
110 revealed a predominant assemblage of quartz, plagioclase, chlorite, muscovite,  
111 illite and other clay minerals (Arculus et al., 2015a). Below ~93 mbsf (in the  
112 lower siltstones) quartz content decreases, while the content of zeolite and clay  
113 minerals increases.

114

### 115 **3. Methods**

116 Anisotropy of magnetic susceptibility (AMS) is a petrofabric tool used to  
117 determine the preferred orientation of minerals (e.g., Jelínek, 1981; Hrouda,  
118 1982; Borradaile, 1988; Rochette et al., 1992; Tarling and Hrouda, 1993;  
119 Borradaile and Jackson, 2004). AMS is geometrically described by an ellipsoid  
120 with principal axes corresponding to the minimum ( $k_{\min}$ ), intermediate ( $k_{\text{int}}$ ), and  
121 maximum ( $k_{\max}$ ) susceptibilities (Hrouda, 1982). The relative magnitude of the  
122 susceptibility axes defines the shape of the AMS ellipsoid, which can be: (1)  
123 isotropic ( $k_{\min} = k_{\text{int}} = k_{\max}$ ) when grains are not aligned preferentially; (2) oblate

124 ( $k_{\min} \ll k_{\text{int}} \approx k_{\max}$ ) when grain alignment defines a foliation plane; (3) triaxial  
125 ( $k_{\min} < k_{\text{int}} < k_{\max}$ ) when grain alignment results in a well-defined foliation and a  
126 lineation; or (4) prolate ( $k_{\min} \approx k_{\text{int}} \ll k_{\max}$ ) when grain alignment defines a  
127 lineation. Here we describe the strength of anisotropy using the corrected  
128 anisotropy degree ( $P_j$ ; Jelínek, 1978), where  $P_j = 1.0$  indicates an isotropic fabric  
129 and, e.g.,  $P_j = 1.05$  indicates 5% anisotropy. The shape of the ellipsoid is  
130 described by the shape parameter ( $T$ ), where  $-1.0 < T < 1.0$  and positive/negative  
131 values of  $T$  indicate oblate/prolate fabrics respectively (Jelínek 1978).  
132 Sedimentary fabrics are characterised by  $P_j > 1.0$ , oblate AMS ellipsoids ( $0 < T <$   
133  $1$ ), vertical  $k_{\min}$  axes, and  $k_{\text{int}}$  and  $k_{\max}$  axes dispersed within the horizontal  
134 foliation plane.

135 We measured the anisotropy of low-field magnetic susceptibility of 173 discrete,  
136 8 cm<sup>3</sup> cubic samples from the top 140 m of Unit I. Measurements were carried  
137 out using an AGICO KLY-3S Kappabridge (Plymouth University), and  
138 susceptibility tensors and associated eigenvectors and eigenvalues calculated  
139 using AGICO Anisoft 4.2 software. Instrument precision for AMS measurements  
140 for the KLY-3S system is > 99%, yielding meaningful principal directions in  
141 weakly anisotropic rocks when  $P_j > 1.003$  (Burmeister et al., 2009). Sample  
142 orientations were determined by correcting cores recovered from Hole U1438B  
143 to the geographic reference frame using “FlexIT” tool data obtained during  
144 advanced piston corer (APC) deployments (Arculus et al., 2015a).

145 Rock magnetic experiments were performed to investigate the nature of the  
146 mineral fractions contributing to the AMS. Curie temperatures were determined  
147 from the high-temperature (20-700°C) variation of magnetic susceptibility of  
148 representative samples, measured using a KLY-3S coupled with a CS-3 high-

149 temperature furnace apparatus. Isothermal remanent magnetization (IRM)  
150 acquisition experiments were conducted on representative samples using a  
151 Molspin pulse magnetizer to apply peak fields up to 800 mT with resulting IRMs  
152 measured using an AGICO JR6A spinner magnetometer.  
153 Finally, we also employ shipboard paleomagnetic and physical property data  
154 from Hole U1438B, collected using methods documented in Arculus et al.  
155 (2015a).

156

#### 157 **4. Results**

158 AMS parameters from the investigated interval of Unit I are extremely variable  
159 with depth, with a clear change occurring at 83 mbsf (Figure 2; Supplementary  
160 material Table 1). The anisotropy degree is low ( $P_j \approx 1.006$ ) indicating weakly  
161 anisotropic sediments in the uppermost 83 m of the sequence, but progressively  
162 increases over an interval of  $\sim 30$  m below 83 mbsf, reaching a maximum of  $P_j$   
163  $\approx 1.060$  at 113 mbsf (Figure 2a). Similarly the shape parameter ( $T$ ) is extremely  
164 variable from -1.0 to +1.0 in the upper interval, becoming consistently oblate  
165 ( $0.6 < T < 1.0$ ) immediately below 83 mbsf, and then strongly oblate ( $0.8 < T <$   
166  $1.0$ ) below 113 mbsf (Figure 2b). In addition, the orientation of the principal  
167 susceptibility axes changes with depth. Above 83 mbsf, magnetic fabrics are  
168 randomly oriented, with only the  $k_{min}$  axes showing a faint alignment near the  
169 vertical (Figure 3a). Below 83 mbsf,  $k_{min}$  axes become tightly clustered along the  
170 vertical, and the  $k_{int}$  and  $k_{max}$  axes align with the horizontal plane (Figure 3b). A  
171 weak clustering of  $k_{max}$  axes along an ESE-NWN orientation is also observed,  
172 potentially reflecting weak bottom water currents during deposition (Hrouda

173 and Jezek, 1999). Mean magnetic susceptibility shows a slight increase with  
174 depth from  $\sim 30$  to  $\sim 80 \times 10^{-5}$  SI (Figure 2c).  
175 Results from high-temperature variation of magnetic susceptibility experiments  
176 revealed consistent maximum Curie temperatures of  $\sim 580^\circ\text{C}$  (Figure 4a),  
177 indicating that the ferromagnetic fraction in these sediments is dominated by  
178 near-stoichiometric magnetite (Dunlop and Özdemir, 1997). This is consistent  
179 with IRM acquisition experiments, where saturation was reached at applied  
180 fields of 200-300 mT (Figure 4b), consistent with the presence of low-coercivity,  
181 fine-grained (single-domain to pseudo-single-domain) magnetite. However, the  
182 vertical alignment of  $k_{min}$  axes (perpendicular to horizontal bedding in the  
183 sediments) indicates a dominance of normal sedimentary fabrics, and excludes  
184 the presence of significant inverse magnetic fabric components due to single  
185 domain effects in these sediments (Potter and Stephenson, 1988).

186

## 187 **5. Discussion**

### 188 *5.1. Comparison of AMS fabric variations with physical properties*

189 Interpretation of AMS data requires understanding of the source of the  
190 susceptibility signal and its potential variability downhole. A linear relationship  
191 between low field magnetic susceptibility and the intensity of natural remanent  
192 magnetization (NRM) suggests constancy of ferromagnetic mineralogy and  
193 grain-size (domain state) through Unit I (Figure 4c), implying that the  
194 contribution of magnetite to the susceptibility signal is controlled by  
195 concentration alone. The relatively low bulk magnetic susceptibility (Figure 2c),  
196 however, indicates that magnetite content is low and that the paramagnetic  
197 phyllosilicate matrix is likely the main contributor to susceptibility in these

198 sediments, as typically observed in clay-rich sediments (e.g., Borradaile and  
199 Jackson, 2004; Sagnotti and Speranza, 1993; Maffione et al., 2012, 2015; Parés,  
200 2015).

201 The rapid increase of the anisotropy degree, together with the acquisition of a  
202 consistently oblate fabric below 83 mbsf within Unit I (Figure 2) marks the onset  
203 of development of a horizontal foliation typical of sedimentary fabrics (Figure  
204 3b). This foliation is progressively enhanced down to 113 mbsf, where the  
205 anisotropy degree reaches a maximum and AMS ellipsoids become strongly  
206 oblate (Figure 2b). This depth interval of fabric development coincides with a  
207 distinct decrease in porosity from 75 to 65% and increase in bulk density from  
208  $\sim 1.45$  to  $\sim 1.55$  g/cm<sup>3</sup> (Figure 2d), seen in shipboard physical property data  
209 acquired on discrete samples analysed during IODP Expedition 351 (Arculus et  
210 al., 2015a). This suggests that fabrics have developed in response to mechanical  
211 compaction and dewatering occurring during burial diagenesis due to increasing  
212 effective overburden stress. Chemical compaction is considered insignificant in  
213 Unit I, as this generally only becomes a significant porosity-occluding process at  
214 depths greater than  $\sim 2$  km (Bjørlykke and Høeg, 1997; Mondol et al., 2008). An  
215 apparent increase in sedimentation rate from  $\sim 0.5$  to  $\sim 2.0$  cm/ka at 83 mbsf  
216 (Figure 2e) is likely to also result from mechanical compaction rather than a  
217 change in rate of sediment supply.

218

## 219 *5.2. Paleomagnetic evidence for compaction*

220 As an additional test for the compaction-related origin of fabric development we  
221 analyzed the occurrence of inclination shallowing of paleomagnetic remanences,  
222 which characteristically results from compaction (e.g., Anson and Kodama, 1987;

223 Arason and Levi, 1990; Tauxe and Kent, 2004; Huang et al., 2013). Shipboard  
224 remanence directions are presented in Figure 5 (based on measurements of  
225 archive half core sections following alternating field demagnetization at 25 mT to  
226 remove significant drilling-induced components; Figure 6). Inclinations above 83  
227 mbsf are close to the value expected at IODP Site U1438, but are generally  
228 shallower than expected below this (Figure 5a). To test this statistically, we  
229 selected only data from the bottom 1.0 m of each core in order to exclude  
230 intervals affected by drilling-induced shearing (commonly encountered in IODP  
231 piston cores (Acton et al., 2002) and evident in the declination data of Figure 5b).  
232 Mean inclinations were calculated after transposing all data to normal polarity  
233 and applying a fixed 45° cut-off to the distribution of corresponding virtual  
234 geomagnetic poles (following the methodology of Johnson et al. (2008)). For the  
235 uppermost 83 m of Unit I, the mean inclination of 44.1° ( $\alpha_{95} = 1.2^\circ$ ,  $n = 408$ ) is  
236 not statistically different from the expected geocentric axial dipole (GAD)  
237 inclination at this site ( $I = 46^\circ$ ). In contrast, the mean inclination below 83 mbsf  
238 ( $I = 39.7^\circ$ ;  $\alpha_{95} = 1.8^\circ$ ,  $n = 266$ ) is statistically significantly shallower. Using the  
239 elongation/inclination (E/I) correction method (Tauxe and Kent, 2004) we  
240 obtained an unflattened inclination for the interval 83-140 mbsf of 42.8° (95%  
241 confidence limits: 40.2°-53.4°) consistent with the expected inclination. These  
242 results further confirm that mechanical compaction has affected the lower  
243 interval of Unit I but is absent or not statistically significant at shallower depths.

244

245 *5.3. Potential causes of the sudden onset of fabric development at 83 mbsf*

246 *5.3.1. Comparison with compaction trends for marine mudrocks*

247 Effective stress-controlled mechanical compaction at the relative shallow depths  
248 of Unit I in Hole U1438B might be expected to produce continuous changes in  
249 physical properties, i.e. with a progressive decrease in porosity and increase in  
250 anisotropy degree with depth. Following Kopf and Behrmann (1997), we  
251 compare downhole porosity variations with published mechanical compaction  
252 functions by converting porosity determinations to uniaxial vertical shortening  
253 values,  $e_v$  using:

$$e_v = \frac{P_0 - P}{P - 100}$$

254 where  $P$  is a given sample porosity (in %) and  $P_0$  is an initial porosity near the  
255 mudline (taken as 75%). Figure 7 compares the resulting values to converted  
256 porosity-depth trends for clays and claystones of Terzaghi (1925), Athy (1930),  
257 Hamilton (1976) and Sclater and Christie (1980). Though variable, each model  
258 predicts a smooth, monotonic increase in shortening with depth, in contrast to  
259 the stepwise increase at 83 mbsf in Unit I from  $e_v$  values close to zero (mean = -  
260 0.03) to a mean value of -0.21 below. This is consistent with the AMS data that  
261 show a uniformly low anisotropy degree down to 83 mbsf and then a sharp  
262 change in character to pronouncedly oblate fabrics. Hence, factors other than  
263 continuous compaction across the whole depth interval of Unit I must influence  
264 fabric development in these sediments.

265

### 266 *5.3.2. Apparent overconsolidation*

267 The sharp increase of the anisotropy degree in the studied unit resembles the  
268 pattern expected for “apparent overconsolidation” (Schwehr et al., 2006),  
269 whereby sediments appear more consolidated than predicted by simple burial

270 for a given depth. According to Schwehr et al. (2006) apparent overconsolidation  
271 can result from unconformities produced by underwater landslides, which  
272 unload pre-consolidated sediments that then become buried by new sediments.  
273 Alternatively, overconsolidation can be caused by repeated shaking events  
274 (Locat and Lee, 2002). The continuous magnetostratigraphic record at Unit I  
275 (Arculus et al., 2015a), however, excludes the existence of unconformities within  
276 the sequence, and it is unlikely that seismic events only caused  
277 overconsolidation of the interval of Unit I below 83 mbsf without also affecting  
278 the top of the section in this tectonically active region.

279

### 280 *5.3.3. Compositional control*

281 Compaction of sediments can be controlled by variations in the composition and  
282 size of sedimentary grains, with, for example, platy phyllosilicate minerals tending  
283 to facilitate compaction. However, no major change in phyllosilicate mineralogy,  
284 grain size or degree of bioturbation of the mudrocks of Unit I (Arculus et al.,  
285 2015a) corresponds with the sharp onset of fabric development at 83 mbsf.  
286 Curtis et al. (1980) have demonstrated that although preferred orientations in  
287 clay-rich sediments commonly result from compactional strain, fabric  
288 development may be affected by the presence of non-platy silicate particles  
289 (such as quartz grains) which inhibit planar fabric development in their  
290 immediate vicinity. In this context, we note a downhole decrease in the volume  
291 percentage of fine-grained quartz particles in the mudrocks, from an average of  
292 8% to 4% (based on smear-slide data collected during shipboard analyses;  
293 Arculus et al., 2015a). This minor compositional change is illustrated semi-  
294 quantitatively by variations in the intensity of quartz peaks at  $26.6^\circ 2\theta$  in XRD

295 analyses of Unit I mud samples (Figure 2g; data from Arculus et al., 2015a).  
296 Quartz content, however, mainly decreases between 50 and 65 mbsf (Figure 2g),  
297 and remains constant throughout the interval where sharp variations of AMS  
298 parameters are observed. Furthermore, the volume percentage change in quartz  
299 affecting fabric development in the mudrocks analysed by Curtis et al. (1980)  
300 was much more dramatic (28% to 15%). Finally, the combined volume  
301 percentage of the main non-platy silicates (quartz plus feldspar) in Hole U1438B  
302 increases slightly from 12% to 15% across the interval where AMS fabrics  
303 develop (Arculus et al., 2015a), suggesting that this mechanism cannot account  
304 for the stepwise change in anisotropy observed here.

305

#### 306 *5.3.4. An effective stress threshold for the 'initial compaction window'*

307 The stepwise variation of anisotropy parameters, porosity/vertical shortening  
308 and bulk density with depth documented here suggests that compaction and  
309 sedimentary fabric development is not progressive, but only starts once the  
310 burial load produces a sufficient effective stress to exceed the internal resistance  
311 to compaction represented by the friction along grain contacts during  
312 reorientation. In this case, the density data of Figure 2d, a water depth of 4700 m  
313 and a seawater density of 1029 kg/m<sup>3</sup> combine to yield an effective stress at 83  
314 mbsf of 0.35 MPa (where effective stress = total stress – pore water pressure  
315 (assumed to be hydrostatic); Terzaghi, 1925). Sedimentary fabric in the studied  
316 section then develops over a restricted interval between 83 and 113 mbsf here  
317 defined as the 'initial compaction window', where pore reduction and preferred  
318 horizontal alignment of phyllosilicate flakes is progressively enhanced over a 30  
319 m-thick interval. A similar compaction window can also be inferred in pelagic

320 sediments from ODP Site 1149 in the NW Pacific Ocean, where a stepwise change  
321 in physical properties occurs between 118 and 150 mbsf (Kawamura and Ogawa,  
322 2002; 2004).

323

#### 324 *5.4 Implications for fabric studies on deformed ancient mudrocks*

325 The excellent age control at Site U1438, based on the IODP Expedition 351  
326 shipboard magnetostratigraphy that successfully identified every geomagnetic  
327 chron in core sections back to 36 Ma (Arculus et al., 2015a), provides constraints  
328 on the timing of initial development of sedimentary fabrics in Unit I mudrocks.  
329 These indicate that the full process of fabric development documented here  
330 started 5.7 Ma after initial deposition and then took 6.5 Ma to complete (Figure  
331 2e). This has implications for studies using AMS to document the progressive  
332 tectonic overprinting of sedimentary fabrics in ancient mudrocks (e.g., Sagnotti  
333 et al., 1998; Larrasoana et al., 2004). Such studies typically assume that initial  
334 sedimentary fabrics (which are subsequently partially overprinted by tectonic  
335 fabrics) correspond to the biostratigraphic age of the sampled rocks. Instead  
336 results here suggest that a ~6 Ma hiatus between deposition and initial fabric  
337 development may need to be considered when documenting the evolution of  
338 anisotropy in deep marine mudrocks.

339

## 340 **6. Conclusions**

341 A detailed record of progressive fabric development in deep marine sediments  
342 sampled at IODP Site U1438 with robust age control indicates that a compaction-  
343 related horizontal foliation started to form 5.7 Ma after initial sedimentation (83  
344 mbsf) and developed over the subsequent 6.5 Ma (down to 113 mbsf), within a

345 30 m-thick interval here termed the 'initial compaction window'. These new  
346 results provide the first high-resolution time/depth constraints on the onset of  
347 fabric development in deep marine environments, with implications for the  
348 tectonic application of AMS in pelagic sedimentary rocks. More data are now  
349 required from such deep marine environments to determine the variability in  
350 the onset of the initial compaction window, including additional quantification of  
351 clay microfabrics through this window using scanning electron microscope  
352 (SEM) and transmission electron microscope (TEM) techniques.

353

#### 354 **Acknowledgements**

355 This research used samples and data provided by International Ocean Discovery  
356 Program (IODP). We thank the members of IODP expedition 351 Science Party  
357 and the captain and crew of the JOIDES Resolution. Shipboard bulk density and  
358 porosity measurements were performed by Mike Gurnis and Morihisa Hamada,  
359 and the anisotropy of magnetic susceptibility measurements on discrete samples  
360 by Matthew Meyer (Plymouth University).

361 Funding: This work was supported by the Natural Environment Research  
362 Council [grant number NE/M007367/1 awarded to AM]; and the European  
363 Research Council [grant number 306810, awarded to D.J.J. van Hinsbergen]. The  
364 paper benefitted from thoughtful reviews by Eric Ferré and an anonymous  
365 reviewer.

366

#### 367 **References**

368 Acton, G. D., 2002. Paleomagnetic overprints in ocean sediment cores and their  
369 relationship to shear deformation caused by piston coring. *J. Geophys. Res.*  
370 107, doi: 10.1029/2001JB000518.

371 Anson, G. L. and Kodama, K. P., 1987. Compaction-induced inclination shallowing  
372 of the post-depositional remanent magnetization in a synthetic sediment.  
373 *Geophys. J. Int.* 88, 673-692.

374 Arason, P. and Levi, S., 1990. Compaction and inclination shallowing in deep-sea  
375 sediments from the Pacific Ocean, *J. Geophys. Res.*, 95(B4), 4501-4510,  
376 doi:10.1029/JB095iB04p04501.

377 Arculus, R. J., Ishizuka, O., Bogus, K. and the Expedition 351 Scientists, 2015a.  
378 *Proceedings of the International Ocean Discovery Program Volume 351*,  
379 doi:10.14379/iodp.proc.351.103.2015.

380 Arculus, R. J., Ishizuka, O., Bogus, K., Gurnis, M., Hickey-Vargas, R., Aljehdali, M. H.,  
381 Bandini-Maeder, A. N., Barth, A. P., Brandl, P., Drab, L., do Monte Guerra, R.,  
382 Hamada, M., Jiang, F., Kanayama, K., Kender, S., Kusano, Y., Li, H., Loudin, L.  
383 C., Maffione, M., Marsaglia, K. M., McCarthy, A., Meffre, S., Morris, A., Neuhaus,  
384 M., Savov, I. P., Sena, C., Tepley III, F. J., van der Land, C., Yogodzinski, G. M.  
385 and Zhang, Z., 2015b. A record of spontaneous subduction initiation in the  
386 Izu–Bonin–Mariana arc. *Nat. Geosci.* 8, 728–733, doi: 10.1038/ngeo2515.

387 Athy, L. F., 1930. Density, porosity and compaction of sedimentary rocks  
388 (abstract). *American Association of Petroleum Geologists, Bulletin*, 14, 24.

389 Bennett, R. H., Bryant, W. R. and Keller, G. H., 1981. Clay fabric of selected  
390 submarine sediments: fundamental properties and models. *J. Sediment.*  
391 *Petrol.* 51, 217-232.

392 Bennett, R. H., O'Brien, N. R. and Hulbert, M. H., 1991. Determinants of clay and  
393 shale microfabric signatures: processes and mechanisms. In: Bennett, R. H.,  
394 Bryant, W. R. and Hulbert, M. H. (Eds.), *Microstructures of Fine-grained*  
395 *Sediments from Mud to Shale*. Springer-Verlag, New York, pp. 5 –32.

396 Bennett, R. H. and Hulbert, M. H., 1986. *Clay Microstructure*. Boston  
397 (International Human Resources Development Corp. Publ.).

398 Bjørlykke, K. and Høeg, K., 1997. Effects of burial diagenesis on stresses,  
399 compaction and fluid flow in sedimentary basins. *Marine and Petroleum*  
400 *Geology*, 14, 267-276.

401 Borradaile, G. J., 1988. Magnetic-susceptibility, petrofabrics and strain:  
402 *Tectonophysics* 156, 1–20, doi: 10.1016/0040-1951(88)90279-x.

403 Borradaile, G. J. and Jackson, M., 2004. Anisotropy of magnetic susceptibility  
404 (AMS): magnetic petrofabrics of deformed rocks. *Geol. Soc. London Sp. Publ.*  
405 238, 299–360, doi: 10.1144/GSL.SP.2004.238.01.18.

406 Burmeister, K. C., Harrison, M. J., Marshak, S., Ferré, E. C., Bannister, R. A. and  
407 Kodama, K. P., 2009. Comparison of Fry strain ellipse and AMS ellipsoid  
408 trends to tectonic fabric trends in very low-strain sandstone of the  
409 Appalachian fold–thrust belt. *J. Struct. Geol.* 31, 1028-1038.

410 Curtis, C. D., Lipshie, S. R., Oertel, G. and Pearson, M. J., 1980. Clay orientation in  
411 some Upper Carboniferous mudrocks, its relationship to quartz content  
412 and some inferences about fissility, porosity and compactional history.  
413 *Sedimentology* 27, 333-339.

414 Deschamps, A. and Lallemand, S., 2003. Geodynamic setting of Izu-Bonin-  
415 Mariana boninites. *Geol. Soc. London Sp. Publ.* 219, 163–185, doi:  
416 10.1144/GSL.SP.2003.219.01.08.

417 Dunlop, D. J. and Özdemir, Ö., 1997. *Rock magnetism: fundamentals and*  
418 *frontiers*. Cambridge University Press, Cambridge, 573 pp.

419 Hamilton, E. L., 1976. Variations of density and porosity with depth in deep sea

420 sediments. *Journal of Sedimentary Petrology*, 46, 280-300.

421 Hirano, S., Ogawa, Y. and Kawamura, K., 2001. Deformation of unlithified  
422 sediments in an early stage of the compaction process deduced from  
423 microtextures and magnetic fabrics: ODP Leg 174B, Hole 1074A. In: Becker,  
424 K., Malone, M.J. (Eds.), *Proc. ODPSci. Results*, vol. 174B, 1 –13.

425 Hrouda, F., 1982. Magnetic anisotropy of rocks and its application in geology and  
426 geophysics. *Geophys. Surveys* 5, 37–82, doi: 10.1007/BF01450244.

427 Hrouda, F. and Jezek, J., 1999. Magnetic anisotropy indications of deformations  
428 associated with diagenesis. In: Tarling, D. H. and Turner, P. (eds),  
429 *Palaeomagnetism and Diagenesis in Sediments*, Geological Society, London,  
430 Special Publication, 151, 127-137.

431 Huang, W., Dupont-Nivet, G., Lippert, P. C., van Hinsbergen, D. J. J. and Hallot, E.,  
432 2013. Inclination shallowing in the Eocene Linzizong sediments from Tibet:  
433 correction, possible causes and implications for reconstructing the India-  
434 Asia collision. *Geophys. J. Int.* 194, 1390-1411.

435 Ishizuka, O., Taylor, R. N., Yuasa, M. and Ohara, Y., 2011a. Making and breaking an  
436 island arc: A new perspective from the Oligocene Kyushu-Palau arc,  
437 Philippine Sea. *Geochem. Geophys. Geosys.* 12, 1–40, doi:  
438 10.1029/2010GC003440.

439 Ishizuka, O., Tani, K., Reagan, M. K., Kanayama, K., Umino, S., Harigane, Y.,  
440 Sakamoto, I., Miyajima, Y., Yuasa, M. and Dunkley, D. J., 2011b. The  
441 timescales of subduction initiation and subsequent evolution of an oceanic  
442 island arc. *Earth Planet. Sci. Lett.* 306, 229–240, doi:  
443 10.1016/j.epsl.2011.04.006

444 Ishizuka, O., Taylor, R. N., Ohara, Y. and Yuasa, M. , 2013. Upwelling, rifting, and  
445 age-progressive magmatism from the Oki-Daito mantle plume. *Geology*, 41,  
446 1011–1014.

447 Jelínek, V., 1978. Statistical processing of anisotropy of magnetic susceptibility  
448 measured on groups of specimens. *Studia Geophys. Geod.* 22, 50–62, doi:  
449 10.1007/BF01613632.

450 Jelínek, V., 1981. Characterization of the magnetic fabric of rocks. *Tectonophysics*  
451 79, 63-67.

452 Johnson, C. L., Constable, C. G., Tauxe, L., Barendregt, R., Brown, L. L., Coe, R. S.,  
453 Layer, P., Mejia, V., Opdyke, N. D., Singer, B. S., Staudigel, H. and Stone, D. B.,  
454 2008. Recent investigations of the 0-5 Ma geomagnetic field recorded by  
455 lava flows. *Geochemistry, Geophys. Geosystems*, 9(4),  
456 doi:10.1029/2007GC001696.

457 Kawamura, K. and Ogawa, Y., 2002. Progressive microfabric changes in  
458 unconsolidated pelagic and hemipelagic sediments down to 180 mbsf,  
459 northwest Pacific, ODP Leg 185, Site 1149. In Ludden, J.N., Plank, T., and  
460 Escutia, C. (Eds.), *Proc. ODP, Sci. Results*, 185.

461 Kawamura, K. and Ogawa, Y., 2004. Progressive change of pelagic clay  
462 microstructure during burial process: examples from piston cores and ODP  
463 cores. *Marine Geol.* 207, 131-144. doi:10.1016/j.margeo.2004.03.016.

464 Kopf, A. and Behrmann, J. H., 1997. Fabric evolution and mechanisms of  
465 diagenesis in fine-grained sediments from the Kita-Yamato Trough, Japan  
466 Sea. *J. Sedim. Res.* 67, 590–600.

467 Larrasoana, J. C., Pueyo, E. L. and Pares, J. M., 2004. An integrated AMS, structural,  
468 palaeo- and rock-magnetic study of Eocene marine marls from the Jaca-

469 Pamplona basin (Pyrenees, N Spain); new insights into the timing of  
470 magnetic fabric acquisition in weakly deformed mudrocks. *Geol. Soc.*  
471 *London Sp. Publ.* 238, 127–143, doi: 10.1144/GSL.SP.2004.238.01.10.  
472 Locat, J. and Lee, H., 2002. Submarine landslides: Advances and challenges.  
473 *Canadian Geotech. J.* 39, 193–212.  
474 Maffione, M., Pucci, S., Sagnotti, L. and Speranza, F., 2012. Magnetic fabric of  
475 Pleistocene continental clays from the hanging-wall of an active low-angle  
476 normal fault (Altotiberina Fault, Italy). *Int. J. Earth Sci.* 101, 849–861, doi:  
477 10.1007/s00531-011-0704-9.  
478 Maffione, M., Hernandez-Moreno, C., Ghiglione, M. C., Speranza, F., van  
479 Hinsbergen, D. J. J. and Lodolo, E., 2015. Constraints on deformation of the  
480 Southern Andes since the Cretaceous from anisotropy of magnetic  
481 susceptibility. *Tectonophysics* 665, 236–250, doi:  
482 10.1016/j.tecto.2015.10.008.  
483 Mondol, N. H., Fawad, M., Jahren, J. and Bjørlykke, K., 2008. Synthetic mudstone  
484 compaction trends and their use in pore pressure prediction. *First Break*,  
485 26, 43-51.  
486 Okino, K. and Fujioka, K., 2003. Central basin spreading center, Philippine Sea:  
487 Structure of an extinct spreading center and implications for marginal basin  
488 information. *J. Geophys. Res.* 108, 2040, doi:10.1029/2001JB001095.  
489 Okino, Y., Shimakawa, Y. and Nagaoka, S., 1994. Evolution of the Shikoku Basin. *J.*  
490 *Geomag. Geoelect.* 46, 463–479.  
491 Parés, J. M., 2015. Sixty years of anisotropy of magnetic susceptibility in  
492 deformed sedimentary rocks. *Frontiers Earth Sci.* 3, 1–13, doi:  
493 10.3389/feart.2015.00004.  
494 Potter, D. K. and Stephenson, A., 1988. Single-domain particles in rocks and  
495 magnetic fabric analysis. *Geophysical Research Letters*, 15, 1097-1100, doi:  
496 10.1029/GL015i010p01097.  
497 Reagan, M. K., Ishizuka, O., Stern, R. J., Kelley, K. A., Ohara, Y., Blichert-Toft, J.,  
498 Bloomer, S. H., Cash, J., Fryer, P., Hanan, B. B., Hickey-Vargas, R., Ishii, T.,  
499 Kimura, J. I., Peate, D. W., Rowe, M. C. and Woods, M., 2010. Fore-arc basalts  
500 and subduction initiation in the Izu-Bonin-Mariana system. *Geochem.*  
501 *Geophys. Geosys.*, 11, 1–17, doi: 10.1029/2009GC002871.  
502 Reynolds, S. and Gorsline, D.S., 1992. Clay microfibrils of deep sea mud(stones),  
503 California Continental Borderland. *J. Sed. Petr.* 62, 41-53.  
504 Rochette, P., Jackson, M. and Aubourg, C., 1992. Rock Magnetism and the  
505 interpretation of magnetic susceptibility. *Reviews Geophys.* 30, 209–226.  
506 Sagnotti, L. and Speranza, F., 1993. Magnetic fabric analysis of the Plio-  
507 Pleistocene clayey units of the Sant'Arcangelo basin, southern Italy. *Physics*  
508 *of the Earth and Planetary Interiors*, 77, 165-176.  
509 Sagnotti, L., Speranza, F., Winkler, A., Mattei, M. and Funicello, R., 1998. Magnetic  
510 fabric of clay sediments from the external northern Apennines (Italy). *Phys.*  
511 *Earth Planet. Int.* 105, 73–93, doi: 10.1016/S0031-9201(97)00071-X.  
512 Schwehr, K., Tauxe, L., Driscoll, N. and Lee, H., 2006. Detecting compaction  
513 disequilibrium with anisotropy of magnetic susceptibility. *Geochem.*  
514 *Geophys. Geosys.* 7, doi: 10.1029/2006GC001378.  
515 Schlater, J. G. and Christie, P. A. F., 1980. Continental stretching: an explanation of  
516 the post-mid-Cretaceous subsidence of the Central North Sea basin. *Journal*  
517 *of Geophysical Research*, 85, 3711-3739.

518 Tarling, D. and Hrouda, F. (Eds.), 1993. Magnetic anisotropy of rocks. Springer  
519 Science & Business Media.  
520 Tauxe, L. and Kent, D. V., 2004. A simplified statistical model for the geomagnetic  
521 field and the detection of shallow bias in paleomagnetic inclinations: was  
522 the ancient magnetic field dipolar? *Timescales Paleomag. Field*, 145, 101–  
523 115.  
524 Terzaghi, K., 1925. *Erdbaumechanik auf Bodenphysikalischer Grundlage*. Franz  
525 Deuticke, Leipzig-Vienna, 274 p.

526  
527

## 528 **Figure captions**

529

530 Figure 1. (a) Location map of the Philippine Sea Plate (PSP) showing the main  
531 geological structures and domains inferred from bathymetry, and the location of  
532 IODP Site U1438. (b) Lithostratigraphic log from Site U1438. The grain size is  
533 averaged over 5 m thick intervals. (cl) clay; (si) silt; (vfs-fs) very fine to fine sand;  
534 (ms-vcs) medium to very coarse sand; (gr), granules. Unit subdivision is  
535 indicated.

536

537 Figure 2. (a-c) Anisotropy of magnetic susceptibility results from 173 discrete  
538 cubic samples from Unit I, IODP Hole U1438B, showing the variation with depth  
539 of (a) the corrected anisotropy degree,  $P_f$  (b) the shape parameter,  $T$ , and (c)  
540 mean susceptibility. (d-g) Variation with depth of physical property and  
541 lithostratigraphic data from IODP Expedition 351 shipboard measurements  
542 (Arculus et al., 2015a). (d) Density and porosity data from discrete samples. (e)  
543 Age model based on magnetostratigraphy. Each point on the age-depth curve  
544 represents a reversal of the magnetic polarity in the core dated using the  
545 reference geomagnetic polarity time scale (GPTS). Sedimentation rates are  
546 calculated on the basis of the magnetostratigraphy. (f) Simplified  
547 lithostratigraphic log of Unit I. (g) Proxy for quartz content derived from peak  
548 intensities on X-Ray diffraction spectra (note arbitrary units). The two dotted  
549 gray lines bracket the interval in which anisotropy and physical parameters  
550 progressively change (here defined as the ‘initial compaction window’).

551

552 Figure 3. Stereographic equal area projections of the principal susceptibility axes  
553 from samples (a) above and (b) below 83 mbsf.

554

555 Figure 4. (a) High-temperature variation of low-field magnetic susceptibility  
556 during a complete heating-cooling cycle, showing a clear Curie temperature of  
557  $\sim 580^\circ\text{C}$  (heating curve). The cooling curve shows a slightly higher Curie  
558 temperature probably due to the oxidation of original magnetite to maghemite.  
559 (b) Isothermal remanent magnetization (IRM) acquisition curve from a  
560 representative sample showing presence of low coercivity magnetite. (c)  
561 Susceptibility vs. intensity of natural remanent magnetization (NRM) showing a  
562 linear relationship suggesting constancy of ferromagnetic mineralogy downhole.

563

564 Figure 5. Variation with depth of (a) inclination and (b) declination of the  
565 magnetic remanence in Unit I after alternating field (AF) demagnetization at 25

566 mT of archive half core sections (2 cm measurement interval; see Arculus et al.  
567 (2015b) for full description of methodology). Data are in geographic coordinates,  
568 after correction using FlexIT core orientation tool data. Declination and  
569 inclination vary downward due to the reversals of the geomagnetic field. (a)  
570 Inclination decreases significantly below core 9H (the effect is more visible in the  
571 normally magnetized intervals). (b) Declination varies within each core likely  
572 due to drilling-induced shear effects, particularly evident in cores 3H, 5H, 8H,  
573 and 13H. The shear effect is expected to be minimum at the bottom of each core.  
574 (c) Core distribution with depth from Hole U1438B.

575

576 Figure 6. Representative orthogonal vector plots of demagnetization data from  
577 intervals used for the calculation of the mean inclination. We select only those  
578 intervals in which 25 mT AF steps provide a reliable measure of the isolated  
579 characteristic remanent magnetization directions (i.e., declination and  
580 inclination of the 25 mT step is at an angle  $< 10^\circ$  from the best fit line calculated  
581 using the 20-40 mT interval). Black (white) dots represent the declination  
582 (inclination) projection.

583

584 Figure 7. Variation of uniaxial vertical shortening derived from the density data  
585 of Figure 2d (open circles) compared to published theoretical compaction trends  
586 for clays and claystones. The two dotted gray lines bracket the interval in which  
587 anisotropy and physical parameters progressively change (here defined as the  
588 'initial compaction window').

589

590 Table S1. Summary of anisotropy of magnetic susceptibility data from Unit I of  
591 IODP Hole U1438B.

592

Figure 1

[Click here to download Figure: Figure\\_1\\_Map\\_Lithos.pdf](#)




Figure 2

[Click here to download Figure: Figure\\_2\\_Downhole\\_plots.pdf](#)





Figure 3

[Click here to download Figure: Figure\\_3\\_AMS\\_stereonet.pdf](#)


a) < 83 mbsf

N



b) > 83 mbsf

N



■  $k_{max}$     △  $k_{int}$     ●  $k_{min}$

Figure 4

[Click here to download Figure: Figure\\_4\\_Rock\\_magnetism.pdf](#)




Figure 5

[Click here to download Figure: Figure\\_5\\_Dec-Inc\\_plot.pdf](#)




Figure 6

[Click here to download Figure: Figure\\_6\\_Z-plots.pdf](#)





Figure 7

[Click here to download Figure: Figure\\_7\\_Shortening\\_v\\_depth.pdf](#)



**Supplementary material for online publication only**

[Click here to download Supplementary material for online publication only: Table 1.xlsx](#)