

# MB-85. Non-invasive temperature measurements by MRI as a predictor of the survival of medulloblastoma patients

Babourina-Brooks, Ben; Kohe, Sarah; Davies, Nigel; Peet, Andrew

DOI:

[10.1093/neuonc/now076.81](https://doi.org/10.1093/neuonc/now076.81)

License:

Other (please specify with Rights Statement)

*Document Version*

Peer reviewed version

*Citation for published version (Harvard):*

Babourina-Brooks, B, Kohe, S, Davies, N & Peet, A 2016, 'MB-85. Non-invasive temperature measurements by MRI as a predictor of the survival of medulloblastoma patients', *Neuro-Oncology*, vol. 18, no. Suppl 3, MB-85, pp. 116. <https://doi.org/10.1093/neuonc/now076.81>

[Link to publication on Research at Birmingham portal](#)

## **Publisher Rights Statement:**

The accepted manuscript (AM) is the final draft author manuscript, as accepted for publication by a journal, including modifications based on referees' suggestions, before it has undergone copyediting, typesetting and proof correction. This is sometimes referred to as the post-print version.

## **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## **Take down policy**

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

- Abstracts are limited to 250 words (not including title, authors/institutional affiliations)
- Figures and tables are not allowed
- Abstract titles should use sentence case (i.e., only the first letter of the title should be capitalized)
- The body of the abstract should be in a single paragraph. If subheaders are used, they should be within that single paragraph in all-caps, not bolded (e.g., METHODS)

## Words 249

Title: Non-invasive temperature measurements by MRI as a predictor of the survival of medulloblastoma patients.

B.Babourina-Brooks<sup>1,2</sup>, S.Kohe<sup>1,2</sup>, N.Davies<sup>3</sup>, A.Peet<sup>1,2</sup>.

<sup>1</sup>Institute of Cancer and Genomic Sciences, University of Birmingham, Edgbaston, Birmingham,

<sup>2</sup>Birmingham Childrens Hospital, Birmingham. <sup>3</sup> University Hospitals Birmingham, Birmingham.

Although five-year survival for paediatric medulloblastoma is now approaching 75%, there remains a large variability in individual patient outcome that can be difficult to predict at diagnosis. It is becoming increasingly important to develop methods of stratifying patients for improved disease management. Tumour temperature has been proposed as a novel marker of prognosis in other tumours and can be measured non-invasively by the technique of magnetic resonance spectroscopy (MRS) thermometry. MRS thermometry has previously detected differences between childhood brain tumour types. In this single centre study we investigated its potential to predict survival in medulloblastoma. Pre-treatment MRS thermometry data were acquired for 40 histology confirmed medulloblastoma patients, 18 of which had died by the end of the study. Treatment was according to national guidelines or clinical trials and included surgery, followed by chemotherapy and/or radiotherapy. Histopathology reports and standard radiological imaging were available for all cases. Using Kaplan-Meier analysis with a median cut-off value, MRS thermometry identified patients with poor survival within the first 5 years ( $p < 0.01$ ). The mean apparent temperature in the poor prognosis group was  $0.7^{\circ}\text{C}$  higher than the good prognosis group, which may reflect increased metabolic rate in this group. No correlations were found between MRS thermometry and Chang stage, histological subtype or radiological features, which individually were not predictors of survival. Age and sex did not significantly influence patient survival. All reported myc status for the cohort were negative. MRS thermometry can provide a novel non-invasive prognostic marker for medulloblastoma and potentially aid treatment stratification.