

Exploring parents' screen-viewing behaviours and sedentary time in association with their attitudes toward their young child's screen-viewing

Solomon-Moore, Emma; Sebire, Simon J; Macdonald-Wallis, Corrie; Thompson, Janice; Lawlor, Debbie A; Jago, Russ

DOI:

[10.1016/j.pmedr.2017.06.011](https://doi.org/10.1016/j.pmedr.2017.06.011)

License:

Creative Commons: Attribution (CC BY)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Solomon-Moore, E, Sebire, SJ, Macdonald-Wallis, C, Thompson, J, Lawlor, DA & Jago, R 2017, 'Exploring parents' screen-viewing behaviours and sedentary time in association with their attitudes toward their young child's screen-viewing', *Preventive Medicine Reports*, vol. 7, pp. 198-205.
<https://doi.org/10.1016/j.pmedr.2017.06.011>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Checked for eligibility: 05/07/2017

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Exploring parents' screen-viewing behaviours and sedentary time in association with their attitudes toward their young child's screen-viewing

Authors:

Emma Solomon-Moore¹,

Simon J. Sebire¹,

Corrie Macdonald-Wallis¹,

Janice L. Thompson²,

Deborah, A. Lawlor^{3,4},

Russell Jago¹,

¹Centre for Exercise, Nutrition and Health Sciences, School for Policy Studies, University of Bristol, Bristol, UK.

²School of Sport, Exercise and Rehabilitation Sciences, University of Birmingham, Birmingham, UK.

³School for Social and Community Medicine, University of Bristol, Bristol, UK.

⁴MRC Integrative Epidemiology Unity, University of Bristol, Bristol, UK.

Corresponding author: Dr Emma Solomon-Moore, Centre for Exercise, Nutrition and Health Sciences, School for Policy Studies, 8 Priory Road, Bristol, BS8 1TZ, UK; 0117 3311099; Emma.Solomon@bristol.ac.uk.

ABSTRACT

Sedentary time and screen-viewing (SV) are associated with chronic disease risk in adults. Parent and child sedentary time and SV are associated. Parents influence children's SV through parenting styles and role modelling. Understanding whether parents' attitudes toward child SV are associated with their own SV and sedentary time will aid development of family interventions to reduce sedentary behaviours. Cross-sectional data with 809 parents from Bristol, UK were collected in 2012-2013 and analysed in 2016. Parental total sedentary time was derived from accelerometer data. Parents self-reported daily television viewing, use of computers, games consoles, and smartphone/tablets (none, 1-59 mins, 1-2 hrs, >2 hrs) and attitudes toward child SV. Adjusted linear and logistic regression models were used to examine associations, separately for weekdays and weekend days. Having negative attitudes toward child SV was associated with lower weekend sedentary time (Coeff: -6.41 [95% CI: -12.37 to -0.45] mins/day). Limiting behaviours and having negative attitudes toward child SV were associated with lower weekday television viewing (OR: 0.72 [0.57-0.90] and 0.57 [0.47-0.70] respectively), weekend television viewing (0.75 [0.59-0.95] and 0.61 [0.50-0.75]), and weekend computer use (0.73 [0.58-0.92] and 0.80 [0.66-0.97]). Negative attitudes were also associated with lower smartphone use on weekdays (0.70 [0.57-0.85]) and weekends (0.70 [0.58-0.86]). Parent self-efficacy for limiting child SV and setting SV rules were not associated with sedentary time or SV. Reporting negative attitudes toward child SV was associated with lower accelerometer-assessed weekend total sedentary time and self-reported SV behaviours, while limiting child SV was also associated with lower self-reported SV.

INTRODUCTION¹

Sedentary behaviours are defined as any waking behaviours characterised by an energy expenditure of ≤ 1.5 METS, where sitting or lying is the dominant mode of posture (e.g., screen-viewing (SV), motorised transport, office work).^{1,2} National data from England in 2012 suggest that adults spend approximately five hours daily being sedentary on both weekdays and weekend days.³ Moreover, half of English adults in 2012 spent two or more hours watching television (TV) or other screens daily, and a third watched TV for over three hours,⁴ with TV viewing the most prevalent leisure-time activity for UK adults in 2005.⁵

Sedentary time and SV (TV, computers, tablets, smartphones, video games) have been found to be associated with increased risk of obesity,⁶⁻⁹ cardiovascular disease,¹⁰⁻¹⁵ diabetes,⁸ cancer,¹⁶⁻¹⁷ all-cause mortality,^{10,12-14} mental disorders,⁴ and poor self-rated health⁴ in adults. A study of Finnish adults found that each additional self-reported daily TV hour was associated with a 1.81 ± 0.44 cm larger waist circumference in women and 2.0 ± 0.44 cm in men (reference category: < 1 hour; $p < 0.0001$).⁶ However, both cross-sectional and prospective studies in children and adults show little association between objectively-assessed time spent sedentary with adiposity or adverse cardio-metabolic health.¹⁸⁻²¹ This lack of association suggests that reporting bias may explain some of the associations with adverse outcomes seen in studies that only use self-report. An alternative explanation may be that SV is more strongly associated with negative health, for example due to an increase in snack consumption during SV,²² with measures of SV currently relying on self-reported data because objective SV measures for use in population studies do not exist. While some sedentary activities are associated with positive educational, mental and social

¹ Abbreviations: SV= screen-viewing, TV= television, IMD= indices of multiple deprivation

benefits (e.g., reading, connecting with loved ones, imaginative play),²³ the links with adverse health outcomes, at least from self-reported data, cannot be ignored. As such, there is a need to develop effective interventions to reduce SV and sedentary time for the whole family. While reductions in sedentary time at work are desirable, it is more likely that major reductions in sedentary behaviour will come from addressing leisure-time behaviours, such as SV, and shifts toward more active travel.¹

To develop effective interventions to reduce SV and sedentary time among families, we must first understand how parent and child sedentary behaviours are associated, and how parents can influence their child's behaviours. Parent TV-viewing time has been found to be strongly associated with child TV-viewing across the week.^{24,25} Parents who report low restriction of sedentary activities, low self-efficacy, and permissive parenting styles have children with greater levels of SV on average.^{26,27} Findings from a previous study using the B-Proact1v dataset, found parental self-efficacy to limit child SV was associated with child weekday TV-viewing and mediated associations between parental control and child SV.²⁸ Beyond these observational studies, a RCT of a school-based intervention aimed at improving 9-10 year olds' physical activity and diet, reduced child-reported SV (though not their accelerometer-assessed sedentary behaviour or any of the primary outcomes) and this effect appeared to be mediated by an effect on child-reported maternal limitation of SV.^{29,30}

These studies demonstrate that associations exist between parent and child SV time, and that parenting styles and preference for limiting child SV are associated with child SV. However, it is yet unknown whether parents' attitudes toward their child's SV are associated with their own SV

and sedentary time. For instance, if parents who report more negative attitudes toward their child's SV also report less SV and spend less time being sedentary themselves, there is potential to develop interventions to encourage parents to have negative attitudes toward their child's SV with the aim of reducing both parent and child SV and sedentary time. Therefore, it is important to understand which aspects of parents' attitudes toward child SV (e.g., self-efficacy for limiting SV, preference for limiting SV, negative attitudes toward SV, setting rules about SV) are associated to parents' own SV and sedentary behaviour.

The aim of this study was to examine whether parents' attitudes toward their young child's SV behaviour was associated with their (the parents) objectively-assessed total sedentary time and self-reported SV behaviours. Specifically, it is hypothesised that parents with a more restrictive attitude toward their young child's SV (i.e., higher preference and efficacy for limiting child SV, more rules and negative attitudes towards SV) would engage in less accelerometer-assessed sedentary time and self-reported SV themselves.

METHODS

Study sample

Data are from the cross-sectional B-Proact1v study, which aimed to identify factors associated with young children's (5-6 years) and parents' physical activity and SV. Details of the study design have been reported previously.³¹ Between February 2012 and May 2013, data were collected from 57 primary schools in the greater Bristol area. In total, 1267 child-parent dyads wore and returned an accelerometer and were included in the final dataset. For the current study, we were interested in parent objectively-assessed sedentary time and self-reported SV

behaviours, and therefore only parents that both wore and returned an accelerometer and completed all the SV measures were included in the analyses (n=809). Figure 1 shows the study flow of participants. Ethical approval was granted by the School for Policy Studies research ethics committee at the University of Bristol, and written informed consent was obtained for all participants.³²

Figure 1. *Study flow of participants*

Measures

Sedentary time

Participants were asked to wear an ActiGraph GT3X waist-worn accelerometer for five consecutive days, including two weekend days, during all waking hours. Data were recorded in 10-second epochs, and uniaxial data were processed using Kinesoft (v3.3.75; Kinesoft, Saskatchewan, Canada). Accelerometer data were considered valid if participants provided at least two weekdays and one weekend day of at least 500 minutes of data. Three days of monitoring have previously been demonstrated to produce reliable estimates of sedentary time in adults.³³ Accelerometer “non-wear” time was defined as periods of ≥ 60 minutes of consecutive zero values, with an allowance of up to 2 minutes of interruptions, and were removed from analyses.³⁴ Sedentary time was determined from accelerometer data using a threshold of < 100 counts per minute.³⁵ Total sedentary time, including both work and leisure time, was analysed separately for weekdays and weekend days. A previous study by Clemes et al. found that objectively-assessed sedentary time was higher on workdays than non-workdays.³⁶

Self-report measures

Parents completed a questionnaire about family characteristics, personal demographics, health aspirations, home media environment, SV time, and their attitudes towards their child's SV behaviour. The Index of Multiple Deprivation (IMD) scores, based upon the English Indices of Deprivation (<http://data.gov.uk/dataset/index-of-multiple-deprivation>), were assigned to each family based on their reported home postcode. Home media environment was assessed by parents indicating how many of each of 10 media devices they have within the home ('TV', 'DVD player', 'digital TV recorder', 'music player', 'desktop computer', 'laptop computer', 'tablet computer', 'games console', 'smartphone', 'handheld console'). The number of devices were summed to create a single score. Health aspirations were assessed on a five-item scale ('to be physically active'; 'to feel good about my level of physical fitness'; 'to keep myself healthy and well'; 'to be relatively free from sickness'; 'to have a physically healthy lifestyle'), where parents indicated the importance of each factor using a seven-point Likert scale, from 1 '*not at all*', through 4 '*moderately*', to 7 '*very*'.^{37,38} Responses were combined and the mean score used in analyses. Parent SV time was assessed via separate questions for the following SV devices: TVs, computers/laptops, games consoles, and smartphones/tablets (except for the time spent talking or texting). For each device, parents reported the time they spent using it outside of work for; a) a normal weekday, and b) a normal weekend day, with response options: '*none*'; '*1-30 minutes*'; '*31 minutes – 1 hour*'; '*1-2 hours*'; '*2-3 hours*'; '*3-4 hours*'; and '*4 hours or more*'. This method of self-reporting SV time has previously been used to assess SV in parents and children.^{26,39,40} A review found that self-reported measures of sedentary time generally showed moderate-to-high correlations for test-retest reliability and that validity correlations were higher in domain-specific measures (e.g., TV viewing, computer use) than for overall sedentary

measures across an entire day.⁴¹ Weekday and weekend SV were assessed independently due to previous evidence that suggests parents report greater SV on weekends than weekdays.⁴²

Parents' self-efficacy to limit their child's SV was assessed via three items (how much can you do to; a) 'control the time your child spends screen-viewing'; b) 'help your children have alternatives to screen-viewing'; c) 'reduce the time your child spends screen-viewing'), using a five-point Likert scale ranging from 1 '*nothing*' to 5 '*a great deal*', adapted from Bandura's Self Efficacy Scale.⁴³ Parents' preference for limiting their child's SV time was measured via three items (I limit how long my child; a) 'plays video games'; b) 'can watch TV and DVDs each day'; c) 'can use the computer for things other than homework'), using a four-point Likert scale ranging from 1 '*strongly disagree*' to 4 '*strongly agree*'.⁴⁴ Parents' attitudes towards their child's SV were measured by asking their perspective on two statements 'children spending several hours per day watching television or playing video games' and 'children spending several hours per day during leisure time using a computer or surfing the Internet' by using four 5-point Likert scales (1-5) with anchor points: '*beneficial=>harmful*,' '*healthy=>unhealthy*,' '*useful=>of no use*,' and '*of no concern=>of concern*'.⁴⁵ Parental rules governing children's SV activities were determined by asking: 'limiting my child's amount of TV viewing, games console or computer use time is' (response options: 1 '*necessary*' to 5 '*unnecessary*'), and 'I let my child decide how much TV he/she watches' (response options: 1 '*never*' to 5 '*always*').⁴⁵ For each of the four SV exposure variables, responses to items were combined and mean scores used for analyses.

Statistical analysis

Distributions of exposures, outcomes and co-variables were compared between participants included in this study and those who were excluded because of key missing data (e.g., not wearing or not having sufficient valid days of accelerometer data) using means, proportions and Chi Square statistics. To explore associations between objectively-assessed total sedentary time and index of multiple deprivation, parents' health aspirations, and home media environment Spearman's correlation coefficients were used. For the associations between SV behaviours and demographic variables means and one-way ANOVA statistics were used. The vast majority of parents did not use a games console on weekdays or weekend days (>90% and 83%, respectively), therefore this behaviour was not included in further analyses. The four exposure variables (self-efficacy for limiting child SV, preference for limiting child SV, negative attitudes towards child SV, and rules about child SV) were treated as continuous variables in all analyses. The responses to the 'rules about child SV' variable were flipped so that higher scores represented more restrictive parenting practices in line with the other exposure variables.

Two of the outcome variables (accelerometer-assessed total sedentary time on weekdays and weekend days) were continuous, as such multivariable linear regression models were used to examine the associations of the four exposure variables with these two sedentary time outcomes. Participant responses to the other six outcome variables (TV viewing, computer use, and smartphone/tablet use on weekdays and weekend days) were collapsed into four time categories: 'none', '1-59 minutes', '1-2 hours' and '2 hours or more'. As these variables were ordinal, multivariable ordered logistic regression models were used to examine the associations of the four exposure variables with each outcome variable. Ordered logistic regression assumes that the coefficients that describe the relationship between, for example, the lowest versus all higher

<u>Included (N=809)</u>	Excluded
-------------------------	----------

categories of the response variable are the same as those that describe the relationship between the next lowest category and all higher categories, known as the proportional odds assumption. To test for proportional odds, likelihood-ratio tests were conducted, and the margins command used. Any models that violated this assumption were analysed separately using generalised ordered logistic regression analyses.⁴⁶

To take account of the parents being recruited via schools, robust standard errors were used.

Adjusted models were adjusted for gender, IMD score, home media environment and health aspirations, as these have previously been associated with sedentary behaviours in adults.⁴⁷⁻⁴⁹

Adjusted linear regression models with total sedentary time were also adjusted for accelerometer wear time. All analyses were performed in Stata version 14.0.⁵⁰

RESULTS

Descriptive statistics are shown in Table 1 for participants included and excluded from the dataset. Parents excluded due to missing data were more likely to be deprived, have lower health aspirations, spend less time sedentary, use computers less but smartphones/tablets more, and have lower self-efficacy for limiting child SV. Mean accelerometer wear-time for parents was 801.9 (SD=101.0) minutes on weekdays and 745.2 (111.6) minutes on weekend days.

	Mean (SD) or %	N	Mean (SD) or %	<i>p</i>
Parent gender (% mothers)	74.4%	427	79.4%	0.05
Index of Multiple Deprivation ^a	13.2 (11.1)	359	18.9 (15.3)	<0.001
Health aspirations	5.9 (1.0)	243	5.6 (1.2)	0.002
Number of media devices	10.8 (4.6)	231	10.9 (4.5)	0.85
Accelerometer-assessed total weekday sedentary time (mins/day)	542.6 (91.7)	337	490.0 (101.8)	<0.001
Accelerometer-assessed total weekend sedentary time (mins/day)	497.6 (94.1)	207	453.4 (101.6)	<0.001
Weekday television viewing		265		0.09
<i>None</i>	3.5%		2.3%	
<i>1-59 minutes</i>	27.7%		22.6%	
<i>1-2 hours</i>	41.0%		40.0%	
<i>2+ hours</i>	27.8%		35.1%	
Weekend television viewing		262		0.05
<i>None</i>	2.2%		3.1%	
<i>1-59 minutes</i>	13.6%		9.2%	
<i>1-2 hours</i>	34.0%		29.0%	
<i>2+ hours</i>	50.2%		58.8%	
Weekday leisure computer use		263		0.006
<i>None</i>	11.4%		19.8%	
<i>1-59 minutes</i>	48.2%		44.9%	
<i>1-2 hours</i>	16.8%		13.3%	
<i>2+ hours</i>	23.6%		22.1%	
Weekend leisure computer use		256		0.001
<i>None</i>	17.3%		27.3%	
<i>1-59 minutes</i>	51.9%		42.2%	
<i>1-2 hours</i>	19.8%		16.0%	
<i>2+ hours</i>	11.0%		14.5%	
Weekday smartphone use		262		<0.001
<i>None</i>	36.6%		34.4%	
<i>1-59 minutes</i>	45.5%		41.6%	
<i>1-2 hours</i>	13.5%		10.7%	
<i>2+ hours</i>	4.5%		13.4%	
Weekend smartphone use		264		0.001
<i>None</i>	36.8%		33.0%	
<i>1-59 minutes</i>	42.3%		41.7%	
<i>1-2 hours</i>	13.8%		10.2%	
<i>2+ hours</i>	7.1%		15.2%	
Self-efficacy for limiting SV	4.6 (0.5)	258	4.5 (0.7)	0.001
Preference for limiting SV	3.4 (0.6)	251	3.3 (0.7)	0.06
Negative attitudes towards SV	3.8 (0.7)	191	3.7 (0.7)	0.15
Rules about SV	4.1 (0.8)	257	4.1 (0.9)	0.45

Table 1. Descriptive characteristics of the study sample (N=809)

^a Index of multiple deprivation: a higher value indicates greater deprivation.

Table 2 displays how the outcome variables vary across demographic variables. For instance, compared to mothers, fathers spent more time being sedentary, reported more computer use on weekdays, and used smartphones/tablets for longer across the week. Participants who spent more time watching TV on weekdays or using computers across the week had lower health aspirations, while participants who spent more time watching TV or using smartphones/tablets across the week had more home media devices on average.

	Index of Multiple Deprivation ^b		Health Aspirations ^b		Home media environment ^b	
	Mean (SD)	r or F (p)	Mean (SD)	r or F (p)	Mean (SD)	r or F (p)
Accelerometer-assessed total weekday sedentary time (mins/day)		-0.07 (0.06)		-0.01 (0.78)		-0.002 (0.94)
Accelerometer-assessed total weekend sedentary time (mins/day)		-0.01 (0.74)		-0.01 (0.74)		-0.02 (0.57)
Weekday television viewing		2.41 (0.07)		4.34 (0.005)		12.63 (<0.001)
None	9.9 (6.4)		6.1 (0.9)		8.1 (4.7)	
1-59 minutes	12.0 (9.5)		6.0 (0.9)		9.7 (4.3)	
1-2 hours	13.6 (12.0)		5.9 (0.9)		11.1 (4.5)	
2+ hours	14.2 (11.6)		5.7 (1.1)		11.8 (4.7)	
Weekend television viewing		0.92 (0.43)		1.83 (0.14)		11.55 (<0.001)
None	14.7 (10.3)		5.7 (1.1)		7.4 (4.3)	
1-59 minutes	12.7 (10.0)		6.0 (0.9)		9.5 (4.1)	
1-2 hours	12.5 (11.0)		5.9 (0.9)		10.4 (4.6)	
2+ hours	13.8 (11.5)		5.8 (1.0)		11.57 (4.5)	
Weekday leisure computer use		0.68 (0.56)		3.44 (0.02)		0.49 (0.69)
None	14.4 (12.3)		6.1 (0.8)		10.8 (4.6)	
1-59 minutes	13.1 (11.1)		5.9 (1.0)		10.6 (4.5)	
1-2 hours	13.7 (10.7)		5.7 (1.0)		11.1 (4.6)	
2+ hours	12.6 (10.9)		5.8 (0.9)		10.9 (4.7)	
Weekend leisure computer use		0.68 (0.57)		4.31 (0.005)		0.97 (0.41)
None	13.4 (10.9)		6.1 (0.8)		11.0 (4.5)	
1-59 minutes	12.9 (11.3)		5.8 (1.0)		10.8 (4.4)	
1-2 hours	13.1 (10.0)		5.8 (1.0)		10.4 (4.4)	
2+ hours	14.7 (12.5)		5.7 (0.9)		11.4 (5.6)	
Weekday smartphone use		0.58 (0.63)		1.79 (0.15)		19.41 (<0.001)
None	13.3 (10.9)		5.9 (1.0)		9.3 (4.3)	
1-59 minutes	13.2 (11.9)		5.9 (1.0)		11.5 (4.5)	
1-2 hours	12.3 (9.1)		5.7 (0.9)		12.2 (4.4)	
2+ hours	15.1 (10.5)		5.7 (1.0)		12.2 (4.5)	
Weekend smartphone use		0.17 (0.91)		1.38 (0.25)		22.28 (<0.001)
None	13.4 (11.1)		5.9 (1.0)		9.2 (4.3)	
1-59 minutes	12.9 (11.7)		5.9 (1.0)		11.4 (4.4)	
1-2 hours	13.6 (10.6)		5.7 (1.0)		12.2 (4.7)	
2+ hours	12.9 (9.2)		5.7 (0.9)		12.6 (4.1)	

Table 2. Intercorrelations, means, proportions, and ANOVA statistics for the study outcome variables and adjustment variables

^a Mean sedentary minutes and T-test statistics presented for continuous outcome variables, proportions in each outcome category and X² statistics presented for categorical outcome variables.

^b Intercorrelations presented for continuous outcome variables, and proportions in each outcome category and F-test statistics presented for categorical outcome variables.

Table 3 presents the associations between parents' attitudes toward their child's SV and their own accelerometer-assessed sedentary time. Having negative attitudes toward child SV was associated with a reduction in parents' weekend accelerometer-assessed total sedentary time, but there were no clear associations between the other three exposure variables and weekend sedentary time, and nor were there associations between any of the four exposures and weekday accelerometer-assessed total sedentary time.

	Unadjusted		Fully adjusted ^a	
	<i>Difference in mean sedentary time per 1 unit of each exposure [95% CI]</i>	<i>p</i>	<i>Difference in mean sedentary time per 1 unit of each exposure [95% CI]</i>	<i>p</i>
Accelerometer-assessed total weekday sedentary time (mins/day)				
Self-efficacy for limiting SV	11.80 [-0.01 to 23.61]	0.05	6.64 [-1.38 to 14.66]	0.10
Preference for limiting SV	3.31 [-7.39 to 14.01]	0.54	-0.71 [-8.01 to 6.60]	0.85
Negative attitudes towards SV	-5.08 [-14.32 to 4.15]	0.28	-2.48 [-8.69 to 3.72]	0.43
Rules about SV	1.16 [-6.81 to 9.12]	0.78	-0.33 [-5.70 to 5.04]	0.90
Accelerometer-assessed total weekend sedentary time (mins/day)				
Self-efficacy for limiting SV	5.68 [-6.47 to 17.84]	0.36	3.86 [-3.86 to 11.58]	0.33
Preference for limiting SV	5.48 [-5.48 to 16.45]	0.33	-0.41 [-7.45 to 6.62]	0.91
Negative attitudes towards SV	-8.50 [-17.95 to 0.94]	0.08	-6.41 [-12.37 to -0.45]	0.04
Rules about SV	-1.43 [-9.59 to 6.74]	0.73	-1.48 [-6.63 to 3.68]	0.57

Table 3. Linear regression analyses showing associations between parents' attitudes toward child screen-viewing and their sedentary time

^a Adjusted for parent gender, index of multiple deprivation score, health aspirations, home media environment, and accelerometer wear-time on weekdays and weekend days respectively.

^b The coefficients represent a per unit increase in the scores for each of the SV exposure variables. Categories for each of the SV variables were: None, 0-59 minutes, 1-2 hours, >2 hours.

All analyses take account of clustering at the school level by using robust standard errors.

The associations between parents' attitudes toward their child's SV and their own SV behaviours are presented in Table 4. Parental self-report of having a preference for limiting child SV and having negative attitudes towards their child's SV were both associated with lower levels of reported weekday and weekend TV viewing, and lower levels of weekend computer use; negative attitudes towards child SV were also associated with lower levels of reported weekday

and weekend smartphone/tablet use. Parental report of self-efficacy for limiting their child's SV and setting rules for child SV were not associated with parents' report of their own SV.

	Unadjusted		Fully adjusted ^a	
	<i>OR for an increase in the level of the SV outcome variables per 1 unit of each exposure [95% CI]</i>	<i>p</i>	<i>OR for an increase in the level of the SV outcome variables per 1 unit of each exposure [95% CI]</i>	<i>p</i>
Weekday television viewing				
Self-efficacy for limiting SV	0.91 [0.71 to 1.15]	0.43	1.03 [0.81 to 1.32]	0.80
Preference for limiting SV	0.63 [0.50 to 0.79]	<0.001	0.72 [0.57 to 0.90]	0.005
Negative attitudes towards SV	0.54 [0.45 to 0.66]	<0.001	0.57 [0.47 to 0.70]	<0.001
Rules about SV	0.85 [0.72 to 1.01]	0.06	0.93 [0.78 to 1.10]	0.39
Weekend television viewing				
Self-efficacy for limiting SV	0.95 [0.74 to 1.22]	0.68	1.03 [0.80 to 1.34]	0.80
Preference for limiting SV	0.67 [0.53 to 0.84]	0.001	0.75 [0.59 to 0.95]	0.02
Negative attitudes towards SV	0.58 [0.48 to 0.71]	<0.001	0.61 [0.50 to 0.75]	<0.001
Rules about SV	0.84 [0.71 to 0.99]	0.05	0.91 [0.76 to 1.09]	0.31
Weekday leisure computer use				
Self-efficacy for limiting SV	0.95 [0.74 to 1.21]	0.67	1.02 [0.79 to 1.31]	0.89
Preference for limiting SV	0.84 [0.68 to 1.05]	0.12	0.90 [0.72 to 1.13]	0.37
Negative attitudes towards SV	0.83 [0.69 to 1.00]	0.05	0.87 [0.72 to 1.05]	0.15
Rules about SV	0.95 [0.81 to 1.12]	0.56	0.98 [0.83 to 1.16]	0.84
Weekend leisure computer use				
Self-efficacy for limiting SV	0.78 [0.61 to 1.00]	0.05	0.87 [0.67 to 1.11]	0.26
Preference for limiting SV	0.68 [0.55 to 0.86]	0.001	0.73 [0.58 to 0.92]	0.009
Negative attitudes towards SV	0.75 [0.62 to 0.91]	0.003	0.80 [0.66 to 0.97]	0.02
Rules about SV	0.87 [0.73 to 1.02]	0.09	0.88 [0.75 to 1.05]	0.16
Weekday smartphone/tablet use				
Self-efficacy for limiting SV	1.05 [0.81 to 1.34]	0.73	1.14 [0.88 to 1.48]	0.33
Preference for limiting SV	0.77 [0.62 to 0.96]	0.02	0.89 [0.70 to 1.12]	0.31
Negative attitudes towards SV	0.67 [0.55 to 0.81]	<0.001	0.70 [0.57 to 0.85]	<0.001
Rules about SV	0.98 [0.83 to 1.15]	0.77	1.09 [0.92 to 1.29]	0.31
Weekend smartphone/tablet use				
Self-efficacy for limiting SV	1.03 [0.80 to 1.32]	0.82	1.12 [0.86 to 1.45]	0.40
Preference for limiting SV	0.80 [0.64 to 0.99]	0.05	0.93 [0.74 to 1.17]	0.53
Negative attitudes towards SV	0.67 [0.55 to 0.81]	<0.001	0.70 [0.58 to 0.86]	0.001
Rules about SV	0.97 [0.82 to 1.14]	0.69	1.09 [0.92 to 1.29]	0.32

Table 4. Ordered logistic regression showing associations between parents' attitudes toward child screen-viewing and their SV behaviour

^a Adjusted for parent gender, index of multiple deprivation score, health aspirations and home media environment.

^b The odds ratios represent the multiplicative change in the odds of belonging to a higher category of SV associated with a unit increase in each of the attitudes to SV variables. Categories for each of the SV variables were: None, 0-59 minutes, 1-2 hours, >2 hours.

All analyses take account of clustering at the school level by using robust standard errors.

Testing of proportional odds assumption

Five models violated the proportional odds assumption (weekend television viewing with negative attitudes towards SV; weekday smartphone use with preference for limiting SV; weekday smartphone use with negative attitudes toward SV; weekend smartphone use with self-efficacy for limiting SV; weekend smartphone use with negative attitudes toward SV), and thus the generalised ordered logistic regression results are presented in Table 5. For the majority of category-specific odds ratios, the associations were in the same direction as the main analysis. For parental report of negative attitudes toward child SV no associations were present with reported weekend television viewing for more than one hour compared with less than one hour, and with using smartphones for more than one minute during the week and weekend compared with no use. There was an inverse association between reported preference for limiting child SV and using a smartphone/tablet for more than one hour on a weekday, compared to less than one hour. Similarly, self-efficacy for limiting SV was inversely associated with using a smartphone/tablet for more than two hours on a weekend day, compared to using a smartphone for less than two hours.

	<i>Unadjusted</i>		<i>Fully adjusted^a</i>	
	<i>OR for each level of the SV outcome variables per 1 unit the exposure variable [95% CI]</i>	<i>p</i>	<i>OR for each level of the SV outcome variables per 1 unit the exposure variable [95% CI]</i>	<i>p</i>
Weekend television viewing & negative attitudes towards SV				
≥1 minute vs. None	0.26 [0.11 to 0.58]	0.001	0.27 [0.12 to 0.61]	0.002
≥1 hour vs. <1 hour	0.75 [0.56 to 0.99]	0.04	0.79 [0.59 to 1.05]	0.11
>2 hours vs. ≤2 hours	0.54 [0.44 to 0.67]	<0.001	0.57 [0.46 to 0.71]	<0.001
Weekday smartphone use & preference for limiting SV^b				
≥1 minute vs. None			1.01 [0.78 to 1.30]	0.95
≥1 hour vs. <1 hour			0.71 [0.52 to 0.97]	0.03
>2 hours vs. ≤2 hours			0.84 [0.47 to 1.51]	0.56
Weekday smartphone use & negative attitudes towards SV				
≥1 minute vs. None	0.78 [0.63 to 0.96]	0.02	0.83 [0.67 to 1.04]	0.10
≥1 hour vs. <1 hour	0.56 [0.43 to 0.72]	<0.001	0.58 [0.44 to 0.76]	<0.001
>2 hours vs. ≤2 hours	0.32 [0.20 to 0.52]	<0.001	0.33 [0.20 to 0.54]	<0.001
Weekend smartphone use & self-efficacy for limiting SV				
≥1 minute vs. None	1.14 [0.87 to 1.48]	0.34	1.25 [0.94 to 1.66]	0.13
≥1 hour vs. <1 hour	0.92 [0.67 to 1.27]	0.62	1.01 [0.72 to 1.43]	0.94
>2 hours vs. ≤2 hours	0.62 [0.43 to 0.89]	0.01	0.67 [0.45 to 0.99]	0.04
Weekend smartphone use & negative attitudes towards SV				
≥1 minute vs. None	0.82 [0.66 to 1.01]	0.06	0.88 [0.71 to 1.10]	0.27
≥1 hour vs. <1 hour	0.55 [0.42 to 0.70]	<0.001	0.57 [0.44 to 0.74]	<0.001
>2 hours vs. ≤2 hours	0.35 [0.24 to 0.51]	<0.001	0.36 [0.25 to 0.53]	<0.001

Table 5. Generalised ordered logistic regression analyses for the variables that violated the proportional odds assumption.

^a Adjusted for parent gender, index of multiple deprivation score, health aspirations, and home media environment.

^b Unadjusted analyses for weekday smartphone use and preference for limiting SV did not violate the proportional odds assumption.

DISCUSSION

Parents who reported more negative attitudes toward their child's SV spent less time being sedentary on weekend days, but not on weekdays. One potential explanation for the null finding on weekdays is that parents with high sedentary time may be engaged in sedentary work, which could be indicative of higher levels of education, and thus confound the association between their attitudes toward child SV and their own weekday sedentary time. As sedentary time was measured via accelerometers, and parents were not asked to report their work hours or occupation, it is not possible to know what activities parents engaged in while being sedentary.

Parents, who had greater preferences for limiting child SV and more negative attitudes toward it, reportedly watched less TV throughout the week and used computers less on weekends.

Additionally, parents with more negative attitudes toward child SV used their smartphone/tablet for less time across the week. The null finding for weekday computer use may be explained by the growing popularity of portable SV devices (tablets/smartphones) and thus computers may be more commonly used for more necessary tasks that may be less influenced by attitude beliefs.⁵¹

This is the first study to compare parents' attitudes towards child SV with parents' own sedentary time and SV behaviour, and so it was important to understand whether parents were adopting a 'Do as I say, not as I do' approach to parenting, or whether they also practice what they preach. Previous studies found that parents who place greater limitations on child SV also reported lower levels of child SV,²⁶ therefore, it seems logical that similar associations would exist with parent SV behaviour, given that parent and child SV are associated.^{24,25} It may be that permissive parents do not limit their child's SV behaviour because they are unwilling to cut down their own SV or sedentary time, or because they are not concerned about SV, while more authoritative parents may engage in less SV and sedentary time themselves in order to role-model 'healthy behaviours' for their child, or because they have negative attitudes toward SV in regards to their own health, and thus have similar attitudes toward SV for their child.

This study found no association between parents' self-efficacy to limit their child's SV or setting SV rules with either parents' own sedentary time or self-reported SV. It is plausible that some

parents felt confident limiting child SV, while not even considering their own SV or sedentary time to be an issue (cognitive dissonance).⁵² It is recommended that future studies explore this association further to examine whether parents' confidence for limiting child SV is associated with their concern and/or awareness of their own behaviours.

Five of the models assessing parents' attitudes with self-reported SV violated the proportional odds assumption, therefore, generalised ordered logistic regression analyses were conducted to provide a more comprehensive model of how associations differed across levels of SV.⁵³ For instance, parents' preference for limiting child SV was associated with lower weekday smartphone use for parents who used their smartphone for at least an hour per day (compared to less than an hour), a finding that was not present in the ordered logistic models. Similarly, parents' self-efficacy for limiting child SV was associated with lower reported weekend smartphone use for parents who used their smartphone for more than two hours per day (compared to less than two hours). These findings demonstrate that associations between these self-reported variables are complicated, and that more advanced models, such as generalised ordered logistic regression models, are necessary to tease out the differences across outcome levels.

The Family Ecological Model illustrates the processes by which parents influence children's diet, activity, and SV behaviours,⁵⁴ however other studies have shown that reciprocal reinforcing relationships exist among family members, and children can influence the health behaviours of their parents.⁵⁵⁻⁵⁸ As such, the family can be a mutually reinforcing environment in which healthy behaviours can be introduced, accepted, and maintained.^{59,60} Therefore, more family

models are needed that account for the complexities of the reciprocal relationship between parent and child health behaviours.

The findings in this study suggest that interventions to educate parents on the ill-effects of SV in order to instil negative attitudes towards child SV and limits for such behaviours could be a potential strategy to reduce both child and parent SV and sedentary time. Indeed, interventions to reduce sedentary behaviours in young people are more likely to be effective if they involve a family component;⁶¹ therefore more family-based interventions to reduce SV and sedentary time are needed. One example of an intervention that successfully reduced sedentary behaviours was PACE+; a primary care-based goal-setting and counselling intervention for adolescents in the United States.⁶² Parents were educated to encourage behaviour change attempts through active support, positive role modelling and praise. Self-reported sedentary time decreased from baseline to one-year follow-up to a greater extent in intervention participants versus control (-77.7 min/day; 95% CI: -105.8 to -49.5).⁶² Therefore, a key target for future research would be to conduct similar interventions encouraging parents to have more negative attitudes toward their child's SV, to limit such behaviours, and be positive SV role models for their child.

Strengths and limitations

The strengths of this study are the availability of data from a reasonably-sized sample of parents, including both mothers and fathers, and that we collected data on both self-reported SV and objectively-assessed sedentary time across both weekdays and weekend days. This, in combination with questionnaire data on family demographics, parenting styles and attitudes

towards child SV allows the dataset to make a novel contribution to the literature. Limitations of the study include its cross-sectional nature so causality could not be examined. ActiGraph accelerometers are waist-worn, thus are unable to distinguish between sitting and standing still, therefore devices that utilise a thigh placement would be more accurate at recording key markers of sedentary behaviour (e.g., sitting or lying posture). 458 participants were excluded from the study due to missing data (N=458), which may have resulted in sampling bias, because these participants differed from included participants in terms of their time spent sedentary, use of screen devices and self-efficacy. The SV measures were self-reported, because there are no objective measures of SV available for use in large cohort studies, however this does mean that reporting bias may explain some of the study findings, where parents who reported more negative attitudes towards their child's SV may have also felt obliged to report less SV behaviour for themselves (irrespective of their actual behaviour).⁶³ Additionally, the ordinal nature of the SV behaviour questionnaire enabled participants to report behaviours easily, however this also necessitated the use of more complex statistical analyses with less interpretable coefficients than a standard linear or logistic regression model. It also meant that it was not possible to calculate a combined SV score.

CONCLUSIONS

Parental report of placing limitations on child SV and having negative attitudes towards it were associated with lower levels of reported TV viewing and weekend computer use among parents. Having negative attitudes towards child SV were also associated with lower levels of smartphone/tablet use and weekend total sedentary time among parents. However, parents' self-

efficacy for limiting child SV and setting SV rules were not associated with either self-reported SV or accelerometer-assessed sedentary time.

ACKNOWLEDGEMENTS

Data presented in this paper were funded by British Heart Foundation project grants (ref PG/11/51/28986 and SP/14/4/31123).

Conflicts of Interest: None

REFERENCES

1. The Sedentary Behaviour and Obesity Expert Working Group. Sedentary Behaviour and Obesity: Review of the Current Scientific Evidence. Department of Health, 2010.
2. Sedentary Behaviour Research Network. Letter to the Editor: Standardized use of the terms “sedentary” and “sedentary behaviours”. *Appl Physiol Nutr Metab*. 2012;37(3):540-542. <http://doi.org/10.1139/h2012-024>.
3. Health & Social Care Information Centre. Health Survey for England 2012: Health, social care and lifestyles. London, England: Health and Social Care Information Centre, 2013.
4. Shiue I. Modeling indoor TV/screen viewing and adult physical and mental health: Health Survey for England, 2012. *Environ Sci Pollut Res*. 2016;23(12):11708-11715. <http://dx.doi.org/10.1007/s11356-016-6354-5>.
5. Office for National Statistics. The Time Use Survey, 2005. How we spend our time. London: Office for National Statistics; 2006.

6. Blanck HM, McCullough ML, Patel AV, et al. Sedentary behavior, recreational physical activity, and 7-year weight gain among postmenopausal U.S. women. *Obesity*. 2007;15:1578-1588. <http://dx.doi.org/10.1038/oby.2007.187>.
7. Heinonen I, Helajärvi H, Pahkala K, et al. Sedentary behaviours and obesity in adults: the Cardiovascular Risk in Young Finns Study. *BMJ Open*. 2013;3:e002901. <http://dx.doi.org/10.1136/bmjopen-2013-002901>.
8. Hu F, Li TY, Colditz GA, et al. Television watching and other sedentary behaviours in relation to risk of obesity and type 2 diabetes mellitus in women. *JAMA*. 2003;289:1785-1791. <http://dx.doi.org/10.1001/jama.289.14.1785>.
9. Shields M, Tremblay MS. Sedentary behaviour and obesity. *Health Reports*. 2008;19:19-30.
10. Dunstan DW, Barr EL, Healy GN, et al. Television viewing time and mortality: the Australian Diabetes, Obesity and Lifestyle Study (AusDiab). *Circulation*. 2010;121(3):384-391. <http://dx.doi.org/10.1161/CIRCULATIONAHA.109.8944824>.
11. Ford ES, Caspersen CJ. Sedentary behaviour and cardiovascular disease: a review of prospective studies. *Int J Epidemiol*. 2012;1-16. <http://dx.doi.org/10.1093/ije/dys078>.
12. Inoue M, Iso H, Yamamoto S, et al. Daily total physical activity level and premature death in men and women: results from a large-scale population-based cohort study in Japan (JPHC study). *Annals of Epidemiology*. 2008;18:522-530. <http://dx.doi.org/10.1016/j.annepidem.2008.03.008>.
13. Katzmarzyk PT, Church T, Craig CL, et al. Sitting time and mortality from all causes, cardiovascular disease and cancer. *Med Sci Sports Exerc*. 2009;41:998-1005. <http://dx.doi.org/10.1249/MSS.0b13e3181930355>.

14. Stamatakis E, Hamer M, Dunstan DW. Screen-Based Entertainment Time, All-Cause Mortality, and Cardiovascular Events. *J Am Coll Cardiol.* 2011;57:292-299.
<http://dx.doi.org/10/1016/j.jacc.2010.05.065>.
15. Wijndaele K, Brage S, Besson H, et al. Television Viewing and Incident Cardiovascular Disease: Prospective Associations and Mediation Analysis in the EPIC Norfolk Study. *Plos One.* 2011;6(5):e20058. <http://dx.doi.org/10.1371/journal.pone.0020058>.
16. Friberg E, Mantzoros CS, Wolk, A. Physical activity and risk of endometrial cancer: A population-based prospective cohort study. *Cancer Epidemiology, Biomarkers and Prevention.* 2006;15:2136-2140. <http://dx.doi.org/10.1158/1055-9965.EPI-06-0465>.
17. Howard RA, Freedman DM, Park Y, et al. Physical activity, sedentary behaviour and the risk of rectal cancer in the NIH-AARP diet and health study. *Cancer Causes and Control.* 2008;19:939-953. <http://dx.doi.org/10.1007/s10552-008-9159-0>.
18. van Ekris E, Altenburg TM, Singh AS, et al. An evidence-update on the prospective relationship between childhood sedentary behaviour and biomedical health indicators: a systematic review and meta-analysis. *Obes Rev.* 2016;17:833-849.
<http://dx.doi.org/10.1111.obr.12426>.
19. Ekelund U, Luan J, Sherar LB, et al. Moderate to vigorous physical activity and sedentary time and cardiometabolic risk factors in children and adolescents. *JAMA.* 2012;307(7):704-712. <http://dx.doi.org/10.1001/jama.2012.156>.
20. Stamatakis E, Hamer M, Tilling K, et al. Sedentary time in relation to cardiometabolic risk factors in adults: Differential associations for self-reported versus accelerometry in working age adults. *Int J Epidemiol.* 2012;41:1328-1337.
<http://dx.doi.org/10.1093/ije/dys077>.

21. Stamatakis E, Coombs N, Tilling K, et al. Sedentary time in late childhood and cardiometabolic health in adolescence. *Pediatrics*. 2015;135:e1432-1441.
<http://dx.doi.org/10.1542/peds.2014-3750>.
22. Pearson N, Biddle SJH. Sedentary Behavior and Dietary Intake in Children, Adolescents, and Adults: A Systematic Review. *Am J Prev Med*. 2011;41(2):178-188.
<http://dx.doi.org/10.1016/j.amepre.2011.05.002>.
23. Jacobs JM, Hammerman-Rozenberg R, Cohen A, et al. Reading Daily Predicts Reduced Mortality Among Men From a Cohort of Community-Dwelling 70-Year-Olds. *J Gerontol B Psychol Sci Soc Sci*. 2008;63(2):S73-80.
24. Jago R, Stamatakis E, Gama A, et al. Parent and Child Screen-Viewing Time and Home Media Environment. *Am J Prev Med*. 2012;43(2):150-158.
<http://dx.doi.org/10.1016/j.amepre.2012.04.012>.
25. Jago R, Thompson JL, Sebire SJ, et al. Cross-sectional associations between the screen-time of parents and young children: differences by parent and child gender and day of the week. *Int J Behav Nutr Phys Act*. 2014;11:54. <http://dx.doi.org/10.1186/1479-5868-11-54>.
26. Jago R, Davison KK, Thompson JL, et al. Parental Sedentary Restriction, Maternal Parenting Style, and Television Viewing Among 10- to 11-Year-Olds. *Pediatr*. 2011;128:e572-578. <http://dx.doi.org/10.1542/peds.2010-3664>.
27. Smith BJ, Grunseit A, Hardy LL, et al. Parental influences on child physical activity and screen viewing time: a population based study. *BMC Public Health*. 2010;10:593.
<http://dx.doi.org/10.1186/1471-2458-10-593>.

28. Jago R, Wood L, Zahra L, et al. Parental Control, Nurturance, Self-Efficacy, and Screen Viewing among 5- to 6-Year-Old Children: A Cross-Sectional Mediation Analysis To Inform Potential Behavior Change Strategies. *Child Obes.* 2015;11(2):139-147.
<http://dx.doi.org/10.1089/chi.2014.0110>.
29. Kipping RR, Howe LD, Jago R, et al. Effect of intervention aimed at increasing physical activity, reducing sedentary behaviour, and increasing fruit and vegetable consumption in children: Active for Life Year 5 (AFLY5) school based cluster randomised controlled trial. *BMJ.* 2014;348:g3256. <http://dx.doi.org/10.1136/bmj.g3256>.
30. Lawlor DA, Howe LD, Anderson EL, et al. The Active for Life Year 5 (AFLY5) school-based cluster randomised controlled trial: effect on potential mediators. *BMC Public Health.* 2016;6(1):68. <http://dx.doi.org/10.1186/s12889-016-2734-5>.
31. Jago R, Sebire SJ, Wood L, et al. Associations between objectively assessed child and parental physical activity: a cross-sectional study of families with 5-6 year old children. *BMC Public Health.* 2014;14:655. <http://dx.doi.org/10.1186/1471-2458-14-655>.
32. Jago R, Bailey R. Ethics and paediatric exercise science: issues and making a submission to a local ethics and research committee. *J Sports Sci.* 2001;19(7):527-535.
<http://dx.doi.org/10.1080/026404101750238980>.
33. Dillon CB, Fitzgerald AP, Kearney PM, et al. Number of Days Required to Estimate Habitual Activity Using Wrist-Worn GENEActiv Accelerometer: A Cross-Sectional Study. *PLOS One.* 2016;11(5):e0109913.
<http://dx.doi.org/10.1371/journal.pone.0109913>.
34. Troiano RP, Berrigan D, Dodd KW, et al. Physical activity in the United States measured by accelerometer. *Med Sci Sports Exerc.* 2008;40(1):181-188.

35. Tudor-Locke C, Brashear MM, Johnson WD, et al. Accelerometer profiles of physical activity and inactivity in normal weight, overweight, and obese US men and women. *Int J Behav Nutr Phys Act.* 2010;7:60. <http://dx.doi.org/10.1186/1479-5868-7-60>.
36. Clemes SA, O'Connell SE, Edwardson CL. Office workers' objectively measured sedentary behaviour and physical activity during and outside working hours. *J Occup Environ Med.* 2014;56(3):298-303. <http://dx.doi.org/10.1097/JOM.0000000000000101>.
37. Kasser T, Ryan RM. A dark side of the American dream: Correlates of financial success as a central life aspiration. *J Pers Soc Psychol.* 1993;65:410-422.
<http://dx.doi.org/10.1037/0022-3514.65.2.410>.
38. Kasser T, Ryan RM. Further examining the American dream: Differential correlates of intrinsic and extrinsic goals. *Pers Soc Psychol B.* 1996;22:80-87.
<http://dx.doi.org/10.1177/0146167296223006>.
39. Jago RP, Fox KR, Page AS, et al. Parent and child physical activity and sedentary time: Do active parents foster active children? *BMC Public Health.* 2010;10;194.
<http://dx.doi.org/10.1186/1471-2458-10-194>.
40. Jago RP, Page AS, Froberg K, et al. Screen-viewing and the home TV environment: The European Youth Heart Study. *Prev Med.* 2008;47(5);525-529.
<https://doi.org/10.1016/j.ypmed.2008.07.016>.
41. Healy GN, Clark BK, Winkler EAH, et al. Measurement of Adults' Sedentary Time in Population-Based Studies. *Am J Prev Med.* 2011;41(2);216-227.
<http://doi.org/10.1016/j.amepre.2011.05.005>.

42. Sigmundova D, Sigmund E, Badura P, et al. Weekday-weekend patterns of physical activity and screen time in parents and their pre-schoolers. *BMC Public Health*. 2016;16(1):898. <http://doi.org/10.1186/s12889-016-3586-8>.
43. Bandura A. Guide for constructing self-efficacy scales. In: Parajes F, Urdan TC, eds. *Self-efficacy beliefs of adolescents*. USA: IAP- Information Age Publishing, 2006:307-337.
44. Davison KK, Li K, Baskin ML, et al. Measuring parental support for children's physical activity in white and African American parents: The Activity Support Scale for Multiple Groups (ACTS-MG). *Prev Med*. 2011;52:39-43. <http://dx.doi.org/10.1016/j.yjpm.2010.11.008>.
45. He M, Piche L, Beynon C, et al. Screen-related Sedentary Behaviors: Children's and Parents' Attitudes, Motivations, and Practices. *J Nutr Educ Behav* 2010;42:17-25. <http://dx.doi.org/10.1016/j.jneb.2008.11.011>.
46. Williams R. Generalized Ordered Logit/ Partial Proportional Odds Models for Ordinal Dependent Variables. *The Stata Journal*. 2006;6(1):58-82.
47. O'Donoghue G, Perchoux C, Mensah K, et al. A systematic review of correlates of sedentary behaviour in adults aged 18-65 years: a socio-ecological approach. *BMC Public Health*. 2016;16:163. <http://dx.doi.org/10.1186/212889-016-2841-3>.
48. Rhodes RE, Mark RS, Temmel CP. Adult Sedentary Behavior: A Systematic Review. *Am J Prev Med*. 2012;42(3):e3-e28. <http://dx.doi.org/10.1016/j.amepre.2011.10.020>.
49. Wood L, Jago R, Sebire SJ, et al. Sedentary time among spouses: a cross-sectional study exploring associations in sedentary time and behaviour in parents of 5 and 6 year old children. *BMC Res Notes*. 2015;8:787. doi: 10.1186/s13104-015-1758-8.

50. STATA Version 14. Statacorp. TX: College Station, 2011.
51. Ofcom. Adults' media use and attitudes: Report 2015. Ofcom, May 2015.
52. Festinger L, Carlsmith M. Cognitive consequences of forced compliance. *J Abnorm Soc Psychol.* 1959;58:203-210.
53. Williams R. Understanding and interpreting generalized ordered logit models. *J Math Sociol.* 2016;40(1):7-20. doi: 10.1080/0022250X.2015.1112384.
54. Davison K, Campbell K. Opportunities to prevent obesity in children within families: an ecological approach. In: Crawford D, Jeffery R, ed. *Obesity Prevention and Public Health.* Oxford: Oxford University Press; 2005:207-230.
55. Crockett SJ, Mullis RM, Perry CL. Parent nutrition education: a conceptual model. *J Sch Health.* 1988;58(2):53–57. <http://dx.doi.org/10.1111/j.1746-1561.1988.tb05822.x>.
56. Nader PR, Sallis JF, Patterson TL, et al. A family approach to cardiovascular risk reduction: results from the San Diego Family Health Project. *Health Educ Q.* 1989;16(2):229–244.
57. Perry CL, Crockett SJ, Pirie P. Influencing parental health behavior: implications of community assessments. *Health Educ.* 1987;18:68–77.
<http://dx.doi.org/10.1080/00970050.1987.10614520>.
58. Perry CL, Luepker RV, Murray DM, et al. Parent involvement with children's health promotion: the Minnesota home team. *Am J Public Health.* 1988;78(9):1156–1160.
59. Gruber KJ, Haldeman LA. Using the Family to Combat Childhood and Adult Obesity. *Prev Chronic Dis.* 2009;6(3):A106.

60. Wrotniak BH, Epstein LH, Paluch RA, et al. Parent weight change as a predictor of child weight change in family-based behavioral obesity treatment. *Arch Pediatr Adolesc Med*. 2004;158(4):342–347. <http://dx.doi.org/10.1001/archpedi.158.4.342>.
61. Biddle SJH, Petrolini I, Pearson N. Interventions designed to reduce sedentary behaviours in young people: a review of reviews. *Br J Sports Med*. 2014;48:182-186. <http://dx.doi.org/10.1136/bjsports-2013-093078>.
62. Patrick K, Calfas KJ, Norman GJ, et al. Randomized Controlled Trial of a Primary Care and Home-Based Intervention for Physical Activity and Nutrition Behaviors. *Arch Pediatr Adolesc Med*. 2006;160:128-136. <http://dx.doi.org/10.1001/archpedi.160.2.128>.
63. Matthews CE, Moore SC, George SM, et al. Improving Self-reports of Active and Sedentary Behaviors in Large Epidemiologic Studies. *Exerc Sport Sci Rev*. 2012;40(3):118-126.