UNIVERSITY^{OF} BIRMINGHAM University of Birmingham Research at Birmingham

Disentangling unclear nuclear breakup channels of beryllium-9 using the three-axis Dalitz plot

Smith, R.; Freer, M.; Wheldon, C.; Curtis, N.; Almaraz-Calderon, S.; Aprahamian, A.; Ashwood, N. I.; Barr, M.; Bucher, B.; Copp, P.; Couder, M.; Fang, X.; Goldring, G.; Jung, F.; Kokalova, Tz; Lesher, S. R.; Lu, W.; Malcolm, J. D.; Roberts, A.; Tan, W. P.

DOI: 10.1088/1742-6596/863/1/012032

License: Creative Commons: Attribution (CC BY)

Document Version Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Smith, R, Freer, M, Wheldon, C, Curtis, N, Almaraz-Calderon, S, Aprahamian, A, Ashwood, NI, Barr, M, Bucher, B, Copp, P, Couder, M, Fang, X, Goldring, G, Jung, F, Kokalova, T, Lesher, SR, Lu, W, Malcolm, JD, Roberts, A, Tan, WP & Ziman, VA 2017, 'Disentangling unclear nuclear breakup channels of beryllium-9 using the three-axis Dalitz plot', *Journal of Physics: Conference Series*, vol. 863, no. 1, 012032. https://doi.org/10.1088/1742-6596/863/1/012032

Link to publication on Research at Birmingham portal

Publisher Rights Statement: Checked for eligibility: 31072017

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

•Users may freely distribute the URL that is used to identify this publication.

•Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.

•User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?) •Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Home Search Collections Journals About Contact us My IOPscience

Disentangling unclear nuclear breakup channels of beryllium-9 using the three-axis Dalitz plot

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2017 J. Phys.: Conf. Ser. 863 012032

(http://iopscience.iop.org/1742-6596/863/1/012032)

View the table of contents for this issue, or go to the journal homepage for more

Download details:

IP Address: 147.188.254.76 This content was downloaded on 12/07/2017 at 17:40

Please note that terms and conditions apply.

You may also be interested in:

New double ionization channel uncovered in four-particle Dalitz plots M Schulz, M F Ciappina, T Kirchner et al.

Study of clustering structures through breakup reactions Pierre Capel

Decay dynamics of N2O under the impact of fast electrons Arnab Khan and Deepankar Misra

Three-axis MEMS Accelerometer for Structural Inspection E Barbin, A Koleda, T Nesterenko et al.

Pioneer of quark model dies Martin Griffiths

Ultrafast molecular dynamics of dissociative ionization in OCS probed by soft X-ray synchrotron radiation Ali Ramadhan, Benji Wales, Reza Karimi et al.

Systematic Analysis of Four-Particle Dalitz Plots for Double Ionization M Schulz, D Fischer, K Schneider et al.

Kinematically Complete studies on Mutual Projectile and Target Ionization X Wang, D Fischer, A Kelkar et al.

Disentangling unclear nuclear breakup channels of beryllium-9 using the three-axis Dalitz plot

R Smith¹, M Freer¹, C Wheldon¹, N Curtis¹, S Almaraz-Calderon², A Aprahamian², N I Ashwood¹, M Barr¹, B Bucher², P Copp³, M Couder², X Fang², G Goldring⁴, F Jung², Tz Kokalova¹, S R Lesher³, W Lu², J D Malcolm¹, A Roberts², W P Tan² and V A Ziman¹

¹School of Physics and Astronomy, University of Birmingham, Edgbaston, B15 2TT, UK. ²Institute for Structure and Nuclear Astrophysics, Department of Physics, University of Notre Dame, Notre Dame, IN 46556, USA.

³Department of Physics, University of Wisconsin - La Crosse, La Crosse, WI 54601, USA.

⁴Department of Particle Physics, Weizmann Institute, 76100 Rehovot, Israel.

E-mail: r.smith.30pgr.bham.ac.uk

Abstract. The three-axis Dalitz plot has been applied to the breakup of a nucleus into unequal mass fragments for the first time. The Dalitz plot allows clear identification of the various breakup channels of $^9\mathrm{Be} \to 2\alpha + n$ process. The method has allowed the branching ratio for the 6.38 MeV level in ⁹Be to be provisionally calculated when examining the ⁹Be(⁴He, α) $\alpha\alpha n$ reaction. The effects of non-uniform angular distributions on the Dalitz plot must still be properly investigated along with the effects of contaminant reaction channels. It is proposed that this method could be used to determine the breakup branching ratio of a newly-measured level in this nucleus.

1. Introduction

A ternary plot is a barycentric plot of three quantities, which sum to a constant value [1]. Using the ternary plot representation, the three quantities are visualised simultaneously by a twodimensional position on an equilateral triangle. While working at the University of Birmingham in 1953, R. H. Dalitz was the first person to apply the ternary plot technique to visualising the three-body decay of sub-atomic particles [2]. Therefore, the ternary plot will henceforth be referred to as a *Dalitz plot* in this context. In the centre of mass (CoM) of a decay $X \to a+b+c$, the energies of a, b and c must sum to the decay Q value (in the non-relativistic limit). The Dalitz plot technique is widely used in high energy physics [3].

Only in recent years has the technique been applied in nuclear physics. Such Dalitz plots are used to determine if the Hoyle state in ${}^{12}C$ decays through a ${}^{8}Be_{q.s.}$ intermediate level or through a direct three-body process [4]. Further, interference patterns in these Dalitz plots are used to determine the angular momentum and parity of ${}^{12}C$ resonances [5]. There is no reason why it cannot also be applied to breakups into unequal mass fragments such as ${}^{9}\text{Be} \rightarrow \alpha + \alpha + n$.

Measuring the breakup branching ratios for states in ⁹Be is important, since the breakup through particular channels is correlated with the strength of the corresponding cluster

Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI. Published under licence by IOP Publishing Ltd 1



Figure 1. Construction of the Dalitz plot for a decay of a nucleus into three particles. ϵ_i corresponds to the fractional energy of particle *i* in the CoM of the decaying system. (a) illustrates that if $\sum \epsilon_i$ = constant then points must lie on the shaded triangle. (b) shows the same plot from a different perspective and (c) illustrates how the allowed area changes from a circle to an ellipse for decays into unequal mass fragments (see text for details).

configurations in the wave functions, which can have implications for astrophysical reaction rates [6,7]. The $n + {}^{8}\text{Be}_{g.s}$, $n + {}^{8}\text{Be}_{2^+}$, $\alpha + {}^{5}\text{He}_{g.s}$ and $\alpha + \alpha + \alpha$ channels are expected to dominate for the low-lying levels. This paper demonstrates, through the use of Monte-Carlo simulations and experimental data, that the three-axis Dalitz plot can be applied to analyse the breakup systematics of the ${}^{9}\text{Be} \rightarrow \alpha + \alpha + n$ system.

2. The Dalitz plot

Figure 1 illustrates how to plot three quantities (which sum to a constant) on a single 2D plot. In this case, ϵ_i corresponds to the fractional energy of particle *i* in the CoM of the decaying system ($\epsilon_i = E_i / \sum_{j=1}^3 E_j$). Figure 1 (*a*) shows that by plotting three quantities in 3D cartesian coordinates, the resulting point must lie on the shaded triangle. By shifting the viewing perspective along a line normal to the plane of the triangle and passing through the origin, figure 1 (*b*) is extracted. A point on the triangle is uniquely defined by ϵ_1 , ϵ_2 and ϵ_3 which are the distances of each point from each edge of the triangle. If particles 1, 2 and 3 are of equal mass, conservation of linear momentum ensures that points can only lie within the circle of figure 1 (*c*). For unequal mass fragments, the circle becomes an inscribed ellipse. The points where the circle or ellipse touch the edge of the triangle are denoted by a, b and c and are fixed such that 2b:b3 equals $m_2:m_3$ (and cyclical for sides 12 and 31). The *x*, *y* coordinates of a point on this triangle are related to ϵ_i by $x = \sqrt{3}(\epsilon_2 - \epsilon_1)/2$ and $y = (2\epsilon_3 - \epsilon_2 - \epsilon_1)/2$.

3. Application to beryllium-9 breakup

An experiment was performed to measure the ${}^{9}\text{Be}({}^{4}\text{He},\alpha)\alpha\alpha n$ reaction, at 22 and 26 MeV beam energies (detailed in Ref. [8]). A new resonance was recorded at 3.8 MeV, and the necessity for reliable branching ratio measurements in order to determine the angular momentum of the state, was highlighted. The Dalitz plot method was applied to this experimental data in order to test if it was a valid method of calculating reliable branching ratios. The $7/2^{-}$ state at 6.38 MeV was chosen as a test subject because branching ratios are already well constrained [6]. The experimental analysis is similar to that of Ref. [8], however, the specific condition that decays proceeded via the ⁸Be ground state was not required. Hence, ⁹Be excitation spectra were reconstructed for events which break up via any of the open channels. Events in the excitation region between 6.1 and 6.7 MeV were selected to consider breakups of the 6.38 MeV level.

It was possible to select which final state α particles arise from the breakup of ⁹Be. The



Figure 2. Dalitz plots for the breakup of the 6.38 MeV level in ⁹Be. Darker regions indicate more counts. (a) depicts the experimental data within an excitation cut of $6.1 < E_x < 6.7$, (b) depicts a simulation of the ⁸Be_{g.s} (upper line) and ⁸Be₂₊ (lower line) channels and (c) shows the simulation of decays through the ⁵He_{g.s} channel. In the case of decays through ⁸Be_{g.s}/⁵He_{g.s}, the first emitted n/α takes the majority of the decay energy, hence the event concentrations in these regions (top centre/ bottom left and right, respectively). Sequential decays through the broad ⁸Be₂₊ state lead to a more even distribution of energies.

energy and momentum of the undetected neutron was calculated from momentum conservation. The momenta of the breakup particles were converted into the CoM frame of the decaying ⁹Be. Their fractional energies in this frame (ϵ_i) were used as the axes of a Dalitz plot. To force the Dalitz plot into a circular, rather than elliptical, shape, the axes were chosen to be ($4\epsilon_{\alpha_1}$, $4\epsilon_{\alpha_2}$, ϵ_n). The resulting Dalitz plot is shown in figure 2 (a). Simulations of the three decay processes with proper threshold dependencies, experimental resolution, but assuming isotropic particle emission, are shown in figures 2 (b) and 2 (c).

4. Conclusions

Figure 2 (a) shows contributions from all three sequential breakup channels as simulated in figures 2 (b) and 2 (c). Each decay contribution is well separated when compared with similar studies using other types of Dalitz plots [6]. One-dimensional projections of the Dalitz plot permitted a quantitative comparison between simulation and experiment. Provisional branching ratios were extracted which are comparable with previous measurements [6]. The effects of non-uniform angular correlations on each breakup stage and contaminant reaction channels need to be considered in order to be confident of these values. Lower levels in ⁹Be such as the new 3.8 MeV state are much closer to the decay thresholds and it is predicted that the signatures from each channel will strongly overlap on the Dalitz plot. Their exact manifestations will depend on the angular distributions and it remains to be seen how separable each channel will be.

- [1] Brazzle B and Tapp A 2016 The Physics Teacher 54(4) 216
- [2] Dalitz R H 1953 Philos. Mag. 44 1068
- [3] Alff C, Berley D, Colley D, Gelfand N, Nauenberg U, Miller D, Schultz J, Steinberger J, Tan T H, Brugger H, Kramer P and Plano R 1962 Phys. Rev. Lett. 9 325
- [4] Itoh M, Ando S, Aoki T, Arikawa H, Ezure S, Harada K, Hayamizu T, Inoue T, Ishikawa T, Kato K, Kawamura H, Sakemi Y and Uchiyama A 2014 Phys. Rev. Lett 113 102501
- [5] Kirsebom O S, Alcorta M, Borge M J G, Cubero M, Diget C A, Dominguez-Reyes R, Fraile L M, Fulton B R, Fynbo H O U, Hyldegaard S et al 2010 Phys. Rev. C 81 064313
- [6] Brown T A D, Papka P, Fulton B R, Watson D L, Fox S P, Groombridge D, Freer M, Clarke N M, Ashwood N I, Curtis N et al 2007 Phys. Rev. C 76 054605
- [7] Papka P, Brown T A D, Fulton B R, Watson D L, Fox S P, Groombridge D, Freer M, Clarke N M, Ashwood N I, Curtis N et al 2007 Phys. Rev. C 75 045803
- [8] Smith R, Wheldon C, Freer M, Curtis N, Kokalova Tz, Almaraz-Calderon S, Aprahamian A, Ashwood N I, Barr M, Bucher B, et al 2016 Phys. Rev. C 94 014320