UNIVERSITY^{OF} BIRMINGHAM University of Birmingham Research at Birmingham

Assessing urban population vulnerability and environmental risks across an urban area during heatwaves – Implications for health protection Macintyre, H. L.; Heaviside, Clare; Taylor, J.; Picetti, R; Symonds, P; Cai, Xiaoming; Vardoulakis, S

DOI: 10.1016/j.scitotenv.2017.08.062

License: Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version Peer reviewed version

Citation for published version (Harvard):

Macintyre, HL, Heaviside, C, Taylor, J, Picetti, R, Symonds, P, Cai, X & Vardoulakis, S 2018, 'Assessing urban population vulnerability and environmental risks across an urban area during heatwaves – Implications for health protection', *Science of the Total Environment*, vol. 610-11, pp. 678-690. https://doi.org/10.1016/j.scitotenv.2017.08.062

Link to publication on Research at Birmingham portal

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

•Users may freely distribute the URL that is used to identify this publication.

•Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.

•User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?) •Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

1

2

3

4

10

11

12

Assessing urban population vulnerability and environmental risks across an urban area during heatwaves – implications for health protection

H. L. Macintyre^a, C. Heaviside^{a,b,c}, J. Taylor^d, R. Picetti^b, P. Symonds^d, Xiaoming Cai^c, S. Vardoulakis^{b,c,e}

a. Chemical and Environmental Effects Department, Centre for Radiation Chemical and Environmental Hazards, Public Health England, Chilton, Oxon, OX11 0RQ, UK. b. London School of Hygiene and Tropical Medicine, Keppel Street, London WC1E 7HT, UK. c. School of Geography, Earth and Environmental Sciences, University of Birmingham, Edgbaston, Birmingham, B15 2TT,

UK.

d. University College London, Gower Street, Kings Cross, London WC1E 6BT, UK.

e. Institute of Occupational Medicine, Research Avenue North, Riccarton, Edinburgh, Midlothian, EH14 4AP, UK.

Correspondence: <u>Helen.Macintyre@phe.gov.uk</u>

13 ABSTRACT

14 Heatwaves can lead to a range of adverse impacts including increased risk of illness and mortality; 15 the heatwave in August 2003 has been associated with ~70,000 deaths across Europe. Due to 16 climate change, heatwaves are likely to become more intense, more frequent and last longer in the 17 future. A number of factors may influence risks associated with heat exposure, such as population 18 age, housing type, and location within the Urban Heat Island, and such factors may not be evenly 19 distributed spatially across a region. We simulated and analysed two major heatwaves in the UK, in 20 August 2003 and July 2006, to assess spatial vulnerability to heat exposure across the West 21 Midlands, an area containing ~5 million people, and how ambient temperature varies in relation to 22 factors that influence heat-related health effects, through weighting of ambient temperatures 23 according to distributions of these factors across an urban area. Additionally we present 24 quantification of how particular centres such as hospitals are exposed to the UHI, by comparing 25 temperatures at these locations with average temperatures across the region, and presenting these 26 results for both day and night times. We find that UHI intensity was substantial during both 27 heatwaves, reaching a maximum of +9.6°C in Birmingham in July 2006. Previous work has shown 28 some housing types, such as flats and terraced houses, are associated with increased risk of 29 overheating, and our results show that these housing types are generally located within the 30 warmest parts of the city. Older age groups are more susceptible to the effects of heat. Our analysis of distribution of population based on age group showed there is only small spatial variation in 31 32 ambient temperature that different age groups are exposed to. Analysis of relative deprivation across the region indicates more deprived populations are located in the warmest parts of the city. 33

34 **KEYWORDS:** urban heat island, spatial vulnerability, heat waves, health effects.

35

36 1. INTRODUCTION

37 1.1 Heat exposure and health

1 Heatwaves, or extended periods of hot weather, are associated with various risks to health 2 including heat exhaustion, heatstroke, emergency hospitalisations, and death. A severe 3 heatwave in August 2003 has been associated with up to 70,000 excess deaths across Europe 4 (Robine et al., 2008), with over 2,000 excess deaths estimated in England (Johnson et al., 2005), 5 and a record breaking maximum temperature of 38.5°C reached in south east England. High 6 temperatures were also recorded throughout much of the UK during the summer of 2006. In 7 the West Midlands, the two heatwave events of 2003 and 2006 were comparable in terms of 8 estimated excess mortality, being around 10% higher than baseline rates at this time of year in 9 both cases (Health Statistics Quarterly, 2006; Johnson et al., 2005). In the future, heatwaves are 10 projected to become more frequent, more intense, and last longer, due to climate change (Kirtman 11 et al., 2013), which will likely lead to increases in heat-related mortality (Hajat et al., 2014; Mitchell 12 et al., 2016; Vardoulakis et al., 2014). Some evidence suggests there is an upper limit to which 13 humans can adapt to temperature (Arbuthnott et al., 2016; Sherwood and Huber, 2010).

In the UK, the risk of temperature-related mortality is projected to increase steeply in the UK over the 21st century under climate change and demographic change scenarios, reaching 260% by 2050, and 540% by 2080 (compared with the 2000s heat-related mortality baseline of around 2,000 premature deaths), with the elderly being most at risk (Hajat et al., 2014; HPA, 2012).

18 Future climate projections are often produced at relatively coarse spatial resolution, due to the 19 cost of computing power required. Factors relevant for the study of heat-exposure and human 20 health, including population age, socioeconomic factors, and the built environment such as 21 dwelling type and the urban heat island, are at a much finer spatial scale. Our study aims to 22 simulate ambient temperature across an urban area during a heatwave period, and subsequently 23 quantify the variation in ambient temperature with other factors that relate to and influence heat-24 related health effects. This includes population-weighting of ambient temperatures, as well as 25 calculating the ambient temperature weighted according to distributions of different housing types, 26 population age, and deprivation score, all factors that influence heat-health relationships. Using 27 environmental modelling techniques in this way to look at human health in relation to heat 28 exposure during heatwaves (which will become increasingly important in the future with climate 29 change) is a novel way of analysing spatial distribution of risks across a large urbanised region. 30 While we have used a detailed case study here, the technique and metrics are applicable to any 31 scenario, which we feel is useful for a wider scientific community.

32 1.2 The Urban Heat island (UHI)

Populations may be particularly at risk from heat due to the Urban Heat Island (UHI) effect (Heaviside et al., 2017), whereby ambient temperatures are often observed to be higher than those in surrounding less-urbanised areas, particularly at night. The main cause of the UHI is the modification of land surfaces, for example, replacing natural surfaces (e.g. vegetation which provide natural shading and cooling via evaporation) by paving or construction of buildings. Urban construction materials (such as concrete, tarmac and asphalt) generally absorb, retain and re-radiate heat more than natural surfaces. Buildings also provide multiple surfaces to

1 reflect and absorb sunlight, increasing urban heating, and impeding air circulation. Heat from 2 human activities (such as air conditioning, vehicles, and industrial processes) can also add to 3 the UHI effect. The UHI effect is often most extreme during anticyclonic summer weather 4 conditions, which are associated with heatwaves. In England and Wales, 82% of the population 5 reside in urban areas (ONS, 2011), leaving them vulnerable to the impacts of heat exposure due 6 to the UHI effect. The West Midlands is a highly urbanised area of the UK, which includes the city 7 of Birmingham with a population of 1.1 million, and has a notable UHI (Bassett et al., 2016; 8 Heaviside et al., 2015; Tomlinson et al., 2012). A health impact assessment for the heatwave of 9 2003 based on high resolution meteorological modelling suggested that in the West Midlands, 10 around half of the heat-related mortality during the heatwave could be attributed to the UHI 11 (Heaviside et al., 2016). The UHI intensity is often defined as the difference in temperature 12 between urban and rural areas, and can be quantified by comparing ambient air temperature 13 observations at a location in the centre of an urban area, and at a location in surrounding rural 14 areas (Bassett et al., 2016; Hatchett et al., 2016; Ketterer and Matzarakis, 2014; Kim and Baik, 2002; 15 Oke, 1973, 1982), or from satellite measurements (Azevedo et al., 2016; Benz et al., 2017; Du et al., 16 2016). However, observation stations are limited in number, are often not sited within urban 17 centres, and may only cover certain time periods, while satellite measurements record land surface 18 temperature (rather than air temperature), and are often temporally limited and may have missing 19 data if it is cloudy. The use of meteorological computer simulations makes it possible to investigate 20 spatial variations in temperature across urban areas, and to quantify the UHI intensity, by 21 comparing temperatures simulated both with and without urban surfaces such as buildings and 22 roads.

23 1.3. Mapping spatial variation in heat-exposure and risk

24 Heat exposure for urban populations will vary across urbanised areas, due to spatial variations in 25 physical infrastructure that influences the UHI. Susceptibility to health risks associated with heat 26 exposure is also influenced by other factors, including population age, housing type, socio-27 economic factors, pre-existing health conditions, and location within the UHI (Taylor et al., 2015; 28 Wolf and McGregor, 2013). It is therefore important in terms of health protection to understand 29 the pattern of risk across a region, which may help target resources to reduce heat risk in the most 30 vulnerable areas, since it is possible that environmental risks and vulnerable population groups are 31 co-located within urban areas.

A number of studies have investigated heat risk across the Greater London area. Wolf and 32 33 McGregor (2013) developed a Heat Vulnerability Index (HVI) for London based on principal component analysis of socio-demographic factors relating to heat vulnerability, combined with land 34 35 surface temperatures (derived from a satellite measurement), finding clustering of high vulnerability in the east and central areas of the city. Heat risk has also been mapped across London 36 37 using a building physics model and monitored weather data, together with information on 38 modelled UHI, housing type and population age, finding that building type and UHI have a 39 significant influence on the distribution of risk across the city for summer 2006 (Taylor et al., 2015).

3

1 Building fabric types and characteristics, and thus thermal properties, will depend on the age of the 2 building, and may be an important modifier for thermal comfort and energy efficiency, in addition 3 to behavioural aspects of building occupancy that can also modify heat exposure risk (Vardoulakis 4 et al., 2015). An existing spatial heat risk assessment of Birmingham combined information derived 5 from credit reference agencies on groups such as the elderly, those with ill health, high population 6 density, and high-rise living, finding that population sub-groups with ill health, and who reside in 7 flats were located in the overall warmest and therefore highest heat risk areas of Birmingham 8 (Tomlinson et al., 2011). This study used remote satellite sensing of the land surface temperature at 9 high spatial resolution (1 km), although only a single snapshot over one night during a heatwave is 10 used, and as such, does not fully represent the range of temperatures that population is exposed to 11 during the heatwave period. Similar studies in large urban areas worldwide have found spatial 12 overlap of sensitive populations and inequity in exposure to the UHI, according to age, occupation, 13 income, and other socio-demographic indicators, and that mapping can help target efforts to 14 reduce heat risk (e.g. via planning the location of parks, cool or green roofs, and social care efforts) 15 (Mitchell and Chakraborty, 2015; Weber et al., 2015; Wong et al., 2016).

16 While mapping of risks can be a useful tool for raising public awareness of risks, it is not clear if 17 producing maps in this way leads to policy or mitigation actions, possibly due to uncertainty in 18 relation to policy and decision making that are difficult to address in such studies (Wolf et al., 19 2015). There is an increasing amount of temperature data available from satellite measurements, 20 but this dataset only gives the temperature of the surface the satellite detects (e.g. the ground, tree 21 tops or building roofs and walls), rather than air temperature which is more relevant for health, 22 while the relationship between ground and air temperature is difficult to quantify. We address 23 these issues by using simulated temperatures and quantifying heat exposure due to the UHI for 24 populations which could be seen to be at risk.

25 1.4. Centres for health and social care

26 Care homes and hospitals are places where sensitive populations may reside. During the 2003 27 heatwave across Europe, of the recorded the recorded mortality among the under 75s in the UK, 28 more of the excess deaths were distributed in residential and nursing homes than would have been 29 expected, and among the over 75s, more of the excess deaths were distributed in hospitals and 30 nursing homes (Kovats et al., 2006). In these settings there may be issues surrounding dependency 31 on care-givers in nursing and residential homes, and the impact this may have on patients' and 32 residents' behaviour in responding to environmental changes (e.g. disempowerment or loss of 33 autonomy), as well as availability of assistance for daily living tasks in institutions (Belmin et al., 34 2007; Brown and Walker, 2008). Reports suggest care schemes have a 'culture of warmth', a 35 perception that the elderly are vulnerable only to cold (not excessive heat), and that design for 36 overheating is rare, with low prioritisation for future climate change and overheating (Gupta et al., 37 2016).

Those in prisons or custodial facilities are not able to directly access health care services, and effective ways to reduce exposure to heat may be limited. While there are few studies examining

1 heat risk in prisons, lawsuits were successfully brought against prisons in Texas and Louisiana in the 2 US, where inmates had died or suffered as a result of heat exposure (Human Rights Clinic, 2015; 3 The Promise of Justice Initiative, 2013). The population in the UK is ageing, as is the prison 4 population. Not only has the prison population in the UK increased from 56,000 to over 85,000 in 5 the last ten years, the number of those aged 60 and over in custody are the fastest growing group, 6 increasing by 120% from 2002-2013 (House of Commons, 2013; Offender management statistics 7 quarterly, 2016). Other factors such as drug use, as well as prevalence of mental health conditions 8 amongst those in custody may also contribute to health risk from exposure to heat.

9 1.5. Social and demographic factors

As well as the elderly, babies and young children are also vulnerable to effects associated with heat exposure (PHE, 2015). Studies have shown that warmer classrooms (25°C compared to 20°C) are detrimental to educational attainment (Wargocki and Wyon, 2007), and school children feel lethargic when temperature and relative humidity are high (Salthammer et al., 2016). Urbanisation and climate change adversely affect thermal comfort and air quality in schools, and the impact of such exposure may go beyond discomfort or ill health, leading to a lifelong burden of disease (Salthammer et al., 2016).

17 Relationships between heat-related mortality and socio-economic and demographic factors vary 18 regionally across the globe. Relationships have been found in some regions between heat-related 19 mortality and factors such as age, gender, those classed as urban-poor, and areas with a lack of 20 high-income earners (with income believed to be a good proxy for access to air conditioning), 21 however, these are often not consistent between countries, in some cases showing no clear 22 relationship for other socio-economic indicators (Hondula and Barnett, 2014; Johnson and Wilson, 23 2009; Yu et al., 2010). In the UK, regional long time-series analysis showed the strongest heat-24 mortality effects were in London, and the strongest cold effects in the East of England. A stronger 25 heat effect was found for those living in urban areas, with no difference for cold effects. Indices of 26 deprivation were not found to be a modifier for heat effects, and only slightly for cold effects in 27 some rural areas, though this is influenced by how well individual circumstances are represented by 28 area-level indicators (Hajat et al., 2007).

29 The World Health Organization (WHO) identified common heat risk factors including being elderly, 30 having pre-existing cardiovascular or respiratory disease, living alone, working outdoors or being 31 involved in heavy labour indoors close to industrial heat sources (World Health Organization, 2015). 32 In some places, gender, nature of dwelling (e.g. hospital or care home), being urban and poor, and 33 having certain medical conditions such as diabetes may also be linked with greater temperature-34 health effects, though causality is complex to determine as certain types of factors may interact 35 with other determinants of health, as well as access to health-care systems (World Health 36 Organization, 2015). Some indicators of socioeconomic status may be associated with heat-related 37 health effects, though inconsistencies exist between studies, with different associations reported 38 between countries, regions and cities within. There is also some evidence that temperature 1 modifies the health effects of some air pollutants, such as ozone (Jerrett et al., 2009; World Health

- 2 Organization, 2015).
- 3 1.6 Summary and aims

We have identified a wide range of factors that are likely to influence the risk of heat-related health effects, including housing type, population age, deprivation indices, and those residing in care and health facilities. People who experience multiple risk factors may be more vulnerable during periods of hot weather. An analysis of the spatial distribution of multiple risk factors across a region may reveal co-location of such factors, and identify places when strategies to mitigate heat risk might be most effective.

10 We address these issues as part of an urban case study for two recent significant heatwave events. 11 We investigate firstly variations in environmental risk factors across the city (temperature, UHI 12 intensity, and thermal comfort indices), and secondly, distribution of other factors which may relate 13 to vulnerability from environmental risks (age, housing type, and deprivation index). A spatial 14 analysis allows us to determine whether these sets of risks are co-located within our study city of 15 Birmingham and the West Midlands region of the UK. Thirdly, we investigate the position of specific 16 locations where sensitive populations are likely to be based, such as hospitals, care-homes, schools, 17 and prisons, with respect to the intensity and position of the UHI, with the hypothesis that since 18 these types of institutions are often situated in urban centres, populations may be at risk from 19 exposure to high temperatures as a consequence of the effects of the UHI.

20

21 2. METHODS AND DATA

In this study we first modelled the ambient air temperature at a height of 2 metres across the West Midlands, and quantified the UHI intensity across the region for the heatwave periods (August 2003 and July 2006) using a mesoscale meteorological model. Using GIS techniques we then extracted (from the modelled data) and analysed the ambient temperatures that correspond to different risk factors such as population age, dwelling type, and deprivation indices.

27 <u>2.1 Modelling temperature</u>

28 The Weather Research and Forecasting (WRF) model is a regional weather model that can be run at 29 high resolution (1 km) to simulate the meteorology across a region (Chen et al., 2011). We used the 30 WRF model with four nested domains (Figure 1a) each with grid resolutions of 36 km, 12 km, 3 km, 31 and 1 km in the smallest (central) domain, with feedback of variables from smaller grids to parent 32 grids. The model time-step was 180, 60, 20 and 5 s for the four domains, respectively, with output 33 every hour. Initial and lateral boundary meteorological conditions were provided by the European 34 Centre for Medium-range Weather Forecasts (ECMWF) ERA-interim reanalysis (Dee et al., 2011) at 35 a spatial resolution of 0.5° every 6 h. There were 39 pressure levels above the surface to 1 hPa. We 36 used an urban canopy scheme, Building Energy Parameterisation (BEP), which models the effect of

buildings on energy and momentum fluxes inside and immediately above the urban street canyons 1 2 (Heaviside et al., 2015; Martilli et al., 2002). Information on building and road properties (e.g. 3 building height, street canyon width, material properties such as albedo, thermal conductivity and 4 heat capacity) is included for three urban categories: Industrial/commercial, high-intensity 5 residential, and low-intensity residential across the West Midlands region (Figure 1b). Details of 6 parameters used for each urban category are given in Table 1a. Land-surface data used as an input 7 to WRF for all domains were based on the US Geological Survey (USGS) 24-category land-use data, 8 and for the inner domain we used two local datasets to generate the urban categories. We used the 9 Noah Land Surface Model, a relatively complex community model, which is often coupled with an

10 urban canopy scheme, and has four layers of soil moisture and temperature (Tewari et al., 2004).



1(b); (b) Urban classifications used in the urban canopy scheme in the WRF model in the central domain. The area shown covers approximately 80 x 80 km.

11 12

Table 1 : (a) Details of urban categories used in BEP. (b) General WRF model setup details.

Category		1: Industrial/ commercial	2: High-intensity residential	3: Low-intensity residential	
	Roof	0.1989	0.1997	0.2027	
Albedo	Wall	0.1989	0.1997	0.2027	
	Ground	0.1989	0.1997	0.2027	
Surface emissivity	Roof	0.9239	0.9274	0.9292	
	Wall	0.9239	0.9274	0.9292	
	Ground	0.9239	0.9274	0.9292	
Average building height		25 m	15 m	10 m	
(b)					
Model setting		Option		Reference	
Long wave radiation		Rapid Radiati	ve Trabsfer Model (RRTM)	Mlawer et al. (1997)	
Short-wave	radiation	Dudhia scher	ne	Dudhia (1989)	
Boundary la	ayer physic	s Bougeault–La	acarrere (designed for use	Bougeault and	
		with BPE urb	an scheme)	Lacarrere (1989)	
Urban physics		BEP urban sc	heme	Martilli et al. (2002)	

13

14 The model has been previously run for this region for the August 2003 heatwave; details of the 15 model simulation and validation are detailed in Heaviside et al. (2015). We have extended the analysis by re-running the 2003 heatwave using an updated version of WRF (v3.6.1), and including a further simulation run during a heatwave period in July 2006. In order to test performance, the model output is compared with observations from MIDAS observational weather stations across the West Midlands (Met Office, 2012). During heatwave periods, air temperature becomes more sensitive to surface moisture properties, which can be difficult for the model to capture. Soil moisture was adjusted in the model to better account for the unusually dry conditions, which led to improvements in the model performance (Figure 2 and Table 2).



Figure 2 : (a) Modelled and observed ambient temperatures at MIDAS sites across the West Midlands in July 2006. (b) Taylor diagram showing statistical comparison between adjusted and unadjusted model. Unadjusted simulation is shown in red, and simulation with reduced initial moisture is shown in blue. A better comparison with observations is indicated by proximity to 'REF' on the horizontal axis.

Table 2 : Statistical comparison between observed air temperature, and modelled temperatures (with soil moisture
 adjustment).

	Edgbaston		Coventry		<u>Coleshill</u>		Church Lawford	
	Observed	Modelled	Observed	Modelled	Observed	Modelled	Observed	Modelled
Mean (°C)	22.54	22.59	22.31	22.64	22.12	22.44	22.32	22.24
Standard deviation	4.54	4.95	4.78	5.11	5.19	5.15	5.21	5.40
(°C)								
RMSD (°C)	_	1.62	-	1.81	-	1.79	-	1.79
Correlation	_	0.94	-	0.94	-	0.94	-	0.94

We ran the model for two heatwave periods (2nd – 10th August 2003, and 16th – 27th July 2006), and in order to quantify the UHI intensity, two simulations were run for each heatwave period. An 'urban' simulation was run as described above, using the urban categories shown in Figure 1b, and is intended to represent the urban morphology of Birmingham and the West Midlands. The simulation is then run again, but with urban land cover replaced by rural cropland and pasture in the innermost domain, as a theoretical 'rural' case. By comparing results between the 'urban' and 'rural' simulations, the UHI intensity can be quantified.

The temperature that people experience or 'feel' can be influenced by a number of factors such as humidity, wind speed, exposure to direct solar radiation, and the amount of clothing they are wearing (Höppe, 1999; Steadman, 1984). National weather services sometimes provide a 'heat index' or 'apparent temperature' as part of weather forecasts during periods of warmer weather, to account for the effect of humidity, with assumptions about human body mass, height, clothing, amount of physical activity, sunlight and ultraviolet radiation exposure, and the wind speed. The heat index is calculated from temperature and relative humidity following the method of the NOAA 1 Weather Prediction Center (www.wpc.ncep.noaa.gov/html/heatindex_equation.shtml). This 2 calculation is a refinement of a result obtained by multiple regression detailed in National Weather 3 Service (NWS) Technical Attachment (SR 90-23). Accounting for the humidity only significantly 4 influences the heat index above about 26°C, and this calculation is not valid for extremes of 5 temperature or humidity (Steadman, 1984).

6 <u>2.2 Population and built environment characteristics</u>

Population data are taken from the National Population Database (2015) gridded at 100 m
resolution, which gives total population at each grid point. Information on age groups is available at
Output Area¹ (OA) level from the most recent census (2011), and attached to each 100 m
population point based on which OA the point falls within. The information is then summed across
each WRF 1 km grid box.

Data on housing type was obtained at Lower Super Output Area² (LSOA) level from the most recent 12 13 census (2011, available from infuse.ukdataservice.ac.uk). Data is available on the number of 14 dwellings of each type within each LSOA. The number of dwellings of each housing type in each 15 WRF model grid box is calculated by summing the fractions of each LSOA that intersect with each grid box, with housing assumed to be distributed evenly across each LSOA. This is combined with 16 17 the 2 m air temperature simulated in the model during the heatwaves of 2003 and 2006, to 18 estimate the average ambient temperature that different housing types across the West Midlands 19 are co-located with. Information on housing type was available (e.g. detached, terrace, flat, etc.) 20 but information on housing age was not recorded spatially.

21 The locations of hospitals, care homes, child care centres, schools, and prisons were considered 22 where 'sensitive populations' are more likely to reside. The National Population Database provides 23 information on these locations as they may have high densities of residents, many of whom may 24 not be able to protect themselves (others have a duty of care towards them). People at these 25 locations may be more vulnerable to harm and potentially harder to evacuate in an emergency 26 situation. The temperature at the location of each of the centres of interest is extracted from the 27 (hourly) model output using bilinear interpolation to the nearest modelled grid points. The mean 28 temperature anomaly for each location is calculated by taking the mean temperature at the 29 selected location over the heatwave period, and subtracting the mean temperature across the 30 entire domain for the same period.

The English Indices of Multiple Deprivation (IMD) scores and ranks are produced periodically by the Department for Communities and Local Government, and are the official measure of relative deprivation for small areas (LSOA based on the 2011 census) in England. The IMD brings together 37 different indicators which cover specific aspects or dimensions of deprivation, such as income,

¹ Output Areas are the lowest geographical level at which census estimates are provided. There are 181,408 Output Areas in England and Wales. The average population per OA is 309.

² Lower Super Output Areas have an average of ~1500 residents and 650 households. Measures of proximity (to give a reasonably compact shape) and social homogeneity (to encourage areas of similar social background) are also included. There are 34,753 LSOAs in England and Wales. http://neighbourhood.statistics.gov.uk/

employment, health and disability, education, skills and training, barriers to housing and services, 1 2 living environment, and crime, which are then weighted and combined to create the overall IMD 3 scores for each LSOA (ONS, 2007). For this work, IMDs for 2007 were used, as this is the closest 4 available year to those modelled using WRF. The IMD score at LSOA level is attached to each 100 m 5 population grid point, and then these are used to calculate population-weighted IMD scores for 6 each WRF 1 km grid-box. Once the IMD scores for each WRF grid box are calculated, these are then 7 associated with the temperatures from the WRF model, and ranked relative to each other, from 8 least- to most-deprived. The data is then split into population deciles (in order of least to most 9 deprived), and the associated temperatures plotted.

- 10
- 11 **3. RESULTS**
- 12 <u>3.1 Environmental risks</u>

13 <u>3.1.1 Temperature/Heat index.</u>

Figure 3a shows the modelled mean 2 m air temperature from 2nd – 10th August 2003, and Figure 3b shows the same for 16th – 27th July 2006. The observed average and the overall minimum temperatures are higher by 1.3°C in 2006 compared with 2003. Average daily maximum ambient temperatures are similar (difference of 0.3°C), though temperatures reached at least 30°C on more days in 2006 than 2003 (see Figure S1 in supplementary material). Comparison of Figure 3 with the land use in Figure 1b shows that urban areas are generally warmer than the surrounding rural areas in the West Midlands modelled domain.



21

1 The average calculated heat index for each of the heatwave periods is shown in supplementary

2 Figure S2. The pattern and magnitude are very similar to the 2 m temperatures shown in Figure 3.

3 Additional discussion of the heat index may be found in the supplementary material.

4 <u>3.1.2. Urban Heat Island</u>

5 The average UHI intensity across the whole period (defined as the difference in 2 m temperature 6 between a simulation with urban land cover and a simulation with rural land cover) for both 7 heatwave periods is shown in Figure 4. Statistics for the two heatwaves are shown in Table 2.

Table 2: Statistics for modelled temperature and the UHI intensity for the two heatwave periods in August
 2003 and July 2006.

			2005 0			
		Δυργασο	Δνοτασο	Daytime†	Nighttime†	
		tomn (°C)		average	average	Max UHI (°C)
		temp (°C)		UHI (°C)	UHI (°C)	
Aug 2003	Regional [‡]	20.5	+0.7	+0.5	+1.0	+6.1 (5pm 10 th ; 11pm 5 th ;
	City centre [*]	21.7	+2.2	+2.0	+2.7	9pm 9 th)
July 2006	Regional [‡]	21.7	+0.6	+0.1	+1.2	$10.6(11nm 17^{th})$
	City centre [*]	23.0	+2.0	+1.0	+3.1	+9.0 (11pm 17)
+Daytime is	defined as 08:00	0 – 19:59; Nig	httime is 20	:00 – 07:59		

‡ Regional is the whole area shown in Figure 1a. * City centre is a point in the centre of Birmingham city.

10

11 While higher average temperatures were reached across the domain during the 2006 heatwave 12 than the 2003 heatwave (Figure 3), the magnitude and spatial distribution of the UHI intensity 13 remains remarkably similar (Figure 4). There are broad similarities in the spatial and temporal 14 diurnal patterns of the UHI intensity between the two heatwaves (Figure 4, Figure S2), although 15 some localised differences exist. The average UHI intensity in the city centre was slightly higher in 16 2003 (+2.2°C in 2003 compared with +2.0°C in 2006), but the UHI intensity at night time was larger 17 in 2006 (+3.1°C in 2006 compared with +2.5°C in 2003).





1

- 2 By combining the data from the two heatwave periods, we have a larger dataset available to
- 3 analyse the distribution of temperatures, and the UHI effects typical during heatwaves across the
- 4 region. Figure 5 shows the spatial distribution of the UHI intensity across the West Midlands,
- 5 determined from both the August 2003 and July 2006 heatwaves combined, and illustrates the
- 6 more intense UHI effect at night time compared with during the day.



1

The heat index and the difference in heat index between urban and rural simulations are shown in supplementary material (Figures S2 and S3), and very similar patterns to the UHI are shown. The specific humidity (i.e. water vapour mixing ratio) is lower in the urban simulation (supplementary Figure S4) due to the presence of urban surfaces replacing vegetated ones that would otherwise provide a supply of moisture to the surrounding air. Overall the average impact of urban surfaces on 2 m temperature and the average impact on the heat index are almost identical (Figure 5 and Figure S3).

9

10 <u>3.2. Social factors</u>

11 <u>3.2.1 – Housing</u>

Buildings are an important modifier to population exposure to heat, with different types of housing being more or less susceptible to overheating, for example top-floor flats are more likely to overheat than detached houses (Beizaee et al., 2013; Symonds et al., 2017; Taylor et al., 2015). Figure 6 shows the average 2 m air temperature that different housing types across the West

1 Midlands are co-located with during heatwave periods, compared to the average temperature 2 across the whole region. The anomaly represents the difference from the average temperature 3 across the entire region for the time period considered (this temperature, representing zero 4 anomaly, is labelled at the base of the vertical axis on the right). The average temperature across all 5 housing types in the region is shown by the horizontal bar (again with absolute value labelled at the 6 right axis). Detached houses, which tend to be located towards more suburban areas, are co-7 located with ambient temperatures almost 0.2°C lower than the average for all housing types. By 8 contrast, terraced houses and flats are co-located with ambient temperatures over 0.1°C higher 9 than the average for all housing types in the region, and 0.6°C warmer than the whole domain 10 average temperature. Our analysis shows that housing types which are more likely to overheat (e.g. 11 flats) are often located in the warmest parts of the city, and the effects are much larger at night 12 (Figure 6b). The differences between ambient temperatures for each housing category are all 13 statistically significant (p<0.05; see supplementary material for details). While the temperature 14 variation between different housing types is less than 1°C, it should be noted that the average UHI 15 intensity for the region was about 0.6 to 0.7°C (and just over 2°C in the centre of the urban area) 16 during heatwaves.



17

18 <u>3.2.2 – Population age</u>

Figure 7 shows the average outdoor temperature anomalies different age groups are exposed to, based on residential address. The anomaly represents the difference from the average temperature across the entire region for the time period considered (this temperature, representing zero anomaly, is labelled at the base of the vertical axis on the right). The average across all age groups

is shown by the horizontal bar (again with absolute value labelled at the right axis). The whole West 1 2 Midlands population is exposed to temperatures about +0.6°C higher than the regional average (as 3 populations reside in more urbanised areas) (Fig. 9a). Young adults and the very young are exposed 4 to slightly higher temperatures than the population average (e.g. 20-24 year olds reside in areas 5 +0.7°C warmer than the regional average of +0.6°C), while the elderly are exposed to slightly lower 6 temperatures than the population average (e.g. 65-75 year olds reside in areas +0.5°C). During the 7 daytime, there is only a modest difference in ambient temperatures exposure in comparison to the 8 regional average; however at night time, population-weighted temperatures are over 1°C higher 9 than the regional average (Figure 7b,c). It should be noted that these figures are based on 10 residential population, so those present in health and care centres are not considered here. Overall, 11 the differences in exposure to ambient temperatures for different population groups are modest, 12 likely a reflection that population age groups are less heterogeneously distributed across this 13 region. The differences between ambient temperatures for each age group are all statistically 14 significant (p<0.05; see supplementary material for details), with the exception of the 15 and the 15 16-17 year old age groups (p=0.25), the 60-64 and 65-69 year old age groups (p=0.069), and the 85-16 89 and 90+ year old age groups (p=0.055).



Figure 7: (a) Average outdoor temperature anomaly (compared with the average across the domain during the heatwave period) for different age groups across the West Midlands, with standard deviations plotted. The analysis is also broken down into (b) night time and (c) day time.

17

18 <u>3.2.3 – Deprivation.</u>

Figure 8 shows the average ambient temperature exposure for increasingly deprived population 1 2 deciles, calculated from population weighted scores across the domain ranked relative to each 3 other. The anomaly represents the difference from the average temperature across the entire 4 region for the time period considered (i.e. day, night, or all times). The horizontal line shows the 5 population weighted temperature for the time period. The most deprived decile of population 6 across the region is exposed to temperatures +1.0°C higher than the average for the region (+0.5°C 7 during the day, and +1.5°C at night). The most deprived decile of population for the region 8 experiences temperatures consistently higher than the average for the region, even during the 9 daytime (each decile represents ~0.5 million people). The overall deprivation scores for all LSOAs in 10 England range from 0.37 to 85.46. For comparison, the deprivation scores calculated for each decile 11 in Figure 8 range from 6.08 to 56.17, placing them around the 12% and 96% most deprived relative 12 to England. The differences between ambient temperatures for each deprivation decile are all 13 statistically significant (p<0.05; see supplementary material for details), with the exception of 14 deciles 4 and 5 (p=0.989).



Figure 8: (a) Average outdoor temperature anomaly (compared with the average across the domain during the heatwave period) for population deciles ranked by deprivation across the West Midlands, with standard deviations plotted. The analysis is also broken down into (b) night time and (c) day time.

15

16

17 <u>3.3 Location based assessment of vulnerability from heat.</u>

1 Sensitive receptors such as hospitals, care homes, child care facilities, schools, and prisons tend to 2 be located in the most highly urbanised areas of the region, with the exception of prisons which are 3 often located on the outskirts of the city. The distribution of temperature anomalies for each 4 location type is shown in Figure 9. The anomalies are based on the number of receptors (i.e. 5 buildings), and not the population at each location, as this will fluctuate. The majority of hospitals, 6 care homes, schools and child care centres are located in areas warmer than the domain average 7 (94% of hospitals, 90% of care homes, 87% of schools, and 88% of child care centres). There are 8 seven prisons located within the study area, with only two of these being warmer on average 9 (Figure 9e).



Figure 9 : Distribution of average ambient 2 m temperature at (a) hospitals, (b) care homes, (c) schools, (d) child care centres, and (e) prisons across the West Midlands. The first column shows the distribution of these locations across the West Midlands, and the next three columns represent the average anomaly averaged first across all times of day, and then broken down into average daytime anomaly (8am – 8pm), and average night time anomaly (8pm – 8am). The average temperature across the domain (used as the reference temperature) is 21.8°C. Red = positive anomaly (warmer); Blue = negative (colder).

1 4. DISCUSSION

2 In this study we used hourly ambient temperature data at high spatial resolution (1 km) across a 3 densely populated region of the UK, capturing temperature variation across the period of two 4 different heatwave episodes. We employ environmental modelling techniques to examine human 5 health in relation to heat exposure during heatwaves, which will become increasingly important in 6 the future as climate change is projected to increase heat-health impacts. Previous studies have 7 examined the spatial distribution of vulnerability to heat risk across a region, often using ambient 8 temperatures that are time-varying but have no spatial variation, or that are well represented 9 spatially but only for a single or limited time frame (Taylor et al., 2015; Tomlinson et al., 2011; Wolf and McGregor, 2013). By using well-validated meteorological modelling techniques, we have 10 11 generated an ambient temperature dataset, based on modelled output, at high temporal and 12 spatial resolution from which to calculate ambient temperatures exposure co-located with various 13 risk factors that relate to and influence heat-related health effects. These include population-14 weighted temperature exposure and the relationship between UHI intensity and demographic 15 factors such as age group. Previous studies have mapped such risks, but our work extends this 16 analysis by calculating the ambient temperature weighted according to distributions of different 17 housing types, population (including age), and deprivation score, all being factors that influence 18 heat-health effects. We have also broken down the analysis by day and night; while the whole 19 region is warmest during the daytime (and urban areas are still warmer than rural ones at this 20 time), the spatial variation in temperature across the region is much greater at night, due to the 21 influence of the UHI (Figure 5). While there is no clear indicator of which temperature metric is 22 most strongly correlated with mortality, and there is variation between countries and cities (Davis 23 et al., 2016), high mortality ratios have been associated with high night time land surface 24 temperatures during the August 2003 heatwave in Paris (Dousset et al., 2011), and there is 25 evidence that higher night time temperatures have a detrimental impact on sleep quality, which is 26 linked with other negative health outcomes (Lack et al., 2008).

We found that almost all locations where sensitive populations are likely to be located (hospitals, care homes, schools, colleges) were warmer than the regional average by up to 2°C, and the number that are warmer is greater at night time (Figure 9). Of the seven prison locations within the study area, most were located in more rural or suburban areas, and therefore were not exposed to higher average ambient temperatures for this region.

32 Locations classified as hospitals here include a few small treatment centres such as family planning, 33 dialysis centres, etc., and therefore do not necessarily have resident populations who might be 34 exposed to the larger temperature anomalies at night. Locations classed as care homes have a high 35 density of residents, present day and night, and also include centres with specialised facilities for 36 those with mental health issues, sensory impairment, learning difficulties, and those who misuse 37 drugs and alcohol. Child care centres have a duty of care to protect people who cannot protect 38 themselves, with potentially high populations, including nurseries, crèches, toddler groups, 39 afterschool clubs, holiday schemes, and youth clubs. School locations are unlikely to be occupied at night, when temperature anomalies are higher, and those of school age are exposed to ambient
temperatures close to the average for the population (though still above regional average). While
colleges have very high populations, they again are unlikely to be occupied at night.

4 The anomalies presented here are based on ambient temperature, although buildings play an 5 important role in determining actual exposure, particularly at night time. Studies using building 6 simulations to estimate indoor temperatures find a significant influence of building characteristics 7 on indoor temperatures (Taylor et al., 2015). However these studies often use a single ambient 8 temperature across all dwelling types, whereas this study shows that different building types are 9 likely to be exposed to different ambient temperatures across a region. Our study could help inform 10 development of inputs to building models by providing a modifier to ambient temperature based 11 on the likely outdoor temperatures for a particular building type.

12 While we have included the highest spatially resolved data that was available at the time, the 13 analysis makes some assumptions about distribution of certain factors. Data for housing types at 14 LSOA level, and population ages at OA level was the finest level of data that was available. Any 15 variations that may exist within an individual LSOA or OA respectively could not be captured by this 16 analysis. The BEP multi-layer urban canopy scheme used with WRF can capture the effects of the 17 built environment on energy and momentum fluxes (accounting for shading and reflections by 18 buildings) at sub-grid scale, providing a suitable representation of how urban areas influence 19 meteorology in a weather model such as WRF. However, microclimatic effects at individual building 20 level are not explicitly captured, as regional weather models can only be run down to 1 km 21 horizontal grid scale constrained by the valid sub-grid scale turbulence schemes, and therefore 22 results for individual locations interpolated from the 1 km grid may have additional variation. The 23 IMDs are a measure of relative deprivation, calculated by weighting data on different indicators 24 that can be used as measures of deprivation. The IMDs are calculated periodically, and although it 25 is a good measure for comparing areas with each other, this measure cannot be easily compared 26 with other years or used to identify trends, as the ranking is performed separately each time a new 27 dataset is released.

This study focuses primarily on the effects of heat, although other environmental or socioeconomic factors in cities that may impact health could be considered in future studies. Such factors may include air pollution, flood risk, social isolation or greenspace. Cold effects could also be investigated. These and other factors identified as influencing health may be included in future studies to develop and overall environmental risk map for cities. Finally, this study examines past heatwave events in two years across the UK, allowing model validation against observations. Future scenarios under climate change could be considered in further work.

35

36 **5. CONCLUSIONS**

1 Previous research suggests that the UHI may contribute around half of the heat related mortality 2 experienced during heatwaves (Heaviside et al., 2016). Increasing urbanisation and climate change 3 will increase heat related health risks in urban areas. We have developed a novel risk mapping 4 methodology that combines high spatial resolution modelling of temperature, population age, and 5 building types to identify locations and population sub-groups at higher health risk during 6 heatwaves. We have simulated and quantified the magnitude and characteristics of the UHI in the 7 West Midlands during two heatwave periods (August 2003 and July 2006), using the WRF-BEP 8 model. The UHI intensity across the region is on average 2.1°C, reaching up to 9.4°C, and there is 9 strong spatial heterogeneity in ambient temperature across the region, particularly at night. Spatial 10 analysis using GIS techniques shows that locations where people already more vulnerable to the 11 effects of exposure to heat (e.g. the elderly, those with pre-existing health conditions, and those 12 dependent on others for care) may reside, tend to be located in the hotter parts of the region. 13 Sensitive receptors (94% of hospitals, 90% of care homes, 87% of schools, and 88% of child care 14 centres) are co-located with higher than the average air temperatures across the West Midlands. 15 Buildings that are more susceptible to overheating, such as flats, are exposed to higher ambient 16 temperatures than other housing types.

17 Interventions such as urban greening or building modifications such as cool roofs may help offset 18 some of the UHI intensity, and reduce the spatial disparity in ambient temperatures, particularly at 19 night. The results from this work could help identify which factors are most strongly correlated with 20 ambient temperature, and help target resources and interventions, as well as focus health 21 messaging during heatwaves to the greatest effect.

Further research is needed to estimate the spatial heterogeneity of factors influencing heat risk in cities. In addition, future work should consider the health impacts and potential benefits that UHI mitigation techniques in a UK city would have on building overheating, as well as the co-benefits or other unintended environmental consequences that may occur in a city.

26

27 ACKNOWLEDGEMENTS

The research was funded by the National Institute for Health Research Health Protection Research Unit (NIHR HPRU) in Environmental Change and Health at the London School of Hygiene and Tropical Medicine in partnership with Public Health England (PHE), and in collaboration with the University of Exeter, University College London, and the Met Office. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR, the Department of Health or Public Health England. The authors gratefully acknowledge the reviews by two anonymous reviewers and the editor, which have helped improve the manuscript.

35

36 **REFERENCES**

- Arbuthnott, K., Hajat, S., Heaviside, C., Vardoulakis, S. (2016) Changes in population susceptibility to heat
 and cold over time: assessing adaptation to climate change. Environmental Health 15, S33.
- 3 Azevedo, J., Chapman, L., Muller, C. (2016) Quantifying the Daytime and Night-Time Urban Heat Island in
- Birmingham, UK: A Comparison of Satellite Derived Land Surface Temperature and High Resolution Air
 Temperature Observations. Remote Sensing 8, 153.
- 6 Bassett, R., Cai, X., Chapman, L., Heaviside, C., Thornes, J.E., Muller, C.L., Young, D.T., Warren, E.L. (2016)
- 7 Observations of urban heat island advection from a high-density monitoring network. Quarterly Journal of
- 8 the Royal Meteorological Society 142, 2434-2441.
- 9 Beizaee, A., Lomas, K.J., Firth, S.K. (2013) National survey of summertime temperatures and overheating risk
 10 in English homes. Building and Environment 65, 1-17.
- Belmin, J., Auffray, J.-C., Berbezier, C., Boirin, P., Mercier, S., de Reviers, B., Golmard, J.-L. (2007) Level of
 dependency: a simple marker associated with mortality during the 2003 heatwave among French dependent
 elderly people living in the community or in institutions. Age and Ageing 36, 298-303.
- Benz, S.A., Bayer, P., Blum, P. (2017) Identifying anthropogenic anomalies in air, surface and groundwater
 temperatures in Germany. Science of The Total Environment 584–585, 145-153.
- Bougeault, P., Lacarrere, P. (1989) Parameterization of Orography-Induced Turbulence in a Mesobeta--Scale
 Model. Monthly Weather Review 117, 1872-1890.
- Brown, S., Walker, G. (2008) Understanding heat wave vulnerability in nursing and residential homes.
 Building Research & Information 36, 363-372.
- 20 Chen, F., Kusaka, H., Bornstein, R., Ching, J., Grimmond, C.S.B., Grossman-Clarke, S., Loridan, T., Manning,
- K.M., Martilli, A., Miao, S., Sailor, D., Salamanca, F.P., Taha, H., Tewari, M., Wang, X., Wyszogrodzki, A.A.,
- Zhang, C. (2011) The integrated WRF/urban modelling system: development, evaluation, and applications to
 urban environmental problems. International Journal of Climatology 31, 273-288.
- Davis, R.E., Hondula, D.M., Patel, A.P. (2016) Temperature Observation Time and Type Influence Estimates of
 Heat-Related Mortality in Seven U.S. Cities. Environmental Health Perspectives 124, 795-804.
- 26 Dee, D.P., Uppala, S.M., Simmons, A.J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M.A.,
- 27 Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A.C.M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C.,
- 28 Dragani, R., Fuentes, M., Geer, A.J., Haimberger, L., Healy, S.B., Hersbach, H., Hólm, E.V., Isaksen, L., Kållberg,
- 29 P., Köhler, M., Matricardi, M., McNally, A.P., Monge-Sanz, B.M., Morcrette, J.J., Park, B.K., Peubey, C., de
- Rosnay, P., Tavolato, C., Thépaut, J.N., Vitart, F. (2011) The ERA-Interim reanalysis: configuration and
 performance of the data assimilation system. Quarterly Journal of the Royal Meteorological Society 137,
- . 32 553-597.
- 33 Dousset, B., Gourmelon, F., Laaidi, K., Zeghnoun, A., Giraudet, E., Bretin, P., Mauri, E., Vandentorren, S.
- (2011) Satellite monitoring of summer heat waves in the Paris metropolitan area. International Journal of
 Climatology 31, 313-323.
- Du, H., Wang, D., Wang, Y., Zhao, X., Qin, F., Jiang, H., Cai, Y. (2016) Influences of land cover types,
 meteorological conditions, anthropogenic heat and urban area on surface urban heat island in the Yangtze
 River Delta Urban Agglomeration. Science of The Total Environment.
- Dudhia, J. (1989) Numerical Study of Convection Observed during the Winter Monsoon Experiment Using a
 Mesoscale Two-Dimensional Model. Journal of the Atmospheric Sciences 46, 3077-3107.

- 1 Gupta, R., Walker, G., Lewis, A., Barnfield, L., Gregg, M., Neven, L., (2016) Care provision fit for a future
- climate. Oxford Brookes University, Published by the Joseph Rowntree Foundation (available from
 jrf.org.uk).
- Hajat, S., Kovats, R.S., Lachowycz, K. (2007) Heat-related and cold-related deaths in England and Wales: who
 is at risk? Occup Environ Med 64, 93-100.
- Hajat, S., Vardoulakis, S., Heaviside, C., Eggen, B. (2014) Climate change effects on human health: Projections
 of temperature-related mortality for the UK during the 2020s, 2050s and 2080s. Journal of Epidemiology and
- 8 Community Health 68, 641-648.
- Hatchett, B.J., Koračin, D., Mejía, J.F., Boyle, D.P. (2016) Assimilating urban heat island effects into climate
 projections. Journal of Arid Environments 128, 59-64.
- Health Statistics Quarterly (2006) Estimated daily mortality during July 2006 in England and Wales. Health
 statistics quarterly / Office for National Statistics, 107-111.
- 13 Heaviside, C., Cai, X.-M., Vardoulakis, S. (2015) The effects of horizontal advection on the urban heat island
- 14 in Birmingham and the West Midlands, United Kingdom during a heatwave. Quarterly Journal of the Royal
- 15 Meteorological Society 141, 1429-1441.
- Heaviside, C., Macintyre, H.L., Vardoulakis, S. (2017) The Urban Heat Island: Implications for health in a
 changing environment. Current Environmental Health Reports, (early online)
- 18 <u>https://link.springer.com/article/10.1007%1002Fs40572-40017-40150-40573</u>.
- Heaviside, C., Vardoulakis, S., Cai, X. (2016) Attribution of mortality to the Urban Heat Island during
 heatwaves in the West Midlands, UK. Environmental Health 15 Suppl 1:27.
- Hondula, D.M., Barnett, A.G. (2014) Heat-Related Morbidity in Brisbane, Australia: Spatial Variation and
 Area-Level Predictors. Environ Health Perspect 122, 831–836.
- Höppe, P. (1999) The physiological equivalent temperature a universal index for the biometeorological
 assessment of the thermal environment. International Journal of Biometeorology 43, 71-75.
- House of Commons, (2013) Older prisoners: Fifth Report of Session 2013–14, Volume I, in: Committee,
 H.o.C.J. (Ed.).
- HPA, (2012) Health Effects of Climate Change in the UK 2012, in: Vardoulakis, S., Heaviside, C. (Eds.). Health
 Protection Agency.
- Human Rights Clinic, (2015) Reckless Indifference: Deadly Heat in Texas Prisons, The University of Texas
 School of Law <u>https://law.utexas.edu/clinics/2015/05/24/usa-extreme-heat-in-texas-prisons-fall-2013/</u>.
- Jerrett , M., Burnett , R.T., Pope , C.A.I., Ito , K., Thurston , G., Krewski , D., Shi , Y., Calle , E., Thun , M. (2009)
 Long-Term Ozone Exposure and Mortality. New England Journal of Medicine 360, 1085-1095.
- Johnson, D.P., Wilson, J.S. (2009) The socio-spatial dynamics of extreme urban heat events: The case of heat related deaths in Philadelphia. Applied Geography 29, 419-434.
- Johnson, H., Kovats, R.S., McGregor, G., Stedman, J., Gibbs, M., Walton, H., Cook, L., Black, E. (2005) The
- impact of the 2003 heat wave on mortality and hospital admissions in England. Health statistics quarterly /
 Office for National Statistics, 6-11.

- 1 Ketterer, C., Matzarakis, A. (2014) Comparison of different methods for the assessment of the urban heat
- 2 island in Stuttgart, Germany. International Journal of Biometeorology.
- Kim, Y.-H., Baik, J.-J. (2002) Maximum Urban Heat Island Intensity in Seoul. Journal of Applied Meteorology
 41, 651-659.
- 5 Kirtman, B., Power, S.B., Adedoyin, J.A., Boer, G.J., Bojariu, R., Camilloni, I., Doblas-Reyes, F.J., Fiore, A.M.,
- 6 Kimoto, M., Meehl, G.A., Prather, M., Sarr, A., Schär, C., Sutton, R., van Oldenborgh, G.J., Vecchi, G., Wang,
- 7 H.J., (2013) Near-term Climate Change: Projections and Predictability, in: Stocker, T.F., Qin, D., Plattner, G.-
- 8 K., Tignor, M., Allen, S.K., Boschung, J., Nauels, A., Xia, Y., Bex, V., Midgley, P.M. (Eds.), Climate Change 2013:
- 9 The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the
- 10 Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, United Kingdom and
- 11 New York, NY, USA, pp. 953-1028.
- Kovats, R.S., Johnson, H., Griffiths, C. (2006) Mortality in southern England during the 2003 heat wave by
 place of death. Health statistics quarterly / Office for National Statistics, 6-8.
- Lack, L.C., Gradisar, M., Van Someren, E.J.W., Wright, H.R., Lushington, K. (2008) The relationship between
 insomnia and body temperatures. Sleep Medicine Reviews 12, 307-317.
- Martilli, A., Clappier, A., Rotach, M. (2002) An Urban Surface Exchange Parameterisation for Mesoscale
 Models. Boundary-Layer Meteorology 104, 261-304.
- Met Office, (2012) Met Office Integrated Data Archive System (MIDAS) Land and Marine Surface Stations
 Data (1853-current). NCAS British Atmospheric Data Centre.
- Mitchell, B.C., Chakraborty, J. (2015) Landscapes of thermal inequity: disproportionate exposure to urban
 heat in the three largest US cities. Environmental Research Letters 10, 115005.
- Mitchell, D., Heaviside, C., Vardoulakis, S., Huntingford, C., Masato, G., Guillod, B.P., Frumhoff, P., Bowery,
 A., Wallom, D., Allen, M. (2016) Attributing human mortality during extreme heat waves to anthropogenic
- 24 climate change. Environmental Research Letters 11, 074006.
- 25 Mlawer, E.J., Taubman, S.J., Brown, P.D., Iacono, M.J., Clough, S.A. (1997) Radiative transfer for
- inhomogeneous atmospheres: RRTM, a validated correlated-k model for the longwave. Journal of
 Geophysical Research: Atmospheres 102, 16663-16682.
- 28 National Population Database, (2015) National Population Database Tool. Health and Safety Laboratory,
- 29 comissioned by HSE. Licensed to PHE under PSMA Contractor Licence (HSL).,
- 30 <u>https://data.gov.uk/dataset/national-population-database4</u>, April 2015.
- 31 Offender management statistics quarterly, (2016) Offender management statistics quarterly: April to June
- 32 2016. Ministry of Justice © Crown copywright 2016, pp.
- 33 <u>https://www.gov.uk/government/collections/offender-management-statistics-quarterly</u>.
- 34 Oke, T.R. (1973) City size and the uran heat island. Atmospheric Environment 7, 769-779.
- Oke, T.R. (1982) The energetic basis of the urban heat island. Quarterly Journal of the Royal Meteorological
 Society 108, 1-24.
- 37 ONS, (2007) The English Indices of Deprivation. Office for National Statistics,
- 38 <u>https://www.gov.uk/government/collections/english-indices-of-deprivation</u>.

- 1 ONS (2011) Office for National Statistics, Census.
- PHE, (2015) Heatwave Plan for England. <u>https://www.gov.uk/government/publications/heatwave-plan-for-</u>
 <u>england</u>.
- Robine, J.-M., Cheung, S.L.K., Le Roy, S., Van Oyen, H., Griffiths, C., Michel, J.-P., Herrmann, F.R. (2008) Death
 toll exceeded 70,000 in Europe during the summer of 2003. Comptes Rendus Biologies 331, 171-178.
- Salthammer, T., Uhde, E., Schripp, T., Schieweck, A., Morawska, L., Mazaheri, M., Clifford, S., He, C.,
 Buonanno, G., Querol, X., Viana, M., Kumar, P. (2016) Children's well-being at schools: Impact of climatic
- 8 conditions and air pollution. Environment International 94, 196-210.
- 9 Sherwood, S.C., Huber, M. (2010) An adaptability limit to climate change due to heat stress. Proceedings of
 10 the National Academy of Sciences 107, 9552-9555.
- Steadman, R.G. (1984) A Universal Scale of Apparent Temperature. Journal of Climate and Applied
 Meteorology 23, 1674-1687.
- 13 Symonds, P., Taylor, J., Mavrogianni, A., Davies, M., Shrubsole, C., Hamilton, I., Chalabi, Z. (2017)
- Overheating in English dwellings: comparing modelled and monitored large-scale datasets. Building Research
 & Information 45, 195-208.
- 16 Taylor, J., Wilkinson, P., Davies, M., Armstrong, B., Chalabi, Z., Mavrogianni, A., Symonds, P., Oikonomou, E.,
- Bohnenstengel, S.I. (2015) Mapping the effects of urban heat island, housing, and age on excess heat-related
 mortality in London. Urban Climate 14, Part 4, 517-528.
- 19 Tewari, M., Chen, F., Wang, W., Dudhia, J., LeMone, M.A., Mitchell, K., Ek, M., Gayno, G., Wegiel, J., Cuenca,
- 20 R.H., (2004) Implementation and verification of the unified Noah land surface model in the WRF model, 20th
- 21 Conference on Weather Analysis and Forecasting/16th Conference on Numerical Weather Prediction,
- 22 Seattle, Washington., p. Abstract 14.12A.
- 23 The Promise of Justice Initiative, (2013) Court rules in Louisiana death row conditions case: Extreme heat on
- 24 death row found to violate eighth amendment to U.S. constitution, <u>http://justicespromise.org/about-</u>
- us/news-press/82-court-rules-in-louisiana-death-row-conditions-case-extreme-heat-on-death-row-found-to violate-eighth-amendment-to-u-s-constitution. The Promise of Justice Initiative.
- Tomlinson, C.J., Chapman, L., Thornes, J.E., Baker, C.J. (2011) Including the urban heat island in spatial heat
 helth risk assessment strategies: a case study for Birmingham, UK. International Journal of Health
 Geographics 10, 42.
- Tomlinson, C.J., Chapman, L., Thornes, J.E., Baker, C.J. (2012) Derivation of Birmingham's summer surface
 urban heat island from MODIS satellite images. International Journal of Climatology 32, 214-224.
- Vardoulakis, S., Dear, K., Hajat, S., Heaviside, C., Eggen, B., McMichael, A. (2014) Comparative assessment of
 the effects of climate change on heat- and cold-related mortality in the United Kingdom and Australia.
 Environmental Health Perspectives 122, 1285-1292.
- Vardoulakis, S., Dimitroulopoulou, C., Thornes, J., Lai, K.-M., Taylor, J., Myers, I., Heaviside, C., Mavrogianni,
 A., Shrubsole, C., Chalabi, Z., Davies, M., Wilkinson, P. (2015) Impact of climate change on the domestic
 indoor environment and associated health risks in the UK. Environment International 85, 299-313.
- Wargocki, P., Wyon, D.P. (2007) The Effects of Moderately Raised Classroom Temperatures and Classroom
 Ventilation Rate on the Performance of Schoolwork by Children (RP-1257). HVAC&R Research 13, 193-220.

- Weber, S., Sadoff, N., Zell, E., de Sherbinin, A. (2015) Policy-relevant indicators for mapping the vulnerability
 of urban populations to extreme heat events: A case study of Philadelphia. Applied Geography 63, 231-243.
- 3 Wolf, T., Chuang, W.-C., McGregor, G. (2015) On the Science-Policy Bridge: Do Spatial Heat Vulnerability
- Assessment Studies Influence Policy? International Journal of Environmental Research and Public Health 12,
 13321.
- Wolf, T., McGregor, G. (2013) The development of a heat wave vulnerability index for London, United
 Kingdom. Weather and Climate Extremes 1, 59-68.
- 8 Wong, M., Peng, F., Zou, B., Shi, W., Wilson, G. (2016) Spatially Analyzing the Inequity of the Hong Kong
 9 Urban Heat Island by Socio-Demographic Characteristics. International Journal of Environmental Research
 10 and Public Health 13, 317.
- 11 World Health Organization, (2015) Heatwaves and Health: Guidance on Warning-System Development, in:
- McGregor, G.R., Bessemoulin, P., Ebi, K., Menne, B. (Eds.). World Health Organization and World
 Meteorological Office.
- 14 Yu, W., Vaneckova, P., Mengersen, K., Pan, X., Tong, S. (2010) Is the association between temperature and
- mortality modified by age, gender and socio-economic status? Science of The Total Environment 408, 3513 3518.
- 17
- 18