

Exercise to preserve β -cell function in recent-onset Type 1 diabetes mellitus (EXTOD) - a randomized controlled pilot trial

Narendran, Parth; Jackson, Nikki; Daley, Amanda; Thompson, Dylan; Stokes, Keith; Greenfield, Sheila; Charlton, Mary; Curran, Michelle; Solomon, Thomas; Nouwen, Arie; Lee, Siang I; Cooper, Ashley R ; Mostazir, Mohammod; Taylor, Rod S ; Kennedy, Amy; Andrews, Rob C

DOI:

[10.1111/dme.13439](https://doi.org/10.1111/dme.13439)

License:

Other (please specify with Rights Statement)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Narendran, P, Jackson, N, Daley, A, Thompson, D, Stokes, K, Greenfield, S, Charlton, M, Curran, M, Solomon, T, Nouwen, A, Lee, SI, Cooper, AR, Mostazir, M, Taylor, RS, Kennedy, A & Andrews, RC 2017, 'Exercise to preserve β -cell function in recent-onset Type 1 diabetes mellitus (EXTOD) - a randomized controlled pilot trial', *Diabetic Medicine*, vol. 34, no. 11, pp. 1521-1531. <https://doi.org/10.1111/dme.13439>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Checked for eligibility: 21/09/2017

This is the peer reviewed version of the following article: Narendran P, Jackson N, Daley A, Thompson D, Stokes K, Greenfield S, Charlton M, Curran M, Solomon TP, Nouwen A, Lee SI. Exercise to preserve β -cell function in recent-onset Type 1 diabetes mellitus (EXTOD)—a randomized controlled pilot trial. *Diabetic Medicine*. 2017. which has been published in final form at [10.1111/dme.13439](https://doi.org/10.1111/dme.13439). This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.


When citing, please reference the published version.

Take down policy


While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Relationship between Δ AUC vs. Δ HOMA-IR



Disposition Index



*6m/12m means are adjusted for their baseline scores