

## Attention training in autism as a potential approach to improving academic performance:

Muller Spaniol, Mayra; Shalev, Lilach; Kossovaki, Lila; Mevorach, Carmel

DOI:

[10.1007/s10803-017-3371-2](https://doi.org/10.1007/s10803-017-3371-2)

License:

None: All rights reserved

*Document Version*

Peer reviewed version

*Citation for published version (Harvard):*

Muller Spaniol, M, Shalev, L, Kossovaki, L & Mevorach, C 2018, 'Attention training in autism as a potential approach to improving academic performance: a school-based pilot study', *Journal of Autism and Developmental Disorders*, vol. 48, pp. 592–610. <https://doi.org/10.1007/s10803-017-3371-2>

[Link to publication on Research at Birmingham portal](#)

### **Publisher Rights Statement:**

The final publication is available at [link.springer.com](http://link.springer.com) via <http://dx.doi.org/10.1007/s10803-017-3371-2>

### **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

### **Take down policy**

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

**Attention training in autism as a potential approach to improving academic performance: a school-based pilot study**

**ABSTRACT**

This study assessed the effectiveness of an attention intervention program (Computerized Progressive Attention Training; CPAT) in improving academic performance of children with ASD. Fifteen 6-10 year olds with ASD attending a mainstream and a special school were assigned to an experimental (CPAT; n=8) and active control (computer games; n=7) group. Children were assessed pre- and post-intervention on measures of behavioural symptoms, cognitive skills and academic performance. The intervention was conducted in school twice a week for 8 weeks. Children in the CPAT group showed cognitive and academic improvements over and above the active control group, while children in both groups showed improvements in behaviour. Results suggest that attention training is a feasible approach to improving academic performance in this population.

Autism Spectrum Disorder (ASD) or autism is a neurodevelopmental disorder characterized by difficulties in social interaction and communication, repetitive and stereotyped interests and behaviours (American Psychiatric Association 2013). ASD is a heterogeneous disorder with complex aetiology (Mandy & Lai 2016), and although impairments in attention are not found amongst its core symptoms, atypical attention is often linked to autism in research, and even associated with the development of ASD symptoms (Keehn, Müller & Townsend 2013).

### *Attention functioning in Autism*

Atypical attentional processes have been noted in individuals with autism from early infancy. One such example is sustained attention (i.e. the ability to remain focused on task over time), which was found to be poorer in children and adolescents with autism in comparison to typically developing (TD) controls (Chien et al. 2015; Chien et al. 2014; Murphy et al. 2014). In a behavioural study measuring sustained attention, participants had to report whenever a specific sequence of numbers appeared within a serial visual processing stream (Chien et al. 2015). Differences in sustained attention were apparent in participants with autism as they showed fewer hits and correct rejections and more misses in comparison to TD children within the same IQ range ( $IQ < 115$ ) (Chien et al. 2015). Similar findings were also reported in a previous study using a Continuous Performance Task (CPT) where children with autism showed worse performance in focused and sustained attention and increased ADHD-like (attentional) symptoms (Chien et al. 2014). Poor performance in sustained attention was also found to be accompanied by reduced brain activation in relevant brain regions in autism compared to controls including lower activation in inferior prefrontal, medial prefrontal, striato-thalamic, and lateral cerebellar regions (Murphy et al. 2014). Even when behavioural differences are not present in sustained attention, brain activation still differs in ASD (Belmonte & Yurgelun-Todd 2003a; Ciesielski, Courchesne, Elmasian 1990; Murphy et al.

2014). While contradicting results were also reported indicating intact sustained attention in ASD (Sanders, Johnson, Garavan, Gill & Gallagher 2008), this may be attributed to scenarios in which the demand for sustained attention is lower (such as in the context of sustained attention to preferred objects).

Evidence for impaired selective-spatial attention (i.e. the ability to select relevant information and suppress irrelevant stimuli) has also been documented (Burack 1994; Plaisted, Swettenham & Rees 1999), with early brain activation differences found for participants with autism (Belmonte & Yurgelun-Todd 2003a; Ciesielski et al. 1990). Belmonte and Yurgelun-Todd (2003a) used functional magnetic resonance imaging (fMRI) while participants performed a task consisting of two oddball streams, presented side-by-side. Participants had to attend to one or the other stream according to target colour. Typically, brain activation patterns represent the locus of spatial attention in such tasks so that activity in ventral visual cortex is enhanced for the attended stream and suppressed for the unattended one (Belmonte & Yurgelun-Todd 2003b). In contrast, brain activation patterns in adults with ASD indicated no spatial attention modulation of ventral visual areas (i.e. there was no enhancement or suppression). This was also accompanied by performance difference where the ASD participants showed reduced selection of targets in the attended stream and reduced compensatory suppression of items in the unattended stream (Belmonte & Yurgelun-Todd 2003a).

Finally, perhaps the most frequently documented attention-related difficulty in autism is in executive control, which is often linked to autism as a primary or secondary deficit (Geurts, Luman & Van Meel 2009; Hill 2004; Pennington & Ozonoff 1996). Executive control (EC) comprises a large set of functions such as working memory, set shifting, inhibition, cognitive flexibility, reasoning and planning, all of which are needed for goal-directed behaviours.

Specifically, set shifting or flexibility was found to be problematic in children with ASD (Happé, Booth, Charlton & Hughes 2006; Verte, Geurts, Roeyers, Oosterlaan & Sergeant 2006). Cognitive flexibility or set-shifting includes processes such as disengaging attention from stimuli and redirecting it towards different locations. Indeed, atypical disengagement of attention is documented in infants with ASD as early as the first year of life (Elsabbagh et al. 2009; Landry & Bryson 2004; Zwaigenbaum et al. 2005). Inhibition and working memory also appear to be considerably delayed in autistic children in early ages, but may recover when entering adulthood (Luna, Doll, Hegedus, Minshew & Sweeney 2007). While some studies have found no problems in inhibition and working memory in autism (Dawson et al. 2002; Luna et al. 2007; Yerys, Hepburn, Pennington & Rogers 2007), increased task complexity and consequently increased demand for executive control tend to bring out such differences. For instance, Rinehart, Bradshaw, Tonge, Brereton & Bellgrove, (2002) measured inhibition in children with autism and TD controls using a Stroop-like task. Levels of cognitive load were systematically increased, gradually adding congruency, conditionality and choice to the task. Results showed that only with increased levels of cognitive load the ASD group showed a decrease in performance (Rinehart et al., 2002).

#### *Academic performance in Autism*

Academic attainment may vary considerably across children on the autism spectrum with a substantial percentage showing additional learning disabilities. In fact, in a review of 10 empirical studies on the prevalence of learning disabilities among children with autism by Emerson, Baines, Allerton & Welch (2010), a very wide range of prevalence rates was reported from 15% to 84%. However, 8 out of the 10 reviewed studies show a prevalence of comorbidity of autism and learning disabilities of above 40%. Overall, poor performance in a range of academic tests such as maths, reading and writing is well documented in ASD

(Keen, Webster & Ridley 2016). Nevertheless, while difficulties in acquiring academic skills are common among individuals with ASD, not all people with autism struggle academically (Estes, Rivera, Bryan, Cali & Dawson 2011). Evidence for difficulties in maths for instance, can be accompanied by reports of average performance in individuals with high functioning autism (HFA) and in some cases performance advantage (Chiang & Lin 2007). Reading skills also seem to be heterogeneous, as children with autism can show average reading ability but they may have difficulties in reading comprehension, and variability in results is large from floor to ceiling levels (Nation, Clarke, Wright & Williams 2006). Writing skills in HFA are reported to be poor, with frequent writing learning disabilities (Whitby & Mancil 2009).

While attention represents a core cognitive process, critical for the development of a number of skills, its importance in the context of classroom settings and academic attainment is of particular interest. Recent research (Erickson, Thiessen, Godwin, Dickerson & Fisher 2015) highlights the importance of selective and sustained attention in supporting learning in a classroom setting since early infancy. In particular, selective attention has been found to play an important role in the development of both literacy and numeracy (Stevens & Bavelier, 2012). Furthermore, Stern and Shalev (2013), found that poor performance in reading and reading comprehension was related to difficulties in sustained attention. Yet, another function of attention - executive attention - was linked to children's ability in Maths (Bull & Scerif, 2001). Evidence linking difficulties in academic skills to attentional difficulties in ASD come from studies by May and colleagues (May, Rinehart, Wilding & Cornish 2013; 2015). Specifically, poor maths performance in children with ASD was found to be related to problems in attention switching in a visual search task (May et al. 2013). In a follow-up study, May et al. (2015) also found that attention switching correlated with both maths and reading performance in ASD. In fact, children with autism might show some similarities in

learning and attentional profile to children with ADHD (Mayes & Calhoun 2007), and children with learning disabilities (LD) (Calhoun & Mayes 2005). In these studies, all clinical groups (ASD, ADHD, LD) were reported to have difficulties in attention and processing speed, which relate to poor performance in academic tests, particularly writing (Calhoun & Mayes 2005; Mayes & Calhoun 2007).

Thus, the behavioural and brain atypicalities seen in the performance of ASD participants in different attention tasks might not only have implications for the severity of ASD symptoms (Keehn et al. 2013), but also for skills needed in learning new abilities, and academic attainment in school settings. Improvement of these attention skills in ASD is therefore of crucial importance in promoting successful learning for children with ASD in a school environment.

#### *Cognitive Interventions in Autism*

A growing number of studies utilising computer-based training programs were carried out with children with autism. These programs typically target specific cognitive functions supporting social interactions such as emotion and face recognition (Golan et al. 2010), language and literacy (Pennington 2010), and social skills (Bernardini, Porayska-Pomsta, Smith & Avramides 2012). However, attempts at targeting more domain-general processes have also been carried out (Hilton et al. 2014; Vries, Prins, Schmand & Geurts 2015). Working memory and cognitive flexibility were separately taught using a computerized program in children with ASD (Vries et al. 2015). In this case, marginal post training effects and generalization occurred, with improvements in working memory and near-transfer from working memory training to attention, and cognitive flexibility training showing improvements in flexibility (Vries et al. 2015). Hilton et al. (2014) also tested cognitive

training for executive function and motor skills in children with autism, using an exergame (i.e. videogame that uses body movement) and found improvements in working memory, metacognition, strength and agility. While certainly not all prior computer-based intervention studies have proven successful as some failed to generalize to naturalistic settings (e.g. Golan & Baron-Cohen 2006; Wass & Porayska-Pomsta 2014), results tend to be overall positive, showing improvements of trained skills and transfer effects (Bernardini et al. 2012; Golan et al. 2010; Pennington 2010), even when more domain-general processes such as working memory and attention are targeted (Vries et al. 2015). However, thus far comprehensive attempts to train a variety of basic attention functions are sparse, especially when applied in a school setting with the aim to impact academic attainment. One attempt to train cognitive flexibility and executive function in a school setting was carried out by Kenworthy et al. (2014) using the "Unstuck and On Target" (UOT) intervention. Using a behavioural protocol to train executive abilities and flexibility, the researchers found post-training improvements in problem-solving, flexibility, and planning/organizing as well as behaviour in classroom, for children with ASD taking part in the UOT training group, in comparison to a social skills intervention group.

The notion of the potential benefit of attention training is not unique to ASD. Posner and Rothbart (2005) argued that attentional training, through repetition can change brain functionality, and bring generalized improvements not only to the attentional network, but also to intelligence and consequently, academic performance. A computerized attention program has been used with children with attention difficulties in a school setting, showing long-term improvements in academic and attention skills (Rabiner et al. 2010). In fact, a number of studies in different populations support the notion that attention can be modified with training (Gagnon & Belleville 2012; Kinnealey et al. 2012; Rabiner et al. 2010;



Sampanis, Mevorach, Shalev, Mohammed & Humphreys 2015; Shalev, Tsal & Mevorach 2007; Wass, Scerif & Johnson 2012).

One attention-training program that has been shown in the past to have generalised training effects in different populations is the Computerised Progressive Attention Training (CPAT) program developed by Shalev et al. (2007). A study assessing the effectiveness of CPAT for 6 to 13 year old children with attention deficits resulted in a decreased level of inattention symptoms as well as improved non-trained academic skills (Shalev et al. 2007). Similarly, a study assessing CPAT in sub-acute stroke patients demonstrated improvements in both specific attention functions (particularly sustained attention) and transfer to other cognitive domains, such as language, memory and number skills which were not directly trained (Sampanis et al. 2015). As CPAT attempts to train different attention functions – sustained, selective-spatial and executive attention, all of which could exhibit atypicalities in ASD, it may be particularly appropriate in order to train attention in children with ASD, especially as its previous uses demonstrated transfer across different cognitive domains as well as to non-trained academic skills.

While various intervention programs for ASD have been studied (Bernardini et al. 2012; Golan et al. 2010; Hilton et al. 2014; Pennington 2010; Vries et al. 2015), to our knowledge none of them has specifically targeted attention training in primary-aged children with ASD in a school setting with the aim of improving academic skills. Moreover, there is a growing need of research and interventions developed for and with schools, considering staff and school resources for effective program implementation (Kasari & Smith 2013; Parsons et al. 2013). Therefore, in the present study, the CPAT program developed by Shalev et al. (2007) was carried out in two school settings (mainstream and special education) with children with ASD, in an attempt to evaluate its viability for such population. In particular, we ask whether

CPAT is an appropriate program to use with children with ASD and whether any changes triggered by training specific attention functions can transfer to non-trained academic skills and cognition.

## METHODS

The study took place at two different times over the school years of 2015-2016, in two primary schools in the UK (one mainstream and one special). 15 children (6-10 years old) across the two schools with a diagnosis of ASD took part and were divided into two groups, one undergoing the CPAT intervention and the other engaging in standard computer games (active control; see details below). To measure the effects of attention training, pre- and post-intervention assessments were taken in the two groups which included an intelligence test (Computerized Progressive Matrices, CPM; Raven, Raven & Court 2008), academic tests in maths, reading and copying, as well as the Childhood Autism Rating Scale (CARS; Schopler, Reichler, DeVellis & Daly, 1980) to measure the severity of autism, and semi-structured interviews with class teachers and teaching assistants (TAs) to get their views on children's performance. The entire study protocol took two months to complete. The intervention protocol and data collection were administered by a principal researcher (PR) and three research assistants (RAs), using personal and school laptops. Pupils were blind in regards of the main purpose of the study, and were all told that playing the computer games may help them improve in school. School staff and parents were told that all children were undergoing a computerized attention training. The possibility of using the CPAT with the active control group was offered to the participating schools following the completion of the study, providing the program as well as training the school staff on its application.

Institutional ethical approval for the study was obtained from the Science, Technology, Engineering and Mathematics Ethic Review Committee of the University of Birmingham. Written consent was obtained from parents and/or legal guardians of all participants included in the study.

### *Participants*

Participants were recruited from the two aforementioned schools. All participants had a diagnosis of autism and a statement of special educational needs (SEN) and none of them had an additional diagnosis (such as ADHD). Diagnostic criteria and ASD statements were checked and reviewed before children could take part in the study. A total of 15 children participated in the study, seven from the mainstream school and eight children from the special school (catering only for students with ASD). Pupils from the mainstream school came from different school years (i.e. 6 to 10 years old: 2 children in Year 1, 1 child in Year 3, 2 in Year 4 and 2 in Year 5) whereas pupils from the special school all attended the same classroom (i.e. 7 to 8 years old). In each school setting pupils were assigned either to an active control (computer games; CG) or to a training (CPAT) group. Participants were paired and matched in age (chronological age and cognitive age equivalent from the CPM, Raven et al. 2008) gender and intelligence (from the CPM; Raven et al. 2008) and randomly assigned to active control or training group. Thus, overall the CPAT group was comprised of 8 children (2 females) and the CG group included 7 children (1 female, see Table 1) across schools. To eliminate possible bias both groups were treated as intervention groups and this information was conveyed to teachers, parents and participants. One child (female) in the active control group did not participate in the computer games sessions, and instead had education as usual (also missing post-intervention assessments for maths and copying). The

age ( $t_{(13)}=.47, p=0.646$ ), age equivalent ( $t_{(13)}=-.772, p=0.454$ ) and the non-verbal intelligence ( $t_{(13)}=-.73, p=0.48$ ) did not differ statistically between the groups.

-----Insert Table 1 page 51 about here -----

*The intervention protocol (CPAT, Shalev, Tsal & Mevorach 2007)*

Attention training was employed using the CPAT, which was developed based on Tsal, Shalev and Mevorach (2005) four-functions of attention model, which proposes that deficits in attention can be present in four functions within the human brain: sustained attention (responsible for maintaining attention for a prolonged time), selective-spatial attention (focusing on relevant information while ignoring adjacent distracting stimuli), executive attention (solving conflicts, inhibition of irrelevant information) and orienting attention (directing and reorienting attention). In this study, three training tasks from the CPAT protocol were used, each separately focusing on either, sustained, selective-spatial or executive attention. The original orienting task (Shalev et al., 2007) was not included in this study as the task included characters in Hebrew and had proven to be too demanding for children with ADHD of similar age to the sample of the current study (Shalev et al., 2007). However, the function of orienting is included to some extent in both the selective spatial and executive attention training tasks (especially as levels of difficulty increase). Tasks were developed in a fun and interactive manner to be used with children from 6 years of age and above who show difficulties in attention (Shalev et al. 2007).

The first of the three training task is *The Computerized Continuous Performance Task (CCPT)*, which was developed to improve the function of sustained attention, that is, to be able to maintain the focus of attention on a given task especially during monotonous activities. The *Conjunctive Search Task (CST)*, based on Treisman & Gelade 1980) was

designed to improve selective-spatial attention, and the *Shift Stroop-like Task* (ST, based on Navon 1977), was designed to improve the function of executive attention and cognitive control. In addition, orienting of attention was trained as a part of the CST, whereby children had to disengage from the distractors and reorient attention towards the target, and also in higher levels within the ST task, when participants had to switch between the global and local levels of the figures (either between or within trials). Snapshots of the tasks are presented in Figure 1. These training tasks were designed with progressive levels of difficulty which are tuned to the individual performance of the trainee. All the training tasks use easy to understand visual and auditory feedbacks. The tasks and different levels of difficulty have been previously tested with children with ADHD (Shalev et al. 2007), children with foetal alcohol spectrum disorder (Kerns et al. 2010) and with patients after stroke (Sampanis et al. 2015). In all these different clinical groups the CPAT program produced positive outcomes.

(Insert Figure 1 page 49)

In the present study, participants in the CPAT group had an average of 13 training sessions (12-14) of approximately 45 minutes, twice a week across a two-month period. Sessions occurred in a distraction-free room, inside the school and were scheduled always at the same time and day of the week, and started as soon as children finished the activity they were working on in the classroom, in the pre-scheduled time set. Experimenters provided one-to-one supervision to each child. Each training task was divided into blocks of different number of trials. The CST and ST had 40 trials per block, while the CCPT had 80 trials per block for low levels of difficulty, or 160 trials for high levels of difficulty. All participants started at the lowest level of difficulty for each training task. Participants then progressed to higher levels of difficulty according to pre-specified criteria within the program, based on maintaining high levels of accuracy and individual improvements in reaction time. The shift between levels of

difficulty in each task was automatically controlled by the program. Training sessions had an average of 6 blocks from the three training tasks (4 to 9). This number varied according to children's age and individual differences. Generally, younger children performed fewer blocks per session. Also, each child took a different amount of time to finish blocks, according to their individual speed in responding to the tasks, which led to different number of blocks they could perform in a 45-minute session. In the CST and ST feedback was highly structured and immediate, with an auditory beep for incorrect answers and immediate positive visual feedback appearing on the screen for correct responses with average latency ("very good") and correct responses showing improvements in reaction time ("excellent"). The CCPT did not show immediate visual positive feedback to maintain high demand of sustained attention. At the end of each block, feedback was translated into points that reflected participants' performance, including the number of correct and incorrect answers and quantity of positive feedback received (in accordance with Shalev et al. 2007 and Sampanis et al. 2015).

#### *Active control protocol (computer games)*

For the active control group, three readily available computer games were used: Plants vs Zombies, Bejewelled 3 and Pacman. Participants in this group took part in a similar number of sessions as the CPAT group (i.e. 13 sessions) which were of the same length, frequency and format as the CPAT group. The control games also had feedback and had different levels of difficulty. Thus, both intervention groups used interactive games with positive feedbacks and rewards (stickers, reward game of their choice at the end of the session), over a 2-month period, twice a week in the mainstream and special school. Importantly, the active control group had the same one-to-one interaction with the experimenters as the training group.

*Pre- and post-intervention measures*

To assess the outcome of the attention training (CPAT) performance was measured in a number of domains (see detailed description below) before and after the intervention for both the CPAT and active control group. These included severity of autism symptoms (CARS), cognitive ability (Raven CPM), and children's academic performance in maths, reading and copying. The views of the class teaching staff were also captured via semi-structured interviews.

Assessments were carried out between one week and two weeks before and after the intervention, with an average of 3 to 4 sessions needed for completion of all assessments (with a frequency of two sessions per week). The assessments took place in the same rooms as the intervention (intervention rooms had sometimes to be changed due to limited room availability in schools but alternated between meeting room, library, computer room or empty classrooms). The PR and RAs had individual sessions with each child and were randomly assigned to work with different children to avoid possible bias. Results were compared within and across the training and control groups.

*CARS*

To assess behavioural symptoms of autism the classroom teachers filled in the Childhood Autism Rating Scale (CARS; Schopler et al. 1980; Schopler, Reichler & Renner 1988) for the participating pupils. CARS is a behaviour observation scale where the child's behaviour is scored against 15 different dimensions, each question corresponding to an ASD symptom. CARS gives a total score measure ranging from non-autistic to severe autism (from a minimum of 15 to a maximum of 60). Each dimension is scored in a scale from 1 to 4. While originally scores below 30 were thought to correspond to the non-autistic range (Schopler et

al. 1988), more recent studies point towards a cut-off score of 25 (Chlebowski, Green, Barton & Fein 2010; Tachimori, Osada & Kurita 2003). Scores above 37 indicate severe autism (Schopler et al. 1988). The reliability has been well documented, showing good internal consistency ( $\alpha=.94$ ,  $n=537$ ), high inter-rater agreement (correlations ranging from .55 to .93,  $M=.71$ ,  $n=280$ ), and test-retest stability over a 1 year period is .88 for the total score ( $n=91$ ; Schopler et al. 1988; CARS manual). High concordance between the CARS and the DSM-III-R and DSM-IV diagnostics is also documented (Perry, Condillac, Freeman, Dunn-Geier & Belair, 2005; Bebko, Perry & Bryson 1996).

#### *Raven CPM*

The Raven's – Educational: Coloured Progressive Matrices (CPM) was developed to provide brief non-verbal screening measure of general ability. Using a set including visual patterns and shapes, it measures nonverbal skills that involve making meaning out of confusion and the ability to form nonverbal constructs that aid handling complex information (Raven et al. 2008). The test is sensitive to intellectual differences, and demonstrates good test-retest reliability (.80; Raven, Court & Raven 1990). It is regarded as a measure of fluid intelligence, especially for children with learning difficulties and disorders of language ability, such as ASD (Carver 1990; Cotton et al. 2005; Raven et al. 2008).

The CPM consists of 36 items, divided in three sets of 12 each: set A, AB, and B. The sets are formed by visual shapes constituted by different patterns that need to be matched, requiring reasoning by analogy, and the ability to take this as a consistent way of thinking and method of inference for upcoming items (Raven et al. 2008). Children can score a maximum of 36 in the raw score, which is converted into standard score, according to points obtained, and chronological age from normative data. The test also provides percentile ranks and age



equivalents. In this study, the PR and RAs administered the CPM individually, prior and after training, following the administration procedure prescribed by Raven et al. (2008), with no time limitations.

#### *Academic assessment*

In order to assess academic performance and the potential impact the intervention has on it, children in both groups completed two series of short tests in maths, reading comprehension and passage copying, one before and one after the intervention program. The tests were the same for all children, prior and post training, and children did not receive any feedback on their performance. The tests were selected from an online database of tests used by schools in the UK. Children had 10 minutes to complete each test, the maths test had a total of 11 questions to be completed, reading comprehension had 17 questions and children had 3 paragraphs of 143 words in total to copy. These academic assessments were chosen for this study because they are commonly used by schools in the UK, providing high ecological validity. Our objective was to evaluate if performance of children improved within the school, using instruments utilized to measure academic attainment by schools. The academic assessments were timed to closely match tests in a school environment, as many aspects of academic performance are also timed in schools (e.g. time to finish tests and complete activities). The tests scores were converted into accuracy rates (percentage correct out of total number of questions) and the copying was scored using the number of words written per minute (total number of words divided by the time provided).

#### *Semi-structured interviews*

Four different class teachers and two TAs took part in individual semi-structured interviews (3 class teachers from different classrooms from the mainstream school, 1 class teacher and 2

TAs from the same classroom from the special school, see Table 2). The interviews were carried out at the same time as the pre- and post-intervention assessments. Teachers and TAs were asked about social, academic and personal performance of children participating in the project, as well as attention skills prior and after the training program. All interviewees were blinded to the existence of a control group, and were told that the pupils who participated in the research project would go through a computerised training on attention skills.

-----Insert Table 2 page 51 about here -----

Prior to the start of the study, it was briefly explained to them what the project consisted of (i.e. rationale, duration and timeline). They were asked about their expectations from the project and asked to describe how they perceived the children in school: social aspects, personal skills, academic performance, attention and concentration in class. The same questions about the children were asked a week after the end of training (see Appendix 1 page 43 for a blank copy of the interview schedule). Interviews were recorded using an audio recorder and additional written notes were kept. Written consents were obtained for the audio recordings.

#### *Data analysis*

A two-way repeated measures ANOVA using time (pre vs post) as within subject factor and group (training vs control) as between subjects factor was used to analyse pre and post data. Significance level was set at  $p < 0.05$ . Effect sizes are reported as partial eta squared ( $\eta_p^2$ ) for ANOVAs and Cohen's  $d$  (same group size) or Hedges  $g$  (different group size) for t-tests. For Raven CPM (which is a standardised measure) we also conducted an analysis of the reliability of the change following the intervention using Reliable Change Indices (RCI; Jacobson & Truax, 1991). All data are reported as mean  $\pm$  standard error of the mean (across

subjects) (SEM). CARS was reported using total scores; Raven CPM was reported and analysed using standard scores; maths and reading were reported in accuracy (percentage correct out of total questions); copying was reported and analysed using copying rate (the average number of words copied per minute). The qualitative data from the interviews were analysed using thematic analysis (Braun & Clark, 2006). Thematic analysis is a widely used method to identify, analyse and report patterns in qualitative data, organising and describing the data in detail, providing flexibility for interpretation of many aspects of the research (Braun & Clark, 2006). This study utilized both quantitative and qualitative data, although the qualitative data were analysed using a chi-square test of independence to verify significant differences between the groups. This decision was taken as a mixed method analysis combining quantitative and qualitative data is considered ideal in order to come to conclusions about human behaviour and effectiveness of interventions, complementing and enhancing the credibility of the research findings (Hesse-Biber, 2010).

## RESULTS

### *Baseline measures*

Scores were distributed normally in all pre and post measures, apart from the maths pre-test (Shapiro-Wilk; control group:  $p=0.039$ ; CPAT group:  $p=0.044$ ). Importantly, the groups did not significantly differ in any of the assessments prior to the commencement of the intervention (Table 3: CPM:  $t_{(13)}=-0.730$ ,  $p=0.48$ ,  $g_s=0.35$ ; maths accuracy:  $t_{(13)}=-1.14$ ,  $p=0.27$ ,  $g_s=0.56$ ; reading accuracy:  $t_{(13)}=-0.78$ ,  $p=0.45$ ,  $g_s=0.38$ ; copying rate:  $t_{(13)}=-1.40$ ,  $p=0.18$ ,  $g_s=0.68$ ; CARS:  $t_{(13)}=1.02$ ,  $p=0.32$ ,  $g_s=0.5$ ). Mean and SEMs of the measures are presented by group in Table 3.

-----Insert Table 3 page 51 about here -----

School membership (whether children were attending the mainstream or the special school) also did not affect pre-intervention baseline measures (CPM:  $t_{(13)}=-0.01$ ,  $p=0.99$ ; maths:  $t_{(13)}=1.39$ ,  $p=0.19$ , reading:  $t_{(13)}=0.97$ ,  $p=0.35$  or copying:  $t_{(13)}=-0.13$ ,  $p=0.99$ ), although CARS scores were higher in the mainstream ( $33.07\pm 2.06$ ) in comparison to the special school ( $26.62\pm 1.01$ ;  $t_{(13)}=2.92$ ,  $p=0.012$ ).

#### *Pre vs. post intervention analyses*

##### *CPM standard score*

CPM scores varied in our sample, ranging from extremely low to average, high average and superior, in both control and training groups. All children from both groups completed all of the CPM test questions on both pre- and post-intervention assessments. A two-way ANOVA was carried on CPM scores with time (pre vs post training) as a within subject factor and group (CPAT vs. CG) as a between subject factor. There was a main effect of time ( $F_{(1,13)}=7.08$ ,  $p=0.020$ ,  $\eta_p^2=0.353$ ), where pre CPM scores were significantly lower ( $89.55\pm 4.53$ ) in comparison to the post CPM scores ( $97.5\pm 5.17$ ). The interaction between group and time was marginally significant ( $F_{(1,13)}=3.78$ ,  $p=0.074$ ,  $\eta_p^2=0.225$ ). Further analysis of simple effects revealed a significant increase in the CPM scores from pre to post assessment for the CPAT group (pre= $86.25\pm 6.73$ ; post= $100\pm 5.98$ ;  $t_{(7)}=-3.67$ ,  $p=0.008$ , *Cohen's d*=-1.32), but not for the CG group (pre= $92.86\pm 5.86$ ; post= $95\pm 8.72$ ;  $t_{(6)}=-0.45$ ,  $p=0.67$ , *Cohen's d*=.011; see Figure 2a). To further assess the likelihood of change in this measure following the intervention, we conducted a Reliable Change Indices (RCI) analysis on the CPM scores (Jacobson & Truax, 1991). RCI is an appropriate measure for small samples which clinically examines significant improvements at the individual level. 6 out of

the 8 children in the CPAT group showed a RCI above the cut-off point of 1.96 (considered to indicate a reliable change), while only 2 out of 7 children from the CG group exhibited a RCI above the cut-off. In addition, 1 child in the CG group showed a reliable decrease in scores. Individual change from pre to post assessment for each participant is depicted in Figure 3a. The data shows that while in the control group both improvement and deterioration were recorded, in the CPAT all participants exhibited improvement.

### *Academic assessments*

#### *Maths*

Data analysis was conducted over the percent correct (correctly answered questions out of total test questions) for each participant. One child (CH7) from the control group was excluded from data analysis since she did not perform the test in the post assessment.

A two-way ANOVA was carried on maths accuracy with time (pre vs post training) as a within subject factor and group (CPAT vs. CG) as a between subject factor. There was a main effect for time ( $F_{(1,12)}=13.92$ ,  $p=0.003$ ,  $\eta_p^2=0.537$ ), showing that post-test scores were generally higher than pre-test scores (pre:  $38.92\pm 9.17$  and post:  $53.13\pm 8.92$ ). More importantly, there was a significant interaction between time and group ( $F_{(1,12)}=6.43$ ,  $p=0.026$ ,  $\eta_p^2=0.349$ ). There was a significant improvement in math scores for the CPAT group (pre= $27.8\pm 10.14$ ; post= $51.7\pm 9.98$ ;  $t_{(7)}=-4.52$ ,  $p=0.003$ ;  $d=-1.60$ ), but not for the CG group (pre= $47.4\pm 14.1$ ; post= $54.5\pm 15.8$ ;  $t_{(5)}=-0.866$ ,  $p=0.426$ ,  $d=-0.18$ ; see Figure 2b). There were no significant differences in the number of attempted answered questions between the pre- and post-tests, and across CPAT and CG group (average of 10 questions attempted out of the total 11). Individual change from pre to post assessment for each participant is depicted in

Figure 3b. Again, the data demonstrate that while in the control group both improvement and deterioration were recorded, in the CPAT group all participants showed improvement.

### *Reading comprehension*

A two-way ANOVA was carried on reading accuracy (% correct out of the total 17 questions in the test) with time (pre vs post training) as a within subject factor and group (CPAT vs. CG) as a between subject factor. Again, there was a main effect of time ( $F_{(1,13)}=8.54$ ,  $p=0.012$ ,  $\eta_p^2=0.397$ ), showing that post training reading scores were generally higher than pre training scores (pre:  $35.63\pm 4.74$  and post:  $47.45\pm 5.15$ ). Again, the interaction between time and group approached significance ( $F_{(1,13)}=3.95$ ,  $p=0.068$ ,  $\eta_p^2=0.233$ ). There was a significant improvement in reading scores for the CPAT group (pre= $39.34\pm 6.19$ ; post= $59.19\pm 5.34$ ;  $t_{(7)}=-4.28$ ,  $p=0.004$ ,  $d=-1.53$ ) but not for the CG group (pre= $31.9\pm 7.3$ ; post= $35.7\pm 9.2$ ;  $t_{(6)}=-0.553$ ,  $p=0.60$ ,  $d=-0.17$ ; see Figure 2c). Here too, there were no significant differences in attempted answered questions between pre and post assessment between the groups (average of 10 questions answered out of 17). Individual change from pre to post assessment for each participant is depicted in Figure 3c. The data once again show a variable picture for the control group with both improvements and deteriorations. In contrast, for the CPAT group a more unified picture of improvement emerges.

(Insert Figure 2 page 49)

### *Copying*

Word copying performance was analysed by calculating the copying rate (words copied per minute). One participant (CH7) from the control group was excluded from the analysis, as she did not perform the post intervention assessment of this test. The copying rate was submitted to a two-way ANOVA on number of copied words per minute with time (pre vs

post training) as a within subject factor and group (CPAT vs. CG) as a between subject factor. There was a main effect of time ( $F_{(1,12)}=7.78$ ,  $p=0.016$ ,  $n_p^2=0.393$ ), showing that post-training scores were generally higher than pre-training scores (pre:  $3.7\pm 0.6$  and post:  $4.6\pm 0.5$ ). More importantly, there was a significant interaction between time and group ( $F_{(1,12)}=11.97$ ,  $p=0.005$ ,  $n_p^2=0.499$ ). For the CPAT group there was a significant improvement in copying rate as a function of time (pre= $2.7\pm 0.46$ ; post= $4.9\pm 0.59$ ;  $t_{(7)}=-7.37$ ,  $p<0.001$ ;  $d=-1.6$ ). In contrast, there was no evidence for improvement in the copying rate for the CG group (pre= $4.6\pm 1.2$ ; post= $4.4\pm 0.9$ ;  $t(5)=3.29$ ,  $p=0.755$ ;  $d=0.09$ ; See Figure 2d). Individual change from pre to post assessment for each participant is depicted in Figure 3d. Here again, a similar pattern appears whereby the control group shows both improvements and deteriorations while the CPAT group shows primarily improvement.

(Insert Figure 3 page 50)

### *CARS*

A two-way ANOVA was carried on CARS scores with time (pre vs post training) as a within subject factor and group (CPAT vs. control) as a between subject factor. There were no significant main effects or interaction for either factor. Thus, as expected, it appears that CARS scores did not differ between the groups ( $F_{(1,13)}=1.43$ ,  $p=0.252$ ,  $n_p^2=0.1$ ) or the passing of time in general ( $F_{(1,13)}=0.71$ ,  $p=0.415$ ,  $n_p^2=0.052$ ).

### *Interviews*

A thematic analysis of the pre-intervention interviews shows that before training, teachers reported difficulties in social interactions, learning academic skills and concentration for all children who participated in the study (both for CG and CPAT groups). In the post-intervention interviews for both CG and CPAT groups, teaching staff noted improvements in

confidence and independence, increased social interactions and a general improved mood and enthusiasm. In order to investigate potential changes following the intervention, teachers' and TAs' accounts were subjected to thematic analysis (Braun & Clarke 2006; interviews were organised and divided in themes covering different topics and patterns, and compared between CPAT and CG groups). Specifically, the common themes were identified and grouped into improvements in: attention, academic and behavioural changes. Quotes from the interviews can be found in Appendix 2 page 44.

#### *CPAT group*

Generally, teaching staff perceived children who took part in the CPAT program as showing academic progress in maths and English, and as being able to concentrate in tasks for longer periods of time, requiring less prompting. They were also perceived to be more confident and independent, although this was not unique to the CPAT group. Some teachers and TAs also reported more 'seeking' behaviours where children seemed to look more for social interactions following training. Behavioural changes were coded from the post intervention interview transcripts with school staff. Results from the pre- and post-intervention assessments, and fragments from the interviews were organised in recurrent themes (attention, academic and behavioural changes) individually and summarized in Table 4. Some extracts from the interviews can be found in Table 4 but a more detailed account of the staff quotations can be found in Appendix 2.

-----Insert Table 4 page 52 about here -----

#### *Control group*

Overall, teaching staff reported positive changes in behaviour, confidence, independence and some progress in academic areas for some of the children in the control group. The same



improvements in attention and concentration were not commonly found in the reports from teaching staff for the control group. Quotes from the interviews are listed in Table 5.

As detailed in Tables 4 and 5, the interviews revealed that improvements in academic skills and attention were more frequently reported for participants in the CPAT group, even though all evaluators believed that all children were taking part in an attention-training program. Positive changes were reported in both control and CPAT groups concerning personal skills such as confidence, independence and motivation. To verify this, a chi-square test of independence was performed on the frequency teaching staff reported positive changes in attention, academic performance and behaviour, in order to examine the relation between attention improvements and group (CPAT and CG), academic improvements and group, and behavioural improvements and group. The relation between attention improvement and group was marginally significant, with more reports of improvements in attention within the CPAT group (87.5%, 7 out of 8) in comparison to the CG group (33.3%, 2 out of 6; Pearson Chi-Square  $\chi^2= 4.38$ ,  $df=1$ ,  $p=.036$ ; Fisher's exact test  $p=.091$ ). The relation between academic improvement and group was significant (Pearson Chi-Square  $\chi^2= 7.47$ ,  $df=1$ ,  $p=.006$ ; Fisher's exact test  $p=.015$ ). Children who underwent CPAT training had significantly more reports of improved academic performance (8 out of 8, 100%) in comparison to children in the CG group (2 out of 6, 33.3%). Behaviour improvement was not significantly different between the two groups ( $p=1$ ), as teaching reported similar improvements in behaviour in children from both groups. In general, the qualitative data analyses corroborated the pattern that was observed in the quantitative data analyses.

-----Insert Table 5 page 53 about here -----

## DISCUSSION

### *Summary*

Previous studies have shown that children with ASD are characterized with atypical attention performance, and as attention functioning is thought to be linked to academic performance, in the present study we attempted to train attention in primary-aged children with ASD using a previously developed attention training protocol (CPAT; Shalev et al. 2007). We assessed the potential efficacy of CPAT in improving academic attainment of children with ASD by obtaining measures of non-verbal cognition, academic skills, autistic symptomatology as well as semi-structured interviews with teaching staff before and after the training protocol was employed in two schools (one mainstream and one special). Importantly, the functions we measured were not trained within the intervention protocol, which only included simple visual attention tasks (sustained-, selective-spatial- and executive attention). Furthermore, we also included an active control group of children with ASD that underwent a similar protocol of intervention with readily available computer games, using the same format and the same length and frequency of sessions (as well as a one-to-one interaction with an experimenter). This enabled us to verify that any changes observed following attention training could not be attributed to the mere passage of time and/or to the children involvement in an engaging computer-based activity during the school day. Overall, the data indicated promising comprehensive improvements for the children in the CPAT intervention group, which were over and above any improvements obtained in the control group in attention and academic performance.

*Intervention effect: quantitative measures*

Particularly, improvements in the CPAT group were evident in a variety of academic tests including maths, reading comprehension and copying speed. Following the CPAT intervention, children with ASD were able to score higher in a time-limited maths test, were more successful in performing a time-limited reading comprehension test and were also able to increase their copying rate (words per minute) of a given text. Furthermore, scores in a non-verbal cognitive assessment (Raven CPM) also improved for the CPAT group following training. Although the small sample size used here should be kept in mind (Button et al. 2013) all of these improvements were either significant or approaching significance at the ANOVA level and all were significant at the simple effects level. This was also supported by an RCI analysis on the cognitive assessment measure where most of the participants in the CPAT group showed indication of a reliable change in performance. Finally, individual change across the different measures appeared to follow a consistent pattern whereby participants in the control group show both improvements and deterioration in performance while the participants in the CPAT group all show improvements across all measures. Thus, our findings give further support to the growing body of interventions utilizing technology in order to improve or teach new skills to children with ASD, including academic, social and cognitive skills (Bernardini et al. 2012; Bosseler & Massaro 2003; Chabani & Hommel 2014; Ganz, Boles, Goodwyn & Flores 2014; Golan et al. 2010; Hetzroni & Shalem 2005; Hilton et al. 2014; Knight, McKissick & Saunders 2013; Pennington 2010; Vries et al. 2015).

While clear academic and cognitive improvements were observed following the attention training, we found no evidence of change in autistic symptomatology. Total scores from the CARS did not change from the pre- to post-intervention assessment and remained equivalent across the two groups (CPAT and CG). It is worth noting that the CARS is designed to be

used by trained clinicians while in this study it was used by class teachers in the two schools. It was also the case that overall CARS scores were relatively low for children with ASD, even more so in the special school. We hypothesise that this is associated with the fact that non-clinicians were completing the forms and therefore based their report on the relative expression of the behaviours in the participants compared to the rest of the school environment. As such, the teacher from the special school judged the behaviours of the participants in this study relative to other ASD children in the school (possibly with more severe ASD and/or learning disabilities) who may well exhibit more extreme autistic behaviours, which resulted with lower scores compared to the mainstream school. It is also noteworthy that while the CARS is widely used for screening and diagnostic purposes as well as a treatment outcome measure, it was not designed to measure behavioural changes (Aman et al., 2004). It is therefore possible that it is not a sensitive tool for such a purpose. Nevertheless, the lack of change in autistic symptomatology may also be related to the dissociation between attention atypicality in autism and core symptoms (similarly to Eaves and Ho 2004 who did not find any differences in CARS scores post-intervention) so that while attention improvement can benefit academic performance it did not affect autistic symptoms. This differential effect provides further support to the efficiency of the CPAT in improving attention and to its specificity.

#### *Intervention effect: qualitative measures*

The academic and cognitive improvements we observed were also reflected in the thematic analysis of the semi-structured interviews we conducted with teachers and TAs in the two participating schools. Teachers and TAs reported improvements in attention and academic performance, which were more frequently reported for children in the CPAT group. Teaching staff evaluations from the interviews match quite closely the academic tests results in maths

and English (reading and writing) for both the CPAT and control group (in the CPAT group teaching staff reported consistent academic improvements for all children whereas such improvement was reported only for one third of the children in the CG group). Teachers also attributed attentional improvements to children from the CPAT but also to children from the CG group (at least to some degree). This may represent a placebo effect as staff in the schools were informed that all children were taking part in an attention training program. Nevertheless, close examination of the interview reports may suggest some degree of improvement in sustained attention for the CPAT group, which is not consistently mentioned for the CG group. For example, behaviours mentioned for children in the CPAT (but not the CG) group include improving concentration in tasks and in the classroom, being able to concentrate for longer periods of time and showing more self-regulation (less prompting needed) when completing routine classroom tasks.

It is also worth noting that the thematic analysis of the interviews did highlight some behavioural changes (albeit in both groups). In particular, teachers and TAs noted that children in both groups seemed more confident, showed high self-esteem and even demonstrated more socially oriented behaviour. Indeed, these behavioural changes may be attributed to the general aspects of the intervention protocol (across both groups); namely, the participation in a special activity with a dedicated member of staff on a regular basis as a tool for engagement and interaction (Blatchford, Bassett, Brown & Webster 2009). Interestingly, the indication that children in the study were showing more socially oriented behaviour may point to an improvement in social interactions, but perhaps not substantial enough to be measured by CARS.

*Attention training as a domain general approach to improving cognition*

The improvements found in maths, reading, copying and non-verbal intelligence following the attention training, are in accordance to previous reports using the CPAT with children with ADHD (Shalev et al. 2007), children with foetal alcohol spectrum disorder (Kerns et al. 2010) and adult stroke patients (Sampanis et al. 2015), where trained attention skills showed transfer effects for non-trained skills. When used with children with ADHD and with children with foetal alcohol spectrum disorder, the CPAT brought improvements for non-trained academic skills (reading, maths and copying; Kerns et al. 2010; Shalev et al. 2007). Similarly, when the CPAT intervention was utilised with sub-acute stroke patients, improvements were documented in a range of non-trained cognitive domains (e.g., language, memory and number skills; Sampanis et al. 2015). Our results are also in accordance with findings regarding improvement in school performance following a different computerized attention-training program, when used with TD children (Rabiner et al. 2010). While it is possible that training with CPAT affected speed of processing in general, which then manifested in better performance in timed tests (c.f. Sampanis et al., 2015 for consideration of general improvement in sustained attention) we note that we found no obvious evidence for such an effect here. First, performance tended to improve both in the timed academic tests and in the non-timed cognitive assessment and furthermore, the number of attempted questions per test did not differ before and after the intervention. It is therefore less likely that a general speed of processing change is the sole reason for improved performance in our study. Thus, the improvements we report in academic skills and cognitive ability following attention-training support the notion that cognitive training of attention has direct impact on learning and general cognition, and can therefore transfer beyond the trained attentional skills.

Another important aspect to consider here is the high comorbidity between ADHD and ASD symptoms and diagnosis. In a recent review, Antshel, Zhang-James, Wagner, Ledesma, & Faraone (2016) pointed out behavioural, biological and cognitive overlaps between the disorders, and that ADHD tends to be the second most common diagnosis comorbid to ASD after social-anxiety disorder (Simonoff et al., 2008). It is therefore still a possibility that the effect of CPAT intervention we document in our participants with ASD is also mediated by comorbidity with ADHD-like symptoms in our cohort. Indeed, although none of our participants had a diagnosis of ADHD, it is difficult to ascertain whether any attention difficulties CPAT has helped to improve, are primarily associated with ASD or are independent of it.

#### *CPAT as a feasible intervention tool for ASD*

Importantly, our findings also support the notion that using CPAT as an intervention within a school setting is feasible. Parsons et al. (2013) and Kasari and Smith (2013) highlighted the importance of the implementation of intervention programs for ASD in real world settings, such as schools. While the CPAT intervention was challenging to implement due to school breaks, children missing classroom time, need of trained staff and private and quiet training rooms, it was still successfully conducted in the two participating schools. Frequent one or two week breaks took place in both schools (i.e. regular school holidays), which required a break in training sessions. Still, children did not show any major differences in performance on the training sessions after the breaks, indicated by similar number of completed blocks in sessions occurring before and after the break. It is also worth noting that we have found no differences in performance (and outcome) across the mainstream and the special school. This further supports the feasibility of the implementation of CPAT in varied school contexts.

Several further promising points regarding the feasibility and suitability of CPAT for children with ASD in school are worth mentioning. First, all children in the CPAT group were able to increase the number of blocks they completed in each session (from 4 initially to 7 or 8 blocks ultimately) and second the levels of difficulty they have reached varied from 2 to 6 (most of the children reached the 3<sup>rd</sup> level of difficulty in all training tasks). This implies that participants were able to perform the training tasks repeatedly and that they improved their performances within the training sessions themselves. Thus, it appears that participants were able to engage with the program in the way intended so that their performance in each level improved before moving on to the next level of difficulty. Importantly, the levels of difficulty participants reached in the current study are not substantially different than previous uses of the CPAT in different populations. Finally, one more important indicator was the fact that no dropouts were recorded in the CPAT groups, and that teachers and TAs reported high motivation in the children when taking part in the training sessions, concomitant to the schools' active participation in the project and its inclusion in the weekly routine. As such, these indicators suggest a good level of compliance and satisfaction that should be investigated more systematically in future studies.

A related issue is the way this study utilized positive feedback and rewards across the two groups, promoting engagement and motivation for both CPAT and CG groups. Improvements in children's independence and self-esteem, as well as high motivation to participate in the program were noted by all class teachers and TAs in the interviews with respect to children in both groups. Previous studies utilising CPAT have highlighted the importance of the tight schedule of online feedbacks embedded within the program which helps the participant to associate the effort they make and the impact it has on performance as well as driving motivation (Geurts, Luman & Van Meel 2008; Kerns et al. 2010; Shalev et al. 2007).



Motivation is an important prerequisite of successful interventions, as motivational incentives tap into reward mechanisms that can aid attentional control (Padmala & Pessoa 2011). Interestingly, in ASD, evidence points to reduced socially-driven motivation, with non-social reward playing a more effective role (Chevallier, Kohls, Troiani, Brodtkin & Schultz 2012; Mundy 1995). In our study a combination of social and non-social motivational tools were used, such as on-screen feedbacks and points acquired, verbal praise, reward games and stickers, making it difficult to discern what type of motivational tool was more effective in this population. Nevertheless, high motivation or the type of feedback in itself could not explain improvements found in the training group, as these were similar across the CPAT and the CG groups.

#### *Limitations and future directions*

It is important to note, however, that the present study investigated a relatively small-sized cohort (8 and 7 children for the CPAT and CG, respectively) which may have impacted on the statistical analyses we have carried out. For instance, although no statistical differences were found for the baseline measures across the two groups such differences may in fact be present in our data. Also, long-term effects of training were not investigated. It is therefore imperative that future studies will continue to investigate the feasibility and efficacy of CPAT (or other attention training protocols) in a larger cohort as well as the long-term effects of training. Testing a larger cohort will also enable to identify specific pre-requisites that facilitate improvement. For instance, looking at the individual change data (Figure 3) may highlight the potential of some participants to improve more than others. It is therefore important to identify the potential causes for these individual differences. Another factor that should be considered is utilizing a double-blind design to avoid any possible bias scoring post-tests as experimenters in this study were aware of the group assignment at the end of

training; however, all possible measures were taken to ascertain children from both groups were treated equally. Parents' evaluations should also be considered for future research, in order to investigate if the effects of training are limited within the school setting or if training and transfer effects are also noted at home. We also recommend that future studies focus on older children (8 years old and above), with no serious behavioural difficulties, as they might benefit more with the CPAT due to the ability to comply with the programme structure and complete more blocks of each task per session.

### *Conclusion*

In conclusion, the results found in this study regarding the application of a computerized progressive attention training program (CPAT) at school are promising. Not only in terms of its effective use in both a mainstream and a special school, but also the near- and far-transfer of attention training to academic and non-verbal cognitive performance of children with ASD. At the same time, the program was engaging and motivating for the children, bringing improvements in a personal level for both the CPAT and CG groups due to highly structured routine and use of rewards and positive feedback.

## REFERENCES

- Aman, M. G., Novotny, S., Samango-Sprouse, C., Lecavalier, L., Leonard, E., Gadow, K. D., ... Chez, M. (2004). Outcome Measures for Clinical Drug Trials in Autism. *CNS Spectrums*, 9(1), 36–47.
- American Psychiatric Association. (2013). Diagnostic and statistical manual of mental disorders. (5<sup>th</sup> ed.). Arlington: VA, *American Psychiatric Publishing*.
- Antshel, K. M., Zhang-James, Y., Wagner, K., Ledesma, A., & Faraone, S. V. (2016). An update on the comorbidity of ASD and ADHD: A focus on clinical management. *Expert Review of Neurotherapeutics*, 16(3), 279-93.
- Bebko J. M., Perry A., Bryson S. E., (1996). Multiple method validation study of facilitated communication: II. Individual differences and subgroup results. *Journal of Autism and Developmental Disorders*, 26, 19-42.
- Belmonte, M. K., & Yurgelun-Todd, D. A. (2003a). Functional anatomy of impaired selective attention and compensatory processing in autism. *Cognitive brain research*, 17(3), 651-664.
- Belmonte, M. K., & Yurgelun-Todd, D. A. (2003b). Anatomic dissociation of selective and suppressive processes in visual attention. *Neuroimage*, 19(1), 180-189.
- Bernardini, S., Porayska-Pomsta, K., Smith, T. J., & Avramides, K. (2012, September). Building autonomous social partners for autistic children. In *International Conference on Intelligent Virtual Agents*, 46-52. Springer Berlin Heidelberg.
- Blatchford, P., Bassett, P., Brown, P., & Webster, R. (2009). The effect of support staff on pupil engagement and individual attention. *British Educational Research Journal*, 35(5), 661-686.
- Bosseler, A., & Massaro, D. W. (2003). Development and evaluation of a computer-animated tutor for vocabulary and language learning in children with autism. *Journal of autism and developmental disorders*, 33(6), 653-672.
- Braun, V. & Clarke, V. (2006) Using thematic analysis in psychology. *Qualitative Research in Psychology*, 3(2), 77-101.

- Bull, R., & Scerif, G. (2001). Executive functioning as a predictor of children's mathematics ability: Inhibition, switching, and working memory. *Developmental neuropsychology*, 19(3), 273-293.
- Burack, J. A. (1994). Selective attention deficits in persons with autism: preliminary evidence of an inefficient attentional lens. *Journal of Abnormal Psychology*, 103(3), 535.
- Button, K. S., Ioannidis, J. P. A., Mokrysz, C., Nosek, B. A., Flint, J., Robinson, E. S. J., & Munafò, M. R. (2013). Power failure: why small sample size undermines the reliability of neuroscience. *Nature Reviews Neuroscience*, 14(5), 365-376.
- Carver, R.P. (1990). Intelligence and reading ability in grades 2–12. *Intelligence*, 1, 449–455.
- Calhoun, S. L., & Mayes, S. D. (2005). Processing speed in children with clinical disorders. *Psychology in the Schools*, 42, 333–343.
- Chabani, E., & Hommel, B. (2014). Visuospatial processing in children with autism: No evidence for (training-resistant) abnormalities. *Journal of autism and developmental disorders*, 44(9), 2230-2243.
- Chevallier, C., Kohls, G., Troiani, V., Brodtkin, E.S. & Schultz, R.T. (2012). The social motivation theory of autism. *Trends in Cognitive Sciences*, 16 (4), 231-239.
- Chiang, H. M., & Lin, Y. H. (2007). Mathematical ability of students with Asperger syndrome and high-functioning autism a review of literature. *Autism*, 11(6), 547-556.
- Chien, Y. L., Gau, S. F., Shang, C. Y., Chiu, Y. N., Tsai, W. C., & Wu, Y. Y. (2015). Visual memory and sustained attention impairment in youths with autism spectrum disorders. *Psychological Medicine*, 45(11), 2263-2273.
- Chien, Y. L., Gau, S. S. F., Chiu, Y. N., Tsai, W. C., Shang, C. Y., & Wu, Y. Y. (2014). Impaired sustained attention, focused attention, and vigilance in youths with autistic disorder and Asperger's disorder. *Research in Autism Spectrum Disorders*, 8(7), 881-889.
- Chlebowski, C., Green, J. A., Barton, M. L., & Fein, D. (2010). Using the childhood autism rating scale to diagnose autism spectrum disorders. *Journal of Autism and Developmental Disorders*, 40(7), 787-799.

- Ciesielski, K. T., Courchesne, E., & Elmasian, R. (1990). Effects of focused selective attention tasks on event-related potentials in autistic and normal individuals. *Electroencephalography and clinical neurophysiology*, 75(3), 207-220.
- Cotton, S. M., Kiely, P. M., Crewther, D. P., Thomson, B., Laycock, R., & Crewther, S. G. (2005). A normative and reliability study for the Raven's Coloured Progressive Matrices for primary school aged children from Victoria, Australia. *Personality and individual differences*, 39(3), 647-659.
- Dawson, G., Munson, J., Estes, A., Osterling, J., McPartland, J., Toth, K. & Abbott, R. (2002). Neurocognitive function and joint attention ability in young children with autism spectrum disorder versus developmental delay. *Child Development*, 73(2), 345-358.
- Eaves, L. C., & Ho, H. H. (2004). The very early identification of autism: Outcome to age 41/2-5. *Journal of Autism and Developmental Disorders*, 34(4), 367-378.
- Elsabbagh, M., Volein, A., Holmboe, K., Tucker, L., Csibra, G., Baron-Cohen, S., & Johnson, M. H. (2009). Visual orienting in the early broader autism phenotype: disengagement and facilitation. *Journal of Child Psychology and Psychiatry*, 50(5), 637-642.
- Emerson, E., Baines, S., Allerton, L., & Welch, V. (2010). Health inequalities and people with learning disabilities in the UK: 2010. *Durham: Improving Health & Lives: Learning Disabilities Observatory*.
- Erickson, L. C., Thiessen, E. D., Godwin, K. E., Dickerson, J. P., & Fisher, A. V. (2015). Endogenously and exogenously driven selective sustained attention: Contributions to learning in kindergarten children. *Journal of experimental child psychology*, 138, 126-134.
- Estes, A., Rivera, V., Bryan, M., Cali, P., & Dawson, G. (2011). Discrepancies between academic achievement and intellectual ability in higher-functioning school-aged children with autism spectrum disorder. *Journal of Autism and Developmental Disorders*, 41(8), 1044-1052.

- Ganz, J. B., Boles, M. B., Goodwyn, F. D., & Flores, M. M. (2014). Efficacy of handheld electronic visual supports to enhance vocabulary in children with ASD. *Focus on Autism and Other Developmental Disabilities*, 29(1), 3-12.
- Gagnon, L. G., & Belleville, S. (2012). Training of attentional control in mild cognitive impairment with executive deficits: results from a double-blind randomised controlled study. *Neuropsychological Rehabilitation*, 22(6), 809-835.
- Geurts, H. M., Luman, M., & Van Meel, C. S. (2008). What's in a game: the effect of social motivation on interference control in boys with ADHD and autism spectrum disorders. *Journal of Child Psychology and Psychiatry*, 49(8), 848-857.
- Geurts, H. M., Corbett, B., & Solomon, M. (2009). The paradox of cognitive flexibility in autism. *Trends in Cognitive Sciences*, 13(2), 74-82.
- Golan, O., Ashwin, E., Granader, Y., McClintock, S., Day, K., Leggett, V., & Baron-Cohen, S. (2010). Enhancing emotion recognition in children with autism spectrum conditions: An intervention using animated vehicles with real emotional faces. *Journal of Autism and Developmental Disorders*, 40(3), 269-279.
- Golan, O., & Baron-Cohen, S. (2006). Systemizing empathy: teaching adults with Asperger syndrome or high-functioning autism to recognize complex emotions using interactive multimedia. *Development and Psychopathology*, 18(2), 591-617.
- Happé, F., Booth, R., Charlton, R., & Hughes, C. (2006). Executive function deficits in autism spectrum disorders and attention-deficit/hyperactivity disorder: examining profiles across domains and ages. *Brain and Cognition*, 61(1), 25-39.
- Hesse-Biber, S. N. (2010). Mixed methods research: Merging theory with practice. *Guilford Press*.
- Hetzroni, O. E., & Shalem, U. (2005). From logos to orthographic symbols: A multilevel fading computer program for teaching nonverbal children with autism. *Focus on Autism and Other Developmental Disabilities*, 20(4), 201-212.
- Hill, E. L. (2004). Executive dysfunction in autism. *Trends in cognitive sciences*, 8(1), 26-32.

- Hilton, C. L., Cumpata, K., Klohr, C., Gaetke, S., Artner, A., Johnson, H., & Dobbs, S. (2014). Effects of exergaming on executive function and motor skills in children with autism spectrum disorder: A pilot study. *American Journal of Occupational Therapy*, 68(1), 57-65.
- Kasari C., & Smith T. (2013). Interventions in schools for children with autism spectrum disorder: Methods and recommendations. *Autism*, 17, 254-267.
- Keehn, B., Müller, R. A., & Townsend, J. (2013). Atypical attentional networks and the emergence of autism. *Neuroscience & Biobehavioral Reviews*, 37(2), 164-183.
- Keen, D., Webster, A., & Ridley, G. (2016). How well are children with autism spectrum disorder doing academically at school? An overview of the literature. *Autism*, 20(3), 276-294.
- Kenworthy, L., Anthony, L. G., Naiman, D. Q., Cannon, L., Wills, M. C., Luong-Tran, C., ... Wallace, G. L. (2014). Randomized controlled effectiveness trial of executive function intervention for children on the autism spectrum. *Journal of Child Psychology and Psychiatry, and Allied Disciplines*, 55(4), 374-383.
- Knight, V., McKissick, B. R., & Saunders, A. (2013). A review of technology-based interventions to teach academic skills to students with autism spectrum disorder. *Journal of Autism and Developmental Disorders*, 43(11), 2628-2648.
- Landry, R., & Bryson, S. (2004). Impaired disengagement of attention in young children with autism. *Journal of Child Psychology and Psychiatry*, 45, 1115–1122.
- Luna, B., Doll, S. K., Hegedus, S. J., Minshew, N. J., & Sweeney, J. A. (2007). Maturation of executive function in autism. *Biological psychiatry*, 61(4), 474-481.
- Mandy, W., & Lai, M. C. (2016). Annual Research Review: The role of the environment in the developmental psychopathology of autism spectrum condition. *Journal of Child Psychology and Psychiatry*, 57(3), 271-292.
- May, T., Rinehart, N., Wilding, J., & Cornish, K. (2013). The role of attention in the academic attainment of children with autism spectrum disorder. *Journal of autism and developmental disorders*, 43(9), 2147-2158.

- May, T., Rinehart, N. J., Wilding, J., & Cornish, K. (2015). Attention and basic literacy and numeracy in children with Autism Spectrum Disorder: A one-year follow-up study. *Research in Autism Spectrum Disorders*, 9, 193-201.
- Mayes, S. D., & Calhoun, S. L. (2007). Learning, attention, writing, and processing speed in typical children and children with ADHD, autism, anxiety, depression, and oppositional-defiant disorder. *Child Neuropsychology*, 13(6), 469-493.
- Mundy, P. (1995) Joint attention and social emotional approach behaviour in children with autism. *Development and Psychopathology*, 7(1), 63-82.
- Murphy, C. M., Christakou, A., Daly, E. M., Ecker, C., Giampietro, V., Brammer, M., & Rubia, K. (2014). Abnormal functional activation and maturation of fronto-striato-temporal and cerebellar regions during sustained attention in autism spectrum disorder. *American Journal of Psychiatry*, 171(10), 1107-1116.
- Nation, K., Clarke, P., Wright, B., & Williams, C. (2006). Patterns of reading ability in children with autism spectrum disorder. *Journal of autism and developmental disorders*, 36(7), 911-919.
- Navon, D. (1977). Forest before trees: The precedence of global features in visual perception. *Cognitive Psychology*, 9, 353-383.
- Padmala, S., & Pessoa, L. (2011). Reward reduces conflict by enhancing attentional control and biasing visual cortical processing. *Journal of cognitive neuroscience*, 23(11), 3419-3432.
- Parsons, S., Charman, T., Faulkner, R., Ragan, J., Wallace, S., & Wittemeyer, K. (2013). Commentary—bridging the research and practice gap in autism: The importance of creating research partnerships with schools. *Autism*, 17(3), 268-280.
- Pennington, B. F., & Ozonoff, S. (1996). Executive functions and developmental psychopathology. *Journal of child psychology and psychiatry*, 37(1), 51-87.
- Pennington, R. C. (2010). Computer-assisted instruction for teaching academic skills to students with autism spectrum disorders: a review of literature. *Focus on Autism and Other Developmental Disabilities*. 25(4), 239-248.



- Perry, A., Condillac, R. A., Freeman, N. L., Dunn-Geier, J., & Belair, J. (2005). Multi-site study of the Childhood Autism Rating Scale (CARS) in five clinical groups of young children. *Journal of autism and developmental disorders*, 35(5), 625-634.
- Plaisted, K., Swettenham, J., & Rees L. (1999). Children with autism show local processing in a divided attention task and global precedence in a selective attention task. *Journal of Child Psychology and Psychiatry*, 40, 733-742.
- Posner, M. I., & Rothbart, M. K. (2005). Influencing brain networks: Implications for education. *Trends in Cognitive Sciences*, 9, 99–10.
- Raven, J.C., Court, J.H., & Raven, J. (1990). Section 2: Coloured Progressive Matrices (with US Norms) Manual for the Raven's Progressive Matrices and Vocabulary Scales, Oxford, *Oxford Psychologist Press*.
- Raven, J., Raven, J.C., & Court, J.H. (2008). Raven's Educational: Coloured Progressive Matrices and Crichton Vocabulary Scale Manual. *London, NCS Pearson*.
- Rinehart, N. J., Bradshaw, J. L., Tonge, B. J., Brereton, A. V., & Bellgrove, M. A. (2002). A neurobehavioral examination of individuals with high-functioning autism and Asperger's disorder using a fronto-striatal model of dysfunction. *Behavioral and Cognitive Neuroscience Reviews*, 1(2), 164-177.
- Sanders, J., Johnson, K. A., Garavan, H., Gill, M., & Gallagher, L. (2008). A review of neuropsychological and neuroimaging research in autistic spectrum disorders: Attention, inhibition and cognitive flexibility. *Research in Autism Spectrum Disorders*, 2(1), 1-16.
- Sampanis, D. S., Mevorach, C., Shalev, L., Mohammed, S., & Humphreys, G. W. (2015). Reducing Cognitive Deficits after Stroke through Computerized Progressive Attentional Training (CPAT): A Pilot Study. *Physical Medicine and Rehabilitation – International*, 2 (7).
- Shalev, L., Tsal, Y., & Mevorach, C. (2007). Computerized progressive attentional training (CPAT) program: effective direct intervention for children with ADHD. *Child neuropsychology*, 13(4), 382-388.



randomized controlled trial. *Journal of Child Psychology and Psychiatry*, 56(5), 566-576.

Wass, S. V., Scerif, G. & Johnson, M. H. (2012). Training attentional control and working memory: is younger, better? *Developmental Review*, 32(4), 360-387.

Wass, S. V., & Porayska-Pomsta, K. (2014). The uses of cognitive training technologies in the treatment of autism spectrum disorders. *Autism*, 18(8), 851-871.

Whitby, P. J. S., & Mancil, G. R. (2009). Academic achievement profiles of children with high functioning autism and Asperger syndrome: A review of the literature. *Education and Training in Developmental Disabilities*, 551-560.

Yerys, B. E., Hepburn, S. L., Pennington, B. F., & Rogers, S. J. (2007). Executive function in preschoolers with autism: Evidence consistent with a secondary deficit. *Journal of autism and developmental disorders*, 37(6), 1068-1079.

Zwaigenbaum, L., Bryson, S., Rogers, T., Roberts, W., Brian, J., & Szatmari, P. (2005). Behavioral manifestations of autism in the first year of life. *International journal of developmental neuroscience*, 23(2), 143-152.

## **APPENDIX 1**

### **Interview with teachers**

#### **Preamble**

Talk about:

- Quickly about project – aimed at improving basic attentional processes involved in school activities
- Academic assessment (content they are seeing in class – math, reading, writing and copying)
- CARS
- Preferences for break activities

#### **Pre interview**

- What do you think are their difficulties regarding academic, social, attentional and personal domains?
- What are your expectations from the training?

#### **Post interview**

- How are they doing academically, socially, personally now?
- How is their attention in the classroom?
- Any other comments about the project?

## APPENDIX 2

Quotes from the interviews:

### CPAT

*“CH11. and CH14 improved in maths, one level up since the beginning of the project” (ST). CH11. is better at doing tasks, it is easier to get him to work on tasks, and attention has improved” (TA1). “He pays attention to details a lot more” (TA2). “CH14. shows improvement in work boxes, she used to flip, but now seems more attentive and more interested in doing the tasks.” (TA1).*

*“CH16. still shows some disruptive behaviours, not as much after the project. Making progress in maths, better concentration but still needs some prompting sometimes. Can do more now on his own” (ST). “CH16 has improved; it’s easier to get him to focus in activities” (TA1).*

*“CH12 has improved concentration and attention skills, he used to get distracted. Maths improved, numbers, English improved, he is progressing really well” (ST). “He is aware of his own behaviour, and is showing better listening skills now” (TA2).*

*“CH3 is more confident and independent... He does things by himself now. CH3 is showing progress in numeracy, improved focus and concentration but still has good and bad days” (MT1).*

*“CH1 shows greater academic achievement; more likely to go and talk to people; more focus and attention to the task. More independent and confident, can do tasks on her own and asks questions, better communication and social interactions” (MT2).*

Attention training in autism as a potential approach to improving academic performance

*“CH5 is concentrating in tasks a bit more, for longer. CH5 is making academic progress; he can concentrate in tasks in class for longer, before it used to be 2 minutes” (MT3).*

*“CH6 made some academic progress, just not as much as one could possibly hope for. He’s not quite there yet, so inconsistent, some days are better than others. Lower levels of learning, numbers need to keep going back to adding, taking away, and number recognition. Talks to himself more. Has many social interactions, but always on his terms”. (MT1)*

**Control**

*“CH8 is more participative in class, still shows difficulties in doing tasks independently and needs prompting. He is progressing academically in maths but not so much in literacy” (MT3).*

*“CH4 understands pure maths but has difficulties with shapes and meaning of words and mathematical problems. He is more confident and independent. Shows more stereotyped behaviours, has problems concentrating and still needs prompting to do tasks” (MT1).*

*“CH9 still needs prompting to pay attention... Same academic performance” (ST).*

*“Attention slips a bit” (TA1). “He gained a lot of confidence, good computer skills, important for life: learning to interact with the computer” (TA2).*

*“CH10 and CH15. have always been good” (TA1). “CH10 shows good behaviour, progressing in maths, English, good concentration” (ST). “CH15’s behaviour got worse, he is expressing more, communicating more (started to hit). Can concentrate in maths, English not so much, depends on the task needs prompting. Keeps good academic progress not major jump” (ST).*

*“CH13 improved behaviour in activities, completely different during activities” (TA1). “He can sit and work through tasks but not major progress academically” (ST).*

#### *Across groups*

Interestingly, there was a direct comparison between two boys, one from the CPAT and one from the control group, who were from the same classroom, according to MT3: *“CH5 is concentrating in tasks a bit more, for longer.... CH5 (CPAT) is more focused than CH8 (control). CH8 gets more distracted, still needs prompting and has difficulties in doing tasks independently. CH5 is making academic progress, he can concentrate in tasks in class for longer, before it used to be 2 minutes”*. In particular, TA2 mentioned that all children were showing *“smiles on their faces: children have been very enthusiastic to participate” (TA2).*

### FIGURE CAPTIONS

*Figure 1.* Example of displays from the three tasks from the CPAT. To the left, the Shift Stroop-like task (executive attention): in this task, participants were instructed to look for the large (global) smiley face (level 1) pressing the keyboard letter L if it was present, and A when it was absent. In level 2, participants had to attend to the small (local) smiley faces, and in advanced levels they had to switch and attend to both large and small smiley faces, according to pre exposed cues. Middle: Conjunctive Search Task (selective attention). In this task participants had to decide if the display contained a target (red smiley boy), pressing L if present and A if target was absent (the display depicts level 1 where visual load is still low). Right: The Computerized Continuous Performance Task (sustained attention). In this task, participants had to respond to the appearance of a red car (target) while maintaining focus and inhibiting responses to other appearing objects (level 1). The car was only present in 30% of the trials. In advanced levels participants only had to respond when the car appeared inside a black box, and ignore its appearance anywhere else on the screen.

*Figure 2.* Performance before and after intervention for the CPAT and CG groups. *a.* Standard scores from the cognitive assessment (CPM) for pre- and post-intervention tests, for the CPAT (black) and CG (grey) groups. *b.* Scores (accuracy) from the maths assessment for pre- and post-intervention tests, for the CPAT (black) and CG group (grey). *c.* Scores (accuracy) from the reading assessment for pre- and post-tests, for the training (black) and control group (grey). *d.* Number of copied words per minute (copying rate) from the copying assessment for pre-and post-tests, for the training (black) and control group (grey).

*Figure 3.* Individual change performance (post *minus* pre intervention scores) for participants from the CPAT and CG groups. Participants are numbered individually, from 1 to 16. *a.*



Attention training in autism as a potential approach to improving academic performance

Change scores for the cognitive assessment (CPM) showing individual performance for the training (CPAT - black) and control groups (CG - grey). *b.* Change scores for the maths assessment, for all participants in the training (CPAT - black) and control (CG - grey) groups. *c.* Individual change scores for the reading assessment, showing all participants in the training (CPAT - black) and control (CG - grey) groups. *d.* Individual change scores for the copying assessment for all participants in the training (CPAT - black) and control (CG - grey) groups.

**FIGURES**

Figure 1 top



Figure 2 top





Figure 3 Top



## TABLES

*Table 1:* Control (CG) and CPAT groups, showing averages, standard deviation (stdv) and range for age, gender, mental age equivalent, and cognitive scores from the CPM in the pre training assessment.

<b>Groups</b>	<b>All</b>	<b>Control (CG)</b>	<b>CPAT Training</b>
<b>N</b>	15	7	8
<b>Gender</b>	3F, 12M	1F, 6M	2F, 6M
<b>Age mean (Stdv) Range</b>	8 (1.07) 6-10	7.86 (1.21) 6-10	8.13 (0.99) 7-10
<b>CPM age equivalent (Stdv)</b>	7.28 (2.07)	7.73 (1.3)	6.89 (2.4)
<b>CPM standard (Stdv) Range</b>	89.33 (17.2) 55	92.86 (15.5) 45	86.25 (19.04) 55

*Table 2:* teachers and TAs taking part on the interviews, from mainstream and special school.

<b>Mainstream school</b>	<b>Special school</b>
(3 different classrooms)	(same classroom)
3 teachers (MT1, MT2, MT3)	1 teacher (ST)
	2 teaching assistants (TA1, TA2)

*Table 3:* Data (means $\pm$ SEM) for the pre-intervention assessments: CPM, maths, reading, copying and CARS. Scores are presented across all participants, as well as separately for the CG and CPAT groups.

<b>Groups</b>	<b>All</b>	<b>Control (CG)</b>	<b>CPAT training</b>
<b>CPM</b>	89.3 $\pm$ 4.4	92.8 $\pm$ 5.8	86.2 $\pm$ 6.7
<b>Maths</b>	37 $\pm$ 8.6	47.4 $\pm$ 14.1	27.8 $\pm$ 10.1
<b>Reading</b>	35.8 $\pm$ 4.6	31.9 $\pm$ 7.3	39.3 $\pm$ 6.2
<b>Copying</b>	3.5 $\pm$ 0.6	4.3 $\pm$ 1.1	2.7 $\pm$ 0.4
<b>CARS</b>	29.6 $\pm$ 1.3	28.1 $\pm$ 1.3	30.9 $\pm$ 2.2

## Attention training in autism as a potential approach to improving academic performance

Table 4: Pre- and post-intervention assessment measures for all children from the training (CPAT) group. M: male; F: female. SS: special school; MS: mainstream school. Improvements in attention, academic skills and behaviour were based on interview extracts for each child.

Children (age, gender, school)	CPM standard (pre/post)	Maths accuracy (pre/post)	Reading accuracy (pre/post)	Copying words/min (pre/post)	Examples from the interview passages	Improvements in:		Behaviour
						Attention	Academic performance	
<b>CH1 (9, F, MS)</b>	75/95	32/59	50/65	3.9/4.5	<i>“Greater academic achievement. More focus and attention to the task (MT2)”</i>	Yes	Yes	Yes
<b>CH3 (7, M, MS)</b>	75/90	14/32	32/47	2.7/5.2	<i>“Progress in numeracy, improved focus and concentration (MT1)”</i>	Yes	Yes	Yes
<b>CH5 (10, M, S)</b>	120/120	91/91	74/79	3.1/5.2	<i>“Academic progress; can concentrate in tasks for longer (MT3)”</i>	Yes	Yes	No
<b>CH6 (8, M, MS)</b>	85/100	0/14	26/38	1.1/2.5	<i>“Some academic progress; inconsistent. Lower levels of learning (MT1)”</i>	No	Yes	Yes
<b>CH11 (7, M, SS)</b>	75/85	9/32	21/56	3.1/6	<i>“Improved in maths (ST). It is easier to get him to work on tasks, attention has improved (TA1)”</i>	Yes	Yes	Yes
<b>CH12 (8, M, SS)</b>	110/115	32/82	50/62	4.9/8.1	<i>“Improved concentration and attention skills. Maths, English improved (ST)”</i>	Yes	Yes	Yes
<b>CH14 (8, F, SS)</b>	65/75	9/32	26/47	1.6/3.8	<i>“Improved in maths (ST). Seems more attentive and interested in doing the tasks (TA1)”</i>	Yes	Yes	Yes
<b>CH16 (8, M, SS)</b>	85/120	36/73	35/79	1.5/4	<i>“Making progress in maths, better concentration.(ST)”</i>	Yes	Yes	Yes
<b>Number of children who improved</b>	7/8	7/8	8/8	8/8		7/8	8/8	7/8

## Attention training in autism as a potential approach to improving academic performance

Table 5: Pre- and post-intervention assessment measures for all children from the control (CG) group. M: male; F: female. SS: special school; MS: mainstream school. Improvements in attention, academic skills and behaviour were based on interview extracts for each child.

Children (age, gender, school)	CPM standard (pre/post)	Maths accuracy (pre/post)	Reading accuracy (pre/post)	Copying words/min (pre/post)	Examples from the interview passages	Improvements in:		Behaviour
						Attention	Academic performance	
<b>CH4 (8, M, MS)</b>	95/90	86/77	35/35	3/2.6	<i>"Difficulties with shapes and words. Still needs prompting, problems concentrating" (MT1)</i>	No	No	Yes
<b>CH8 (10, M, MS)</b>	90/95	91/100	62/82	8.2/5.2	<i>"Needs prompting. Progressing academically in maths" (MT3)</i>	No	Yes	Yes
<b>CH9 (8, M, SS)</b>	85/90	9/14	35/15	2.6/4.6	<i>"Still needs prompting to pay attention. Same academic performance" (ST)</i>	No	No	Yes
<b>CH10 (8, M, SS)</b>	110/120	23/50	41/26	4.9/3.9	<i>"Always been good. (TA1). Progressing in maths, English, good concentration" (ST)</i>	Yes	Yes	Yes
<b>CH13 (8, M, SS)</b>	70/50	9/5	9/9	1/ 2.1	<i>"Improved behaviour (TA1). Not major progress academically" (ST)</i>	No	No	Yes
<b>CH15 (7, M, SS)</b>	115/115	82/82	35/47	8.3/8.2	<i>"Always been good (TA1). Can concentrate in maths. Keeps good academic progress not major jump" (ST)</i>	Yes	No	No
<b>Number of children who improved</b>	3/6	3/6	2/6	2/6		2/6	2/6	5/6

