

University of Birmingham Research at Birmingham

Can we identify older people most vulnerable to living in cold homes during winter?

Sartini, Claudio; Tammes, Peter; Hay, Alastair D.; Preston, Ian; Lasserson, Daniel; Whincup, Peter H.; Goya Wannamethee, S.; Morris, Richard W.

DOI.

10.1016/j.annepidem.2017.11.008

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version
Peer reviewed version

Citation for published version (Harvard):

Sartini, C, Tammes, P, Hay, AD, Prestón, I, Lasserson, D, Whincup, PH, Goya Wannamethee, S & Morris, RW 2017, 'Can we identify older people most vulnerable to living in cold homes during winter?', *Annals of Epidemiology*. https://doi.org/10.1016/j.annepidem.2017.11.008

Link to publication on Research at Birmingham portal

Publisher Rights Statement:

Accepted for publication in Annals of Epidemiology, publication on 10.1016/j.annepidem.2017.11.008

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

•Users may freely distribute the URL that is used to identify this publication.

- •Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- •User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)

•Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Download date: 20. Apr. 2024

1 **Title**: Can we identify older people most vulnerable to living in cold homes during winter? 2 3 Authors' names and affiliations Claudio Sartini^{1*}, Peter Tammes², Alastair D Hay², Ian Preston³, Daniel Lasserson⁴, Peter H 4 Whincup⁵, S Goya Wannamethee¹, and Richard W Morris² 5 6 ¹ University College London, Department of Primary Care & Population Health, London, UK 7 8 ² Centre for Academic Primary Care, School of Social and Community Medicine, University of 9 Bristol, Bristol, UK ³Centre for Sustainable Energy, Bristol, UK 10 ⁴ Nuffield Department of Medicine, University of Oxford, UK 11 ⁵ Population Health Research Institute, St George's University of London, UK 12 13 * Corresponding author 14 15

Email: c.sartini@ucl.ac.uk

16 Abstract 17 Words: (now 188, max 200) 18 19 **Purpose**: Living in a cold home increases the risk of dying in winter, especially in older 20 people. However, it is unclear which individual factors predict whether older people are 21 living in cold homes. 22 23 Methods: 1402 men aged 74-95 from a UK population-based study reported difficulties in 24 keeping warm during winter answering four simple "yes/no" questions. Associations 25 between individual's characteristics and each of the four self-reported measures of cold 26 homes were estimated using logistic regression models. Next, we investigated whether 27 measures of cold homes predict mortality over the subsequent 2.1 years. 28 29 Results: Manual social class, difficulties making ends meet, and not being married were 30 each associated (p<0.05) with each of the four measures of cold homes (adjusted odds 31 ratios ranged from 1.61 to 4.68). Social isolation, poor respiratory health and grip strength 32 were also associated with reports of cold homes. 126 men died; those who reported the 33 presence of at least three measures cold homes had increased mortality [adjusted hazard 34 ratios 2.85 (95%CI 1.11-7.30, p=0.029)]. 35 36 Conclusions: Older people who find it hard to keep warm in winter, and have an elevated 37 mortality, could be identified using a self-report questionnaire. 38 39 Word count of manuscript text: 2442 words (+ 3 tables embedded in this file)

Background

Excess winter mortality in the United Kingdom (UK) has been partially attributed to cold housing [1, 2], with an extra 5500 more deaths occurring annually in the coldest homes than would occur if those homes were warm [3]. Greater susceptibility of older people to cold has been suggested [4] as they have worse cardiovascular and respiratory profiles at lower indoor [5] and outdoor [6] temperatures. However, it is unclear how to identify older people who particularly find it hard to keep warm in winter [7]. Existing evidence including qualitative [8-10] and quantitative [1, 9, 11, 12] study designs and different measures of cold homes (e.g. self-reported [8, 9] and indoor temperature [5, 11]), do not identify factors related to living in cold homes.

Using data from a UK population-based study of older men, we aim to highlight factors independently associated with living in cold homes to identify vulnerable older people who find it hard to keep warm [13]. We investigate the associations between (i) sociodemographic measures, (ii) health factors, (iii) behavioural factors, (iv) other personal circumstances, and (v) house characteristics with four self-reported measures of cold homes during winter. As it is unclear which measures of cold homes best predict those at risk of death [11], we also investigated whether reports of cold homes in our study relate to mortality.

Material and methods

62 Sample

The British Regional Heart Study (BRHS) is a prospective, population-based cohort study following up 7735 men (99% Caucasian) recruited from primary care practices in 24 British towns in 1978–80. In 2014, 2820 surviving men aged 74-96 years were invited to complete a comprehensive health status and life style questionnaire including self-reported measures of cold homes [14]. 1655 men responded (99% between April and October) and 1402 had complete data on all covariables of interest. Of the 1402, the number of observations available varied from 1385 to 1402. The National Research Ethics Service (NRES) Committee London provided ethical approval. Participants provided informed written consent to the investigation, in accordance with the Declaration of Helsinki.

72 73 Self-reported measures of cold homes 74 Men were asked whether they were (i) having difficulties in meeting the heating/fuel costs; 75 (ii) staying in bed longer in order to stay warm during the previous winter; (iii) unable to 76 keep the living room comfortably warm during the cold winter weather, and (iv) turning the 77 heating off even when cold because of worries about the costs during the previous winter. 78 Having difficulties in meeting the heating/fuel costs was chosen as our main outcome, as an 79 overall proxy measure of cold housing. 80 81 *Individual factors* 82 Individual factors selected in this study were selected on the basis of previously reported 83 associations with cold homes in qualitative [8-10] and quantitative [1, 9, 11, 12] studies: for 84 example, manual social class, difficulties in making ends meet and being not married. 85 Factors investigated were categorised consistently with previous published work from the 86 BRHS [14-19], and represented five different domains: (i) socio-demographic (age, social 87 class, and region of residence), (ii) general health (number of chronic conditions, respiratory 88 health, mobility limitations outdoors, grip strength, depression, and feeling of social 89 isolation), (iii) behavioural factors (smoking and alcohol consumption), (iv) personal 90 circumstances (having increasing financial difficulties and house ownership), and (v) house 91 characteristics (types of home insulation, heating system). Also, a proxy measure of the 92 house energy efficiency (Energy Efficiency rating [20], aggregated from households within 93 participants' Lower Super Output Area [LSOA]) was linked to each of the BRHS men. Energy 94 Efficiency (EE) rating was investigated using descriptive statistics only; as a graded trend in 95 association with the main outcome of interest was not found, this variable was not included 96 in final models. 97 98 Statistical analyses 99 We examined the distribution of all variables of interest according to self-reported 100 measures of cold homes. As the Energy Efficiency rating did not show a graded linear

relationship with self-reported difficulties in keeping warm, we preferred to include house

characteristics collected at individual level (e.g. types of home insulation and heating) in

101

102

103

further analysis.

104 105 Logistic regression models 106 Logistic regression was used to estimate the associations of individual factors with each of 107 the cold home measures, firstly unadjusted, then mutually adjusted. The mutually adjusted 108 models were performed to demonstrate factors which were independently associated with 109 each of the four self-reported measures of cold homes collected in this study. 110 111 112 Subsidiary analyses 113 As older people are less active in winter [21], they are also likely to spend most time at 114 home during this period [7]; therefore, a subsidiary analysis was carried out to find out 115 which individual factors were associated with men both having difficulties in meeting the 116 heating/fuel costs and also being inactive or occasionally active. Physical activity levels were 117 derived from a self-reported questionnaire validated against objectively measured physical 118 activity in a previous study [15]. 119 120 As it is unclear which measures of cold homes best predict those at risk of death [11], we 121 also investigated whether self-reported measures of cold homes, alone or in combination, 122 predicted mortality using Cox proportional hazard models; estimates (hazard ratios) were 123 adjusted for potential confounding factors such as age, social class, marital status, chronic 124 conditions, respiratory health, and region. We selected a combination of three measures of 125 cold homes: 126 1) Having difficulties in meeting the heating/fuel costs 127 2) Inability to keep the living room warm 128 3) Staying in bed longer in order to stay warm 129 130 Question 1 was regarded as an overall proxy measure of cold housing, question 2 was used 131 as the living room is likely to be used the most by older people in the daytime, question 3 132 included information about the bedroom, representing a distinct part of the house in which 133 older people will spend a large number of hours. We believe the 4th measure - "Turning the 134 heating off because of worries of the costs" was less informative than "having difficulties in

meeting the heating/fuel costs", as it did not give any specific information about the room

135

136 where the heating was turned on or. 137 138 Results 139 We found (i) 288 (20.7%) men had difficulties in meeting the heating/fuel costs; (ii) 173 140 (12.4%) stayed in bed longer in order to stay warm; (iii) 47 (3.3%) could not keep 141 comfortably warm in the living room, and (iv) 130 (9.4%) turned heating off because of 142 worries about the costs. Manual social class, increasing financial difficulties, poor health in 143 general, and being not married were more common in men who were having difficulties in 144 meeting the heating/fuel costs (Table 1). Similar findings were found for other self-reported 145 measures of cold homes (results not shown). 146 147 Logistic regression models 148 In unadjusted models (Table 2, left column), many individual factors were associated with 149 having difficulties in meeting the heating/fuel costs. However, in fully adjusted models 150 (Table 2, right column), fewer associations were found. Having increasing financial 151 difficulties showed the strongest association (OR= 4.68, 95%CI 3.74-5.87, p<0.001). Also, 152 men who were of manual social class, not married, feeling socially isolated, having three or 153 more chronic conditions, persistent sputum production, and who were younger were more 154 likely to have difficulties in meeting the heating/fuel costs (all p<0.05). 155 156 Overall, only three individual factors showed consistent associations with each of the four 157 outcomes of interest: manual social class, having increasing financial difficulties, and being 158 not married (see Table 2 and supplementary Table 2). Other less consistent associations 159 were found in fully adjusted models: poor respiratory health (persistent sputum 160 production), lower grip strength and social isolation were associated with at least two out of 161 four measures of cold homes. Increasing age and drinking alcohol daily (vs occasionally) 162 were also inversely associated with cold homes (Table 2 and supplementary Table 2). The 163 absence of cavity/solid wall insulation in the house increased the odds of living in cold 164 homes, (adjusted OR=1.87, 95%CI 0.98-3.55, p=0.057 for inability to keep the living room comfortably warm, and OR=1.47, 95%CI 0.97-2.22, p=0.072 for turning the heating off when 165

cold because of worries about costs, see supplementary Table 2 – Part C). Overall, other

166

167 factors such as smoking, region of residence, and heating system were not consistently 168 associated with cold home reports. 169 170 Subsidiary analyses 171 129 (9.5%) men had both difficulties in meeting the heating/fuel costs and were inactive or 172 occasionally active. Three individual factors showed statistically significant associations with 173 this outcome: having mobility limitations (OR=3.60, 95%CI 2.15-6.04), persistent sputum 174 production (OR=2.07, 95%CI 1.31-3.27), and having increasing financial difficulties (OR= 175 3.77, 95%CI 2.78-5.11). No associations were found between other individual factors and 176 this outcome (results not shown). 177 178 126 men died after completing the questionnaire (median follow-up period of 2.12 years, 179 interquartile range 2.15-2.25 years). Single self-reported measures of cold homes were not 180 associated with mortality in unadjusted cox proportional hazard models (Table 3). However, 181 assenting to having difficulties in meeting the heating/fuel costs, staying in bed longer in 182 order to stay warm, and inability to keep the living room comfortably warm vs not, n=21, 183 predicted all-cause mortality in unadjusted models (unadjusted Hazard Ratio [HR]=2.90, 184 95%CI 1.18-7.09, p=0.020; adjusted HR=2.85, 95%CI 1.11-7.30, p=0.029). Additional 185 adjustment for social isolation, financial difficulties, grip strength and other factors did not 186 alter the magnitude of this association (results not shown). 187 188 Discussion 189 To our knowledge, this is the most comprehensive investigation of associations of individual 190 factors (socio-demographic, economic, health, and house conditions) with self-reported 191 measures of cold homes in older men, and reports of cold homes related to mortality. 192 193 Overall findings 194 Our findings showed that identifying older people who find it hard to keep warm in winter is 195 possible using a self-report questionnaire designed in the BRHS. First, we narrowed down 196 the list of factors which were independently associated with each of the four self-reported 197 measures of cold homes collected in this study; we thought it was important to assess 198 whether an individual factor remained significantly associated with reports of cold homes

after mutual adjustment for other individual factors. As expected these factors were increasing financial difficulties, manual social class, and being not married (e.g. living alone) which are known determinants of fuel poverty [1]. Nevertheless, men having more chronic conditions (three or more vs none), who persistently produced sputum in winter (a marker of chronic lung disease or respiratory infection), with lower grip strength (a marker of physical frailty [22]), and who were feeling socially isolated (an indicator of reduced quality of life [23]) were also more likely to live in cold homes. There was also a suggestion that presence of mobility limitations particularly increased the odds of having difficulties in meeting the heating/fuel costs if the men were also inactive (or occasionally active). As most participants lived in a centrally heated home, absence of cavity or solid wall insulation in the house appeared more relevant to cold housing. With these analyses we have gone beyond findings reported in previous qualitative and quantitative studies which merely listed factors linked with living in cold homes.

Measures of cold homes and mortality

Those who reported cold homes had also increased mortality rates. However, only a specific combination of three measures of cold homes predicted mortality, while single measures of cold homes did not. This means that to identify an exhaustive measure of cold homes is very complex. Keeping the living room warm was more strongly related to mortality than other single measures, possibly because the living room gets used the most by older people.

Comparison with previous studies

Consistent with our main findings, previous studies identified highest levels of fuel poverty in households occupied by a single person over 60 years old (vs couples over 60) with low income [12]. The English Longitudinal Study of Ageing (ELSA) also found that age in older adults was inversely associated with living in cold homes; the authors reported that ELSA participants under 80 years who were living in cold homes had a worse cardiovascular risk profile including higher blood pressure, and were less likely to have blood pressure checked [5]. Interestingly, we also found that men who were drinking alcohol daily (vs occasionally) were less likely to stay in bed longer in order to stay warm, and less likely to turn the heating off because of worries of the costs. To our knowledge these findings were not previously reported and may indicate reduced sensitivity of alcohol consumers to cold [24]

or an indicator of greater financial resources. It is also plausible that daily drinkers spent more time outside and thus required less home heating.

Strengths and limitations

Previous investigations of cold homes have been at household levels [12], while the current study was a large population-based study of older men, thus applying at individual level. We were able to account for confounding between individual factors, and estimated independent associations of these factors with measures of cold homes. To our knowledge, this analysis was not carried out elsewhere. Moreover, the self-reported measures of cold homes we used were similar to the ones collected in other quantitative studies [4, 25], but never used to predict mortality in survival models. A specific combination of three measures of cold homes was associated with a threefold increased mortality risk. Other factors potentially related to cold homes and mortality, for example respiratory infections, objectively measured respiratory function, and biological markers of inflammation (e.g. Interleukin-6 and C-Reactive protein) were not available during the relevant data collection phase. Further studies, which take these variables into account, will need to be undertaken to better understand the mechanism which relates cold homes to mortality The follow-up period for survival analysis was relatively short and the statistical power reduced due to a low number of deaths observed. Future studies with longer follow-up and repeated measures of cold homes over time are required.

The study lacked an objective measure of cold homes, such as indoor temperature (a better marker of thermal efficiency of the dwellings). Also, we observed that a higher EE rating of the house measured at LSOA level did not correspond to less difficulties in meeting the fuel costs in the BRHS; suggesting that this broad measure of EE is less suitable than the individual level data available in the BRHS. However, we acknowledge its relevance in other studies on cold homes at household or macro-area level.

A further minor limitation is the inclusion of only male participants; in the UK and in comparison to men, a higher proportion of the female population are aged 75 and over (9%, compared with 7% of males in 2013 [26]), so we would expect a higher absolute number of women exposed to cold weather, and so cold housing, than men. We would expect that a

cold homes-mortality relationship could be found in the female population; previous reports found that women were more likely to suffer fatal events in a cold period than men [1, 4]. Lastly, although our measure of grip strength was self-reported, our finding was consistent with one from a previous study [5].

We also acknowledge the potential importance of factors which were not measured nor reported in our study, such as biological markers of inflammation and influenza rates. This is a limitation of our study; to measure those factors could have helped in understanding the biological pathways linking cold homes with mortality [5]. Larger studies may explore this important scientific questions in the future. However, our work still makes an important contribution to the literature and enhance the understanding of which profiles of older men live in cold homes, and the implications for their future mortality.

Implications

Our findings suggest that experiencing increasing financial difficulties and lower social class, known to be strongly associated with fuel poverty [27], are not the only factors which increase older people's difficulties in keeping warm during winter. With an aging population, UK policies should acknowledge the detrimental contribution of multiple risk factors which increase with age and are more common in people living in cold homes such as social isolation, poor respiratory health or lower physical function in general. Interventions developed at address these could also reduce winter mortality, as well as interventions to lower fuel payments.

Our findings also suggest that a few simple questions, such as the ones on grip strength and persistent sputum production, may be a useful tool in identifying those who find it hard to keep warm in winter in primary care. Present studies are already evaluating the feasibility of implementing grip strength measurement into routine clinical practice, because it is inexpensive and simple to measure [22]. Other factors related to cold homes in our study are already collected in primary care (e.g. chronic conditions, marital status, and alcohol consumption), while others can be potentially routinely collected in the future (e.g. spirometry to measure lung function [28], or a single item question rather than a complex score to measure social isolation [29]), as part of an admission procedure during winter. This

would help primary care teams in identifying, or improving the assessment of heating needs of, older people who find it hard to keep warm without visit them at home, as the National Institute for Health and Care Excellence (NICE) have recommended in England [2, 13].

Conclusions

Identifying older people who find it hard to keep warm in winter and have an increased mortality risk is possible. Increasing financial difficulties and lower social class are not the only factors which increase older people's difficulties in keeping warm during winter. With an increasing aging population, UK policies need to tackle the adverse effect of multiple risk factors which increase with age and are more common in people living in cold homes, such as social isolation, poor respiratory health and physical frailty.

AUTHORS CONTRIBUTIONS

CS processed the data, performed statistical analyses, drafted and revised the manuscript, and incorporated revisions of co-authors. RWM contributed to the study design and supervised the statistical analyses. SGW, PHW, and RWM contributed to the BRHS design and the acquisition of data. IP contributed to the acquisition of the data from the Centre of Sustainable Energy. All authors provided an important intellectual contribution to the work, revised the manuscript, and approved the final version.

Acknowledgements

This paper presents independent research funded by the National Institute of Health Research School for Primary Care Research (NIHR SPCR grant reference number 281). The NIHR programme grant was awarded to RWM, and supported CS and PT. DSL is funded by the NIHR Oxford Biomedical Research Centre. The British Regional Heart study is supported by a British Heart Foundation (BHF) programme grant (RG/13/16/30528). The funders had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and the decision to submit the manuscript for publication. The views expressed in this publication are those of the author(s) and not necessarily those of the NIHR, the Department of Health, the NHS or BHF.

328 Conflict of interest statement

327

329 The authors report no relationships that could be construed as a conflict of interest

Table 1 – Descriptive statistics of individual characteristics in BRHS men collected on one occasion during 2014 (left column), and descriptive statistics stratified by having difficulties in meeting the fuel costs.

	All (n=1399) ⁷		ving difficulties in ne heating/fuel costs	
	All (n=1399)	Yes (n=288, 20.7%)	No (n=1111, 79.3%)	p-value
Socio-demographic characteristics				
Age (years), mean (SD)	81.0 (4.3)	80.5 (4.2)	81.1 (4.3)	0.025
Social class:				
Manual, n (%)	633 (45.3)	183 (63.5)	450 (40.5)	<0.001
Non-Manual, n (%)	731 (45.3)	100 (34.7)	631 (56.8)	
HMF, n (%)	35 (2.5)	5 (1.7)	30 (2.7)	
Region, n (%)				0.511
South	493 (35.2)	97 (33.7)	396 (35.6)	
Midlands	196 (14.0)	39 (13.5)	157 (14.1)	
North	555 (39.7)	113 (39.2)	442 (39.8)	
Scotland	155 (11.8)	39 (13.5)	116 (10.4)	
General health				
Chronic conditions ¹ , n (%)				<0.001
None	589 (42.1)	99 (34.4)	490 (44.1)	
One/Two	700 (50.0)	150 (52.1)	550 (49.5)	
Three or more	110 (7.9)	39 (13.5)	71 (6.4)	
Chronic obstructive pulmonary disease (COPD) ² : Yes, n (%)	90 (6.4)	31 (10.7)	59 (5.3)	0.001
Persistent sputum production: Yes, n (%)	349 (25.0)	108 (37.5)	241 (21.7)	<0.001
Mobility limitations in getting about outdoor: Yes, n (%)	539 (38.5)	157 (54.5)	382 (34.4)	<0.001
Grip strength ³ : fair/poor vs good/very good, n (%)	256 (18.4)	82 (28.5)	174 (15.7)	<0.001
Depression: Yes vs no, n (%)	145 (10.4)	46 (16.0)	99 (8.9)	<0.001
Feeling isolated from others: Sometimes/often vs rarely/not, n (%)	311 (22.2)	101 (35.1)	210 (18.9)	<0.001

Behavioural factors				
Smoking: Yes vs no, n (%)	47 (3.4)	11 (3.8)	36 (3.2)	0.627
Alcohol consumption, n (%)				0.008
Occasionally	719 (51.4)	165 (57.3)	554 (49.9)	
None	208 (14.9)	48 (16.7)	160 (14.4)	
Daily	472 (33.7)	75 (26.0)	397 (35.7)	
Personal circumstances				
Finance managing ⁴ , n (%)				<0.001
Very well	748 (53.5)	42 (14.6)	706 (63.6)	
Quite well	456 (32.6)	127 (44.1)	329 (29.6)	
Alright or not well	195 (13.9)	119 (41.3)	76 (6.8)	
House ownership ⁵ : Renting/Other vs owner, n (%)	145 (10.4)	40 (13.9)	105 (9.5)	0.028
Present circumstances, n (%)				0.015
Married	1007 (72.0)	196 (68.1)	811 (73.0)	
Single/Alone/Divorced/Separated	97 (6.9)	31 (10.7)	66 (5.9)	
Widowed	295 (21.1)	61 (21.2)	234 (21.1)	
House characteristics				
House centrally heated: No vs Yes, n (%)	146 (10.4)	34 (11.8)	112 (10.1)	0.394
Cavity/solid wall insulation: No vs Yes, n (%)	461 (33.0)	93 (32.3)	368 (33.1)	0.789
House energy efficiency rating ⁶				0.012
1 – Lowest energy efficiency	103 (10.3)	29 (14.4)	74 (9.2)	
2	173 (17.3)	34 (16.9)	139 (17.3)	
3	371 (37.0)	77 (38.3)	294 (36.7)	
4	143 (14.3)	34 (16.9)	109 (13.6)	
5 – Highest energy efficiency	213 (21.2)	27 (13.4)	186 (23.2)	

¹ Men were asked if their doctor had ever diagnosed chronic conditions including angina, heart attack, heart failure, claudication, stroke, diabetes, cancer, chronic kidney disease, osteoporosis, Parkinson's disease.

² Doctor's diagnosis of COPD

³ Rating in comparison with men of same age

⁴ Men were asked to describe how they were managing financially at present

⁵ Renting from local authorities or privately. Category other included living in residential or nursing home (n=5 men), or living in sheltered accommodation (n=9 men), or unspecified accommodation (n=24 men)

⁶ The rating was not self-reported by the BRHS men, but aggregated from households within participants' Lower Super Output Area [LSOA]).

⁷ Descriptive statistics in this table were from 1399 men with complete data on all variables and who answered the question about difficulties in meeting the heating/fuel costs

Table 2 - Cross-sectional associations from logistic regression models between individual characteristics and difficulties in meeting the heating/fuel costs in BRHS men (aged 74-95) who completed a questionnaire in 2014. Per each of the individual characteristics the associations are reported as Odds Ratios (ORs) in comparison to the reference category. The statistically significant results are reported in bold.

			neating/fuel costs? Y	es vs No
	Unadjusted m	odel ⁶	Full adjusted n	nodel ⁷
	OR (95%CI)	p-value	OR (95%CI)	p-value
Socio-demographic characteristics				
Age (years)	0.97(0.94,1.00)	0.026	0.94(0.90,0.98)	0.003
Social class: Non-manual (ref.)	1		1	
Manual	2.57(1.95,3.37)	<0.001	1.66(1.20,2.31)	0.002
HMF	1.05(0.40,2.77)	0.919	1.00(0.34,2.91)	0.998
Region: South (ref.)	1		1	
Midlands	1.01(0.67,1.54)	0.947	1.00(0.61,1.64)	0.999
North	1.04(0.77,1.41)	0.782	0.89(0.62,1.28)	0.541
Scotland	1.37(0.90,2.10)	0.144	1.03(0.62,1.72)	0.903
General health				
Chronic conditions ¹ : None (ref.)	1		1	
One/Two	1.35(1.02,1.79)	0.037	1.21(0.87,1.70)	0.261
Three or more	2.72(1.74,4.25)	<0.001	2.13(1.19,3.82)	0.011
COPD ² : No (ref.)	1		1	
Yes	2.15(1.36,3.39)	0.001	1.00(0.54,1.86)	0.994
Persistent sputum production: No (ref.)	1		1	
Yes	2.17(1.64,2.86)	<0.001	1.83(1.29,2.59)	0.001
Difficulties in getting outdoor: None (ref.)	1		1	
Yes	2.29(1.76,2.97)	<0.001	1.22(0.85,1.74)	0.274
Grip Strength ³ : Good/Very good (ref.)	1		1	
Fair/Poor	2.13(1.58,2.89)	<0.001	1.25(0.85,1.82)	0.253
Depression (van Marwijk score): No (ref.)	1		1	
Yes	1.94(1.33,2.83)	0.001	1.03(0.62,1.71)	0.896
Feeling isolated from others: No/rarely (ref.)	1		1	
Sometimes/often	2.32(1.74,3.08)	<0.001	1.61(1.10,2.37)	0.014
Behavioural factors				
Smoking: No (ref.)	1		1	
Yes	1.19(0.60,2.36)	0.627	0.60(0.25,1.43)	0.249
Alcohol consumption: Occasionally (ref.)	1		1	
None	1.01(0.70,1.45)	0.969	0.81(0.52,1.26)	0.344
Daily	0.63(0.47,0.86)	0.003	0.82(0.57,1.18)	0.294
Personal circumstances				
Finance managing (score 1-3) 4: Well/Quite well/alright or not well	5.10(4.15,6.28)	<0.001	4.68(3.74,5.87)	<0.001

House ownership: Owner (ref.)	1		1	
Renting/Other ⁵	1.55(1.05,2.28)	0.029	0.64(0.39,1.05)	0.076
Present circumstances: Married (ref.)	1		1	
Single/Alone/Divorced/Separated	1.94(1.23,3.06)	0.004	2.20(1.25,3.89)	0.006
Widowed	1.08(0.78,1.49)	0.645	1.10(0.72,1.68)	0.653
House characteristics				
House centrally heated, Yes (ref.)	1		1	
No	1.19(0.79,1.79)	0.394	0.96(0.57,1.61)	0.874
Cavity/solid wall insulation, Yes (ref.)	1		1	
No	0.96(0.73,1.27)	0.789	1.07(0.76,1.50)	0.690

¹ Men were asked if their doctor had ever diagnosed chronic conditions including angina, heart attack, heart failure, claudication, stroke, diabetes, cancer, chronic kidney disease, osteoporosis, Parkinson's disease.

² Doctor's diagnosis of COPD

³ Rating in comparison with men of same age

⁴ Men were asked to describe how they were managing financially at present

⁵ Renting from local authorities or privately. Category other included living in residential or nursing home (n=5 men), or living in sheltered accommodation (n=9 men), or unspecified accommodation (n=24 men)

⁶ Variables included one at a time. Number of observation in all models = 1399

⁷ All listed variables included in the model. Number of observation = 1399

Table 3 – Prospective associations between self-reported measures of cold homes with all-cause mortality in men aged 74-96 years from the BRHS. Results were reported as Hazard Ratios (HR) with 95% Confidence Intervals (CI) from Cox proportional hazard models. Statistically significant HRs are marker in bold.

	All-caus	e Mortality ¹
	Model 1	Model 2
	Unadjusted Model	Full adjusted model ²
	HR (95% CI)	HR (95%CI)
Self-reported measures of cold homes during previous winter		
(1) Having difficulties in meeting the heating/fuel costs	1.14 (0.75, 1.73) p=0.547	1.04 (0.67,1.60) p=0.861
(2) Staying in bed longer in order to stay warm	1.15 (0.69, 1.91) p=0.601	1.05 (0.62, 1.78) p=0.857
(3) Can't keep the living room comfortably warm	1.81 (0.84, 3.88) p=0.127	1.38 (0.64, 3.01) p=0.406
(4) Turning the heating off because of worries about the costs	0.69 (0.33, 1.40) p=0.302	0.62 (0.30, 1.29) p=0.202
Combination of measures		
Assenting to (1) and (2) vs others ³	1.47 (0.80, 2.74) p=0.217	1.34 (0.71, 2.54) p=0.372
Assenting to (1) and (3) vs others ⁴	2.22 (0.98, 5.04) p=0.056	1.80 (0.77, 4.18) p=0.172
Assenting to (1), (2) and (3) vs others ⁵	2.90 (1.18, 7.09) p=0.020	2.85(1.11, 7.30) p=0.029

¹ Median follow-up period of 2.12 years during years 2014-2016; 126 men died during this period (the total number of men included in each of the survival models was 1385).

² Adjusted models for age, social class, region, marital status, number of chronic conditions, and persistent sputum production

³ Men assenting to (1) and (2) were n=89

⁴ Men assenting to (1) and (3) were n=34

⁵ Men assenting to (1), (2) and (3) were n=21

Supplementary table 1 - PART A- Cross-sectional associations from logistic regression models between individual characteristics and self-reported measures of cold homes measured in BRHS men (aged 74-95) who completed a questionnaire in 2014. Per each of the individual characteristics the associations are reported as Odds Ratio (OR) in comparison to the reference category. The statistically significant results are reported in bold.

	Staying in bed longer in order to stay warm during the previous winter n=1393				Inability to keep the living room comfortably warm during the cold winter weather n=1385				Turning the heating off even when cold because of worries about the costs during the previous winter n= 1402			
	Unadjuste	d Model	Adjusted	Model	Unadjuste	d Model	Adjusted I	Model	Unadjusted	l Model	Adjusted	Model
	OR (95%CI)	p-value	OR (95%CI)	p-value	OR (95%CI)	p-value	OR (95%CI)	p-value	OR (95%CI)	p-value	OR (95%CI)	p-value
Socio-demographic characteristics												
Age (years)	0.99 (0.95,1.03)	0.611	0.95 (0.91,0.99)	0.023	0.99 (0.93,1.06)	0.813	0.99 (0.91,1.07)	0.738	0.95 (0.91,1.00)	0.033	0.92 (0.88,0.97)	0.003
Social class: Non-manual (ref.)	1		1		1		1		1		1	
Manual	2.52 (1.79,3.53)	<0.001	1.61 (1.11,2.35)	0.013	4.20 (2.06,8.54)	<0.001	2.71 (1.25,5.88)	0.012	2.63 (1.78,3.89)	<0.001	1.55 (1.00,2.40)	0.049
HMF	1.97 (0.74,5.27)	0.178	1.60 (0.56,4.62)	0.382	4.38 (0.92,20.8 1)	0.063	4.54 (0.87,23.63)	0.072	2.16 (0.73,6.42)	0.165	1.95 (0.61,6.22)	0.257
Region: South (ref.)	1		1		1		1		1		1	
Midlands	0.87 (0.49,1.53)	0.622	0.78 (0.43,1.45)	0.439	0.94 (0.36,2.43)	0.891	1.00 (0.36,2.79)	0.993	0.89 (0.47,1.67)	0.710	0.84 (0.42,1.66)	0.614
North	1.33 (0.91,1.94)	0.141	1.31 (0.87,1.98)	0.196	1.00 (0.50,1.98)	0.995	1.02 (0.49,2.12)	0.967	1.39 (0.91,2.13)	0.126	1.27 (0.80,2.03)	0.306
Scotland	2.07 (1.27,3.39)	0.004	1.81 (1.05,3.11)	0.032	1.42 (0.57,3.51)	0.452	1.10 (0.41,2.98)	0.853	1.57 (0.87,2.83)	0.136	1.19 (0.62,2.30)	0.595
General health												
Chronic conditions ¹ : None (ref.)	1		1		1		1		1		1	
One/Two	1.15 (0.82,1.62)	0.416	0.88 (0.60,1.29)	0.509	1.46 (0.75,2.85)	0.267	1.17 (0.57,2.41)	0.670	1.26 (0.86,1.86)	0.236	1.20 (0.77,1.85)	0.419
Three or more	1.95 (1.14,3.36)	0.015	0.91 (0.48,1.72)	0.768	3.72 (1.57,8.83)	0.003	1.92 (0.71,5.24)	0.200	1.75 (0.93,3.30)	0.085	1.09 (0.51,2.34)	0.825
COPD ² : No (ref.)	1		1		1		1		1		1	
Yes	2.03 (1.19,3.47)	0.009	0.97 (0.51,1.84)	0.923	4.32 (2.07,9.00)	<0.001	2.13 (0.86,5.29)	0.102	1.96 (1.07,3.58)	0.029	1.08 (0.52,2.26)	0.840

Supplementary table 1 - PART B - Cross-sectional associations from logistic regression models between individual characteristics and self-reported measures of cold homes measured in BRHS men (aged 74-95) who completed a questionnaire in 2014.

	Staying in bed longer in order to stay warm during the previous winter n=1393			Inability to keep the living room comfortably warm during the cold winter weather n=1385				Turning the heating off even when cold because of worries about the costs during the previous winter n= 1402				
	Unadjuste	d Model	Adjusted 1	Model	Unadjuste	d Model	Adjusted l	Model	Unadjusted	l Model	Adjusted	Model
General health	OR (95%CI)	p-value	OR (95%CI)	p-value	OR (95%CI)	p-value	OR (95%CI)	p-value	OR (95%CI)	p-value	OR (95%CI)	p-value
Persistent sputum production: No (ref.)	1		1		1		1		1		1	
Yes	1.79 (1.28,2.52)	0.001	1.34 (0.91,1.98)	0.141	2.51 (1.39,4.52)	0.002	1.62 (0.82,3.21)	0.166	2.13 (1.46,3.10)	<0.001	1.83 (1.18,2.84)	0.007
Difficulties in getting outdoor: None (ref.)	1		1		1		1		1		1	
Yes	3.00 (2.16,4.17)	<0.001	2.02 (1.35,3.02)	0.001	2.93 (1.60,5.36)	0.001	1.44 (0.68,3.05)	0.340	1.75 (1.22,2.52)	0.002	0.86 (0.54,1.37)	0.529
Grip Strength ³ : Good/Very good (ref.)	1		1		1		1		1		1	
Fair/Poor	2.99 (2.11,4.23)	<0.001	2.09 (1.41,3.10)	<0.001	1.76 (0.91,3.38)	0.091	0.88 (0.42,1.87)	0.744	2.29 (1.54,3.41)	<0.001	1.59 (1.00,2.53)	0.050
Depression (van Marwijk score): No (ref.)	1		1		1		1		1		1	
Yes	2.13 (1.38,3.28)	0.001	1.01 (0.61,1.69)	0.964	3.16 (1.60,6.24)	0.001	1.69 (0.76,3.76)	0.202	2.06 (1.26,3.36)	0.004	1.02 (0.57,1.84)	0.943
Feeling isolated from others: No/rarely (ref.)	1		1		1		1		1		1	
Sometimes/often	2.97 (2.13,4.15)	<0.001	1.78 (1.19,2.66)	0.005	2.45 (1.35,4.44)	0.003	1.24 (0.60,2.58)	0.565	3.40 (2.35,4.94)	<0.001	2.44 (1.55,3.83)	<0.001
Behavioural factors												
Smoking: No (ref.)	1		1		1		1		1		1	
Yes	2.24 (1.12,4.49)	0.023	1.15 (0.52,2.57)	0.725	2.85 (0.98,8.29)	0.055	1.29 (0.37,4.48)	0.691	1.47 (0.61,3.55)	0.386	0.70 (0.25,1.94)	0.498
Alcohol consumption: Occasionally (ref.)	1		1		1		1		1		1	
None	1.06 (0.69,1.62)	0.785	0.86 (0.54,1.38)	0.545	0.91 (0.41,2.01)	0.811	0.69 (0.30,1.62)	0.400	1.13 (0.69,1.85)	0.618	0.94 (0.55,1.62)	0.820
Daily	0.41 (0.27,0.63)	<0.001	0.46 (0.29,0.71)	0.001	0.45 (0.21,0.95)	0.036	0.54 (0.24,1.21)	0.135	0.63 (0.41,0.97)	0.038	0.72 (0.45,1.16)	0.180

Supplementary table 1 - *PART C* - Cross-sectional associations from logistic regression models between individual characteristics and self-reported measures of cold homes measured in BRHS men (aged 74-95) who completed a questionnaire in 2014. Per each of the individual characteristics the associations are reported as Odds Ratio (OR) in comparison to the reference category. The statistically significant results are reported in bold.

	Staying in bed longer in order to stay warm during the previous winter n=1393				Inability to keep the living room comfortably warm during the cold winter weather n=1385				Turning the heating off even when cold because of worries about the costs during the previous winter n= 1402			
	TT 1'			v f 1 1	TT 11 .			v f 1 1	TT 1'			3.6.1.1
	Unadjuste	d Model	Adjusted 1	Vlodel	Unadjusted	1 Model	Adjusted I	Vlodel	Unadjusted	Model	Adjusted	Model
	OR (95%CI)	p-value	OR (95%CI)	p-value	OR (95%CI)	p-value	OR (95%CI)	p-value	OR (95%CI)	p-value	OR (95%CI)	p-value
Personal circumstances												
Finance managing ⁴ (score 1-3): Well/Quite well/alright or not well	2.20 (1.78,2.71)	<0.001	1.59 (1.25,2.01)	<0.001	3.02 (2.05,4.44)	<0.001	2.24 (1.46,3.43)	<0.001	3.07 (2.41,3.93)	<0.001	2.62 (2.00,3.44)	<0.001
House ownership: Owner (ref.)	1		1		1		1		1		1	
Renting/Other ⁵	2.11 (1.36,3.25)	0.001	0.98 (0.59,1.63)	0.942	2.13 (1.01,4.50)	0.048	0.66 (0.27,1.60)	0.356	1.66 (1.00,2.77)	0.051	0.71 (0.39,1.30)	0.264
Present circumstances: Married (ref.)	1		1		1		1		1		1	
Single/Alone/ Divorced/Separated	2.67 (1.59,4.48)	<0.001	2.60 (1.44,4.70)	0.002	4.31 (2.01,9.23)	<0.001	3.88 (1.60,9.45)	0.003	2.17 (1.19,3.93)	0.011	2.24 (1.13,4.43)	0.021
Widowed	1.93 (1.34,2.79)	<0.001	1.73 (1.12,2.67)	0.013	1.47 (0.72,3.01)	0.292	1.08 (0.47,2.47)	0.864	1.66 (1.09,2.52)	0.018	1.73 (1.04,2.87)	0.034
House characteristics												
House centrally heated, Yes (ref.)	1		1		1		1		1		1	
No	1.27 (0.78,2.06)	0.334	1.02 (0.59,1.77)	0.938	2.73 (1.36,5.48)	0.005	1.70 (0.76,3.79)	0.195	1.04 (0.58,1.86)	0.897	0.81 (0.42,1.57)	0.536
Cavity/solid wall insulation, Yes (ref.)	1		1		1		1		1		1	
No	1.09 (0.78,1.52)	0.630	1.04 (0.72,1.52)	0.818	1.83 (1.02,3.27)	0.043	1.87 (0.98,3.55)	0.057	1.35 (0.93,1.95)	0.116	1.47 (0.97,2.22)	0.072

¹ Men were asked if their doctor had ever diagnosed chronic conditions including angina, heart attack, heart failure, claudication, stroke, diabetes, cancer, chronic kidney disease, osteoporosis, Parkinson's disease.

² Doctor's diagnosis of COPD

³ Rating in comparison with men of same age

⁴ Men were asked to describe how they were managing financially at present

⁵ Renting from local authorities or privately. Category other included living in residential or nursing home (n=5 men), or living in sheltered accommodation (n=9 men), or unspecified accommodation (n=24 men)

Supplementary table 2 – Descriptive statistics of individual factors in BRHS men aged 74-96 who died before the end of the follow up time (left column) vs men who were still alive (right column).

	•	ticipants at the end of e case analysis used ir	•
	Dead	Alive	
	(n=126, 9.1%)	(n=1259, 89.9%)	p-value
Socio-demographic characteristics			
Age at baseline (years), mean (SD)	80.7 (4.1)	83.9 (5.2)	<0.001
Social class:			
Manual, n (%)	65 (51.6)	559 (44.4)	0.234
Non-Manual, n (%)	57 (45.2)	669 (53.1)	
HMF, n (%)	4 (3.2)	31 (2.5)	
Region, n (%)			0.360
South	40 (31.8)	446 (35.4)	
Midlands	179 (13.5)	178 (14.1)	
North	49 (38.9)	500 (39.7)	
Scotland	20 (15.9)	135 (10.7)	
General health			
Chronic conditions ¹ , n (%)			0.012
None	40 (31.8)	544 (43.2)	
One/Two	70 (55.6)	624 (49.6)	
Three or more	16 (12.7)	91 (7.2)	
Chronic obstructive pulmonary disease (COPD) ² : Yes, n (%)	21 (16.7)	68 (5.4)	0.001

Persistent sputum production: Yes, n (%)	49 (38.9)	292 (23.2)	<0.001
Mobility limitations in getting about outdoor: Yes, n (%)	69 (54.8)	461 (36.6)	<0.001
Grip strength ³ : fair/poor vs good/very good, n (%)	26 (20.7)	227 (18.1)	0.481
Depression: Yes vs no, n (%)	20 (15.9)	125 (9.9)	0.038
Feeling isolated from others: Sometimes/often vs rarely/not, n (%)	31 (24.6)	273 (21.7)	0.299
Behavioural factors			
Smoking: Yes vs no, n (%)	6 (4.8)	41 (3.3)	0.374
Alcohol consumption, n (%)			0.636
Occasionally	67 (53.2)	645 (51.2)	
None	21 (16.7)	185 (14.7)	
Daily	38 (30.2)	429 (34.1)	
Personal circumstances			
Finance managing ⁴ , n (%)			0.211
Very well	60 (47.3)	683 (54.3)	
Quite well	43 (34.1)	409 (32.5)	
Alright or not well	23 (18.3)	167 (13.3)	
House ownership ⁵ : Renting/Other vs owner, n (%)	23 (18.25)	121 (9.6)	0.002
Present circumstances, n (%)			0.070
Married	86 (68.3)	914 (72.6)	
Single/Alone/Divorced/Separated	15 (11.9)	81 (6.4)	
Widowed	25 (19.8)	264 (21.0)	
House characteristics			

House centrally heated: No vs Yes, n (%)	14 (11.1)	131 (10.4)	0.805
Cavity/solid wall insulation: No vs Yes, n (%)	40 (31.7)	418 (33.2)	0.741

Men were asked if their doctor had ever diagnosed chronic conditions including angina, heart attack, heart failure, claudication, stroke, diabetes, cancer, chronic kidney disease, osteoporosis, Parkinson's disease.
 Doctor's diagnosis of COPD
 Rating in comparison with men of same age
 Men were asked to describe how they were managing financially at present
 Renting from local authorities or privately. Category other included living in residential or nursing home, living in sheltered accommodation, or unspecified accommodation

References

- [1] Marmot Review Team. The Health Impacts of Cold Homes and Fuel Poverty. 2011. Available from https://www.foe.co.uk/sites/default/files/downloads/cold-homes-health.pdf.
- [2] National Institute for Heath and Care Excellence (NICE) guidelines. Excess winter deaths and illness and the health risks associated with cold homes. Published date: March 2015.

 Available from https://www.nice.org.uk/guidance/ng6/chapter/1-Recommendations
- [3] Dear KB, McMichael AJ. The health impacts of cold homes and fuel poverty. BMJ. 2011;342:d2807.
- [4] Wilkinson P, Pattenden S, Armstrong B, Fletcher A, Kovats RS, Mangtani P, et al. Vulnerability to winter mortality in elderly people in Britain: population based study. BMJ. 2004;329(7467):647-53.
- [5] Shiue I. Cold homes are associated with poor biomarkers and less blood pressure check-up: English Longitudinal Study of Ageing, 2012-2013. Environmental science and pollution research international. 2016;23(7):7055-9.
- [6] Sartini C, Barry SJ, Whincup PH, Wannamethee SG, Lowe GD, Jefferis BJ, et al. Relationship between outdoor temperature and cardiovascular disease risk factors in older people. European journal of preventive cardiology. 2017;24(4):349-56.
- [7] McCartney M. Margaret McCartney: can doctors fix cold homes? BMJ. 2015;350:h1595.
- [8] Tod A, Lusambili A, Cooke J, Homer C, Abbott J, Stocks A, et al. Barriers to keeping warm in later life. Nursing older people. 2013;25(10):22-9.
- [9] Burholt V, Windle G. Keeping warm? Self-reported housing and home energy efficiency factors impacting on older people heating homes in North Wales. Energy Policy. 2006;34(10):1198-208.
- [10] Grey CN, Jiang S, Nascimento C, Rodgers SE, Johnson R, Lyons RA, et al. The short-term health and psychosocial impacts of domestic energy efficiency investments in low-income areas: a controlled before and after study. BMC Public Health. 2017;17(1):140.
- [11] Wilkinson P, Landon M, Armstrong B, Stevenson S, McKee M. Cold comfort: the social and environmental determinants of excess winter death in England, 1986-1996: Joseph Rowntree Foundation, York; 2001.
- [12] Department for Business, Energy & Industrial Strategy. Annual fuel poverty statistics report: 2016. Office of National Statistics. Available from https://www.gov.uk/government/statistics/annual-fuel-poverty-statistics-report-2016.
- [13] Kmietowicz Z. GPs should identify and visit people at risk from cold homes, says NICE. BMJ. 2015;350:h1183.
- [14] Lennon LT, Ramsay SE, Papacosta O, Shaper AG, Wannamethee SG, Whincup PH. Cohort Profile Update: The British Regional Heart Study 1978–2014: 35 years follow-up of cardiovascular disease and ageing. International Journal of Epidemiology. 2015;44(3):826a-g.
- [15] Jefferis BJ, Sartini C, Ash S, Lennon LT, Wannamethee SG, Whincup PH. Validity of questionnaire-based assessment of sedentary behaviour and physical activity in a population-based cohort of older men; comparisons with objectively measured physical activity data. Int J Behav Nutr Phys Act. 2016;13(1):14.
- [16] Walker M, Whincup P, Shaper A. The British Regional Heart Study 1975–2004. International Journal of Epidemiology. 2004;33(6):1185-92.
- [17] Sartini C, Wannamethee SG, Iliffe S, Morris RW, Ash S, Lennon L, et al. Diurnal patterns of objectively measured physical activity and sedentary behaviour in older men. BMC Public Health. 2015;15:609.
- [18] Shaper AG, Pocock SJ, Walker M, Cohen NM, Wale CJ, Thomson AG. British Regional Heart Study: cardiovascular risk factors in middle-aged men in 24 towns. BMJ. 1981;283(6285):179-86.

- [19] Jefferis BJ, Sartini C, Lee IM, Choi M, Amuzu A, Gutierrez C, et al. Adherence to physical activity guidelines in older adults, using objectively measured physical activity in a population-based study. BMC Public Health. 2014;14:382.
- [20] Centre for Sustainable Energy. Mapping domestic energy efficiency in Great Britain. March 2015. Available from https://www.cse.org.uk/projects/view/1305.
- [21] Sartini C, Morris RW, Whincup PH, Wannamethee SG, Ash S, Lennon L, et al. Association of Maximum Temperature With Sedentary Time in Older British Men. J Phys Act Health. 2017;14(4):265-9.
- [22] Ibrahim K, May C, Patel HP, Baxter M, Sayer AA, Roberts H. A feasibility study of implementing grip strength measurement into routine hospital practice (GRImP): study protocol. Pilot Feasibility Stud. 2016;2(27):2-10.
- [23] Findlay RA. Interventions to reduce social isolation amongst older people: where is the evidence? Ageing and Society. 2003;23(05):647-58.
- [24] Freund BJ, Obrien C, Young AJ. Alcohol Ingestion and Temperature Regulation during Cold-Exposure. Journal of Wilderness Medicine. 1994;5(1):88-98.
- [25] Rudge J, Gilchrist R. Excess winter morbidity among older people at risk of cold homes: a population-based study in a London borough. J Public Health (Oxf). 2005;27(4):353-8.
- [26] Office of National Statistics (ONS) bulletin. Excess Winter Mortality in England and Wales: 2013-14 and 2012-13. Available from https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/bulletins/excesswintermortalityinenglandandwales/2014-11-28.
- [27] Iparraguirre J. Have winter fuel payments reduced excess winter mortality in England and Wales? J Public Health (Oxf). 2015;37(1):26-33.
- [28] Rothnie KJ, Mullerova H, Goss H, Chandan J, Quint JK. Validity and Interpretation of Spirometry for Patients in Primary Care. Thorax. 2015;70(Suppl 3):A188-A90.
- [29] Steptoe A, Shankar A, Demakakos P, Wardle J. Social isolation, loneliness, and all-cause mortality in older men and women. Proc Natl Acad Sci. 2013;110(15):5797-801.