UNIVERSITY^{OF} BIRMINGHAM University of Birmingham Research at Birmingham

TNF- α Autocrine Feedback Loops in Human Monocytes

Gane, Jennie M.; Stockley, Robert A.; Sapey, Elizabeth

DOI: 10.1155/2016/1079851

License: Creative Commons: Attribution (CC BY)

Document Version Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Gane, JM, Stockley, RA & Sapey, E 2016, 'TNF- α Autocrine Feedback Loops in Human Monocytes: The Proand Anti-Inflammatory Roles of the TNF- α Receptors Support the Concept of Selective TNFR1 Blockade in Vivo', *Journal of Immunology Research*, vol. 2016, 1079851. https://doi.org/10.1155/2016/1079851

Link to publication on Research at Birmingham portal

Publisher Rights Statement: Checked for eligibility:08/01/2018

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

•Users may freely distribute the URL that is used to identify this publication.

•Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.

•User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?) •Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.



Research Article

TNF- α Autocrine Feedback Loops in Human Monocytes: The Pro- and Anti-Inflammatory Roles of the TNF- α Receptors Support the Concept of Selective TNFR1 Blockade *In Vivo*

Jennie M. Gane,¹ Robert A. Stockley,² and Elizabeth Sapey¹

¹Department of Clinical and Experimental Medicine, University of Birmingham, Edgbaston, Birmingham, UK ²Lung Investigation Unit, University Hospital Birmingham NHS Foundation Trust, Edgbaston, Birmingham, UK

Correspondence should be addressed to Elizabeth Sapey; e.sapey@bham.ac.uk

Received 7 May 2016; Revised 29 July 2016; Accepted 7 August 2016

Academic Editor: Senthamil Selvan

Copyright © 2016 Jennie M. Gane et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Selective TNFR1 blockade in inflammatory diseases is emerging as a clinical strategy. We studied the roles of the two TNF- α receptors, TNFR1 and TNFR2, in human monocytes, the principal producer of TNF- α and central to many TNF- α driven diseases. We hypothesised that TNF- α has pro- and anti-inflammatory effects on monocytes, occurring differentially via TNFR1 and TNFR2. Monocytes were isolated from healthy human subjects and exposed to LPS, plus/minus the addition of blocking antibodies to TNF- α or its receptors. Pro- and anti-inflammatory cytokine production was quantified using real-time PCR and ELISAs. Cell surface expression of TNFR1/2 was measured by flow cytometry. We demonstrated that monocytes vary in the expression patterns of TNFR1 and TNFR2. Autocrine binding of TNF- α led to sustained upregulation of proinflammatory effect. TNFR2 was responsible for binding soluble TNF- α secreted by monocytes, clearing the cytokine from the pericellular environment. TNFR1 blockade did not change the cell surface expression of TNFR2, leaving this receptor free to upregulate IL-10. These novel results support the concept of selective TNFR1 blockade *in vivo* in order that positive anti-inflammatory effects of TNF- α can be retained via TNFR2 ligation.

1. Introduction

TNF- α is classically regarded as a proinflammatory cytokine, playing an important role in the pathophysiology of inflammatory diseases, such as rheumatoid arthritis and inflammatory bowel disease [1]. Anti-TNF- α molecules, which block both soluble and membrane bound TNF- α , are effective in inflammatory arthropathies [2, 3] and inflammatory bowel disease [4, 5]. Despite this, 30% of patients with these diseases do not respond to anti-TNF- α agents and the reasons are unclear [1]. Intriguingly, some patients receiving anti-TNF- α therapy who had multiple sclerosis as a comorbidity developed exacerbation of their neurological disease, suggesting a proinflammatory consequence of generic TNF- α blockade in certain circumstances [6, 7].

Monocytes are the principal source of TNF- α in humans and are believed to be central in many diseases, including

inflammatory bowel disease, arthropathies, and septic shock [1, 8]. Monocytes are also one of the few cells that express both TNF- α receptors in humans; TNFR1 is expressed on most human cells whereas TNFR2 is found only on immune cells and vascular endothelial cells [9]. TNFR1 is historically considered as the receptor through which the majority of the proinflammatory effects are elicited [8, 10]. The role of TNFR2 in general and its intracellular signalling pathways is less clear. Studies using murine knockout models suggest it plays a predominantly anti-inflammatory role, completely or partially protecting against experimentally induced arthritis, encephalitis, and heart failure [8], and it may have an important role in virus elimination [11]. However, in some studies, a proinflammatory role has also been suggested, as TNFR2 knockout mice developed less emphysema in response to cigarette smoke [12].

The importance of characterising the respective roles of TNFR1 and TNFR2 in human cells is being increasingly recognised. If TNFR1, ubiquitously expressed on almost all cell types, is the "proinflammatory" receptor, with TNFR2 responsible for a more immunomodulatory role, selective TNFR1 blockade would seem a more appropriate strategy in chronic inflammatory diseases. Indeed, a phase one study of Atrosab, a humanized monoclonal antibody that specifically blocks TNFR1, has demonstrated an acceptable safety profile and phase two proof-of-concept trials in rheumatoid arthritis and psoriasis are planned for 2016 [13]. In addition, another monoclonal antibody targeting TNFR1, GSK1995057, has been tested in healthy volunteers with an associated reduction in proinflammatory mediators in bronchoalveolar lavage fluid in response to LPS [14]. Conversely, recent studies in platelets from rheumatoid arthritis have shown proinflammatory effects of TNF- α signalling via TNFR2, specifically upregulation of the adhesion molecule P-selectin leading to platelet-neutrophil complex formation [15].

We hypothesised that TNF- α signalling in monocytes has both pro- and anti-inflammatory effects, occurring differentially through TNFR1 and TNFR2. Our experimental aims were as follows: to study the respective roles of the two receptors on autocrine regulation of pro- and anti-inflammatory cytokine production, to assess the expression patterns of both receptors on monocytes, and to study the effects of TNF- α on their cell surface expression pattern.

2. Materials and Methods

2.1. Study Subjects. Peripheral blood monocytes were obtained from 19 healthy subjects (12 male) who were non-smokers and not receiving regular medication. The median age was 30 years (range: 21 to 45) and all subjects provided informed consent. The studies had received approval from a research ethics committee (Research and Ethics Committee number 3359a).

2.2. Isolation of Blood Monocytes. Whole blood was collected into heparinised tubes, layered over Lymphoprep[™] (Axis Shield, Stockport, UK), and centrifuged to obtain a buffy coat layer of PBMCs. The cells were resuspended in sterile PBS containing 2 mM EDTA and 0.1% BSA and CD14+ CD16– monocytes were extracted using the Dynabeads[®] Untouched[™] Human Monocytes kit (Life Technologies, Paisley, UK), following the product protocol. The monocytes (>95% viable by exclusion of trypan blue) were resuspended in culture medium (sterile RPMI medium 1640; Sigma Chemicals Ltd., Poole, UK) supplemented with 10% foetal calf serum, 10% L-glutamine, and 10% penicillin V and streptomycin. Throughout all stages of the isolation, the cells were kept on ice to minimise cellular activation.

2.3. Cell Culture/Stimulation. Cell culture was conducted in 12 or 24-well tissue culture plates (Costar, Fisher Scientific, Loughborough, UK) at 37°C and in 5% CO_2 . Monocytes were plated at a concentration of between 0.25 and 0.45 million per mL of culture medium. The number of replicates used is specified in individual experiments. Salmonella Enteritidis derived LPS (100 ng/mL; Sigma Chemicals Limited, Poole, UK) was used as the TNF- α stimulant. The blocking mAbs used in relevant experiments are shown in Table 1 and are marketed as preventing binding of each relevant protein to its usual ligand, thereby blocking the usual function of that protein. Each mAb was added at a concentration of 10 μ g/mL to 1 mL of culture medium per well (except in Figure 4 where 25 μ g/mL of TNF- α mAb was used to ensure that all sTNF- α and mTNF- α were blocked), following initial concentration response experiments. An irrelevant mouse IgG₁ antibody was used as a control in validation experiments to exclude any nonspecific effect of the mAbs (data not shown). Culture plates containing cells incubated with mAbs were gently agitated for 20 minutes prior to the addition of LPS.

2.4. Measurement of Inflammatory Mediators in Cell Culture Supernatants. Concentrations of the mediators were measured in monocyte supernatants using specific ELISAs according to manufacturer instructions and read using a Synergy HT microplate reader (Biotech, GMI, Ramsey, USA). All samples and standards were run in duplicate and the average was taken as the sample result. Manufacturers' details and lower and upper limits of quantification are given in Table 1.

2.5. RNA Isolation, Reverse Transcription, and Real-Time Quantitative PCR. Adherent cells were pelleted and stored in RNAlater[®] (Life Technologies, Paisley, UK) at -80°C for future study. Messenger RNA was extracted from each cell pellet using the Isolate RNA Minikit (Bioline, London, UK) and adequate quantity and purity of extracted RNA were determined using a Nanodrop 2000c spectrophotometer (Fisher Scientific, Loughborough, UK). RNA samples were reverse transcribed to cDNA using a High Capacity RNAto-cDNA Kit (Life Technologies, Paisley, UK) and real-time quantitative PCR carried out for each gene of interest using TaqMan[™] gene expression assays (Life Technologies, Paisley, UK), as listed in Table 1. All assays were conducted as described previously [16]. GAPDH was found to be stably expressed in a random selection of samples and was therefore used as the normalising gene in the current studies (data not shown). The $2^{-\Delta CT}$ formula was used to calculate the relative expression of mRNA of the genes of interest [17].

2.6. Analysis of TNF- α Receptors 1 and 2 Expression on the Surface of Monocytes. Monocytes (0.25 × 10⁶ per sample) underwent flow cytometry either immediately after isolation or after a relevant period in culture and were labelled with fluorophore-labelled mAbs to TNFR1 and TNFR2, as described in [18]. Control samples labelled with isotype matched control mAbs were run simultaneously to control for nonspecific binding. Full antibody details are provided in Table 1. Positively labelled samples were run in duplicate where specified and the mean result was taken. Doublet events were excluded. MFI results are expressed as the ratio of the intensity of cells labelled with the flow cytometry antibody of interest to those incubated with isotype control flow cytometry antibodies.

		TABLE 1: Assays.		
mAb	Clone number	Fluorophore	Function	Manufacturer
Mouse IgG_1 TNF- α	28401	N/A	Blocking	R and D Systems
Mouse IgG, TNFRI	16805	N/A	Blocking	R and D Systems
Mouse IgG, TNFR2	22210	N/A	Blocking	R and D Systems
Mouse IgG ₁ isotype control	11711	N/A	Control	R and D Systems
Mouse IgG ₁ TNFR1	16803	PE	Flow cytometry	R and D Systems
Mouse IgG _{2A} TNFR2	22235	FITC	Flow cytometry	R and D Systems
Mouse IgG ₁ isotype control	11711	PE	Flow cytometry	R and D Systems
Mouse IgG _{2A} control	X39	FITC	Flow cytometry	BD Biosciences
ELISA	Minimum detectable dose (pg/mL)	Lower limit of quantification (pg/mL)	Upper limit of quantification (pg/mL)	Manufacturer
TNF-a	1.6	15.6	1000	R and D Systems
CXCL8	3.5	31.3	1000	R and D Systems
TNFRI	50	50	45000	Invitrogen
TNFR2	100	100	138000	Invitrogen
Gene symbol (TaqMan [™])	Assay-on-demand number	NCBI RefSeq transcript number	Amplicon length	Manufacturer
GAPDH	Hs9999905_ml	NM_002046.5	122 bp	Life Technologies
$TNF-\alpha$	Hs00174128_ml	NM_000594.3	80 bp	Life Technologies
CXCL8	Hs00174103_ml	NM_000584.3	101 bp	Life Technologies
$IL-1\beta$	Hs01555410_ml	NM_000576.2	91 bp	Life Technologies
IL-6	Hs00985639_ml	NM_000600.3	66 bp	Life Technologies
$TGF-\beta$	Hs00998133_ml	NM_000660.4	57 bp	Life Technologies
IL-10	Hs00961622_m1	NM_000572.2	74 bp	Life Technologies
Details of blocking and flow cytome	rry antibodies. ELISAs and TaqMan gene expres	sion assays are supplied.		

ď.
lie
ď
In
ê
ar
ys
sa
as
ц
sic
es
pr
ex
Je
Sei
<u> </u>
Ia
2
Lac
ų.
ũ
ŝ
SA
Ë
Ξ
es.
qi
õ
E:
an
Ň
eti
Ę
5
Ċ
N N
flc
q
an
ğ
Ġ
ocl
Ыс
íÌ
s
_

TABLE 2: Effect of TNF- α mAb on pro- and anti-inflammatory cytokine mRNA expression in LPS-stimulated monocytes.

	Area under the curve (AUC) Δ CT. hours(median,range)			2)		
	TNF-α	CXCL8	IL-1 β	IL-6	IL-10	TGF- β
LPS	159 (86–233)	3050 (2170–3697)	1571 (1436–1891)	242 (120–1022)	11 (8–13)	33 (21–50)
LPS and TNF- α mAb	152 (54–197)	2248 (1679–2988)	940 (885–1394)	144 (93–235)	6 (4–10)	29 (18–49)
<i>p</i> value		0.03	0.03	0.03	0.03	

The table outlines area under the curve values for time course profiles of each of the cytokines in response to LPS or LPS stimulation in the presence of TNF- α mAb (n = 4). AUC values were compared with a Wilcoxon Signed-Rank Test. p values are one-tailed.

2.7. Statistical Analysis. Data are presented as median and interquartile range/range or mean and standard deviation where percentages are presented. Friedman's test was employed to detect overall differences between experimental conditions or time points, followed by *post hoc* pairwise comparison testing if appropriate. Individual pairwise comparisons were tested using a Wilcoxon Signed-Rank Test. Area under the curve calculations were made using the trapezoid method. Statistical significance was set at $p \le 0.05$. Two tailed p values were used unless otherwise specified. One-tailed p values were employed where previous experimental work provided a clear expectation of effects. Data was analysed using the SPSS statistical program (version 22.0, Chicago, USA).

3. Results

3.1. Autocrine Regulation of Monocyte Inflammatory Cytokine *Production by TNF-\alpha*. To determine whether autocrine binding of TNF- α had a positive feedback effect on monocytes, cells (from n = 4) were stimulated with LPS in the presence or absence of TNF- α mAb. Figure 1 shows the time course profiles of mRNA expression of TNF- α , CXCL8, IL-1 β , IL-6, IL-10, and TGF- β after stimulation with LPS and LPS in the presence of TNF- α mAb. Table 2 details median AUC values for each cytokine time course profile for both experimental conditions. Significantly lower AUC values in LPS-stimulated monocytes cultured in the presence of TNF- α mAb were observed for CXCL8, IL-1 β , IL-6, and IL-10 (p values are shown in Figure 1 if significant). TNF- α and TGF- β mRNA expression were not affected by blockade of secreted TNF- α mAb. To test the hypothesis that the effect of TNF- α blockade would also influence the secreted protein, the concentration of one mediator, CXCL8, was determined in the supernatants from the experiments shown in Figure 1 (n = 4) at each time point by ELISA. A significant reduction in CXCL8 in the LPS-stimulated cell supernatants was observed from 32 hours onwards in the presence of TNF- α mAb, from 172.7 ng/mL (range: 150.6 to 237.0 ng/mL) to 138.3 ng/mL (range: 122.3 to 159.4 ng/mL) at 32 hours (p = 0.03).

3.2. The Role of Individual Receptors. To determine the role of individual TNF- α receptors following autocrine binding of TNF- α , monocytes (from n = 7) were cultured following stimulation with 100 ng/mL LPS in the presence or absence of 10 μ g/mL TNFR1 mAb or TNFR2 mAb or both. No effect was observed for TNF- α mRNA expression (Figure 2(a)).

However, significant reductions in expression of CXCL8, IL- 1β , and the anti-inflammatory cytokine IL-10 were observed when TNFR1 alone was blocked (Figures 2(b), 2(c), and 2(e)) and a similar trend (though not statistically significant) was observed for IL-6 (Figure 2(d)). TNFR2 blockade alone had no effect on proinflammatory gene expression but did result in a reduction in IL-10 expression (see Figure 2(e)) and dual receptor blockade led to greater reduction in IL-10 mRNA than blocking TNFR2 alone.

3.3. TNFR1 and TNFR2 Surface Expression on Monocytes. Current therapeutic strategies to target TNF- α affect signalling through both TNFR1 and TNFR2. Figure 3 shows the distribution of the pattern of receptor expression on freshly isolated monocytes and monocytes which had been stimulated with LPS. There appeared to be four distinct subpopulations of cells in freshly isolated monocytes with some expressing TNFR1 alone (5.8% \pm 5.0%), TNFR2 alone $(34.0\% \pm 34.0\%)$, both $(46.7\% \pm 34\%)$, or neither receptor $(13.5\% \pm 13.3\%)$. However, both before and following stimulation, most cells expressed either TNFR2 alone or both receptors. A small but significant increase in the percentage of monocytes expressing neither receptor was observed in LPSstimulated cells compared to freshly isolated monocytes. Raw data broken down by subject for freshly isolated monocytes is shown in Table 3 and illustrates the high degree of intrasubject variation in expression pattern observed.

3.4. The Autocrine Effect of TNF-Alpha on Cell Surface TNFR1 and TNFR2. We wished to determine whether TNF- α affects the expression of its own receptors, in addition to upregulating cytokines. To investigate this, monocytes were stimulated with LPS with and without an excess of either TNF- α mAb (to prevent cell-derived TNF- α binding to the cell via either receptor) or a mAb specific to either of the two TNF- α receptors (to ensure TNF- α signalling occurred solely via one of its two receptors).

Figures 4(a) and 4(b) show that blockade of TNF- α led to a significant decrease in TNFR1 surface expression and conversely a small but significant increase in TNFR2. In addition, Figure 4(a) shows that TNFR1 membrane expression was decreased further when TNFR2 was blocked rather than TNF- α itself. The results suggest that there is a positive feedback loop in monocytes, where TNFR1 expression is increased following signalling through TNFR2 by TNF- α . An example of flow cytometry overlay histograms from one subject are shown in Figure 5.



FIGURE 1: Effect of TNF- α mAb on pro- and anti-inflammatory cytokine mRNA expression in LPS-stimulated monocytes. Each figure illustrates a time course experiment showing cytokine mRNA expression by monocytes stimulated with LPS (100 ng/mL) ± TNF- α mAb (10 µg/mL). Data points are expressed as median (range). AUC values were compared using the Wilcoxon Signed-Rank Test. *p* values are one-tailed and are shown if significant. A significant reduction in AUC in the presence of TNF- α mAb was observed for CXCL8, IL-1 β , IL-6, and IL-10 (*p* = 0.03). The experiments represent data from 4 subjects. Time points were conducted singly in culture wells and the resultant cellular material in duplicate for subsequent real-time PCR.



FIGURE 2: Effect of selective TNF- α receptor blockade on cytokine mRNA expression. Monocytes were stimulated with 100 ng/mL of LPS (positive control) for 20 hours ± TNFR1/TNFR2 mAb/both (10 µg/mL), and cytokine mRNA expression was measured. Each column shows median (IQR) values. Overall differences between experimental conditions for each cytokine were assessed with Friedman's test and *post hoc* pairwise comparisons conducted if appropriate (*post hoc* p values are shown in the figures). (a) TNF- α : no overall difference observed. (b) CXCL8: Friedman's test p = 0.03. (c) IL-1 β : Friedman's test p < 0.01. (d) IL-6: no overall difference observed. (e) IL-10: Friedman's test p < 0.01. The experiments represent data from 7 subjects. Experimental conditions were conducted singly in culture wells and the resultant cellular material used in duplicate for subsequent real-time PCR.

Given the findings in Figure 4(a), we hypothesised that we would observe an increase in the TNFR1 MFI ratio over time in stimulated monocytes expressing both TNFR1 and TNFR2, greater than any increase in monocytes solely expressing TNFR1, in which TNF- α autocrine feedback via TNFR2 could

not occur. Figure 4(c) shows a trend for an increase over time of TNFRI expression in monocytes expressing both receptors after LPS stimulation (Friedman's test p = 0.09) and shows that by 6 and 22 hours after stimulation TNFRI MFI ratio was significantly greater in the monocytes expressing both

Freshly isolated monocytes	TNFR1 only (%)	TNFR2 only (%)	Both receptors (%)	Neither receptor (%)
Subject one	0	76	1	22
Subject two	9	10	45	36
Subject three	1	72	20	7
Subject four	4	42	42	12
Subject five	6	4	87	3
Subject six	14	1	84	1

TABLE 3: Expression patterns of TNFR1 and TNFR2 on freshly isolated monocytes.

The table shows the expression patterns of TNFR1 and TNFR2 on freshly isolated monocytes from six subjects.



FIGURE 3: Pattern of TNF- α receptor expression on monocytes. Each column shows the mean percentage of monocytes (with SD bars) expressing each of the four possible cell surface receptor combinations, in freshly isolated cells and after stimulation with 100 ng/mL of LPS for 20 hours. Overall differences in receptor distribution subtype for each of the two conditions were assessed with separate Friedman's tests and post hoc pairwise comparisons conducted if appropriate. In freshly isolated unstimulated monocytes, no overall difference was observed. In monocytes stimulated for 20 hours with 100 ng/mL of LPS, Friedman's test was significant (p = 0.02). Significant post hoc p values are shown in the figure and were significant for comparisons with monocytes expressing TNFR1 only. Comparisons for each of the four receptor distribution subtypes between freshly isolated and LPS-stimulated monocytes were conducted with Wilcoxon Signed-Ranks Testing and demonstrated an increase in the percentage of monocytes expressing neither receptor in LPS-stimulated cells. The experiments represent data from 6 subjects. Experimental conditions for freshly isolated monocytes were conducted singly and for LPS-stimulated monocytes in duplicate (due to limitations to cell numbers available).

TNFR1 and TNFR2 than in those expressing TNFR1 only. In contrast, although there was a significant increase in TNFR2 expression over time in monocytes expressing TNFR2 alone

or both receptors, there was no difference between those two monocyte types (Figure 4(d)).

To determine if the reduction in cell surface TNFR1 following TNF- α blockade was due to increased shedding of the receptor into the pericellular environment, soluble TNFR1/TNFR2 concentrations were measured in cell-free supernatants from monocytes (stimulated with LPS with and without a TNF- α mAb). Figures 4(e) and 4(f) show TNF- α mAb caused a small but significant increase in soluble TNFR1 consistent with an autocrine effect but had no effect on soluble TNFR2 concentration.

3.5. Autocrine Binding of Soluble TNF- α Occurs Predominantly via TNFR2 in Monocytes Cultured in Isolation. It was noted in early experiments that the concentration of sTNF- α in the culture medium of monocytes initially increased, in the absence of (Figure 6(a)) and in response to LPS (Figure 6(b)), but was followed by a subsequent decrease.

The TNF- α ELISA detects both free and soluble receptorbound cytokine indicating that the fall in detectable sTNF- α was not due to binding to its soluble receptors but could represent degradation or exclusion from the supernatant by binding to cell surface receptors. Figure 6(c) shows TNF- α concentration in the cell-free supernatant of monocytes stimulated with LPS in the presence or absence of selective TNF- α receptor blockade. Blockade of TNFR1 did not lead to an increase in sTNF- α , whereas blockade of TNFR2 did, indicating the decrease in sTNF- α over time was likely due to binding to TNFR2.

4. Discussion

The current series of experiments provides novel data concerning TNF- α autocrine feedback loops, the respective roles of TNFR1 and TNFR2 in human monocytes, and the finding that human monocytes have a variable expression pattern of TNF- α receptors.

Global TNF- α blockade *in vivo* is generally anti-inflammatory [8] but is associated with the abrogation of both pro- and anti-inflammatory cytokine production in monocytes. Signalling for proinflammatory mediators in monocytes occurred via TNFR1 whereas anti-inflammatory IL-10 upregulation occurred via either TNFR1 or TNFR2, with an additive effect of dual receptor ligation. Data from the current work suggests that on average over 80% of monocytes studied expressed TNFR2, whilst approximately 50% expressed TNFR1 (with a high degree of intrasubject variation). Global



FIGURE 4: Autocrine effects of TNF- α on TNF- α receptor cell surface expression and soluble receptor concentration. In (a) and (b), the surface expression of TNFR1 and TNFR2 on monocytes stimulated with LPS (100 ng/mL) ± TNF- α mAb or selective TNF- α receptor blockade (25 µg/mL) for 20 hours was determined (e.g., TNFR1 expression was determined in monocytes where TNF- α signalling via TNFR2 had been prevented with a blocking mAb). Concentrations were compared to LPS-stimulated cells, using a Wilcoxon Signed-Ranks Test. In (c) and (d), time course profiles of TNFR1 MFI and TNFR2 MFI, respectively, are shown in LPS-stimulated monocytes expressing one or both types of receptor. No overall difference was observed for time course profiles of TNFR1 expression in cells expressing solely TNFR1 (Friedman's p = 0.33) nor cells expressing both receptors (Friedman's p = 0.09). Friedman's test was significant for TNFR2 expression in cells expressing TNFR2 solely (p = 0.02) and cells expressing both receptors (p = 0.02). Significant *post hoc* pairwise comparisons p values are shown in the figures. Differences in MFI at each time point between cell types were assessed with a Wilcoxon Signed-Ranks Test. In (e) TNFR1 and (f) TNFR2, the columns show the concentration in the supernatant of LPS-stimulated monocytes (for 22 hours) ± TNF- α mAb (10 µg/mL). Concentrations were compared to the positive control, LPS-stimulated cells, using a Wilcoxon Signed-Ranks Test. Columns in all figures show median (IQR) values. The experiments represent data from 6 subjects. Experimental conditions/time points for flow cytometry analyses were conducted in duplicate (a–d) with the exception of freshly isolated monocytes (time: 0 hours) in (c) and (d) due to limitations to cell numbers available. Experimental conditions in (e) and (f) were conducted in culture wells in duplicate and in each sample run in duplicate on the subsequent ELISA.



FIGURE 5: An example of flow cytometry overlay histograms. Overlay histograms are shown for gated monocytes from one subject. (a) Four histograms are shown, with fluorescence intensity in the PE channel on the *x*-axis and cell count on the *y*-axis. The black histogram represents LPS-stimulated monocytes incubated with a PE-labelled isotype control antibody. The red histogram represents monocytes stimulated with LPS and labelled with PE-TNFR1 mAb. Green and pink histograms represent LPS-stimulated monocytes incubated in the presence of TNF- α mAb or TNFR2 mAb, respectively, and then labelled with PE-TNFR1 mAb. The altered positions of the histograms illustrate the effect of TNF- α or TNFR2 blockade on subsequent TNFR1 expression. (b) Four histograms are shown, with fluorescence intensity in the FITC channel on the *x*-axis and cell count on the *y*-axis. The black histogram represents LPS-stimulated monocytes incubated with a FITC-labelled isotype control antibody. The red histogram represents monocytes stimulated with LPS and labelled with FITC-TNFR2 mAb. Green and pink histograms overlay each other exactly illustrating that in this subject's monocytes there was no effect of TNF- α or TNFR1 blockade on subsequent TNFR2 expression. Each of the three experimental conditions was conducted alongside samples subjected to the same conditions but incubated with an isotype control flow cytometry antibody. As all isotype control histograms overlaid each other, only one of three is shown in each graph for ease of visualisation.

TNF- α blockade will therefore have a significant effect on both pro- and anti-inflammatory signalling from monocytes, whilst TNFRI blockade would provide a predominantly anti-inflammatory effect.

Previous authors studying the respective roles of the two receptors in human cells have been limited by the use of exogenous sTNF- α or alternatives such as mutant forms of TNF- α which can activate TNFR2 and/or cell lines where mTNF- α has been overexpressed [19–21]. Whilst useful, these models may not resemble naturally occurring events with mTNF- α . In our experimental system, we used LPS as a potent but relevant inducer of TNF- α [22] as mTNF- α must be produced before its release as sTNF- α , providing confidence that signalling could occur through both forms of the cytokine. This makes our experiments more physiological. In addition, the same batch of LPS was used throughout to overcome potential variations between bacterial sources and even batches from the same strain [23] to ensure the best replication of experiments.

IL-10 is also produced by monocytes in response to bacterial products such as LPS [24]. In our study, we saw an additional positive feedback effect of TNF- α on IL-10 mRNA expression, illustrating the complexity of feedback

loops as IL-10 itself can downregulate TNF- α expression [25]. IL-10 mRNA expression was upregulated by TNF- α binding to either receptor, additively, and importantly TNFR2 ligation alone induced IL-10 upregulation in the absence of any significant upregulation of proinflammatory cytokines. IL-10 is a key immune-regulatory cytokine with anti-inflammatory effects in many diseases. Monocytes, in particular the dominant CD14+ CD16– monocytes studied here, and other cells of the monocyte lineage are a major source of IL-10 [26]. By blocking the TNF- α molecule itself, for example, with infliximab, any autocrine upregulation of IL-10 as a result of TNF- α binding might be lost, disrupting the proinflammatory/anti-inflammatory balance, and may explain the adverse events in patients with multiple sclerosis [6, 7].

These findings provide supporting evidence in favour of TNFR1 blockade *in vivo*, in diseases in which TNFR1 signalling has been shown to play the dominant role in pathogenesis, such as inflammatory arthritis and multiple sclerosis [8, 27]. TNFR1 expression is ubiquitous, whereas TNFR2 expression is limited to leukocytes and vascular endothelial cells [9]. Selective blockade of TNFR1 in a clinical setting might therefore ameliorate the proinflammatory



FIGURE 6: Autocrine binding of sTNF- α secreted by monocytes. Time course profiles for monocytes cultured alone (a) or with 100 ng/mL of LPS (b) are shown. Each datum point is the median value (range). (a) Friedman's test: p < 0.01. Significant post hoc pairwise comparisons of each time point to time zero are shown in the graphs. (b) Friedman's test: p < 0.001. Significant post hoc pairwise comparisons of each time point to time zero are shown in the graphs. (c) Monocytes were cultured with 100 ng/mL of LPS for 20 hours $\pm 10 \,\mu$ g/mL of TNFR1 mAb or TNFR2 mAb or both mAbs. Each column shows median and IQR values. Friedman's test: p < 0.001. *Post hoc* pairwise comparisons p values are shown in the figures. The experiments represent data from 5 subjects (a and b) or 7 subjects (c). Time points in (a) and (b) were conducted in culture wells singly and experimental conditions in (c) in duplicate. Each sample was run in duplicate on the subsequent ELISAs.

effects of TNF- α , whilst allowing ongoing beneficial effects of anti-inflammatory IL-10 produced by monocytes via TNFR2 signalling. To this end, our data supports preliminary *in vivo* studies examining the role of selective TNFR1 blockade in an LPS-model designed to simulate acute lung injury [14].

The reduction in sTNF- α in the supernatants of our experiments with time is likely to reflect internalisation and degradation of the ligand-receptor complex [28]. The majority of sTNF- α binds to TNFR2, an intriguing finding given that mTNF- α rather than sTNF- α is believed to stimulate TNFR2 [19, 29]. This may represent a previously unrecognised homeostatic mechanism for limiting sTNF- α ligation to TNFR1, in the same way that soluble TNF- α receptors are able to exclude active TNF- α in the circulation [9]. Similar

reduction in sTNF- α has also been observed in monocytederived macrophages, indicating this possible homeostatic mechanism is not restricted to monocytes alone but also to other major cellular producers of TNF- α [30]. This concept requires further study as our experiments studied monocytes in isolation and hence did not assess competition for sTNF- α binding by TNFR1 on other cell types.

A further novel finding was that signalling via TNFR2 also upregulated the cell surface expression of TNFR1, most likely through increased production of the receptor rather than reduced shedding into the supernatant, as TNF- α blockade had only a slight effect on soluble TNFR1 concentration. TNFR1 upregulation did not seem to have an appreciable proinflammatory effect, as blockade of TNFR2 did not reduce



FIGURE 7: TNF- α effects on monocytes via TNR1 and TNFR2. The figure shows two adjacent monocytes. (1) LPS induces both proand anti-inflammatory cytokine gene/protein expression. (2) TNF- α , induced by LPS stimulation, is expressed at the monocyte surface as mTNF- α and cleaved to sTNF- α by the enzyme TACE. (3) sTNF- α signals in an autocrine/paracrine fashion via TNFR1 to augment the expression of CXCL8, IL1- β , IL-6, and IL-10 mRNA and (4) mTNF- α via TNFR2 to further augment IL-10 mRNA expression. (5) Autocrine/paracrine binding of membrane TNF- α to TNFR2 also upregulates the expression of TNFR1 at the cell surface. (6) TNFR2 binds sTNF- α . Selective TNFR1 blockade will switch off TNF- α driven proinflammatory cytokine production whilst leaving TNFR2 free to upregulate the immunoregulatory cytokine, IL-10, and possibly clear a proportion of sTNF- α from the pericellular environment.

proinflammatory cytokine output (through reduction of cell surface TNFRI). The relevance of this increase in TNFRI secondary to TNF- α signalling via TNFR2 on monocytes, which can express both receptors, is unknown but would not negate the clinical efficacy of selectively blocking TNFR1 as any increase in TNFR1 expression on cells by TNFR2 ligation would still be blocked by the TNFR1 mAb. It is also important to note that TNFR1 blockade did not reduce TNFR2 expression on the monocyte surface thereby leaving it available to bind TNF- α and induce IL-10 production. The increase in TNFR2 receptors over time is likely to have occurred as a direct result of LPS stimulation (as shown recently [31]). Figure 7 summarises concepts from the data in pictorial form.

There are some limitations to our study. Firstly, as a practical issue, the monocyte yield is low compared to other leukocytes, and hence the number of cellular functions and experimental conditions we could study at one time was limited, thus requiring separate experiments to generate a global view. Secondly, it was not possible to include a negative control arm to experiments, whereby monocytes were not exposed to LPS. However, validation experiments revealed that monocytes cultured in the absence of LPS became apoptotic, especially at later time points. This meant that obtaining a sufficient number of events for each flow cytometry sample for viable unstimulated cells would not have been possible. Thirdly, if targeted TNFR1 blockade in TNF- α disease is to become a viable strategy, it will be necessary to replicate and extend our experiments within the relevant disease cohorts.

Further delineation of the TNFR1 and TNFR2 intracellular signalling pathways in monocytes is also needed. The ability of TNFR2 ligation to induce signalling pathways that initiate IL-10 upregulation in the absence of any upregulation of proinflammatory cytokines, whilst TNFR1 ligation can do both, is of importance in abrogation of the inflammatory/anti-inflammatory balance and hence damage versus resolution. Determining the steps in the pathways should make it possible to separate these effects.

In addition, further work is needed to clarify the role of TNFR2 in monocytes. Whether monocytes that express both TNFR1 and TNFR2 differ substantially in function from those only expressing TNFR1 or TNFR2 and whether expression patterns change in response to a stimulus or as part of an illness are of particular interest. Intriguingly, approximately 15% to 25% of monocytes in both the freshly isolated and stimulated state expressed neither TNFR1 nor TNFR2 and the percentage of monocytes with this subtype increased slightly in response to LPS stimulation. The significance of this finding in the context of monocyte function is unclear and could be investigated further by separating monocytes

by TNFR1/TNFR2 status by flow cytometry and tracking changes to receptor expression within each subgroup. Furthermore, it would be informative to examine the functional effects of selective TNFR2 blockade (both soluble and membrane bound). The cell-free supernatants from monocytes cultured in the presence or absence of TNFR2 mAb could be harvested and used as a stimulus for neutrophils or endothelial cells. This would identify if the excess pericellular soluble TNF- α , present as a direct result of TNFR2 blockade, could enhance release of neutrophil products such as reactive oxygen species or the expression of adhesion molecules on endothelial cells. This would support the hypothesis that membrane bound TNFR2 has an important homeostatic role in clearing TNF- α from the pericellular environment and further support selective TNFR1 blockade *in vivo*.

Lastly, it would be informative to determine whether the imbalance in favour of IL-10 secretion in the presence of TNFR1 blockade has a beneficial effect on macrophage differentiation. IL-10 drives monocytes towards the more immunomodulatory M2 macrophage phenotype (rather than the type 1 proinflammatory M1 phenotype [32]), which in itself could have positive effects *in vivo* in TNF- α associated inflammatory diseases. Both macrophage phenotypes can be produced *in vitro* and hence the relative roles of TNFR1 and TNFR2 could be studied by macrophage subtype. However, this may not be relevant *in vivo* as recent studies suggest that macrophages are rarely derived from monocytes [33]. Thus, it would be more relevant to conduct such studies with tissue derived macrophages.

5. Conclusions

In susceptible individuals, disordered and excessive TNF- α signalling leads to chronic inflammatory disease. TNFR1 is ubiquitous and its ligation by TNF- α has widespread proinflammatory effects, whereas TNFR2 expression is limited mainly to leukocytes and its function in humans has been less clear. The data presented here highlights that monocytes vary in their expression patterns of TNFR1 and TNFR2 but support the concept that proinflammatory effects of TNF- α occur via TNFR1 whilst TNFR2 predominantly plays an anti-inflammatory role by increasing IL-10 output and removing sTNF- α from the pericellular environment. These findings may support the current strategy of selective TNFR1 blockade for TNF- α driven chronic inflammatory disease.

Abbreviations

μg:	Microgram
AUC:	Area under the curve
BSA:	Bovine serum albumin
CD:	Cluster of differentiation
cDNA:	Complementary deoxyribonucleic acid
CXCL8:	Interleukin 8
EDTA:	Ethylenediaminetetraacetic acid
ELISA:	Enzyme-linked immunosorbent assay
FITC:	Fluorescein isothiocyanate
GAPDH:	Glyceraldehyde 3-phosphate dehydrogenase
Ig:	Immunoglobulin

IL:	Interleukin
IQR:	Interquartile range
LPS:	Lipopolysaccharide
mAb:	Monoclonal antibody
(m)RNA:	(Messenger) ribonucleic acid
mL:	Millilitre
MFI:	Median fluorescence intensity
ng:	Nanogram
PBMC:	Peripheral blood mononuclear cell
PBS:	Phosphate buffered saline
PCR:	Polymerase chain reaction
PE:	Phycoerythrin
pg:	Picogram
RPMI:	Roswell Park Memorial Institute
TGF- β :	Tumour growth factor-beta
TNF-α:	Tumour necrosis factor-alpha
TNFR1/2:	Tumour necrosis factor-alpha receptor
	one/two.

Ethical Approval

All procedures performed in studies involving human participants were in accordance with the ethical standards of the University of Birmingham and national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Consent

Informed consent was obtained from all individual participants included in the study.

Competing Interests

The authors declare that they have no conflict of interests.

Authors' Contributions

Jennie M. Gane designed and conducted the experiments and wrote the manuscript. Elizabeth Sapey and Robert A. Stockley supervised the project and reviewed the manuscript.

Acknowledgments

The authors thank Dr. Gillian McNab for laboratory assistance. This study is supported by the West Midlands Chest Fund.

References

- N. Parameswaran and S. Patial, "Tumor necrosis factor-a signaling in macrophages," *Critical Reviews in Eukaryotic Gene Expression*, vol. 20, no. 2, pp. 87–103, 2010.
- [2] J. A. Singh, R. Christensen, G. A. Wells et al., "Biologics for rheumatoid arthritis: an overview of Cochrane reviews," *Cochrane Database of Systematic Reviews*, no. 4, Article ID CD007848, 2009.

- [3] J. Yost and J. E. Gudjonsson, "The role of TNF inhibitors in psoriasis therapy: new implications for associated comorbidities," *F1000 Medicine Reports*, vol. 1, article 30, 2009.
- [4] J. R. Cummings, S. Keshav, and S. P. L. Travis, "Medical management of Crohn's disease," *British Medical Journal*, vol. 336, no. 7652, pp. 1062–1066, 2008.
- [5] S. Danese and C. Fiocchi, "Ulcerative colitis," *The New England Journal of Medicine*, vol. 365, no. 18, pp. 1713–1725, 2011.
- [6] X. Chen and J. J. Oppenheim, "Contrasting effects of TNF and anti-TNF on the activation of effector T cells and regulatory T cells in autoimmunity," *FEBS Letters*, vol. 585, no. 23, pp. 3611– 3618, 2011.
- [7] E. Andreadou, E. Kemanetzoglou, C. Brokalaki et al., "Demyelinating disease following anti-tnfa treatment: a causal or coincidental association? Report of four cases and review of the literature," *Case Reports in Neurological Medicine*, vol. 2013, Article ID 671935, 9 pages, 2013.
- [8] S. Blüml, C. Scheinecker, J. S. Smolen, and K. Redlich, "Targeting TNF receptors in rheumatoid arthritis," *International Immunology*, vol. 24, no. 5, pp. 275–281, 2012.
- [9] B. B. Aggarwal, "Signalling pathways of the TNF superfamily: A double-edged sword," *Nature Reviews Immunology*, vol. 3, no. 9, pp. 745–756, 2003.
- [10] D. J. MacEwan, "TNF receptor subtype signalling: differences and cellular consequences," *Cellular Signalling*, vol. 14, no. 6, pp. 477–492, 2002.
- [11] M. I. Kafrouni, G. R. Brown, and D. L. Thiele, "The role of TNF-TNFR2 interactions in generation of CTL responses and clearance of hepatic adenovirus infection," *Journal of Leukocyte Biology*, vol. 74, no. 4, pp. 564–571, 2003.
- [12] A. I. D'hulst, K. R. Bracke, T. Maes et al., "Role of tumour necrosis factor-α receptor p75 in cigarette smoke-induced pulmonary inflammation and emphysema," *European Respiratory Journal*, vol. 28, no. 1, pp. 102–112, 2006.
- [13] Baliopharm, "Product Pipeline /ATROSAB," 2016, http://www .baliopharm.com/de/product-pipeline/atrosab.html.
- [14] A. Proudfoot, C. O'Kane, A. Bayliffe et al., "A novel TNFRItargeting domain antibody attenuates pulmonary inflammation in a human model of lung injury, via actions on the lung microvascular endothelium," in A47. Critical Illness: Novel Molecules and Models, A6589, American Thoracic Society, 2014.
- [15] A. A. Manfredi, M. Baldini, M. Camera et al., "Anti-TNF α agents curb platelet activation in patients with rheumatoid arthritis," *Annals of the Rheumatic Diseases*, vol. 75, no. 8, pp. 1511–1520, 2016.
- [16] J. M. Gane, R. A. Stockley, and E. Sapey, "The rs361525 polymorphism does not increase production of tumor necrosis factor alpha by monocytes from alpha-1 antitrypsin deficient subjects with chronic obstructive pulmonary disease—a pilot study," *Journal of Negative Results in BioMedicine*, vol. 14, no. 1, article 20, 2015.
- [17] M. L. Wong and J. F. Medrano, "Real-time PCR for mRNA quantitation," *BioTechniques*, vol. 39, no. 1, pp. 75–85, 2005.
- [18] Systems RaD, Flow Cytometry Protocol for Staining Membrane-Associated Proteins in Suspended Cells, 2015, https://www .rndsystems.com/resources/protocols/flow-cytometry-protocolstaining-membrane-associated-proteins-suspended-cells.
- [19] M. Grell, E. Douni, H. Wajant et al., "The transmembrane form of tumor necrosis factor is the prime activating ligand of the 80 kDa tumor necrosis factor receptor," *Cell*, vol. 83, no. 5, pp. 793– 802, 1995.

- [20] J. A. J. Barbara, W. B. Smith, J. R. Gamble et al., "Dissociation of TNF-α cytotoxic and proinflammatory activities by p55 receptor- and p75 receptor-selective TNF-α mutants," *The EMBO Journal*, vol. 13, no. 4, pp. 843–850, 1994.
- [21] S. Bryde, I. Grunwald, A. Hammer et al., "Tumor necrosis factor (TNF)-functionalized nanostructured particles for the stimulation of membrane TNF-specific cell responses," *Bioconjugate Chemistry*, vol. 16, no. 6, pp. 1459–1467, 2005.
- [22] J. H. Youn, Y. J. Oh, E. S. Kim, J. E. Choi, and J.-S. Shin, "High mobility group box 1 protein binding to lipopolysaccharide facilitates transfer of lipopolysaccharide to CD14 and enhances lipopolysaccharide-mediated TNF- α production in human monocytes," *The Journal of Immunology*, vol. 180, no. 7, pp. 5067–5074, 2008.
- [23] Sigma-Aldrich, "Lipopolysaccharides from Salmonella enteriditis-Product Information," http://www.sigmaaldrich.com/catalog/product/sigma/l6011.
- [24] M. Saraiva and A. O'Garra, "The regulation of IL-10 production by immune cells," *Nature Reviews Immunology*, vol. 10, no. 3, pp. 170–181, 2010.
- [25] H. L. Dickensheets, S. L. Freeman, M. F. Smith Jr., and R. P. Donnelly, "Interleukin-10 upregulates tumor necrosis factor receptor type-II (p75) gene expression in endotoxin-stimulated human monocytes," *Blood*, vol. 90, no. 10, pp. 4162–4171, 1997.
- [26] K. Asadullah, W. Sterry, and H. D. Volk, "Interleukin-10 therapy—review of a new approach," *Pharmacological Reviews*, vol. 55, no. 2, pp. 241–269, 2003.
- [27] G. C. Suvannavejh, H.-O. Lee, J. Padilla, M. C. Dal Canto, T. A. Barrett, and S. D. Miller, "Divergent roles for p55 and p75 tumor necrosis factor receptors in the pathogenesis of MOG₃₅₋₅₅induced experimental autoimmune encephalomyelitis," *Cellular Immunology*, vol. 205, no. 1, pp. 24–33, 2000.
- [28] K. Imamura, D. Spriggs, and D. Kufe, "Expression of tumor necrosis factor receptors on human monocytes and internalization of receptor bound ligand," *Journal of Immunology*, vol. 139, no. 9, pp. 2989–2992, 1987.
- [29] H. Rauert, A. Wicovsky, N. Müller et al., "Membrane Tumor Necrosis Factor (TNF) induces p100 processing via TNF receptor-2 (TNFR2)," *The Journal of Biological Chemistry*, vol. 285, no. 10, pp. 7394–7404, 2010.
- [30] F. Gantner, R. Kupferschmidt, C. Schudt, A. Wendel, and A. Hatzelmann, "In vitro differentiation of human monocytes to macrophages: change of PDE profile and its relationship to suppression of tumour necrosis factor-α release by PDE inhibitors," *British Journal of Pharmacology*, vol. 121, no. 2, pp. 221–231, 1997.
- [31] J. A. Lopatnikova, F. F. Vasilyev, A. A. Alshevskaya, and S. V. Sennikov, "Quantitative flow cytometric analysis of expression of tumor necrosis factor receptor types I and II on mononuclear cells," *Journal of Receptors and Signal Transduction*, vol. 33, no. 1, pp. 49–55, 2013.
- [32] N. Wang, H. Liang, and K. Zen, "Molecular mechanisms that influence the macrophage M1-M2 polarization balance," *Frontiers in Immunology*, vol. 5, article 614, 2014.
- [33] F. Ginhoux and S. Jung, "Monocytes and macrophages: developmental pathways and tissue homeostasis," *Nature Reviews Immunology*, vol. 14, no. 6, pp. 392–404, 2014.





The Scientific World Journal



Research and Practice









Computational and Mathematical Methods in Medicine

Behavioural Neurology





Oxidative Medicine and Cellular Longevity