

## Cinacalcet for symptomatic hypercalcemia caused by AP2S1 mutations

Howles, Sarah A.; Hannan, Fadil M.; Babinsky, Valerie N.; Rogers, Angela; Gorvin, Caroline M.; Rust, Nigel; Richardson, Tristan; McKenna, Malachi J.; Nesbit, M. Andrew; Thakker, Rajesh V.

DOI:

[10.1056/NEJMc1511646](https://doi.org/10.1056/NEJMc1511646)

License:

None: All rights reserved

### Document Version

Publisher's PDF, also known as Version of record

### Citation for published version (Harvard):

Howles, SA, Hannan, FM, Babinsky, VN, Rogers, A, Gorvin, CM, Rust, N, Richardson, T, McKenna, MJ, Nesbit, MA & Thakker, RV 2016, 'Cinacalcet for symptomatic hypercalcemia caused by AP2S1 mutations', *The New England Journal of Medicine*, vol. 374, no. 14, pp. 1396-1398. <https://doi.org/10.1056/NEJMc1511646>

[Link to publication on Research at Birmingham portal](#)

### Publisher Rights Statement:

Published in the *New England Journal of Medicine* on 07/04/2016

Copyright © 2016 Massachusetts Medical Society. All rights reserved.

### General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

## Cinacalcet for Symptomatic Hypercalcemia Caused by *AP2S1* Mutations

**TO THE EDITOR:** Familial hypocalciuric hypercalcemia is a genetically heterogeneous disorder with three variants: types 1, 2, and 3. Type 1 is caused by calcium-sensing receptor (*CASR*) mutations,<sup>1</sup> and type 2 is caused by guanine nucleotide-binding protein (G-protein) subunit  $\alpha_{11}$  (*GNA11*) mutations.<sup>2</sup> Type 3, which is the most severe variant clinically, is caused by adaptor-related protein complex 2 (AP2), sigma 1 subunit (*AP2S1*) heterozygous mutations. *AP2S1* mutations cause Arg15Cys, Arg15His, and Arg15Leu substitutions, and the mutant AP2-sigma proteins result in impaired calcium-sensing receptor-mediated signal transduction.<sup>3,4</sup>

There is currently no effective therapy for familial hypocalciuric hypercalcemia type 3. We therefore evaluated the usefulness of cinacalcet, a licensed calcium-sensing receptor allosteric activator,<sup>5</sup> in correcting signaling defects due to *AP2S1* mutations and in ameliorating symptomatic hypercalcemia.

We evaluated three previously unreported cases of familial hypocalciuric hypercalcemia type 3. Each of the three probands had an *AP2S1* mutation causing an Arg15Cys, Arg15His, or Arg15Leu substitution (Fig. 1A). The probands included a 34-year-old woman who had a 12-year history of hypercalcemia. She presented with fatigue, headaches, and persistent generalized aches that did not resolve after parathyroidectomy. In addition, a 22-year-old man presented with hypercalcemia, fatigue, and generalized rib pain, and a 52-year-old woman who had an approximately 20-year history of hypercalcemia presented with headaches, abdominal pain, vomiting, fatigue, and musculoskeletal pain that did not resolve after pamidronate infusion or parathyroidectomy (Fig. 1A and 1B).

The *in vitro* effects of cinacalcet on the signaling responses of cells expressing the calcium-sensing receptor and familial hypocalciuric hypercalcemia type 3-associated AP2-sigma mutants were assessed by measurement of intracellular calcium concentrations and of the activity of the mitogen-activated protein kinase (MAPK)-serum response element reporter in response to alterations in extracellular calcium (see the Methods

section in the Supplementary Appendix, available with the full text of this letter at NEJM.org). Cinacalcet (at a concentration of 10 nM) corrected the abnormal intracellular calcium signaling and MAPK responses to extracellular calcium in cells expressing the Cys15, His15, or Leu15 AP2-sigma mutants (Fig. S1 and S2 in the Supplementary Appendix).

Oral cinacalcet (at a dose of 30 to 60 mg daily) was administered to the three probands with familial hypocalciuric hypercalcemia type 3 over a 33-to-45-month period. This treatment led to more than 20% reductions in serum calcium concentrations (Fig. 1B) and abatement of symptoms. Cinacalcet also increased serum phosphate concentrations and reduced serum parathyroid hormone concentrations, although these values remained within the normal range (Fig. 1B).

Adverse effects such as nausea, vomiting, and hypocalcemia did not develop in any of the probands. However, long-term surveillance will be required to assess safety and monitor for hypocalcemia.

Our results show that cinacalcet-mediated allosteric modulation of the calcium-sensing receptor can correct the loss of function of *AP2S1* mutations. In addition, in the short term, cinacalcet can reduce the symptoms of familial hypocalciuric hypercalcemia type 3 associated with all three *AP2S1* mutations.


### Figure 1 (facing page). Cinacalcet for Hypercalcemia Associated with *AP2S1* Mutations.

Panel A shows the clinical and biochemical findings at presentation in three unrelated probands with familial hypocalciuric hypercalcemia who had heterozygous adaptor-related protein complex 2 (AP2), sigma 1 subunit (*AP2S1*) mutations resulting in Arg15Cys, Arg15His, or Arg15Leu substitutions in the AP2-sigma mutant proteins. Reference ranges are from Nesbit et al.<sup>2</sup> Panel B shows the effect of cinacalcet on serum concentrations of calcium, phosphate, and parathyroid hormone in the three probands. Vertical dashed lines indicate initiation of cinacalcet (CIN) therapy, pamidronate (PMD) infusion, or parathyroidectomy (PTX). Shaded gray areas indicate reference ranges.

A

Variable	Reference Range	Proband 1	Proband 2	Proband 3
Sex		Female	Male	Female
Age at presentation — yr		22	22	32
Family history of hypercalcemia		Yes	Yes	No
Associated clinical features		Fatigue, headaches, generalized musculo-skeletal pain	Fatigue, rib pain	Fatigue, headaches, vomiting, abdominal and musculo-skeletal pain
Serum measurements				
Calcium — mmol/liter				
Ionized	1.10–1.30	1.72	1.53	—
Albumin-adjusted	2.10–2.60	—	—	3.07
Phosphate — mmol/liter				
Total	0.70–1.40	1.00	0.80	0.95
Alkaline phosphatase				
Total — U/liter	30–130	55.0	—	81.0
Bone-specific — $\mu$ g/liter	3.7–20.9	—	20.6	—
Magnesium — mmol/liter				
Total	0.70–1.40	0.97	—	0.98
Parathyroid hormone				
Base units	10–65 ng/liter	28.5	65.1	—
SI units	1.3–7.6 pmol/liter	—	—	6.2
Urinary calcium-to-creatinine clearance ratio				
Ratio	>0.02	0.001	0.002	0.001
AP2S1 mutation				
Mutation		Arg15Cys	Arg15His	Arg15Leu

B



Sarah A. Howles, D.Phil.  
 Fadil M. Hannan, D.Phil.  
 Valerie N. Babinsky, D.Phil.  
 Angela Rogers, M.B., B.S.  
 Caroline M. Gorvin, D.Phil.  
 Nigel Rust, B.Sc.

University of Oxford  
 Oxford, United Kingdom

Tristan Richardson, M.B., B.S.

Royal Bournemouth Hospital  
 Bournemouth, United Kingdom

Malachi J. McKenna, M.D.

St. Vincent's University Hospital  
 Dublin, Ireland

M. Andrew Nesbit, Ph.D.

Rajesh V. Thakker, M.D.

University of Oxford  
 Oxford, United Kingdom  
 rajesh.thakker@ndm.ox.ac.uk

Drs. Howles and Hannan contributed equally to this letter.

A complete list of authors is available with the full text of this letter at [NEJM.org](http://NEJM.org).

Supported by program grants (G9825289 and G1000467, to Drs. Hannan, Gorvin, Nesbit, and Thakker) from the United Kingdom Medical Research Council, a grant (to Drs. Nesbit and Thakker) from the National Institute for Health Research (NIHR) Oxford Biomedical Research Centre Programme, a grant (FP7-264663, to Dr. Babinsky) from the European Commission Seventh Framework Program, Wellcome Trust clinical training fellowships (to Drs. Howles and Rogers), a Wellcome Trust Investigator Award (to Dr. Thakker), and an NIHR Senior Investigator Award (to Dr. Thakker).

Disclosure forms provided by the authors are available with the full text of this letter at [NEJM.org](http://NEJM.org).

1. Hannan FM, Nesbit MA, Zhang C, et al. Identification of 70 calcium-sensing receptor mutations in hyper- and hypo-calcaemic patients: evidence for clustering of extracellular domain mutations at calcium-binding sites. *Hum Mol Genet* 2012;21:2768-78.
2. Nesbit MA, Hannan FM, Howles SA, et al. Mutations affecting G-protein subunit  $\alpha 11$  in hypercalcemia and hypocalcemia. *N Engl J Med* 2013;368:2476-86.
3. Nesbit MA, Hannan FM, Howles SA, et al. Mutations in AP2S1 cause familial hypocalciuric hypercalcemia type 3. *Nat Genet* 2013;45:93-7.
4. Hannan FM, Howles SA, Rogers A, et al. Adaptor protein-2 sigma subunit mutations causing familial hypocalciuric hypercalcaemia type 3 (FHH3) demonstrate genotype-phenotype correlations, codon bias and dominant-negative effects. *Hum Mol Genet* 2015;24:5079-92.
5. Nemeth EF, Shoback D. Calcimimetic and calcilytic drugs for treating bone and mineral-related disorders. *Best Pract Res Clin Endocrinol Metab* 2013;27:373-84.

DOI: [10.1056/NEJMc1511646](https://doi.org/10.1056/NEJMc1511646)

Correspondence Copyright © 2015 Massachusetts Medical Society.