

The coexistence of two ^{bla}NDM-5 genes on an IncF plasmid as revealed by nanopore sequencing

Feng, Yu; Liu, Lu; McNally, Alan; Zong, Zhiyong

DOI:

[10.1128/AAC.00110-18](https://doi.org/10.1128/AAC.00110-18)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Feng, Y, Liu, L, McNally, A & Zong, Z 2018, 'The coexistence of two ^{bla}NDM-5 genes on an IncF plasmid as revealed by nanopore sequencing', *Antimicrobial Agents and Chemotherapy*.
<https://doi.org/10.1128/AAC.00110-18>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

1 **The coexistence of two *bla*_{NDM-5} genes on an IncF plasmid**
2 **as revealed by nanopore sequencing**

3 **Yu Feng^{1,2}, Lu Liu^{1,2}, Alan McNally⁵, Zhiyong Zong^{1,2,3,4*}**

4 *¹Center of Infectious Diseases, West China Hospital, Sichuan University, Chengdu,*
5 *China. ²Division of Infectious Diseases, State Key Laboratory of Biotherapy, Chengdu,*
6 *China. ³Department of Infection Control, West China Hospital, Sichuan University,*
7 *Chengdu, China. ⁴Center for Pathogen Research, West China Hospital, Sichuan*
8 *University, Chengdu, China. ⁵Institute of Microbiology and Infection, College of*
9 *Medical and Dental Sciences, University of Birmingham, Birmingham, UK.*

10
11 **Running title: two *bla*_{NDM-5} on a plasmid**

12
13 **Keywords:** carbapenem resistance; plasmids; Escherichia coli.

14
15 ***Corresponding author. Mailing address:** Center of Infectious Diseases, West China
16 Hospital (Huaxi), Guoxuexiang 37, Chengdu 610041, China. Phone:
17 86-28-8542-2637. Fax: 86-28-8542-3212. E-mail: zongzhiy@scu.edu.cn

18

19 **Abstract**

20 In a carbapenem-resistant *Escherichia coli* clinical isolate of sequence type 167, two
21 copies of *bla*_{NDM-5} were found on a 144,225-bp IncF self-transmissible plasmid of the
22 F36:A4:B- type. Both *bla*_{NDM-5} genes were located in 11,065-bp regions flanked by
23 two copies of IS26. The two regions were identical in sequence but were present at
24 different locations on the plasmid, suggesting a duplication of the same region. This
25 study highlights the complex genetic contexts of *bla*_{NDM-5}.

26 New Delhi metallo-β-lactamase (NDM) is a type of carbapenem-hydrolysing enzymes
27 (carbapenemases) with the ability to hydrolyze all β-lactams except monobactams (1),
28 representing a serious challenge for treatment of bacterial infections, infection control
29 and public health. ~~Up to now~~To date, there are 21 variants of NDM, ~~among which~~with
30 NDM-5 ~~is~~ one of the most common variants encountered in the Enterobacteriaceae
31 (2-5). The NDM-5-encoding gene, *bla*_{NDM-5}, usually exists in a single copy on
32 plasmids. However, we have found the peculiar presence of two copies of *bla*_{NDM-5} on
33 a single plasmid within an *Escherichia coli* clinical isolate, which is reported here.

34
35 *E. coli* strain SCEC020007 was recovered from urine of a female outpatient with
36 urinary tract infection in October 2016 in China. The strain was resistant to amikacin
37 (MIC, >512 µg/ml), ceftazidime (>512 µg/ml), ceftazidime-avibactam (>512/4 µg/ml),
38 ciprofloxacin (256 µg/ml), imipenem (64 µg/ml), meropenem (256 µg/ml),
39 piperacillin-tazobactam (>512/4 µg/ml) and trimethoprim-sulfamethoxazole
40 (128/2,432 µg/ml), but was susceptible to aztreonam (8 µg/ml), colistin (2 µg/ml) and
41 tigecycline (0.25 µg/ml) as determined using the broth dilution method of the Clinical
42 Laboratory Standards Institute (6). As there are no breakpoints of colistin and
43 tigecycline from CLSI, those defined by EUCAST (<http://www.eucast.org/>) were
44 applied.

45
46 A draft genome sequence of the strain was generated on the Illumina HiSeq X10
47 platform, which generated 5,557,833 clean reads and 1.67 Gb clean bases. A total of
48 113 contigs (102 >1,000 bp; *N50* 126,680 bp) with a 50.76% GC content were *de*
49 *novo* assembled using SPAdes (7). Strain SCEC020007 belonged to phylogenetic
50 group A as determined using PCR as described previously (8) and sequence type

51 167 (ST167) as determined using the genomic sequence to query the *E. coli*
52 multi-locus sequence typing database
53 (<http://enterobase.warwick.ac.uk/species/index/ecoli>). Antimicrobial resistance genes
54 were identified from genome sequences using the ABRicate program
55 (<https://github.com/tseemann/abricate>) to query the ResFinder database
56 (<http://genomicepidemiology.org/>). Strain SCEC020007 had 9 antimicrobial
57 resistance genes mediating resistance to aminoglycosides (*aadA2*, *aadA5*, *rmtB*),
58 β -lactams (*bla*_{NDM-5} and *bla*_{TEM-1}), tetracycline (*tet(A)*), sulphonamides (*sul1*) and
59 trimethoprim (*dfrA12* and *dfrA17*). Plasmid replicon types within strain SCEC020007
60 were determined using ~~by~~ the PlasmidFinder tool at <http://genomicepidemiology.org/>.
61 Surprisingly, strain SCEC020007 had an IncFIA, an IncFII and an IncB/O/K/Z replicon
62 but no IncX3, which is the common replicon type of plasmids associated with *bla*_{NDM-5}.
63
64 To untangle the genetic context of *bla*_{NDM-5}, strain SCEC020007 was subjected to
65 sequencing using the long-read ~~real-time~~ MinION Sequencer (Nanopore, Oxford, UK).
66 ~~The A~~ de novo hybrid assembly of both short Illumina reads and long MinION reads
67 was ~~performed-constructed~~ using Unicycler v0.4.3 (9) under conservative mode for
68 ~~an~~ increased accuracy. ~~The C~~ complete circular contigs generated were then
69 corrected using Plion v1.22 (10) with Illumina reads for several rounds until no
70 change was detected. The hybrid assembly of Illumina and MinION reads revealed
71 that strain SCEC020007 had a 4.8-Mb circular chromosome, a 144,225-bp plasmid
72 containing ~~an~~ IncFIA and ~~a~~ FII replicons (designated pNDM5_020007) and an
73 84,952-bp plasmid with an IncB/O/K/Z replicon (designated pBOKZ_020007).
74 Surprisingly, there were two copies of *bla*_{NDM-5} in strain SCEC020007, both of which
75 were present on pNDM5_020007. Both *bla*_{NDM-5} genes were located in 11,065-bp

76 regions flanked by two copies of IS26 and the two regions were identical in sequence
77 but were present at different locations on pNDM5_020007 (Figure 1), suggesting that
78 the 11,065-bp region is duplicated. The presence of the two *bla*_{NDM-5} genes and their
79 locations on pNDM5_020007 were confirmed by PCR. The 11,065-bp region
80 contained a complex class 1 integron with a *dfrA17-aadA5* cassette array and *ISCR1*
81 (insertion sequence common region 1), which is truncated by IS26 at its 5' conserved
82 segment, a 69-bp remnant of *ctuA1* (encoding an ion tolerant protein), *dsbC*
83 (encoding an oxidoreductase), *trpF* (encoding a phosphoribosylanthranilate
84 isomerase), *ble* (mediating bleomycin resistance), *bla*_{NDM-5}, a truncated *ISAbA125*
85 and a truncated *ISEcp1/ISEc9* element (Figure 1). The co-existence of two *bla*_{NDM-5}
86 genes has not been reported before but the co-existence of two *bla*_{NDM-1} genes has
87 been described previously (11, 12). Two tandem copies of *bla*_{NDM-1} genes have been
88 found in the chromosome of an ST167 *E. coli* in China (11) and a *Pseudomonas*
89 *aeruginosa* strain in Serbia (12). In both cases, the tandem copies of *bla*_{NDM-1} are
90 associated with *ISCR1* but not IS26. It is known that *ISCR1* uses the rolling circle
91 mechanism for transposition and may generate tandem duplication of its mobilized
92 sequence via homologous recombination (13). However, the duplication of the
93 11,065-bp region carrying *bla*_{NDM-5} on pNDM5_020007 is not tandem, suggesting that
94 the duplication might not result from the action of *ISCR1* but could be mediated by
95 IS26. The exact mechanism for the duplication of such a large region warrants further
96 studies.

97
98 Assembly based on Illumina reads alone generated only a single contig containing
99 *bla*_{NDM-5} and was unable to reveal that there were actually ~~were~~ two identical copies of
100 the same contig. This imposes difficulties for completing the *bla*_{NDM-5}-carrying plasmid

101 | sequence by convention^{al} methods including PCR and Sanger sequencing to close
102 | gaps between contigs. By contrast, MinION sequencing ~~is~~was able to resolve the
103 | copy numbers of genes and contigs and their exact position on the plasmid relative to
104 | each other.

105

106 | Plasmid multi-locus sequence typing (pMLST) was performed using the pMLST tool
107 | (<https://cge.cbs.dtu.dk/services/pMLST/>). pNDM5_020007 belongs to the F36:A4:B-
108 | type. pNDM5_020007 ~~was~~has closest similarity (97% coverage and 99% identity) to
109 | a 149.5-kb unnamed plasmid (GenBank accession no. CP023871) from *E. coli* strain
110 | FDAARGOS_434, which was recovered from a human rectal swab in British
111 | Columbia, Canada, in 2014. This unnamed plasmid also carries *bla*_{NDM-5} (a single
112 | copy) and belongs to the F36:A4:B- type. Backbones of pNDM5_020007 and the
113 | unnamed plasmid of strain FDAARGOS_434 are almost identical, suggesting that
114 | they might have originated from a common plasmid. Conjugation experiments were
115 | carried out in broth and on filters with the azide-resistant *E. coli* strain J53 as the
116 | recipient. pNDM5_020007 was able to be transferred by conjugation, suggesting that
117 | it is self-transmissible.

118

119 | In conclusion, we identified the presence of two *bla*_{NDM-5} genes on an F36:A4:B-
120 | self-transmissible plasmid. The co-existence of two *bla*_{NDM-5} genes was due to the
121 | duplication of an IS26-bracketed region containing *ISCR1*.

122

123 | **Nucleotide sequence accession numbers.** The complete sequence of
124 | pBOKZ_020007, pNDM5_020007 and the chromosome of strain SCEC020007 has

125 been deposited into GenBank under the accession no. CP025625, CP025626 and
126 CP025627, respectively.

127

128 **Acknowledgments**

129 This work was supported by grants from the National Natural Science Foundation of
130 China (project no. 81222025 and 81572030) and a joint grant from the National
131 Natural Science Foundation of China (project no. 81661130159) and the Newton
132 Advanced Fellowship, Royal Society, UK (NA150363).

133

134 **References**

- 135 1. **Kumarasamy KK, Toleman MA, Walsh TR, Bagaria J, Butt F, Balakrishnan R,**
136 **Chaudhary U, Doumith M, Giske CG, Irfan S, Krishnan P, Kumar AV, Maharjan S,**
137 **Mushtaq S, Noorie T, Paterson DL, Pearson A, Perry C, Pike R, Rao B, Ray U, Sarma**
138 **JB, Sharma M, Sheridan E, Thirunarayan MA, Turton J, Upadhyay S, Warner M,**
139 **Welfare W, Livermore DM, Woodford N.** 2010. Emergence of a new antibiotic
140 resistance mechanism in India, Pakistan, and the UK: a molecular, biological, and
141 epidemiological study. *Lancet Infect Dis* **10**:597-602.
- 142 2. **Hornsey M, Phee L, Wareham DW.** 2011. A novel variant, NDM-5, of the New Delhi
143 metallo- β -lactamase in a multidrug-resistant *Escherichia coli* ST648 isolate recovered
144 from a patient in the United Kingdom. *Antimicrob Agents Chemother* **55**:5952-5954.
- 145 3. **Iovleva A, Doi Y.** 2017. Carbapenem-Resistant Enterobacteriaceae. *Clin Lab Med*
146 **37**:303-315.
- 147 4. **Logan LK, Weinstein RA.** 2017. The Epidemiology of Carbapenem-Resistant
148 Enterobacteriaceae: The Impact and Evolution of a Global Menace. *J Infect Dis*
149 **215**:S28-S36.
- 150 5. **Zhang R, Liu L, Zhou H, Chan EW, Li J, Fang Y, Li Y, Liao K, Chen S.** 2017. Nationwide
151 Surveillance of Clinical Carbapenem-resistant Enterobacteriaceae (CRE) Strains in
152 China. *EBioMedicine* **19**:98-106.
- 153 6. **CLSI.** 2017. Performance Standards for Antimicrobial Susceptibility Testing;
154 Twenty-Third Informational Supplement. M100-S27. Clinical and Laboratory
155 Standards Institute, Wayne, PA, USA.
- 156 7. **Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM,**
157 **Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G,**
158 **Alekseyev MA, Pevzner PA.** 2012. SPAdes: a new genome assembly algorithm and
159 its applications to single-cell sequencing. *J Comput Biol* **19**:455-477.
- 160 8. **Clermont O, Bonacorsi S, Bingen E.** 2000. Rapid and simple determination of the
161 *Escherichia coli* phylogenetic group. *Appl Environ Microbiol* **66**:4555-4558.

- 162 9. **Wick RR, Judd LM, Gorrie CL, Holt KE.** 2017. Unicycler: Resolving bacterial genome
163 assemblies from short and long sequencing reads. PLoS Comput Biol **13**:e1005595.
- 164 10. **Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA, Zeng**
165 **Q, Wortman J, Young SK, Earl AM.** 2014. Pilon: an integrated tool for comprehensive
166 microbial variant detection and genome assembly improvement. PLoS One
167 **9**:e112963.
- 168 11. **Shen P, Yi M, Fu Y, Ruan Z, Du X, Yu Y, Xie X.** 2017. Detection of an *Escherichia coli*
169 Sequence Type 167 Strain with Two Tandem Copies of *bla*_{NDM-1} in the Chromosome. J
170 Clin Microbiol **55**:199-205.
- 171 12. **Jovicic B, Lepsanovic Z, Begovic J, Rakonjac B, Perovanovic J, Topisirovic L, Kojic M.**
172 2013. The clinical isolate *Pseudomonas aeruginosa* MMA83 carries two copies of the
173 *bla*_{NDM-1} gene in a novel genetic context. Antimicrob Agents Chemother
174 **57**:3405-3407.
- 175 13. **Toleman MA, Bennett PM, Walsh TR.** 2006. ISCR elements: novel gene-capturing
176 systems of the 21st century? Microbiol Mol Biol Rev **70**:296-316.
177

178 **Figure legend**

179 **Figure 1.** pNDM5_020007 and the genetic context of *bla*_{NDM-5}. The two
180 11,065-bp *bla*_{NDM-5}-containing regions bracketed by IS26 are indicated by
181 orange circles in the map of pNDM5_020007 and are shown in detail at the
182 bottom. Δ represents truncated genes or mobile genetic elements.