

Prevention of rickets and osteomalacia in the UK

Uday, Suma; Högler, Wolfgang

DOI:

[10.1136/archdischild-2018-314826](https://doi.org/10.1136/archdischild-2018-314826)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Uday, S & Högler, W 2018, 'Prevention of rickets and osteomalacia in the UK: political action overdue', *Archives of Disease in Childhood*, vol. 103, no. 9, pp. 901-906. <https://doi.org/10.1136/archdischild-2018-314826>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Final version of record available at: <http://dx.doi.org/10.1136/archdischild-2018-314826>

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Archives of
Disease in Childhood

Prevention of rickets and osteomalacia in the United Kingdom: Political action overdue

Journal:	<i>Archives of Disease in Childhood</i>
Manuscript ID	archdischild-2018-314826.R1
Article Type:	Review
Edition:	not in use
Date Submitted by the Author:	n/a
Complete List of Authors:	Uday, Suma ; Birmingham Women's and Children's NHS Foundation Trust, Endocrinology and Diabetes; University of Birmingham , Institute of Metabolism and Systems Research Hoegler, Wolfgang; Birmingham Women's and Children's NHS Foundation Trust; University of Birmingham , Institute of Metabolism and Systems Research
Keywords:	immigration, dark skin, south Asian, African, vitamin D deficiency

SCHOLARONE™
 Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 **Title:**

2 Prevention of rickets and osteomalacia in the United Kingdom: Political action overdue

3

4 **Authors:**

5 Suma Uday^{1,2}, Wolfgang Högler^{1,2}

6 1. Department of Endocrinology & Diabetes, Birmingham Women's and Children's Hospital,
7 Birmingham, UK

8 2. Institute of Metabolism and Systems Research, University of Birmingham, Birmingham, UK

9

10 **Corresponding author:**

11 PD Dr Wolfgang Högler,

12 Department of Endocrinology and Diabetes, Birmingham Women's and Children's Hospital,

13 Steelhouse Lane, B4 6NH, Birmingham, UK

14 Tel: +44 121 333 8197

15 Email: wolfgang.hogler@nhs.net

16

17 **Keywords:** Immigration, dark skin, South Asian, African, vitamin D deficiency, calcium, food
18 fortification

19

20 **Word count: 3427**

21 **Figures total: 3**

Colour: 1

22 **Tables: 3**

23

24

25

26

27

28

29

30

31

32

33

1
2
3 34 **Abstract**
4

5 35 The consequences of vitamin D and dietary calcium deficiency have become a huge public health
6 36 concern in the UK. The burden of disease from these deficiencies includes not just rickets, but also
7 37 hypocalcaemic seizures, dilated cardiomyopathy and mostly occult myopathy and osteomalacia. The
8 38 increasing burden of disease is intrinsically linked to ethnicity and the population demographic
9 39 changes in the UK. Three facts have led to the resurfacing of the English disease: 1) The UK has no
10 40 ultraviolet sunlight for at least 6 months of the year 2) Dark skin produces far less vitamin D than
11 41 white skin per unit ultraviolet light exposure, and 3) non-EU immigration over the last century. To date,
12 42 the UK government demonstrates incomplete understanding of these three facts, and its failure to
13 43 adjust its prevention programs to changing demographics is endangering the health and life of UK
14 44 residents **with dark skin**, of whom infants are the most vulnerable. **Establishing accountability through**
15 45 the implementation of monitored antenatal and infantile supplementation programs and mandatory
16 46 food fortification is overdue.

17
18
19
20
21
22
23
24
25
26
27 47
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

48 Introduction

49 Quite in contrast to their continental European counterparts, paediatric A&E doctors in the UK often
50 encounter dark skinned infants with hypocalcaemic seizures, hypocalcaemic cardiomyopathy and
51 rickets during winter and spring. This article will explain the critical role of the UK government in the
52 causation and prevention of what was once called, and still is, the 'English disease'.

53 Rickets and osteomalacia occur due to defective mineralisation of hypertrophic growth plate
54 chondrocytes and pre-formed osteoid in existing bone, respectively. In children, rickets and
55 osteomalacia always co-exist. The most common cause of rickets and osteomalacia worldwide is
56 calcium deprivation, secondary to low dietary calcium ('nutritional') and/or low ultraviolet B (UVB)
57 sunlight exposure [1], the main source of cutaneous vitamin D production ('solar') [2]. There is very
58 little vitamin D in diet (including breast milk) and thus humans are mainly dependent on solar UVB for
59 vitamin D supply. Cholecalciferol (vitamin D3) and ergocalciferol (vitamin D2) undergo hepatic 25-
60 hydroxylation to form calcidiol or 25 hydroxyvitamin D (25OHD, stable form reflecting vitamin D status)
61 and then renal 1-hydroxylation to form the hormone calcitriol or 1,25 dihydroxyvitamin D. In contrast to
62 calcitriol, vitamin D and 25OHD are biologically inert. The main action of calcitriol is to aid intestinal
63 calcium absorption. When dietary calcium or solar vitamin D availability is low, secondary
64 hyperparathyroidism sets in which increases intestinal calcium absorption and bone resorption in the
65 short term to maintain normal serum calcium concentrations. Unopposed, this compensation will fail in
66 the long term and hyperparathyroidism-induced renal phosphate wasting will lead to
67 hypophosphataemia, which ultimately impairs mineralisation on the tissue level causing rickets and
68 osteomalacia [3]. Hypocalcaemic complications of calcium deprivation can occur before these
69 hypophosphataemic complications, especially during phases of rapid growth i.e during infancy and
70 adolescence [4][5].

71 Dietary calcium deficiency rickets (due to nutritional deficits) prevails in low income countries whereas
72 vitamin D deficiency rickets (due to reduced solar UVB exposure/latitude) predominates in high
73 income countries [3]. Calcium deprivation and all its hypocalcaemic or hypophosphataemic
74 complications can be easily prevented by ensuring adequate calcium and vitamin D supply [6].
75 Factors causing vitamin D deficiency in high income countries such as the UK specifically include:

1
2
3 76 1) **Latitude**: lack of UVB radiation for at least 6 months of the year due to its geographic location [2][7]
4
5 77 2) **Ethnicity**: dark skin [Fitzpatrick skin type IV (light brown), V (dark brown) and VI (black)] massively
6
7 78 reduces cutaneous synthesis of vitamin D [7]
8
9 79 3) **Culture**: covered clothing, low dietary vitamin D and calcium intake [3][8]
10
11 80 4) **Sunscreen**: Excessive use of sunscreen [9] due to fear of skin cancer
12
13 81 It is not difficult to see that the majority of the above factors are non-modifiable. Therefore,
14
15 82 supplementation or food fortification with vitamin D is necessary to prevent rickets and osteomalacia
16
17 83 in an ethnically and culturally increasingly diverse society. Non-EU immigration over the last century
18
19 84 has led to a change in population demographics with a growing proportion of the resident population
20
21 85 (14% in the 2011 census) being dark skinned (**Figure 1**) [10]. Whole body clothing for cultural,
22
23 86 medical or other reasons further reduces sunlight exposure. These high-risk groups are unable to
24
25 87 produce sufficient vitamin D in the UK. Unfortunately, the UK has not adapted to its ethnic and cultural
26
27 88 diversity which has led to a resurgence in rickets [11] and hypocalcaemic complications [5], a problem
28
29 89 further augmented by a lack of mandatory food fortification [12] and a complex and poorly
30
31 90 implemented infant vitamin D supplementation policy [13].

91 **Burden of disease from vitamin D deficiency and rickets in the UK**

32
33
34 92 'Nutritional' rickets was indeed originally caused by poor nutrition but was always also of 'solar' origin
35
36 93 due to pollution during the industrial revolution. The fact that the English disease has resurfaced as a
37
38 94 major public health problem in the UK in the last 50 years however is much less 'nutritional', but due
39
40 95 to the increase in the ethnic high-risk proportion of the population. Over the past few decades several
41
42 96 UK studies have highlighted the increasing burden of rickets [5][11][14][15][16] and the increased
43
44 97 susceptibility of the Black, Asian and Minority Ethnic (BAME) groups. The burden of disease from
45
46 98 solar vitamin D and dietary calcium deficiency is much wider and goes beyond the bone pathology.
47
48 99 The British Paediatric Surveillance Unit survey (September 2011-2013) reported a total of 91 cases
49
50 100 (85%, n=77 were infants) of hypocalcaemic seizures due to vitamin D deficiency, amounting to an
51
52 101 annual incidence of 3.49 per million children (0-15 years), with highest incidence in the South Asian
53
54 102 population at 26.04 per million [5]. In contrast to the white population, its high incidence in the BAME
55
56 103 population precludes rickets from being classified as a rare disease (**Table 1**) [17]. A retrospective
57
58 104 study from southeast England identified sixteen BAME cases (6 Asian, 10 Black) of hypocalcaemic

1
2
3 105 dilated cardiomyopathy secondary to severe vitamin D deficiency between 2000 and 2006 [18]. These
4 106 cases displayed significant mortality (n=3) and morbidity requiring mechanical ventilatory and
5 107 circulatory support (n=2 each) and referral for cardiac transplantation (n=2) [18]. In a post-mortem
6 108 study from London (n=52), hypocalcaemia due to vitamin D deficiency was considered accountable
7 109 for death in 3 BAME children; they all showed radiological and histological rickets: 2 babies had
8 110 cardiomyopathy and a 3-year-old had hypocalcaemic seizures [19]. These reports are not collections
9 111 of tragic medical rarities. Infants presenting with complications of hypocalcaemia such as seizures
10 112 [4][5][11], dilated cardiomyopathy [18], cardiac death [19], and overt rickets [14][15] only represent the
11 113 tip of the iceberg of a public health crisis. The extent of hidden pathology remains unknown.
12 114 Histological changes of rickets and osteomalacia occur long before the disease becomes evident on
13 115 radiographs [3]. One way to gain insight into the widespread prevalence of rickets and osteomalacia
14 116 in the UK is from post mortem studies [19] which demonstrate histological changes in nearly 70% of
15 117 infants studied, **which however were not quantified**. Osteomalacia in adults on post mortem
16 118 examination is also highly prevalent (nearly 25% of a low-risk population studied, n=675) in northern
17 119 Europe [20].

18
19
20
21
22
23
24
25
26
27
28
29
30
31 120 The increased prevalence of rickets in the immigrant and resident BAME population in the UK,
32 121 especially the South Asian community, is not new knowledge and was noted as far back as the 1960s
33 122 [21], when the UK government commonly referred to it as the 'Asian rickets' [22]. Although
34 123 immigration of dark-skinned individuals has contributed to the rise in rickets in most developed
35 124 countries [17][23], an additional contributing factor in the UK is its geographic location [24] and
36 125 possibly the changing climate [25]. Hence, vitamin D deficiency is neither limited to the winter months
37 126 nor exclusive to BAME population. A study from Manchester reported median serum 25OHD
38 127 concentrations in summer of 28.6 nmol/L [interquartile range (IQR) 21.3-41.6], falling further to 18.4
39 128 nmol/L (IQR 12.7-25.7) in the winter in adult residents of South Asian origin [8]. These findings were
40 129 resonated in another study in women of child bearing age where severe deficiency (25OHD <25
41 130 nmol/L) was prevalent in 81% and 79.2% of UK South Asians in winter and autumn, respectively [26].
42 131 In a study of white women from North-West England (n=333) 27% and 7% had insufficient (<50
43 132 nmol/L) and deficient (<25 nmol/L) serum 25OHD levels during pregnancy, and 48% and 11% four
44 133 months post-delivery, respectively [27]. Median 25OHD in cord blood samples were only 50% of
45 134 maternal 25OHD levels and the prevalence of deficiency and insufficiency in infants at 4 months

1
2
3 135 (n=322) was 13% and 24%, respectively [27]. No UK study compared vitamin D levels in native and
4
5 136 migrant mother-baby pairs, but an Italian study showed that severe vitamin D deficiency (25OHD < 25
6
7 137 nmol/L) was widely prevalent in the migrant mothers and their newborns (48.4% and 76.2%,
8
9 138 respectively) when compared to natives (38% and 18%, respectively) [28].

10
11 139 The UK National Diet and Nutrition Survey found that the mean intake of vitamin D was below the
12
13 140 reference nutrient intake: only 29% for children aged 1.5 to 3 years and 33% for adults \geq 65 years and
14
15 141 nearly a fifth of all subjects studied were vitamin D deficient (25OHD < 25 nmol/L) [29].

16
17 142 Although vitamin D deficiency is pandemic in Europe it is more prevalent in some mid-latitude
18
19 143 countries including the UK when compared to northerly latitude countries such as Norway, Iceland or
20
21 144 Finland even after accounting for ethnicity [30]. The lower prevalence in the northern latitude is
22
23 145 attributed to better supplementation policies and also food fortification [30].

24 25 146 **Factors contributing to resurgence of rickets in the UK**

26 27 28 147 ***The increasing BAME resident and immigrant population***

29
30
31 148 A question on ethnic group was first introduced in the 1991 Census, not only “to enable monitoring for
32
33 149 equal opportunities / anti-discrimination policies” but also “to plan for the future through resource
34
35 150 allocation and provision of services” [10]. However, this does not seem to have happened in relation
36
37 151 to rickets prevention despite the so obviously increasing proportion of dark skinned individuals. The
38
39 152 most recent census (2011) reported that 14% of the population in England and Wales are of BAME
40
41 153 origin which is a significant rise from previous census in 2001 (8.7%) and 1991 (5.9%) (**Figure 1**) [10].
42
43 154 Every year, nearly 13% of babies born in England and Wales have been of BAME background since
44
45 155 2014 [10] and around 100-150,000 people of BAME background (net) are added to the UK population
46
47 156 through immigration (**Figure 2**) [10]. Therefore, the government needs to adjust its health care
48
49 157 systems to these population demographic changes.

50
51 158 A multitude of factors contribute to the increased prevalence of vitamin D deficiency in BAME groups
52
53 159 residing in high latitude countries. An adult study of South Asian residents in Manchester reported
54
55 160 lower oral intake of dietary vitamin D (mean of 1.32 μ g compared to 3.26 μ g in white subjects), sun
56
57 161 avoidance when outside and lower amounts of skin surface exposure [8]. However by far the greatest

1
2
3 162 contributing factor is reduced skin synthesis of vitamin D [7]. Dark skin, when compared to white skin,
4 163 requires much greater exposure to sunlight to improve circulating 25OHD levels as melanin absorbs a
5 164 substantial proportion of the UVB light [7]. Hence, even the UK recommended summer sunlight
6 165 exposure amounts fail to produce sufficient circulating 25OHD in the dark skinned population [7]. Lack
7 166 of ethnicity-specific guidance on sunlight exposure [7], minimal impact of dietary vitamin D in
8 167 improving serum 25OHD levels [26] and poor supplementation policies in infants [13] and pregnant
9 168 women all contribute to widespread vitamin D deficiency in dark skinned individuals, all year round
10 169 [26].
11
12
13
14
15
16
17

18 ***UK politics and policy: From success to failure***

19
20
21 171 There is an interesting history behind the increasing burden of rickets in the UK. During World War II,
22 172 Britain triumphed eradication of rickets through rationing, food fortification and supplementation of
23 173 infants and pregnant women with cod-liver oil. Strangely, the same country which once demonstrated
24 174 the effectiveness of good public health policies is now struggling to tackle this easily preventable
25 175 disease. In the 1950s, uncontrolled fortification of infant formula and food (4000 IU/day, which is 10
26 176 times the current recommended dose of 400 IU/day) was implicated in the causation of infantile
27 177 hypercalcaemia [31] which led the government to withdraw fortification. Ever since this time, the
28 178 'English disease' has returned to the UK to the extent that today, Britain serves as an example of how
29 179 lack of effective policy implementation can lead to a rise in a preventable disease such as rickets.
30 180 Rickets or 'Asian rickets' has enormous political significance and has been the subject of debate in
31 181 parliament since the 1960s [22]. There is a suggestion that the resurgence of rickets symbolises
32 182 socioeconomic inequality and racial or ethnic health disparity in the UK, which is discussed in detail
33 183 elsewhere [22]. The government continues to shy away from the issue. The senior author submitted a
34 184 parliamentary question to the House of Lords in November 2017 with evidence demonstrating the
35 185 ineffectiveness of the UK's infant vitamin D supplementation program [13] and queried the
36 186 government's future plans to prevent rickets in the UK. The response, detailed in **Table 2**, reflects the
37 187 government's current misconceptions.
38
39
40
41
42
43
44
45
46
47
48
49
50

51 ***The UK prevention program: complex, outdated, unmonitored, ineffective***

1
2
3 189 The Scientific Advisory Committee on Nutrition (SACN) sets vitamin D 'reference nutrient intake' for
4
5 190 adults and children aged >4 years (10 µg or 400 IU/day) and 'safe intake' for infants (8.5-10 µg or
6
7 191 340-400 IU/day) and children aged 1-4 years (10 µg or 400 IU/day) [35]. The SACN does not
8
9 192 recommend supplements or food fortification but acknowledges the difficulty of achieving the
10
11 193 recommended intakes from natural food sources and advised the government to consider strategies
12
13 194 for the UK population to achieve the above intake [35]. Whilst most authorities consider optimal levels
14
15 195 of serum 25OHD to be above 50 nmol/L [6], the SACN recommends that levels should not fall below
16
17 196 25 nmol/L [35]. The National Institute for Health and Care Excellence (NICE) makes certain
18
19 197 recommendations to multiple agencies (including the Department of Health, Public Health England
20
21 198 and healthcare professionals) to increase supplement use in at risk populations [34]. Based on the
22
23 199 latest SACN vitamin D and health report [35], Public Health England recommend that populations at
24
25 200 risk should take supplements containing 10 µg (400 IU) of vitamin D all year round
26
27 201 (<https://www.gov.uk/government/news/phe-publishes-new-advice-on-vitamin-d>). The recommendation
28
29 202 for infants and children however is not only overly complex (**Table 3**) but also now out-dated in
30
31 203 relation to formula feeding and dose for children >1 year. The evidence-based global consensus on
32
33 204 the prevention of nutritional rickets recommends universal supplementation of all infants [6], which is
34
35 205 followed by most European countries (85%, n=24/29) [13]. SACN does not recommend vitamin D
36
37 206 supplements for formula-fed infants whilst new evidence demonstrates that formula feeding does not
38
39 207 protect from rickets [5][6]. In the British Paediatric Surveillance Unit survey, 19% (n=15/77) of infants
40
41 208 presenting with hypocalcaemic seizures were exclusively formula-fed and 9% (n=7) received mixed
42
43 209 breast and formula feed [5].

44
45 210 In contrast to the global consensus recommendation (10 µg or 400 IU for infants and 15 µg or 600 IU
46
47 211 for children > 1 year), SACN and Public Health England recommend 8.5 to 10 µg of vitamin D for
48
49 212 infants and children and the government supplies **Healthy Start** vitamins which only provide 7.5 µg.
50
51 213 The operation of the **Healthy Start** scheme which provides free vitamins to low income families is also
52
53 214 very complex [33], leading to low uptake, poor accessibility and lack of motivation to take daily
54
55 215 supplements [36], not to mention the lack of parental awareness [32]. Apart from the complexity of the
56
57 216 guidance, it is mainly the complete lack of accountability and monitoring, that is responsible for the
58
59 217 failing of the **Healthy Start** program to improve the vitamin D status of the at-risk population [33]. The
60

218 UK's unmonitored infant vitamin D supplementation program has the lowest adherence rate in
219 Europe, with just 5-20% of infants actually receiving vitamin D drops (**Figure 3**) [13].

220 **Preventative strategies**

221 Given the low uptake of supplements the long term solution to vitamin D deficiency is mandatory food
222 fortification [12] which has been successfully implemented in countries such as Canada. However,
223 this may not be easily adopted by the UK government given that it also shies away from adopting folic
224 acid fortification despite overwhelming evidence on its role in preventing neural tube defects [37].
225 Mandatory fortification of flour with folic acid is adopted by 78 countries but not the UK, which has
226 resulted in a continued rise in neural tube defects in the UK [37]. Countries with *mandatory* folic acid
227 fortification have seen a decline in neural tube defects compared to those with voluntary or no
228 fortification [38]. Despite SACN recommendation to adopt mandatory folic acid fortification of flour (in
229 2006 and 2009), the UK government has not taken any positive steps. Not surprisingly in 2017, Food
230 Standards Scotland requested SACN to provide advice on whether its previous recommendations on
231 mandatory fortification still apply, with an intention to proceed unilaterally [39]. Similarly, iodine
232 deficiency which is considered the single most important preventable cause of brain damage by the
233 World Health Organisation, is thought to be re-emerging as a public health concern in the UK due to
234 lack of national salt-iodisation and monitoring of iodine levels in pregnancy [40].

235 Whilst mandatory food fortification may be the best long term solution to improve the vitamin D status
236 at a population level [12], certain measures in the interim are crucial in protecting the most vulnerable
237 group - infants. As most infants acquire vitamin D deficiency from the mother it is equally important to
238 ensure adequate supplementation during pregnancy and to start infant supplementation at birth.
239 Success of supplementation policies requires that authorities adopt multiple, simple strategies. We
240 have previously compared vitamin D supplementation policies across Europe and determined that the
241 following policy implementation features were significantly associated with improved adherence in the
242 first year of life: universal supplementation independent of feeding mode ($p = 0.007$), providing
243 information on supplementation at discharge from neonatal units ($p = 0.02$), providing financial family
244 support ($p = 0.005$) and monitoring adherence at child health surveillance visits ($p = 0.001$) [13]. The
245 UK currently does not adopt any of the above features, except providing financial support only to low
246 income families.

1
2
3 247 Public health policies need to be simple to work. Universal supplementation of all infants [6] enhances
4 248 adherence [13], moreover formula feeds do not protect the infant from rickets [5][6] and hence the
5
6 249 SACN guidance requires updating and simplifying. Most importantly, similar to the vaccination
7
8 250 program, a group of healthcare professionals needs to be made responsible for delivering this
9
10 251 prevention program, in the same way as vaccination program. In the authors' point of view,
11
12 252 establishing accountability is the most essential change required in the UK.

13
14 253 Based on data from the European survey [13], the suggested responsibilities are as follows:

- 15
16
17 254 1) Health care professionals providing antenatal care include monitoring of supplementation at every
18
19 255 routine visit.
- 20 256 2) Neonatal units provide specific information on prevention programs to parents, commence vitamin
21
22 257 D supplementation at birth and provide first bottle at discharge.
- 23
24 258 3) Upon registration at the GP, adherence to vitamin D supplementation is checked and then
25
26 259 monitored at every single vaccination time point and red book visit.
- 27
28 260 4) Inclusion of a monitoring question on adherence to vitamin D supplements in maternity notes and
29
30 261 personal child health record (red book), at all routine time points, which reminds healthcare
31
32 262 professionals and parents.
- 33 263 5) The government should provide financial remuneration to GPs delivering this prevention program
34
35 264 in a similar way as for the immunization program (prevention of infections). The government
36
37 265 should consider linking financial family support to adherence to all prevention programs.

38 39 266 40 41 267 **Future considerations**

42
43
44 268 The majority of the European medical literature on rickets and hypocalcaemic complications comes
45
46 269 from the UK [1]. Current studies are mainly based on clinical presentation or serum 25OHD
47
48 270 measurements. Further research is required to explore the true burden of rickets and specifically
49
50 271 osteomalacia in the UK. Histological changes occur long before the disease becomes clinically,
51
52 272 biochemically or radiologically evident [3]. Therefore, the true prevalence of occult rickets and
53
54 273 osteomalacia can only be identified by **quantitative** growth plate histology and bone

1
2
3 274 histomorphometry. These investigations are invasive in alive children or adults but can be undertaken
4 275 during post-mortem examination.

6
7 276 Taking daily vitamin D supplements prevents rickets but may pose an issue with adherence, similar to
8
9 277 folic acid supplementation, especially if un-monitored. However, vitamin D can also be easily and
10
11 278 safely administered in bolus doses alleviating the need for daily supplementation and parental
12
13 279 compliance. Licensed preparations for bolus administration are readily available in the UK. Feasibility
14
15 280 of administering bolus doses of vitamin D during immunisations and routine child care visits in the UK
16
17 281 should be explored. Likewise, safety and feasibility of administering bolus doses of vitamin D during
18
19 282 antenatal visits should also be explored.

20
21 283 The increasing UK BAME population requires life-long supplementation. Since this can never be
22
23 284 delivered, food fortification is the obvious solution to the problem. Further studies are required to
24
25 285 bridge the gap in knowledge required to facilitate food fortification, and to compare the cost-
26
27 286 effectiveness of fortification and supplementation programs.

28
29 287

30 31 32 288 **Conclusions**

33
34
35 289 The UK population is more ethnically and culturally diverse than ever. The increasing UK BAME
36
37 290 population is fully exposed to the complications of vitamin D deficiency due to their dark skin and
38
39 291 cultural traditions. The UK government demonstrates serious misconception of the actual public
40
41 292 health issue and the implementation of effective prevention programs, and an astounding ignorance
42
43 293 of changing population demographics. National statistics demonstrate a constant rise of the UK
44
45 294 BAME resident population through continued non-EU immigration from countries with predominantly
46
47 295 dark-skinned population (Asia, Africa), with high birth rates. We call for the government to consider
48
49 296 adaptation of their public health policy implementation strategy, and introduce accountability through
50
51 297 monitoring.

52 298

53
54
55 299

1
2
3 **References:**
4

- 5 301 1. Creo AL, Thacher TD, Pettifor JM, Strand MA, Fischer PR. Nutritional rickets around the world:
6 302 an update. *Paediatr Int Child Health*. 2017;37(2):84–98.
- 7 303 2. Högler W. Complications of vitamin D deficiency from the foetus to the infant: One cause, one
8 304 prevention, but who's responsibility? *Best Pract Res Clin Endocrinol Metab*. 2015;29(3):385–
9 305 98.
- 10 306 3. Uday S, Högler W. Nutritional Rickets and Osteomalacia in the Twenty-first Century: Revised
11 307 Concepts, Public Health, and Prevention Strategies. *Curr Osteoporos Rep*. 2017;15(4):293–
12 308 302.
- 13 309 4. Ahmed SF, Franey C, McDevitt H, Somerville L, Butler S, Galloway P, et al. Recent trends and
14 310 clinical features of childhood vitamin D deficiency presenting to a children's hospital in
15 311 Glasgow. *Arch Dis Child*. 2011;96(7):694–6.
- 16 312 5. Basatemur E, Sutcliffe A. Incidence of hypocalcemic seizures due to vitamin D deficiency in
17 313 children in the United Kingdom and Ireland. *J Clin Endocrinol Metab*. 2015;100(1):E91–5.
- 18 314 6. Munns CF, Shaw N, Kiely M, Specker BL, Thacher TD, Ozono K, et al. Global Consensus
19 315 Recommendations on Prevention and Management of Nutritional Rickets. *J Clin Endocrinol*
20 316 *Metab*. 2016;101(2):394–415.
- 21 317 7. Farrar MD, Kift R, Felton SJ, Berry JL, Durkin MT, Allan D, et al. Recommended summer
22 318 sunlight exposure amounts fail to produce sufficient vitamin D status in UK adults of South
23 319 Asian origin. *Am J Clin Nutr*. 2011;94(5):1219–24.
- 24 320 8. Kift R, Berry JL, Vail A, Durkin MT, Rhodes LE, Webb AR. Lifestyle factors including less
25 321 cutaneous sun exposure contribute to starkly lower vitamin D levels in U.K. South Asians
26 322 compared with the white population. *Br J Dermatol*. 2013;169(6):1272–8.
- 27 323 9. Hyppönen E, Power C. Hypovitaminosis D in British adults at age 45y: nationwide cohort study
28 324 on dietary and lifestyle predictors. *Am J Clin Nutr*. 2007;85(3):860–8.
- 29 325 10. Office for National Statistics. Available from: <https://www.ons.gov.uk/>
- 30 326 11. Goldacre M, Hall N, Yeates DGR. Hospitalisation for children with rickets in England: A
31 327 historical perspective. *Lancet*. 2014;383(9917):597–8.
- 32 328 12. Hayes A, Cashman KD. Food-based solutions for vitamin D deficiency: putting policy into
33 329 practice and the key role for research. *Proc Nutr Soc*. 2017;76(1):54–63.
- 34 330 13. Uday S, Kongjonaj A, Aguiar M, Tulchinsky T, Högler W. Variations in infant and childhood
35 331 Vitamin D supplementation programs across Europe and factors influencing adherence.
36 332 *Endocr Connect*. 2017;6(8):667–75.
- 37 333 14. Ashraf S, Mughal MZ. The prevalence of rickets among non-Caucasian children. *Arch Dis*
38 334 *Child*. 2002;87(3):263–4.
- 39 335 15. Ladhani S, Srinivasan L, Buchanan C, Allgrove J. Presentation of vitamin D deficiency. *Arch*
40 336 *Dis Child*. 2004;89(8):781–4.
- 41 337 16. Callaghan AL, Moy RJD, Booth IW, DeBelle G, Shaw NJ. Incidence of symptomatic vitamin D
42 338 deficiency. *Arch Dis Child*. 2006;91(7):606–7.
- 43 339 17. Thacher TD, Pludowski P, Shaw NJ, Mughal MZ, Munns CF, Högler W. Nutritional rickets in
44 340 immigrant and refugee children. *Public Health Rev*. 2016;37(3):1–10.
- 45 341 18. Maiya S, Sullivan I, Allgrove J, Yates R, Malone M, Brain C, et al. Hypocalcaemia and vitamin
46 342 D deficiency: an important, but preventable, cause of life-threatening infant heart failure. *Heart*.
47 343 2008;94(5):581–4.

- 1
2
3 344 19. Scheimberg I, Perry L. Does Low Vitamin D Have a Role in Pediatric Morbidity and Mortality ?
4 345 An Observational Study of Vitamin D in a Cohort of 52 Postmortem Examinations. *Pediatr Dev*
5 346 *Pathol.* 2014;17(6):455–64.
- 6 347 20. Priemel M, Domarus C Von, Klatter TO, Kessler S, Schlie J, Meier S, et al. Bone Mineralization
7 348 Defects and Vitamin D Deficiency : Histomorphometric Analysis of iliac crest bone biopsies
8 349 and circulating 25-hydroxyvitamin D in 675 patients. *J Bone Miner Res.* 2010;25(2):305–12.
- 9
10 350 21. Ford JA, Colhoun EM, McIntosh WB, Dunnigan MG. Rickets and Osteomalacia in the Glasgow
11 351 Pakistani Community, 1961-71. *Br Med J.* 1972;2(5815):677–80.
- 12 352 22. Bivins R. Ideology and disease identity: the politics of rickets, 1929-1982. *Med Humanit.*
13 353 2014;40(1):3–10.
- 14
15 354 23. Högler W, Munns CF. Rickets and osteomalacia: a call for action to protect immigrants and
16 355 ethnic risk groups. *Lancet Glob Heal.* 2016;4(4):e229-30.
- 17 356 24. O'Neill CM, Kazantzidis A, Ryan MJ, Barber N, Sempos CT, Durazo-Arvizu RA, et al.
18 357 Seasonal changes in vitamin D-effective UVB availability in Europe and associations with
19 358 population serum 25-hydroxyvitamin D. *Nutrients.* 2016;8(9):E533.
- 20
21 359 25. Majeed H, Moore GWK. Impact of Multidecadal Climate Variability on United Kingdom Rickets
22 360 Rates. *Sci Rep.* 2017;7(1):1–6.
- 23 361 26. Darling AL, Hart KH, MacDonald HM, Horton K, Kang'Ombe AR, Berry JL, et al. Vitamin D
24 362 deficiency in UK South Asian Women of childbearing age: A comparative longitudinal
25 363 investigation with UK Caucasian women. *Osteoporos Int.* 2013;24(2):477–88.
- 26
27 364 27. Emmerson AJB, Dockery K, Mughal MZ, Roberts SA, Tower CL, Berry JL. Vitamin D status of
28 365 White pregnant women and infants at birth and 4 months in North West England: A cohort
29 366 study. *Matern Child Nutr.* 2018;14(1):e12453.
- 30 367 28. Cadario F, Savastio S, Magnani C, Cena T, Pagliardini V, Bellomo G, et al. High prevalence of
31 368 vitamin D deficiency in native versus migrant mothers and newborns in the North of Italy: A call
32 369 to act with a stronger prevention program. *PLoS One.* 2015;10(6).
- 33
34 370 29. National Diet and Nutrition Survey Results from Years 5 and 6 (combined) of the Rolling
35 371 Programme (2012/2013 – 2013/2014). 2016. Available from:
36 372 [https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/551352/NDNS_](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/551352/NDNS_Y5_6_UK_Main_Text.pdf)
37 373 [Y5_6_UK_Main_Text.pdf](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/551352/NDNS_Y5_6_UK_Main_Text.pdf)
- 38 374 30. Cashman, K D. Dowling KG. Vitamin D deficiency in Europe: pandemic? *Am J Clin Nutr.*
39 375 2016;103(4):1033–44.
- 40
41 376 31. Stewart WK, Mitchell RG, Morgan HG, Lowe KG, Thomson J. The Changing Incidence of
42 377 Rickets and Infantile Hypercalcaemia As Seen in Dundee. *Lancet.* 1964;1(7335):679–82.
- 43 378 32. Drury R, Rehm A, Johal S, Nadler R. Vitamin D supplementation: we must not fail our children!
44 379 *Medicine (Baltimore).* 2015;94(18):e817.
- 45
46 380 33. McFadden A, Green JM, McLeish J, McCormick F, Williams V, Renfrew MJ. Healthy Start
47 381 vitamins—a missed opportunity: findings of a multimethod study. *BMJ Open.*
48 382 2015;5(1):e006917–e006917.
- 49
50 383 34. NICE. Vitamin D: supplement use in specific population groups [Internet]. 2014. Available
51 384 from: <https://www.nice.org.uk/guidance/ph56>
- 52 385 35. Scientific Advisory Committee on Nutrition. Vitamin D and Health 2016. 2016. Available from:
53 386 [https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/537616/SACN_](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/537616/SACN_Vitamin_D_and_Health_report.pdf)
54 387 [Vitamin_D_and_Health_report.pdf](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/537616/SACN_Vitamin_D_and_Health_report.pdf)
- 55
56 388 36. Jessiman T, Cameron A, Wiggins M, Lucas PJ. A qualitative study of uptake of free vitamins in

- 1
2
3 389 England. Arch Dis Child. 2013;98(8):587–91.
- 4 390 37. Morris JK, Rankin J, Draper ES, Kurinczuk JJ, Springett A, Tucker D, et al. Prevention of
5 391 neural tube defects in the UK: A missed opportunity. Arch Dis Child. 2016;101(7):604–7.
- 6
7 392 38. Atta CAM, Fiest KM, Frolkis AD, Jette N, Pringsheim T, St Germaine-Smith C, et al. Global
8 393 birth prevalence of spina bifida by folic acid fortification status: A systematic review and meta-
9 394 analysis. Am J Public Health. 2016;106(1):e24–34.
- 10 395 39. Scientific Advisory Committee on Nutrition. Update on folic acid. 2017. Available from:
11 396 [https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/637111/SACN_](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/637111/SACN_Update_on_folic_acid.pdf)
12 397 [Update_on_folic_acid.pdf](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/637111/SACN_Update_on_folic_acid.pdf)
- 13
14 398 40. Bath SC, Steer CD, Golding J, Emmett P, Rayman MP. Effect of inadequate iodine status in
15 399 UK pregnant women on cognitive outcomes in their children: Results from the Avon
16 400 Longitudinal Study of Parents and Children (ALSPAC). Lancet. Elsevier Ltd;
17 401 2013;382(9889):331–7.
- 18 402
19
20 403
21
22 404
23 405
24
25 406
26
27 407
28
29 408
30
31 409
32 410
33
34 411
35
36 412
37 413
38
39 414
40
41 415
42 416
43
44 417
45
46 418
47
48 419
49 420
50
51 421
52 422
53
54 423
55
56 424
57
58
59
60

1
2
3 425 **Table 1: In high-income countries, rickets overall is a rare disease. However, the vast majority**
4 426 **of cases in these countries are from the BAME community, in which rickets is a common**
5 427 **disease [17]**

6 428

Country	Overall incidence (per 100,000)	Incidence in dark skinned (per 100,000)
UK	8	95
USA	24	220
Australia	5	2300
Denmark	3	60
European Union definition of a rare disease: affecting < 50 in 100,000 of the general population		

7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22 429

23
24
25 430

26 431

27
28 432
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 2: Parliamentary response to future plans on rickets prevention in the UK, which demonstrates serious misconceptions and incomplete knowledge of evidence (available from <http://www.parliament.uk/business/publications/written-questions-answers-statements/written-question/Lords/2017-11-13/HL3098/>)

Question	
To ask Her Majesty's Government whether they have any plans to designate a group of health care professionals to be accountable and responsible for the prevention of rickets and its complications. (HL2925)	
Parliamentary Response:	Revealed misconceptions and comments
<p>The Government has no plans to designate a group of healthcare professionals focused on the prevention of rickets.</p> <p>Rickets can generally be prevented by ensuring that children have a healthy, balanced diet, spend some time outside in the sun and take appropriate supplements containing ten micrograms of vitamin D.</p> <p>There is already a range of resources available to increase awareness of the need for vitamin D supplements, including advice on the NHS Choices website on the importance of vitamin D for bone health, and supplements are readily available over the counter and through the Healthy Start vitamins scheme.</p> <p>In August the National Institute for Health and Care Excellence (NICE) updated its public health guidance on increasing vitamin D supplement use among population groups at risk of vitamin D deficiency, in the light of the 2016 Scientific Advisory Committee on Nutrition report on Vitamin D and health. A copy of Vitamin D: supplement use in specific population groups is attached.</p>	<p>The Government instructs GPs to deliver prevention of serious infections (vaccinations) but assumes prevention of rickets will work by other means.</p> <p>The government does not understand the consequences of the UK's latitude and specific needs of the increasing BAME population, and ignores the rising incidence of symptomatic deficiency in the UK [5][11].</p> <p>This response ignores evidence of the ineffectiveness of those resources, given that nearly 85% of a well learned population in Oxford are unaware of the need for vitamin D supplementation in their children [32] and they fail to acknowledge the poor uptake of these vitamins [33].</p> <p>The NICE only makes recommendations. In this financially strained era, the government leaves the local commissioners with the choice of how to take the recommendations forward, without determining responsibilities or funding. This vitamin D prevention 'program' completely lacks mandatory monitoring [13] and accountability [2] much in contrast to other prevention</p>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

<p>This guideline includes recommendations on how to: increase access to and availability of vitamin D supplements, including for at-risk groups such as the BAME community, ensure health professionals recommend vitamin D supplements, raise awareness of the importance of these supplements amongst the population, and monitor provision and uptake of vitamin D supplements. Local commissioners will wish to consider how best to take forward these recommendations in respect of local BAME populations.</p>	<p>programs in infants (immunizations). The high incidence of rickets (Table 1) [4][5][15][18] and the increasing BAME proportion of the population (Figure 1) [10] requires robust policy implementation strategies. Moreover, the use of supplements in the population, especially BAME is low [8]. European countries which adopt monitoring of vitamin D supplementation during child health surveillance visits have demonstrated a high adherence in the first year of life [13].</p>
<p>The Government has made no formal assessment of the research referred to. Given the widespread availability of vitamin D supplements and clear guidance to health professionals and the public the Government does not believe there is a need for further strategies to prevent rickets.</p>	<p>Folic acid supplements have been widely available and campaigned for a number of years, but the incidence of neural tube defects in the UK continues to rise [37][38]. This simply goes to show that advocating supplementation does not work. The government needs to consider food fortification in order to eradicate rickets and osteomalacia in the UK [11].</p>

Confidential: For Review Only


Table 3: UK supplementation policy in comparison to global consensus recommendation:

United Kingdom Public Health England (PHE) (Following SACN guidance) Available from https://www.gov.uk/government/news/phe-publishes-new-advice-on-vitamin-d	Most of Continental Europe (Following global consensus recommendations [6])
Children aged 1 to 4 years should have a daily 10 µg vitamin D supplement. PHE recommends that babies are exclusively breastfed until around 6 months of age. As a precaution, all babies under 1 year should have a daily 8.5 to 10 µg (340 to 400 IU) vitamin D supplement to ensure they get enough. Children who have more than 500ml of infant formula a day do not need any additional vitamin D as formula is already fortified	Universal vitamin D supplementation of all infants with 10 µg (400 IU) daily, from birth

1
2
3 **Figure 1:** Census data (1991 to 2011) demonstrating an upward trend in the proportion of non-white
4 population and a downward trend in the proportion of white population. Data from national statistics
5 [10].
6
7
8
9


10 **Figure 2: UK net migration data over the last decade demonstrating an increase in the number**
11 **of people with dark skin living in the UK.** There has been a negative net migration of the White
12 British population and a positive net migration in the Black, Asian and Minority Ethnic (BAME)
13 population. BAME net migration was derived from national statistics data summarising the following
14 non-EU global regions of origin/destination. BAME = net Asian migrants + net Rest of the World
15 migrants [Sub-Saharan & North Africa, Central & South America, Oceania; excluding North American
16 migrants] [10]. The assumption was made that migrants from/to EU and North America are exclusively
17 white, and from/to BAME global regions are exclusively dark skinned. Net total migration (grey bar)
18 equals net BAME (black bar) + net white British (white bar) + net North America + net EU migration.
19
20
21
22
23

24 **Figure 3:** Adherence rates for infant vitamin D supplementation in the first year of life in Europe, with
25 UK reporting the lowest rates [13]. Good adherence ($\geq 80\%$ of infants supplemented) is indicated in
26 green, moderate adherence (79–50%) in orange and low adherence ($< 50\%$) in red.
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



Census data (1991 to 2011) demonstrating an upward trend in the proportion of non-white population and a downward trend in the proportion of white population. Data from national statistics [10].


203x162mm (96 x 96 DPI)



UK net migration data over the last decade demonstrating an increase in the number of people with dark skin living in the UK. There has been a negative net migration of the White British population and a positive net migration in the Black, Asian and Minority Ethnic (BAME) population. BAME net migration was derived from national statistics data summarising the following non-EU global regions of origin/destination. BAME = net Asian migrants + net Rest of the World migrants [Sub-Saharan & North Africa, Central & South America, Oceania; excluding North American migrants] [10]. The assumption was made that migrants from/to EU and North America are exclusively white, and from/to BAME global regions exclusively dark skinned. Net total migration (grey bar) equals net BAME (black bar) + net white British (white bar) + net North America + net EU migration.

203x162mm (96 x 96 DPI)

Only



Adherence rates for infant vitamin D supplementation in the first year of life in Europe, with UK reporting the lowest rates [13]. Good adherence ($\geq 80\%$ of infants supplemented) is indicated in green, moderate adherence (79–50%) in orange and low adherence (<50%) in red.

203x181mm (96 x 96 DPI)

Dear Editor

12/03/2018

Thank you for accepting our review article for publication in ADC. We have addressed the comments raised by the reviewer. A point by point response is detailed below and the relevant changes in the manuscript are highlighted in red.

Comments to the Author

This is a strong and important paper. The content is well informed and provides a timely critique of the current state of affairs with regard to the public health problems arising from widespread vitamin D deficiency. In parts it may lack balance in that arguments against the points being made are not always strongly presented e.g. the case against mandatory food fortification. However, that the paper is polemic in nature is clear from the title and tone.

1. Generally the language used in the paper to present evidence and categorise groups in the population at higher risk of vitamin D deficiency is accurate and sensitive. This is important so as not to detract from the key messages. However, on a handful of occasions, I believe that the choice of language might be improved. Skin colour is a continuum and so I have difficulty with the presentation of skin types within the general population as a dichotomy between "white" and "dark skinned".

Response: 'Dark skin' is a term used in SACN vitamin D and health report and the NICE guidelines on supplementation of risk groups. We recognise that skin colour is a continuum and therefore we have made the 'dark skin' (BAME) risk groups more objective by clarifying the Fitzpatrick skin types (IV-VI) we are referring to.

Amendments made: Where we first mention 'dark skin' - we have added the following in brackets [Fitzpatrick type IV-VI (light brown, dark brown, black)]. Page 3, line 77

2. The text accompanying Figure 2 provides some explanation of the authors' presentation of ONS data but the main body of the paper could usefully contain some more detail.

Response: We are very conscious of having already exceeded the word limit for reviews. The text contains the main information (net positive BAME migration) and the figure legend explains how the figure was derived, including underlying assumptions. We feel this is sufficient.

3. In particular, the last paragraph of the Introduction describes 14% of the population in 2011 as "dark skinned" without any qualification or reference to Figure 2.

Response: We believe you are referring to figure 1. We have now anchored Figure 1 in the text demonstrating the census population breakdown. Page 3, line 85

4. The same sentence provides a list of groups "wearing cultural whole body covering (Muslim, African, South Asian)" which runs the risk of giving the impression that everyone from the listed groups shares a single/common culture and that no one from outside the groups listed shares practices such as modest dress. I do not think that this is the intention of the authors.

Response: We have rephrased the sentence to include all groups.

Amendment: Whole body clothing for cultural, medical or other reasons further reduces sunlight exposure. Page 3, line 85-86

5. Some difficulties may be removed with the substitution of "darker skinned" for "dark skinned" in some instances. As an article that strongly advocates for the BAME population, it would be a shame if the message and intention was lost as a result of choices in words/language.

Response: Please see our response above to comment 1

1
2
3
4 6. In the introduction there is a list of four factors causing vitamin D deficiency in countries such as the
5 UK. It is stated that, "It is not difficult to see that none of the above factors are modifiable". In fact, only a
6 couple are not modifiable although I accept that the other two (culture and sunscreen use) may be difficult
7 to change (one more than the other).

8 **Response: Reworded to 'It is not difficult to see that the majority of the above factors are non-**
9 **modifiable'. Page 3, line 81**

10
11
12 7. p5 - at end of first paragraph there is mention of Scheimberg's work (ref 19). I think that it would be good
13 to provide some more detail re limitations of this work and/or at least to qualify the description of
14 "histological change" in order to provide greater clarity.

15
16 **Response: We agree that the work had some limitations and we have therefore made a comment to**
17 **highlight the lack of quantitative histology.**

18
19 **Amendments: added 'which however were not quantified'. Page 5, line 117.**

20
21 **We have also added that future studies should look at quantitative histology. Page 10, line 273**

22
23 8. p8 - Healthy Start should be capitalised, I think.

24 **Response: Amended accordingly**

25
26 9. Under "Preventative Strategies", whilst I do not disagree re mandatory fortification perhaps being a
27 solution, I think that it would be helpful to mention/consider voluntary food fortification as an option. The
28 experience with folic acid supplementation is rightly brought into the paper - surely this would suggest that
29 mandatory fortification with vitamin D is not likely to be imminent in the UK.

30
31
32 **Response: The current voluntary food fortification practice in the UK is not working and is unmonitored.**
33 **Without regulation, there is also the risk of cumulative excess vitamin D intake with consumption of multiple**
34 **fortified products.**

35
36
37 10. The list of "suggested responsibilities"/means of establishing accountability listed on p10 may be worth
38 highlighting more than they are at present.

39 **Response: We feel that we have sufficiently highlighted the suggested responsibilities on page 10.**
40 **However, we have in response, incorporated the need for 'establishing accountability' into the abstract.**
41 **Page 2, line 44**

42
43
44
45 Thanking you

46
47 Yours sincerely

48
49
50 Suma Uday