# UNIVERSITY<sup>OF</sup> BIRMINGHAM University of Birmingham Research at Birmingham

# Prevention of rickets and osteomalacia in the UK

Uday, Suma; Högler, Wolfgang

DOI: 10.1136/archdischild-2018-314826

License: None: All rights reserved

Document Version Peer reviewed version

Citation for published version (Harvard):

Uday, S & Högler, W 2018, 'Prevention of rickets and osteomalacia in the UK: political action overdue', Archives of Disease in Childhood, vol. 103, no. 9, pp. 901-906. https://doi.org/10.1136/archdischild-2018-314826

Link to publication on Research at Birmingham portal

#### **Publisher Rights Statement:**

Final version of record available at: http://dx.doi.org/10.1136/archdischild-2018-314826

#### **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

•Users may freely distribute the URL that is used to identify this publication.

Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)

•Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

#### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

# Archives of **Disease in Childhood**

# Prevention of rickets and osteomalacia in the United Kingdom: Political action overdue

Journal:	Archives of Disease in Childhood
Manuscript ID	archdischild-2018-314826.R1
Article Type:	Review
Edition:	not in use
Date Submitted by the Author:	n/a
Complete List of Authors:	Uday, Suma ; Birmingham Women's and Children's NHS Foundation Trust, Endocrinology and Diabetes; University of Birmingham , Institute of Metabolism and Systems Research Hoegler, Wolfgang; Birmingham Women's and Children's NHS Foundation Trust; University of Birmingham , Institute of Metabolism and Systems Research
Keywords:	immigration, dark skin, south Asian, African, vitamin D deficiency

**SCHOLARONE**<sup>™</sup> Manuscripts

2	1	Title
3	T	The.
5	2	Prevention of rickets and osteomalacia in the United Kingdom: Political action overdue
6 7	3	
8 9	4	Authors:
10 11	5	Suma Uday <sup>1,2</sup> , Wolfgang Högler <sup>1,2</sup>
12 13	6	1. Department of Endocrinology & Diabetes, Birmingham Women's and Children's Hospital,
14	/	Birmingham, UK
15	8	2. Institute of Metabolism and Systems Research, University of Birmingham, Birmingham, UK
16	9	
17		
18 19	10	Corresponding author:
20	11	PD Dr Wolfgang Högler,
21	12	Department of Endocrinology and Diabetes, Birmingham Women's and Children's Hospital,
22	13	Steelhouse Lane, B4 6NH, Birmingham, UK
23	14	Tel: +44 121 333 8197
25	15	Email: wolfgang hogler@nhs net
26	15	
27	16	
28	17	Keywords: Immigration, dark skin, South Asian, African, vitamin D deficiency, calcium, food
29 30	18	fortification
31 32	19	
33 34	20	Word count: 3427
35	21	Figures total: 3 Colour: 1
36 37	22	Tables: 3
38 39	23	
40 41	24	
42 43	25	
44	26	
46	27	
47 48	28	
49 50	29	
51 52	30	
53 54	31	
55 56	32	
57 58	33	
59 60		https://mc.manuscriptcentral.com/adc

#### Abstract

The consequences of vitamin D and dietary calcium deficiency have become a huge public health concern in the UK. The burden of disease from these deficiencies includes not just rickets, but also hypocalcaemic seizures, dilated cardiomyopathy and mostly occult myopathy and osteomalacia. The increasing burden of disease is intrinsically linked to ethnicity and the population demographic changes in the UK. Three facts have led to the resurfacing of the English disease: 1) The UK has no ultraviolet sunlight for at least 6 months of the year 2) Dark skin produces far less vitamin D than white skin per unit ultraviolet light exposure, and 3) non-EU immigration over the last century. To date, the UK government demonstrates incomplete understanding of these three facts, and its failure to adjust its prevention programs to changing demographics is endangering the health and life of UK residents with dark skin, of whom infants are the most vulnerable. Establishing accountability through d antenatar the implementation of monitored antenatal and infantile supplementation programs and mandatory food fortification is overdue.

### 48 Introduction

49 Quite in contrast to their continental European counterparts, paediatric A&E doctors in the UK often 50 encounter dark skinned infants with hypocalcaemic seizures, hypocalcaemic cardiomyopathy and 51 rickets during winter and spring. This article will explain the critical role of the UK government in the 52 causation and prevention of what was once called, and still is, the 'English disease'.

Rickets and osteomalacia occur due to defective mineralisation of hypertrophic growth plate chondrocytes and pre-formed osteoid in existing bone, respectively. In children, rickets and osteomalacia always co-exist. The most common cause of rickets and osteomalacia worldwide is calcium deprivation, secondary to low dietary calcium ('nutritional') and/or low ultraviolet B (UVB) sunlight exposure [1], the main source of cutaneous vitamin D production ('solar') [2]. There is very little vitamin D in diet (including breast milk) and thus humans are mainly dependent on solar UVB for vitamin D supply. Cholecalciferol (vitamin D3) and ergocalciferol (vitamin D2) undergo hepatic 25-hydroxylation to form calcidiol or 25 hydroxyvitamin D (250HD, stable form reflecting vitamin D status) and then renal 1-hydroxylation to form the hormone calcitriol or 1,25 dihydroxyvitamin D. In contrast to calcitriol, vitamin D and 25OHD are biologically inert. The main action of calcitriol is to aid intestinal calcium absorption. When dietary calcium or solar vitamin D availability is low, secondary hyperparathyroidism sets in which increases intestinal calcium absorption and bone resorption in the short term to maintain normal serum calcium concentrations. Unopposed, this compensation will fail in the long term and hyperparathyroidism-induced renal phosphate wasting will lead to hypophosphataemia, which ultimately impairs mineralisation on the tissue level causing rickets and osteomalacia [3]. Hypocalcaemic complications of calcium deprivation can occur before these hypophosphataemic complications, especially during phases of rapid growth i.e during infancy and adolescence [4][5].

71 Dietary calcium deficiency rickets (due to nutritional deficits) prevails in low income countries whereas 72 vitamin D deficiency rickets (due to reduced solar UVB exposure/latitude) predominates in high 73 income countries [3]. Calcium deprivation and all its hypocalcaemic or hypophosphataemic 74 complications can be easily prevented by ensuring adequate calcium and vitamin D supply [6]. 75 Factors causing vitamin D deficiency in high income countries such as the UK specifically include:

1) Latitude: lack of UVB radiation for at least 6 months of the year due to its geographic location [2][7]

2) Ethnicity: dark skin [Fitzpatrick skin type IV (light brown), V (dark brown) and VI (black)] massively

78 reduces cutaneous synthesis of vitamin D [7]

3) **Culture**: covered clothing, low dietary vitamin D and calcium intake [3][8]

4) **Sunscreen:** Excessive use of sunscreen [9] due to fear of skin cancer

It is not difficult to see that the majority of the above factors are non-modifiable. Therefore, supplementation or food fortification with vitamin D is necessary to prevent rickets and osteomalacia in an ethnically and culturally increasingly diverse society. Non-EU immigration over the last century has led to a change in population demographics with a growing proportion of the resident population (14% in the 2011 census) being dark skinned (Figure 1) [10]. Whole body clothing for cultural, medical or other reasons further reduces sunlight exposure. These high-risk groups are unable to produce sufficient vitamin D in the UK. Unfortunately, the UK has not adapted to its ethnic and cultural diversity which has led to a resurgence in rickets [11] and hypocalcaemic complications [5], a problem further augmented by a lack of mandatory food fortification [12] and a complex and poorly implemented infant vitamin D supplementation policy [13].

# 91 Burden of disease from vitamin D deficiency and rickets in the UK

'Nutritional' rickets was indeed originally caused by poor nutrition but was always also of 'solar' origin due to pollution during the industrial revolution. The fact that the English disease has resurfaced as a major public health problem in the UK in the last 50 years however is much less 'nutritional', but due to the increase in the ethnic high-risk proportion of the population. Over the past few decades several UK studies have highlighted the increasing burden of rickets [5][11][14][15][16] and the increased susceptibility of the Black, Asian and Minority Ethnic (BAME) groups. The burden of disease from solar vitamin D and dietary calcium deficiency is much wider and goes beyond the bone pathology. The British Paediatric Surveillance Unit survey (September 2011-2013) reported a total of 91 cases (85%, n=77 were infants) of hypocalcaemic seizures due to vitamin D deficiency, amounting to an annual incidence of 3.49 per million children (0-15 years), with highest incidence in the South Asian population at 26.04 per million [5]. In contrast to the white population, its high incidence in the BAME population precludes rickets from being classified as a rare disease (Table 1) [17]. A retrospective study from southeast England identified sixteen BAME cases (6 Asian, 10 Black) of hypocalcaemic

dilated cardiomyopathy secondary to severe vitamin D deficiency between 2000 and 2006 [18]. These cases displayed significant mortality (n=3) and morbidity requiring mechanical ventilatory and circulatory support (n=2 each) and referral for cardiac transplantation (n=2) [18]. In a post-mortem study from London (n=52), hypocalcaemia due to vitamin D deficiency was considered accountable for death in 3 BAME children; they all showed radiological and histological rickets: 2 babies had cardiomyopathy and a 3-year-old had hypocalcaemic seizures [19]. These reports are not collections of tragic medical rarities. Infants presenting with complications of hypocalcaemia such as seizures [4][5][11], dilated cardiomyopathy [18], cardiac death [19], and overt rickets [14][15] only represent the tip of the iceberg of a public health crisis. The extent of hidden pathology remains unknown. Histological changes of rickets and osteomalacia occur long before the disease becomes evident on radiographs [3]. One way to gain insight into the widespread prevalence of rickets and osteomalacia in the UK is from post mortem studies [19] which demonstrate histological changes in nearly 70% of infants studied, which however were not quantified. Osteomalacia in adults on post mortem examination is also highly prevalent (nearly 25% of a low-risk population studied, n=675) in northern Europe [20].

The increased prevalence of rickets in the immigrant and resident BAME population in the UK, especially the South Asian community, is not new knowledge and was noted as far back as the 1960s [21], when the UK government commonly referred to it as the 'Asian rickets' [22]. Although immigration of dark-skinned individuals has contributed to the rise in rickets in most developed countries [17][23], an additional contributing factor in the UK is its geographic location [24] and possibly the changing climate [25]. Hence, vitamin D deficiency is neither limited to the winter months nor exclusive to BAME population. A study from Manchester reported median serum 25OHD concentrations in summer of 28.6 nmol/L [interquartile range (IQR) 21.3-41.6], falling further to 18.4 nmol/L (IQR 12.7-25.7) in the winter in adult residents of South Asian origin [8]. These findings were resonated in another study in women of child bearing age where severe deficiency (250HD <25 nmol/L) was prevalent in 81% and 79.2% of UK South Asians in winter and autumn, respectively [26]. In a study of white women from North-West England (n=333) 27% and 7% had insufficient (<50 nmol/L) and deficient (<25 nmol/L) serum 25OHD levels during pregnancy, and 48% and 11% four months post-delivery, respectively [27]. Median 25OHD in cord blood samples were only 50% of maternal 25OHD levels and the prevalence of deficiency and insufficiency in infants at 4 months

(n=322) was 13% and 24%, respectively [27]. No UK study compared vitamin D levels in native and
migrant mother-baby pairs, but an Italian study showed that severe vitamin D deficiency (25OHD < 25</li>
nmol/L) was widely prevalent in the migrant mothers and their newborns (48.4% and 76.2%,
respectively) when compared to natives (38% and 18%, respectively) [28].

The UK National Diet and Nutrition Survey found that the mean intake of vitamin D was below the reference nutrient intake: only 29% for children aged 1.5 to 3 years and 33% for adults  $\ge$  65 years and nearly a fifth of all subjects studied were vitamin D deficient (250HD < 25 nmol/L) [29].

Although vitamin D deficiency is pandemic in Europe it is more prevalent in some mid-latitude countries including the UK when compared to northerly latitude countries such as Norway, Iceland or Finland even after accounting for ethnicity [30]. The lower prevalence in the northern latitude is attributed to better supplementation policies and also food fortification [30].

# 146 Factors contributing to resurgence of rickets in the UK

#### 147 The increasing BAME resident and immigrant population

A question on ethnic group was first introduced in the 1991 Census, not only "to enable monitoring for equal opportunities / anti-discrimination policies" but also "to plan for the future through resource allocation and provision of services" [10]. However, this does not seem to have happened in relation to rickets prevention despite the so obviously increasing proportion of dark skinned individuals. The most recent census (2011) reported that 14% of the population in England and Wales are of BAME origin which is a significant rise from previous census in 2001 (8.7%) and 1991 (5.9%) (Figure 1) [10]. Every year, nearly 13% of babies born in England and Wales have been of BAME background since 2014 [10] and around 100-150,000 people of BAME background (net) are added to the UK population through immigration (Figure 2) [10]. Therefore, the government needs to adjust its health care systems to these population demographic changes.

A multitude of factors contribute to the increased prevalence of vitamin D deficiency in BAME groups
residing in high latitude countries. An adult study of South Asian residents in Manchester reported
lower oral intake of dietary vitamin D (mean of 1·32 μg compared to 3·26 μg in white subjects), sun
avoidance when outside and lower amounts of skin surface exposure [8]. However by far the greatest

contributing factor is reduced skin synthesis of vitamin D [7]. Dark skin, when compared to white skin, requires much greater exposure to sunlight to improve circulating 250HD levels as melanin absorbs a substantial proportion of the UVB light [7]. Hence, even the UK recommended summer sunlight exposure amounts fail to produce sufficient circulating 25OHD in the dark skinned population [7]. Lack of ethnicity-specific guidance on sunlight exposure [7], minimal impact of dietary vitamin D in improving serum 25OHD levels [26] and poor supplementation policies in infants [13] and pregnant women all contribute to widespread vitamin D deficiency in dark skinned individuals, all year round [26].

#### 170 UK politics and policy: From success to failure

There is an interesting history behind the increasing burden of rickets in the UK. During World War II, Britain triumphed eradication of rickets through rationing, food fortification and supplementation of infants and pregnant women with cod-liver oil. Strangely, the same country which once demonstrated the effectiveness of good public health policies is now struggling to tackle this easily preventable disease. In the 1950s, uncontrolled fortification of infant formula and food (4000 IU/day, which is 10 times the current recommended dose of 400 IU/day) was implicated in the causation of infantile hypercalcaemia [31] which led the government to withdraw fortification. Ever since this time, the 'English disease' has returned to the UK to the extent that today, Britain serves as an example of how lack of effective policy implementation can lead to a rise in a preventable disease such as rickets. Rickets or 'Asian rickets' has enormous political significance and has been the subject of debate in parliament since the 1960s [22]. There is a suggestion that the resurgence of rickets symbolises socioeconomic inequality and racial or ethnic health disparity in the UK, which is discussed in detail elsewhere [22]. The government continues to shy away from the issue. The senior author submitted a parliamentary question to the House of Lords in November 2017 with evidence demonstrating the ineffectiveness of the UK's infant vitamin D supplementation program [13] and queried the government's future plans to prevent rickets in the UK. The response, detailed in Table 2, reflects the government's current misconceptions.

188 The UK prevention program: complex, outdated, unmonitored, ineffective

The Scientific Advisory Committee on Nutrition (SACN) sets vitamin D 'reference nutrient intake' for adults and children aged >4 years (10 µg or 400 IU/day) and 'safe intake' for infants (8.5-10 µg or 340-400 IU/day) and children aged 1-4 years (10 µg or 400 IU/day) [35]. The SACN does not recommend supplements or food fortification but acknowledges the difficulty of achieving the recommended intakes from natural food sources and advised the government to consider strategies for the UK population to achieve the above intake [35]. Whilst most authorities consider optimal levels of serum 25OHD to be above 50 nmol/L [6], the SACN recommends that levels should not fall below 25 nmol/L [35]. The National Institute for Health and Care Excellence (NICE) makes certain recommendations to multiple agencies (including the Department of Health, Public Health England and healthcare professionals) to increase supplement use in at risk populations [34]. Based on the latest SACN vitamin D and health report [35], Public Health England recommend that populations at risk should take supplements containing 10 µg (400 IU) of vitamin D all year round (https://www.gov.uk/government/news/phe-publishes-new-advice-on-vitamin-d). The recommendation for infants and children however is not only overly complex (Table 3) but also now out-dated in relation to formula feeding and dose for children >1 year. The evidence-based global consensus on the prevention of nutritional rickets recommends universal supplementation of all infants [6], which is followed by most European countries (85%, n=24/29) [13]. SACN does not recommend vitamin D supplements for formula-fed infants whilst new evidence demonstrates that formula feeding does not protect from rickets [5][6]. In the British Paediatric Surveillance Unit survey, 19% (n=15/77) of infants presenting with hypocalcaemic seizures were exclusively formula-fed and 9% (n=7) received mixed breast and formula feed [5].

In contrast to the global consensus recommendation (10 µg or 400 IU for infants and 15 µg or 600 IU for children > 1 year), SACN and Public Health England recommend 8.5 to 10 µg of vitamin D for infants and children and the government supplies Healthy Start vitamins which only provide 7.5 µg. The operation of the Healthy Start scheme which provides free vitamins to low income families is also very complex [33], leading to low uptake, poor accessibility and lack of motivation to take daily supplements [36], not to mention the lack of parental awareness [32]. Apart from the complexity of the guidance, it is mainly the complete lack of accountability and monitoring, that is responsible for the failing of the Healthy Start program to improve the vitamin D status of the at-risk population [33]. The

218 UK's unmonitored infant vitamin D supplementation program has the lowest adherence rate in 219 Europe, with just 5-20% of infants actually receiving vitamin D drops (**Figure 3**) [13].

## **Preventative strategies**

Given the low uptake of supplements the long term solution to vitamin D deficiency is mandatory food fortification [12] which has been successfully implemented in countries such as Canada. However, this may not be easily adopted by the UK government given that it also shies away from adopting folic acid fortification despite overwhelming evidence on its role in preventing neural tube defects [37]. Mandatory fortification of flour with folic acid is adopted by 78 countries but not the UK, which has resulted in a continued rise in neural tube defects in the UK [37]. Countries with mandatory folic acid fortification have seen a decline in neural tube defects compared to those with voluntary or no fortification [38]. Despite SACN recommendation to adopt mandatory folic acid fortification of flour (in 2006 and 2009), the UK government has not taken any positive steps. Not surprisingly in 2017, Food Standards Scotland requested SACN to provide advice on whether its previous recommendations on mandatory fortification still apply, with an intention to proceed unilaterally [39]. Similarly, iodine deficiency which is considered the single most important preventable cause of brain damage by the World Health Organisation, is thought to be re-emerging as a public health concern in the UK due to lack of national salt-iodisation and monitoring of iodine levels in pregnancy [40].

Whilst mandatory food fortification may be the best long term solution to improve the vitamin D status at a population level [12], certain measures in the interim are crucial in protecting the most vulnerable group - infants. As most infants acquire vitamin D deficiency from the mother it is equally important to ensure adequate supplementation during pregnancy and to start infant supplementation at birth. Success of supplementation policies requires that authorities adopt multiple, simple strategies. We have previously compared vitamin D supplementation policies across Europe and determined that the following policy implementation features were significantly associated with improved adherence in the first year of life: universal supplementation independent of feeding mode (p = 0.007), providing information on supplementation at discharge from neonatal units (p = 0.02), providing financial family support (p = 0.005) and monitoring adherence at child health surveillance visits (p = 0.001) [13]. The UK currently does not adopt any of the above features, except providing financial support only to low income families.

Public health policies need to be simple to work. Universal supplementation of all infants [6] enhances adherence [13], moreover formula feeds do not protect the infant from rickets [5][6] and hence the SACN guidance requires updating and simplifying. Most importantly, similar to the vaccination program, a group of healthcare professionals needs to be made responsible for delivering this prevention program, in the same way as vaccination program. In the authors' point of view, establishing accountability is the most essential change required in the UK.

- 253 Based on data from the European survey [13], the suggested responsibilities are as follows:
- Health care professionals providing antenatal care include monitoring of supplementation at every
   routine visit.
  - 2) Neonatal units provide specific information on prevention programs to parents, commence vitamin
     D supplementation at birth and provide first bottle at discharge.
- 258 3) Upon registration at the GP, adherence to vitamin D supplementation is checked and then
  259 monitored at every single vaccination time point and red book visit.
- 4) Inclusion of a monitoring question on adherence to vitamin D supplements in maternity notes and
  personal child health record (red book), at all routine time points, which reminds healthcare
  professionals and parents.
  - 5) The government should provide financial remuneration to GPs delivering this prevention program
    in a similar way as for the immunization program (prevention of infections). The government
    should consider linking financial family support to adherence to all prevention programs.

## **Future considerations**

The majority of the European medical literature on rickets and hypocalcaemic complications comes from the UK [1]. Current studies are mainly based on clinical presentation or serum 25OHD measurements. Further research is required to explore the true burden of rickets and specifically osteomalacia in the UK. Histological changes occur long before the disease becomes clinically, biochemically or radiologically evident [3]. Therefore, the true prevalence of occult rickets and osteomalacia can only be identified by quantitative growth plate histology and bone

histomorphometry. These investigations are invasive in alive children or adults but can be undertakenduring post-mortem examination.

Taking daily vitamin D supplements prevents rickets but may pose an issue with adherence, similar to folic acid supplementation, especially if un-monitored. However, vitamin D can also be easily and safely administered in bolus doses alleviating the need for daily supplementation and parental compliance. Licensed preparations for bolus administration are readily available in the UK. Feasibility of administering bolus doses of vitamin D during immunisations and routine child care visits in the UK should be explored. Likewise, safety and feasibility of administering bolus doses of vitamin D during antenatal visits should also be explored.

The increasing UK BAME population requires life-long supplementation. Since this can never be delivered, food fortification is the obvious solution to the problem. Further studies are required to bridge the gap in knowledge required to facilitate food fortification, and to compare the costeffectiveness of fortification and supplementation programs.

# **Conclusions**

The UK population is more ethnically and culturally diverse than ever. The increasing UK BAME population is fully exposed to the complications of vitamin D deficiency due to their dark skin and cultural traditions. The UK government demonstrates serious misconception of the actual public health issue and the implementation of effective prevention programs, and an astounding ignorance of changing population demographics. National statistics demonstrate a constant rise of the UK BAME resident population through continued non-EU immigration from countries with predominantly dark-skinned population (Asia, Africa), with high birth rates. We call for the government to consider adaptation of their public health policy implementation strategy, and introduce accountability through monitoring.

# 300 **References:**

1 2

3

4 5

6 7

8

9

10 11

12

13 14

15

16

17 18

19

58

59

- Creo AL, Thacher TD, Pettifor JM, Strand MA, Fischer PR. Nutritional rickets around the world: an update. Paediatr Int Child Health. 2017;37(2):84–98.
- Högler W. Complications of vitamin D deficiency from the foetus to the infant: One cause, one
   prevention, but who's responsibility? Best Pract Res Clin Endocrinol Metab. 2015;29(3):385–
   98.
- Uday S, Högler W. Nutritional Rickets and Osteomalacia in the Twenty-first Century: Revised Concepts, Public Health, and Prevention Strategies. Curr Osteoporos Rep. 2017;15(4):293– 308.
- Ahmed SF, Franey C, McDevitt H, Somerville L, Butler S, Galloway P, et al. Recent trends and clinical features of childhood vitamin D deficiency presenting to a children's hospital in Glasgow. Arch Dis Child. 2011;96(7):694–6.
- 3125.Basatemur E, Sutcliffe A. Incidence of hypocalcemic seizures due to vitamin D deficiency in<br/>children in the United Kingdom and Ireland. J Clin Endocrinol Metab. 2015;100(1):E91–5.
- 203146.Munns CF, Shaw N, Kiely M, Specker BL, Thacher TD, Ozono K, et al. Global Consensus22315Recommendations on Prevention and Management of Nutritional Rickets. J Clin Endocrinol23316Metab. 2016;101(2):394–415.
- 243177.Farrar MD, Kift R, Felton SJ, Berry JL, Durkin MT, Allan D, et al. Recommended summer25318sunlight exposure amounts fail to produce sufficient vitamin D status in UK adults of South26319Asian origin. Am J Clin Nutr. 2011;94(5):1219–24.
- 320
  320
  321
  322
  321
  322
  321
  322
  321
  322
  321
  322
  321
  322
  321
  322
  321
  322
  322
  323
  324
  325
  325
  326
  327
  327
  327
  328
  329
  329
  329
  320
  320
  321
  321
  321
  322
  321
  322
  321
  322
  321
  322
  322
  323
  324
  325
  325
  325
  326
  327
  327
  327
  328
  328
  329
  329
  329
  320
  321
  321
  321
  321
  321
  322
  321
  322
  321
  322
  321
  322
  321
  322
  321
  322
  321
  321
  321
  322
  321
  321
  321
  321
  321
  321
  321
  321
  321
  322
  321
  321
  321
  321
  322
  321
  321
  321
  322
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
  321
- 313239.Hyppönen E, Power C. Hypovitaminosis D in British adults at age 45y: nationwide cohort study32324on dietary and lifestyle predictors. Am J Clin Nutr. 2007;85(3):860–8.
- 33
   34 325 10. Office for National Statistics. Available from: https://www.ons.gov.uk/
- 35
   326
   327
   37
   326
   327
   327
   328
   327
   329
   329
   320
   320
   320
   321
   321
   322
   323
   324
   325
   326
   326
   327
   326
   327
   328
   329
   329
   320
   320
   321
   321
   321
   321
   321
   322
   323
   324
   325
   326
   326
   327
   326
   327
   328
   328
   329
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
   321
- 3832812.Hayes A, Cashman KD. Food-based solutions for vitamin D deficiency: putting policy into<br/>practice and the key role for research. Proc Nutr Soc. 2017;76(1):54–63.
- 40
  41
  42
  43
  43
  43
  44
  45
  46
  47
  47
  48
  49
  49
  49
  40
  40
  41
  42
  43
  43
  43
  43
  43
  44
  44
  45
  46
  46
  47
  48
  49
  49
  49
  49
  49
  40
  40
  40
  41
  41
  42
  43
  43
  43
  44
  44
  45
  46
  47
  47
  48
  49
  49
  49
  40
  40
  41
  41
  41
  41
  42
  43
  43
  44
  44
  45
  46
  46
  47
  47
  48
  49
  49
  40
  40
  41
  41
  41
  41
  41
  42
  43
  44
  44
  45
  46
  47
  47
  48
  49
  49
  49
  40
  40
  41
  41
  41
  41
  42
  43
  44
  44
  45
  46
  47
  47
  48
  49
  49
  49
  40
  40
  41
  41
  41
  41
  42
  43
  44
  44
  44
  44
  44
  45
  46
  46
  47
  47
  48
  48
  49
  49
  49
  49
  49
  49
  49
  40
  40
  41
  41
  41
  42
  43
  44
  44
  44
  44
  44
  44
  44
  44
  44
  44
  44
  44
  44
  44
  44
  44
  44
  44
  44
  44
  44
  44
  44
  44
  44
  <
- 4433314.Ashraf S, Mughal MZ. The prevalence of rickets among non-Caucasian children. Arch Dis45334Child. 2002;87(3):263–4.
- 464733515.Ladhani S, Srinivasan L, Buchanan C, Allgrove J. Presentation of vitamin D deficiency. Arch48336Dis Child. 2004;89(8):781–4.
- 49 337 16. Callaghan AL, Moy RJD, Booth IW, Debelle G, Shaw NJ. Incidence of symptomatic vitamin D
  50 338 deficiency. Arch Dis Child. 2006;91(7):606–7.
  51
- 5233917.Thacher TD, Pludowski P, Shaw NJ, Mughal MZ, Munns CF, Högler W. Nutritional rickets in53340immigrant and refugee children. Public Health Rev. 2016;37(3):1–10.
- 5434118.Maiya S, Sullivan I, Allgrove J, Yates R, Malone M, Brain C, et al. Hypocalcaemia and vitamin55342D deficiency: an important, but preventable, cause of life-threatening infant heart failure. Heart.563432008;94(5):581–4.57

1			
2 3 4 5	344 345 346	19.	Scheimberg I, Perry L. Does Low Vitamin D Have a Role in Pediatric Morbidity and Mortality? An Observational Study of Vitamin D in a Cohort of 52 Postmortem Examinations. Pediatr Dev Pathol. 2014;17(6):455–64.
6 7 8	347 348 349	20.	Priemel M, Domarus C Von, Klatte TO, Kessler S, Schlie J, Meier S, et al. Bone Mineralization Defects and Vitamin D Deficiency : Histomorphometric Analysis of iliac crest bone biopsies and circulating 25-hydroxyvitamin D in 675 patients. J Bone Miner Res. 2010;25(2):305–12.
9 10 11	350 351	21.	Ford JA, Colhoun EM, McIntosh WB, Dunnigan MG. Rickets and Osteomalacia in the Glasgow Pakistani Community, 1961-71. Br Med J. 1972;2(5815):677–80.
12 13	352 353	22.	Bivins R. Ideology and disease identity: the politics of rickets, 1929-1982. Med Humanit. 2014;40(1):3–10.
15 16	354 355	23.	Högler W, Munns CF. Rickets and osteomalacia: a call for action to protect immigrants and ethnic risk groups. Lancet Glob Heal. 2016;4(4):e229-30.
17 18 19 20	356 357 358	24.	O'Neill CM, Kazantzidis A, Ryan MJ, Barber N, Sempos CT, Durazo-Arvizu RA, et al. Seasonal changes in vitamin D-effective UVB availability in Europe and associations with population serum 25-hydroxyvitamin D. Nutrients. 2016;8(9):E533.
21 22	359 360	25.	Majeed H, Moore GWK. Impact of Multidecadal Climate Variability on United Kingdom Rickets Rates. Sci Rep. 2017;7(1):1–6.
23 24 25 26	361 362 363	26.	Darling AL, Hart KH, MacDonald HM, Horton K, Kang'Ombe AR, Berry JL, et al. Vitamin D deficiency in UK South Asian Women of childbearing age: A comparative longitudinal investigation with UK Caucasian women. Osteoporos Int. 2013;24(2):477–88.
27 28 29	364 365 366	27.	Emmerson AJB, Dockery K, Mughal MZ, Roberts SA, Tower CL, Berry JL. Vitamin D status of White pregnant women and infants at birth and 4 months in North West England: A cohort study. Matern Child Nutr. 2018;14(1):e12453.
30 31 32 33	367 368 369	28.	Cadario F, Savastio S, Magnani C, Cena T, Pagliardini V, Bellomo G, et al. High prevalence of vitamin D deficiency in native versus migrant mothers and newborns in the North of Italy: A call to act with a stronger prevention program. PLoS One. 2015;10(6).
34 35 36 37	370 371 372 373	29.	National Diet and Nutrition Survey Results from Years 5 and 6 (combined) of the Rolling Programme (2012/2013 – 2013/2014). 2016. Available from: https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/551352/NDNS_ Y5_6_UK_Main_Text.pdf
38 39 40	374 375	30.	Cashman, K D. Dowling KG. Vitamin D deficiency in Europe: pandemic? Am J Clin Nutr. 2016;103(4):1033–44.
41 42	376 377	31.	Stewart WK, Mitchell RG, Morgan HG, Lowe KG, Thomson J. The Changing Incidence of Rickets and Infantile Hypercalcaemia As Seen in Dundee. Lancet. 1964;1(7335):679–82.
43 44 45	378 379	32.	Drury R, Rehm A, Johal S, Nadler R. Vitamin D supplementation: we must not fail our children! Medicine (Baltimore). 2015;94(18):e817.
46 47 48	380 381 382	33.	McFadden A, Green JM, McLeish J, McCormick F, Williams V, Renfrew MJ. Healthy Start vitaminsa missed opportunity: findings of a multimethod study. BMJ Open. 2015;5(1):e006917–e006917.
49 50 51	383 384	34.	NICE. Vitamin D: supplement use in specific population groups [Internet]. 2014. Available from: https://www.nice.org.uk/guidance/ph56
52 53 54 55	385 386 387	35.	Scientific Advisory Committee on Nutrition. Vitamin D and Health 2016. 2016. Available from: https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/537616/SACN_ Vitamin_D_and_Health_report.pdf
56 57	388	36.	Jessiman T, Cameron A, Wiggins M, Lucas PJ. A qualitative study of uptake of free vitamins in
58 59 60			13 https://mc.manuscriptcentral.com/adc

3	389		England. Arch Dis Child. 2013;98(8):587–91.
4 5 6	390 391	37.	Morris JK, Rankin J, Draper ES, Kurinczuk JJ, Springett A, Tucker D, et al. Prevention of neural tube defects in the UK: A missed opportunity. Arch Dis Child. 2016;101(7):604–7.
7 8 9	392 393 394	38.	Atta CAM, Fiest KM, Frolkis AD, Jette N, Pringsheim T, St Germaine-Smith C, et al. Global birth prevalence of spina bifida by folic acid fortification status: A systematic review and meta- analysis. Am J Public Health. 2016;106(1):e24–34.
10 11 12 13	395 396 397	39.	Scientific Advisory Committee on Nutrition. Update on folic acid. 2017. Available from: https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/637111/SACN_ Update_on_folic_acid.pdf
14 15 16 17	398 399 400 401	40.	Bath SC, Steer CD, Golding J, Emmett P, Rayman MP. Effect of inadequate iodine status in UK pregnant women on cognitive outcomes in their children: Results from the Avon Longitudinal Study of Parents and Children (ALSPAC). Lancet. Elsevier Ltd; 2013;382(9889):331–7.
18 19	402		
20	403		
22	404		
23	405		
25 26	406		
27 28	407		
29 30	408		
31	409		
33	410		
34 35	411		
36 37	412		
38 39	415		
40 41	415		
42 43	416		
44	417		
45 46	418		
47 48	419		
49 50	420		
51 52	421		
53	422		
54 55	423		
56 57	424		
58 59			14

disease [17]

Table 1: In high-income countries, rickets overall is a rare disease. However, the vast majority

of cases in these countries are from the BAME community, in which rickets is a common

	Country	Overall incidence (per 100,000)	Incidence in dark skinned (per 100,000)
	UK	8	95
	USA	24	220
	Australia	5	2300
	Denmark	3	60
	European Union defin	nition or a rare disease: affecting < 5	0 in 100,000 of the general population
29		<i>C</i> .	
30			
31			
2			

# Table 2: Parliamentary response to future plans on rickets prevention in the UK, which demonstrates serious misconceptions and incomplete knowledge of evidence (available from <a href="http://www.parliament.uk/business/publications/written-questions-answers-statements/written-question/Lords/2017-11-13/HL3098/">http://www.parliament.uk/business/publications/written-questions-answers-statements/written-question/Lords/2017-11-13/HL3098/</a>)

# Question

To ask Her Majesty's Government whether they have any plans to designate a prevention of rickets and its complications. (HL2925)	group of health care professionals to be accountable and responsible for the
Parliamentary Response:	Revealed misconceptions and comments
The Government has no plans to designate a group of healthcare	The Government instructs GPs to deliver prevention of serious infections
professionals focused on the prevention of rickets.	(vaccinations) but assumes prevention of rickets will work by other means.

Rickets can generally be prevented by ensuring that children have a healthy, balanced diet, spend some time outside in the sun and take appropriate supplements containing ten micrograms of vitamin D.

There is already a range of resources available to increase awareness of the need for vitamin D supplements, including advice on the NHS Choices website on the importance of vitamin D for bone health, and supplements are readily available over the counter and through the Healthy Start vitamins scheme.

In August the National Institute for Health and Care Excellence (NICE) updated The National Institute for Health and Care Excellence (NICE) updated The National public health guidance on increasing vitamin D supplement use among the go population groups at risk of vitamin D deficiency, in the light of the 2016 take to Scientific Advisory Committee on Nutrition report on Vitamin D and health. A funding copy of Vitamin D: supplement use in specific population groups is attached.

a latitude and specific needs of the increasing BAME population, and ignores the rising incidence of symptomatic deficiency in the UK [5][11].
 This response ignores evidence of the ineffectiveness of those resources,

The government does not understand the consequences of the UK's

Inis response ignores evidence of the ineffectiveness of those resources,
given that nearly 85% of a well learned population in Oxford are unaware
ly of the need for vitamin D supplementation in their children [32] and they
fail to acknowledge the poor uptake of these vitamins [33].

The NICE only makes recommendations. In this financially strained era, the government leaves the local commissioners with the choice of how to take the recommendations forward, without determining responsibilities or funding. This vitamin D prevention 'program' completely lacks mandatory monitoring [13] and accountability [2] much in contrast to other prevention

This guideline includes recommendations on how to: increase access to and	programs in infants (immunizations). The high incidence of rickets (Table
availability of vitamin D supplements, including for at-risk groups such as the	1) [4][5][15][18] and the increasing BAME proportion of the population
BAME community, ensure health professionals recommend vitamin D	(Figure 1) [10] requires robust policy implementation strategies. Moreover,
supplements, raise awareness of the importance of these supplements	the use of supplements in the population, especially BAME is low [8].
amongst the population, and monitor provision and uptake of vitamin D	European countries which adopt monitoring of vitamin D supplementation
supplements. Local commissioners will wish to consider how best to take	during child health surveillance visits have demonstrated a high adherence
forward these recommendations in respect of local BAME populations.	in the first year of life [13].
406	
The Government has made no formal assessment of the research referred to.	Folic acid supplements have been widely available and campaigned for a
Given the widespread availability of vitamin D supplements and clear guidance	number of years, but the incidence of neural tube defects in the UK
to health professionals and the public the Government does not believe there	continues to rise [37][38]. This simply goes to show that advocating
is a need for further strategies to prevent rickets.	supplementation does not work. The government needs to consider food
	fortification in order to eradicate rickets and osteomalacia in the UK [11].

#### Table 3: UK supplementation policy in comparison to global consensus recommendation:

United Kingdom Public Health England (PHE)	Most of Continental Europe			
(Following SACN guidance)	(Following global consensus			
Available from	recommendations [6])			
https://www.gov.uk/government/news/phe-				
publishes-new-advice-on-vitamin-d				
Children aged 1 to 4 years should have a daily 10 $\mu$ g vitamin D supplement. PHE recommends that babies are exclusively breastfed until around 6 months of age. As a precaution, all babies under 1 year should have a daily 8.5 to 10 $\mu$ g (340 to 400 IU) vitamin D supplement to ensure they get enough. Children who have more than 500ml of infant formula a day do not need any additional vitamin D as formula is already fortified	Universal vitamin D supplementation of all infants with 10 μg (400 IU) daily, from birth			

**Figure 1:** Census data (1991 to 2011) demonstrating an upward trend in the proportion of non-white population and a downward trend in the proportion of white population. Data from national statistics [10].

**Figure 2: UK net migration data over the last decade demonstrating an increase in the number of people with dark skin living in the UK.** There has been a negative net migration of the White British population and a positive net migration in the Black, Asian and Minority Ethnic (BAME) population. BAME net migration was derived from national statistics data summarising the following non-EU global regions of origin/destination. BAME = net Asian migrants + net Rest of the World migrants [Sub-Saharan & North Africa, Central & South America, Oceania; excluding North American migrants] [10]. The assumption was made that migrants from/to EU and North America are exclusively white, and from/to BAME global regions are exclusively dark skinned. Net total migration (grey bar) equals net BAME (black bar) + net white British (white bar) + net North America + net EU migration.

Figure 3: Adherence rates for infant vitamin D supplementation in the first year of life in Europe, with UK reporting the lowest rates [13]. Good adherence (280% of infants supplemented) is indicated in green, moderate adherence (79–50%) in orange and low adherence (<50%) in red.



60



Census data (1991 to 2011) demonstrating an upward trend in the proportion of non-white population and a downward trend in the proportion of white population. Data from national statistics [10].

203x162mm (96 x 96 DPI)



UK net migration data over the last decade demonstrating an increase in the number of people with dark skin living in the UK. There has been a negative net migration of the White British population and a positive net migration in the Black, Asian and Minority Ethnic (BAME) population. BAME net migration was derived from national statistics data summarising the following non-EU global regions of origin/destination. BAME = net Asian migrants + net Rest of the World migrants [Sub-Saharan & North Africa, Central & South America, Oceania; excluding North American migrants] [10]. The assumption was made that migrants from/to EU and North America are exclusively white, and from/to BAME global regions exclusively dark skinned. Net total migration (grey bar) equals net BAME (black bar) + net white British (white bar) + net North America + net EU migration.

203x162mm (96 x 96 DPI)





Adherence rates for infant vitamin D supplementation in the first year of life in Europe, with UK reporting the lowest rates [13]. Good adherence (≥80% of infants supplemented) is indicated in green, moderate adherence (79–50%) in orange and low adherence (<50%) in red.

203x181mm (96 x 96 DPI)



#### Dear Editor

#### 12/03/2018

Thank you for accepting our review article for publication in ADC. We have addressed the comments raised by the reviewer. A point by point response is detailed below and the relevant changes in the manuscript are highlighted in red.

#### Comments to the Author

This is a strong and important paper. The content is well informed and provides a timely critique of the current state of affairs with regard to the public health problems arising from widespread vitamin D deficiency. In parts it may lack balance in that arguments against the points being made are not always strongly presented e.g. the case against mandatory food fortification. However, that the paper is polemic in nature is clear from the title and tone.

1. Generally the langauage used in the paper to present evidence and categorise groups in the population at higher risk of vitamin D deficiency is accurate and sensitive. This is important so as not to detract from the key meassages. However, on a handful of occasions, I believe that the choice of language might be improved. Skin colour is a continuum and so I have difficulty with the presentation of skin types within the general population as a dichotomy between "white" and "dark skinned".

Response: 'Dark skin' is a term used in SACN vitamin D and health report and the NICE guidelines on supplementation of risk groups. We recognise that skin colour is a continuum and therefore we have made the 'dark skin' (BAME) risk groups more objective by clarifying the Fitzpatrick skin types (IV-VI) we are referring to.

Amendments made: Where we first mention 'dark skin'- we have added the following in brackets [Fitzpatrick type IV-VI (light brown, dark brown, black)]. Page 3, line 77

2. The text accompanying Figure 2 provides some explanation of the authors' presentation of ONS data but the main body of the paper could usefully contain some more detail.

Response: We are very conscious of having already exceeded the word limit for reviews. The text contains the main information (net positive BAME migration) and the figure legend explains how the figure was derived, including underlying assumptions. We feel this is sufficient.

3. In particular, the last paragraph of the Introduction describes 14% of the population in 2011 as "dark skinned" without any qualification or reference to Figure 2.

Response: We believe you are referring to figure 1. We have now anchored Figure 1 in the text demonstrating the census population breakdown. Page 3, line 85

4. The same sentence provides a list of groups "wearing cultural whole body covering (Muslim, African, South Asian)" which runs the risk of giving the impression that everyone from the listed groups shares a single/common culture and that no one from outside the groups listed shares practices such as modest dress. I do not think that this is the intention of the authors.

Response: We have rephrased the sentence to include all groups.

Amendment: Whole body clothing for cultural, medical or other reasons further reduces sunlight exposure. Page 3, line 85-86

5. Some difficulties may be removed with the substitution of "darker skinned" for "dark skinned" in some instances. As an article that strongly advocates for the BAME population, it would be a shame if the message and intention was lost as a result of choices in words/langauge.

Response: Please see our response above to comment 1

6. In the introduction there is a list of four factors causing vitamin D deficiency in countries such as the UK. It is stated that, "It is not difficult to see that none of the above factors are modifiable". In fact, only a couple are not modifiable although I accept that the other two (culture and sunscreen use) may be difficult to change (one more than the other).

Response: Reworded to 'It is not difficult to see that the majority of the above factors are nonmodifiable'. Page 3, line 81

7. p5 - at end of first paragraph ther is mention of Scheimberg's work (ref 19). I think that it would be good to provide some more detail re limitations of this work and/or at least to qualify the description of "histological change" in order to provide greater clarity.

Response: We agree that the work had some limitations and we have therefore made a comment to highlight the lack of quantitative histology.

Amendments: added 'which however were not quantified'. Page 5, line 117.

We have also added that future studies should look at quantitative histology. Page 10, line 273

8. p8 - Healthy Start should be capitalised, I think.

Response: Amended accordingly

9. Under "Preventative Strategies", whilst I do not disagree re mandatory fortification perhaps being a solution, I think that it would be helpful to mention/consider voluntary food fortification as an option. The experience with folic acid supplementation is rightly brought into the paper - surely this would suggest that mandatory fortification with vitamin D is not likely to be imminent in the UK.

Response: The current voluntary food fortification practice in the UK is not working and is unmonitored. Without regulation, there is also the risk of cumulative excess vitamin D intake with consumption of multiple fortified products.

10. The list of "suggested responsibilities"/means of establishing accountability listed on p10 may be worth highlighting more than they are at present.

Response: We feel that we have sufficiently highlighted the suggested responsibilities on page 10. However, we have in response, incorporated the need for 'establishing accountability' into the abstract. Page 2, line 44

Thanking you

Yours sincerely

Suma Uday