

Groups with a p -element acting with a single non-trivial Jordan block on a simple module in characteristic p

Craven, David

DOI:
[10.1515/jgth-2018-0014](https://doi.org/10.1515/jgth-2018-0014)

License:
None: All rights reserved

Document Version
Peer reviewed version

Citation for published version (Harvard):
Craven, D 2018, 'Groups with a p -element acting with a single non-trivial Jordan block on a simple module in characteristic p ', *Journal of Group Theory*, vol. 21, no. 5, pp. 719-787. <https://doi.org/10.1515/jgth-2018-0014>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Craven, Groups with a p -element acting with a single non-trivial Jordan block on a simple module in characteristic p , *J. Group Theory* 21 (2018), 719 – 787

DOI 10.1515/jgth-2018-0014

©de Gruyter 2018

checked on 6/12/2018

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Groups with a p -element acting with a single non-trivial Jordan block on a simple module in characteristic p

David A. Craven

April 12, 2018

Let V be a vector space over a field of characteristic p . In this paper we complete the classification of all irreducible subgroups G of $\mathrm{GL}(V)$ that contain a p -element whose Jordan normal form has exactly one non-trivial block, and possibly multiple trivial blocks. Broadly speaking, such a group acting primitively is a classical group acting on a symmetric power of a natural module, a 7-dimensional orthogonal group acting on the 8-dimensional spin module, a complex reflection group acting on a reflection representation, or one of a small number of other examples, predominantly with a self-centralizing cyclic Sylow p -subgroup.

1 Introduction

The classification of primitive permutation groups that contain a p^a -cycle (see [32] and [4, p.229]) has been of great use in answering a variety of problems in permutation group theory. In a different direction, groups generated by transvections have been studied by many authors, culminating in a complete determination in [15]. A simultaneous generalization of these two concepts, when the transvections are unipotent, is the idea of a *minimally active* element. This is a unipotent element u whose Jordan normal form has at most one block of size greater than 1, and all other blocks of size 1. (This is equivalent to $\dim(C_{M/C_M(u)}(u)) \leq 1$, or $[M, u] \cap C_M(u)$ having dimension at most 1, where M is the underlying vector space.) Such elements also appeared in work of Oliver, Semeraro and the author [3] in the context of exotic fusion systems. The more general concept of *almost cyclic* elements in matrix groups has been looked at for sporadic groups [24] and Weil modules for classical groups [25], and minimally active elements for algebraic groups have been studied in [29] and [30]. (An almost cyclic matrix is a matrix that is similar to the sum of a scalar matrix and a matrix whose minimal and characteristic polynomials coincide.)

In this article we give a general classification theorem for all irreducible subgroups of $\mathrm{GL}(M)$ that contain a minimally active element. In some cases, most notably irreducible but imprimitive subgroups, because of the wide range of examples, we give a general construction of such groups but cannot in any real sense give a full classification.

Theorem 1.1. *Let M be a vector space over a field k of characteristic p , and let $G \leq \mathrm{GL}(M)$ be an irreducible subgroup. If G contains a unipotent element u such that $\dim(C_{M/C_M(u)}(u)) = 1$ then one of the following holds:*

- (i) G acts imprimitively on M , the element u acts on M with a single Jordan block (i.e., $\dim(C_M(u)) = 1$) and if

$$M = M_1 \oplus M_2 \oplus \cdots \oplus M_t$$

is the finest direct sum decomposition stabilized by G , then t is a power of p and u^t stabilizes each M_i and acts with a single Jordan block on each M_i .

- (ii) p is odd and M factorizes as $M_1 \otimes M_2$ with $\dim(M_i) = 2$, and u lies in the central product $\mathrm{SL}_2(k) \circ \mathrm{SL}_2(k)$ with the obvious action on M , with u acting on M with a block of size 3 and a block of size 1;
- (iii) G stabilizes a factorization of M as $M_1 \otimes M_2$ with $\dim(M) = 4, 8, 9$, $p = 2, 3, 2$ respectively, and $o(u) = 4, 9, 8$ respectively, with u acting on M with a single block if $\dim(M) = 4, 8$, and a block of size 8 and one of size 1 if $\dim(M) = 9$;
- (iv) p is a Fermat or Mersenne prime $2^n \pm 1$, $\dim(M) = 2^n$, G is a subgroup of the extraspecial type group $(Z_4 \circ 2^{1+2n}).\mathrm{Sp}_{2n}(2)$, $o(u) = p$ and u acts with a block of size p and one of size 1 if p is Mersenne, and a single block if p is Fermat;
- (v) $p = 3$, $\dim(M) = 2^n$ for $n = 2, 3$, G is a subgroup of the extraspecial type group $(Z_4 \circ 2^{1+2n}).\mathrm{Sp}_{2n}(2)$, $o(u) = 3, 9$ respectively, and u acts with either a block of size 3 and a block of size 1, or a single block of size 8;
- (vi) the image of G in $\mathrm{PGL}(M)$ is almost simple, and G acts absolutely irreducibly.

Cases (i)–(v) are in some sense general, although note that (ii), (iii) and (v) only occur for $\dim(M) < 10$, with (i) and (iv) being the generic case. For (i) in particular, and also for the other cases, we give more information and are more specific about which classes occur in the relevant sections.

Of course, for (vi) we can be much more specific, and this is the content of the next theorem. We say that $G \leq \mathrm{GL}(M)$ is *tensor decomposable* if $M = M_1 \otimes \cdots \otimes M_t$ with G stabilizing the factors, so that G is a subgroup of $\mathrm{GL}(M_1) \wr \mathrm{Sym}_t$.

Theorem 1.2. *Let M be a vector space over a field of characteristic p , and let $G \leq \mathrm{GL}(M)$ be an irreducible subgroup such that the image of G in $\mathrm{PGL}(M)$ is almost simple. Suppose that G acts primitively and tensor indecomposably. If G contains a unipotent element u such that $\dim(C_{M/C_M(u)}(u)) = 1$ then (up to automorphism) G is one of the following:*

- (i) a linear or unitary group acting on a symmetric power of the natural module, a classical group, $G_2(q)$, ${}^2G_2(q)$, ${}^2B_2(q)$, or ${}^3D_4(q)$ acting on the natural module (minimal module for G_2), or $\mathrm{PSL}_3(2^a).2$ in characteristic 2 and M of dimension 8;
- (ii) the group $\mathrm{Spin}_7(q)$ acting on the 8-dimensional spin module;
- (iii) a subgroup of a complex reflection group acting on a non-trivial composition factor of a reflection representation;
- (iv) a group with a self-centralizing cyclic Sylow p -subgroup with $\dim(M) \leq o(u) + 1$;
- (v) one of the groups
 - (a) $\mathrm{Alt}_7 \leq \mathrm{SL}_4(2)$;
 - (b) $2 \cdot \mathrm{Alt}_7 \leq \mathrm{SU}_4(3)$;
 - (c) $3 \cdot M_{22} \leq \mathrm{SL}_6(4)$;
 - (d) $J_2 \leq \mathrm{SL}_6(4)$;

$$(e) \ 3 \cdot M_{10} \leq \mathrm{GL}_9(4);$$

$$(f) \ 3 \cdot J_3 \leq \mathrm{SL}_9(4).$$

(For (i) of this result, we allow C_2 to be viewed as B_2 and A_3 as D_3 , so they have two ‘natural’ modules.)

We give exact descriptions of all pairs of almost simple groups and simple modules such that the group contains an element acting minimally actively in various results throughout the paper, but there are far too many to list here. Alternating groups are in Propositions 4.1 and 4.3, sporadic groups are in Proposition 6.2, Lie type groups in characteristic not p are in Propositions 7.1, 8.3, 8.5 and 9.1, and Lie type in characteristic p are given in Propositions 5.1, 5.2 and 5.3.

Given the results in [24, 25, 29, 30], what remains for almost quasisimple groups is the alternating groups, outer automorphisms of groups of Lie type in defining characteristic, outer automorphisms of sporadic groups, and groups of Lie type in cross characteristic acting on non-Weil modules. After a preliminary section establishing notation and proving some important basic lemmas, in Section 3 we prove Theorem 1.1. After this, we work with almost simple groups, studying alternating groups in Section 4, Lie type groups in defining characteristic in Section 5 and sporadic groups in Section 6. In Section 7 we give some preliminaries about groups of Lie type in cross characteristic, and then Sections 8 and 9 consider classical and exceptional groups respectively. Finally, Section 10 establishes Theorem 1.2.

2 Notation and Preliminaries

Throughout this paper, let p be a prime and let k be an algebraically closed field of characteristic p . Let G be a finite group such that $p \mid |G|$, and let u be a p -element of G .

For specific groups, we write Alt_n for the alternating group of degree n , to distinguish it from the algebraic group of type A , and write Sym_n for consistency. Similarly, a cyclic group of order n will be denoted Z_n rather than C_n . Groups of Lie type are given their standard names of SL , PSL , PSP , and so on.

All modules considered are finite-dimensional and defined over k . We denote the trivial module by k or k_G if the group needs to be emphasized, and if H is a subgroup of G and M is a kG -module then $M \downarrow_H$ is the restriction of M to H . As usual, \oplus and \otimes denote direct sum and tensor product, with $\Lambda^i(M)$ and $S^i(M)$ denoting the exterior and symmetric powers of M .

If M is a kG -module and u is a p -element then the action of u on M is conjugate in $\mathrm{GL}(M)$ to a triangular matrix and has a Jordan normal form, made up of blocks of various sizes. If the action of u is conjugate to a triangular matrix with Jordan blocks of sizes m_1, \dots, m_r , then we say that u has type (m_1, \dots, m_r) on M . We often place the m_i in weakly decreasing order, but this is not necessary.

The modules of interest are as follows.

Definition 2.1. *Let G be a finite group and let k be an algebraically closed field of characteristic $p > 0$. If u is a p -element of G and M is a kG -module, then u acts minimally actively on M if, in the Jordan normal form of u on M , there is at most one Jordan block of size greater than 1, i.e., if u has type $(m, 1, \dots, 1)$ for some $m \geq 1$ on M , or equivalently if $[M, u] \cap C_M(u)$ is at most 1-dimensional. We say that M is minimally active if there exists a non-trivial 2-element acting minimally actively on M .*

Notice that the identity acts minimally actively on all modules, and all p -elements act minimally actively on 1-dimensional modules.

We use the term minimally active here, following [3], rather than almost cyclic, following [24, 30], because almost cyclic elements need not be p -elements, where p is the characteristic of the underlying field, i.e., unipotent elements of the corresponding $\text{GL}(M)$. Since we definitely require this extra hypothesis, we prefer to use this more specific term, to avoid leading the reader to believe we have classified all irreducible groups containing almost cyclic elements.

In our work we often need to know how many conjugates of a given element u generate the normal closure of $\langle u \rangle$ in a given group, so we introduce some notation, following [8].

Definition 2.2. *Let G be a finite group and let u be an element of G . We denote by $\alpha(u)$ the smallest number of conjugates $u_1, \dots, u_{\alpha(u)}$ of u such that*

$$\langle u_1, \dots, u_{\alpha(u)} \rangle = \langle u^G \rangle,$$

i.e., the fewest number of conjugates of u needed to generate the normal closure of $\langle u \rangle$. Write

$$\alpha(G) = \max_{u \in G} \alpha(u).$$

Of course, if a group has even order and is not dihedral then $\alpha(G) \geq 3$, and $\alpha(G)$ is the maximum of $\alpha(u)$ for all elements u of prime order. In [8], various bounds for almost simple groups were obtained, and we will use them frequently to get general constraints on finite groups with elements acting minimally actively on a simple module. For example, in Lemmas 2.9 and 2.10 we show that $\alpha(u) = 2$ for some specific conjugacy classes of permutations inside symmetric groups.

We collect several basic facts about minimally active modules now.

Lemma 2.3. *Let G be a finite group and let M be a faithful kG -module.*

- (i) *If u acts minimally actively on M , then u acts minimally actively on any submodule or quotient of M , and on the dual of M .*
- (ii) *If u is contained in a subgroup H of G and acts minimally actively on M , then u acts minimally actively on $M \downarrow_H$.*
- (iii) *If $M = M_1 \oplus M_2$ and u acts minimally actively on M , then $\langle u^G \rangle$ acts trivially on at least one of the M_i .*
- (iv) *If M is simple and u acts minimally actively on M then $\dim(M) \leq \alpha(u) \cdot (o(u) - 1)$. More generally, if $a = \dim(M) - \dim(C_M(u))$ then $\dim(M) \leq a \cdot \alpha(u)$.*
- (v) *Suppose that M and N are kG -modules with $\dim(M) \leq \dim(N)$, and such that u acts non-trivially on $M \otimes N$. We have that u acts minimally actively on $M \otimes N$ if and only if either $\dim(M) = 1$ and u acts minimally actively on N , or p is odd, $\dim(M) = \dim(N) = 2$, and u acts non-trivially (i.e., has type (2)) on both M and N .*
- (vi) *If u acts non-trivially and minimally actively on $\Lambda^2(M)$, then u has type (2), (2, 1) or (3) on M , or p is odd and u has type (4) on M .*
- (vii) *If u acts non-trivially and minimally actively on $S^2(M)$, then p is odd and u has type (2) on M , or $p \geq 5$ and u has type (3) on M .*

Type	Symmetric square	Exterior square
(2)	(3) ($p \neq 2$), (2, 1) ($p = 2$)	(1) (all p)
(2, 1)	(3, 2, 1) ($p \neq 2$), (2, 2, 1, 1) ($p = 2$)	(2, 1) (all p)
(3)	(5, 1) ($p \neq 2, 3$), (3, 3) ($p = 3$), (4, 2) ($p = 2$)	(3) (all p)
(3, 1)	(5, 3, 1, 1) ($p \neq 2, 3$), (3, 3, 3, 1) ($p = 3$), (4, 3, 2, 1) ($p = 2$)	(3, 3) (all p)
(4)	(7, 3) ($p \neq 2, 5$), (5, 5) ($p = 5$), (4, 4, 2) ($p = 2$)	(5, 1) ($p \neq 2$), (4, 2) ($p = 2$)
(5)	(9, 5, 1) ($p \neq 2, 5$), (5, 5, 5) ($p = 5$), (8, 4, 3) ($p = 2$)	(7, 3) ($p \neq 5$), (5, 5) ($p = 5$)

Table 1: Types of symmetric and exterior squares

Proof. The first three parts are clear. For the fourth part, note that the codimension a of $C_M(u)$ is at most $o(u) - 1$, whence the codimension of $C_M(\langle u, u^{g^2}, \dots, u^{g^r} \rangle)$ is at most $ra \leq r(o(u) - 1)$. If G is generated by r conjugates of u then this is $C_M(G) = 0$, so that $\dim(M) \leq ra \leq r(o(u) - 1)$, as claimed.

For (v), note that if $\dim(M) = 1$ and u acts minimally actively on N then the result is clear, and if p is odd and u acts as a single Jordan block of size 2 on both M and N , then u acts on $M \otimes N$ with type (3, 1), so one direction holds. For the other, if $\dim(M) = \dim(N) = 2$ and $p = 2$ then u acts on $M \otimes N$ with type (2, 2), and otherwise $\dim(N) \geq 3$. If u acts trivially on M then it must act non-trivially on N , and the action of u on $M \otimes N$ contains two copies of the action of u on N (as $\dim(M) \geq 2$) so that u cannot act minimally actively. If u acts non-trivially on M and $\dim(N) \geq 3$, then M contains a u -invariant subspace on which u acts with type (2), and N contains a u -invariant subspace with type either (3) or (2, 1). In the first case, u acts on the tensor product of these subspaces as (4, 2) (or (3, 3) if $p = 3$), and in the second as (3, 2, 1) (or (2, 2, 2) if $p = 2$), so u does not act minimally actively in either case, by applying (i).

For the statements about exterior and symmetric powers, recall that

$$S^2(A \oplus B) = S^2(A) \oplus (A \otimes B) \oplus S^2(B)$$

and similarly for exterior squares. Thus if u has at least three blocks then it contains a submatrix of type (2, 1, 1), and the symmetric and exterior squares of this have two blocks of size 2. From Table 1, we see that u acts minimally on $\Lambda^2(M)$ and $S^2(M)$ when claimed, and that u cannot act minimally on either of these when (3, 1) or (5) is a submatrix of the type of u on M . All other possibilities are in the table above, and this completes the proof. \square

In characteristic 2, we will have to consider modules that are not exterior squares, but exterior squares with one or two trivial composition factors removed.

Lemma 2.4. *Let $p = 2$, let G be a finite group and let M be a faithful, simple module of dimension at least 6. If V is obtained from $\Lambda^2(M)$ by removing at most two trivial composition factors, then V is not minimally active for any non-trivial 2-element of G .*

Proof. The exterior square of a block of size 6 has type (8, 6, 1), so even a submodule of codimension 2 cannot be minimally active for u . Similarly, the exterior square of a matrix of type (4, 1) has type (4, 4, 2), so again we cannot find a minimally active submodule of codimension 2 for u . The exterior square of a matrix of type (3, 1, 1) has type (3, 3, 3, 1) and that of (2, 1⁴) has type (2⁴, 1⁷), so again this cannot work. Every type for u acting on M contains one of these types as a submodule, hence u cannot act minimally actively on V . \square

We now give a lemma on when a power of an element can be minimally active. This uses the classification of groups generated by transvections given in [15].

Lemma 2.5. *Let G be a finite group and let M be a faithful, simple kG -module. Let u be a p -element and suppose that a non-trivial element v of $\langle u^p \rangle$ acts minimally actively on M .*

Then v acts as a transvection on M . Furthermore, G contains a classical group in its natural representation as a normal subgroup, or $p = 2$, k contains \mathbb{F}_4 , and G is either $3 \cdot \text{Alt}_6 \leq \text{GL}_3(k)$ or $3 \cdot \text{PSU}_4(3) \leq \text{GL}_6(k)$.

Proof. The p th power of a single Jordan block of size ap is the sum of p blocks of size a ; from this it is easy to see that the p th power of a single block of size $ap + b$ is the sum of b blocks of size $a + 1$ and $p - b$ blocks of size a . In order for this to be minimally active, we must have $a = b = 1$. Thus u is the sum of one block of size $p^a + 1$ for some a and blocks of size at most p^a , and an element v of order p in $\langle u \rangle$ is a transvection, i.e., has type $(2, 1^{n-2})$ for $n = \dim(M)$.

Since G possesses a faithful simple module in characteristic p , $O_p(G) = 1$. (There are many ways to see this: one is that parabolic subgroups act reducibly in the general linear group, and the normalizer of a p -subgroup is contained in a parabolic.) The subgroup H generated by all conjugates of v is a normal subgroup of G , whence acts semisimply on M as a sum of conjugate modules but also v acts minimally actively, whence H acts irreducibly on M by Lemma 2.3(iii). Thus H is an irreducible subgroup of $\text{GL}(M)$, with $O_p(H) = 1$, containing a transvection, so is one of the groups on Kantor's list in [15, Theorem II].

Of these, we need to check which have a transvection as a proper power of a p -element. Classical groups certainly do (cases (T1) and (T2) in Kantor's list), whereas no 2-element powers to a transposition in Sym_n , (cases (T3) and (T9)), and case (T6) has Sylow p -subgroups of exponent p . Cases (T4) and (T8) are not irreducible, and (T5) and (T7) have a single class of involutions, which must be transpositions, and do not have exponent 2, so are examples. This exhausts the list. \square

If the Sylow p -subgroup of G is cyclic then we can say more about minimally active modules. This next lemma is a generalization of [3, Propositions 3.7 and 3.9], and the proof follows the same method. We do not give all the background on Green correspondence needed for their proof here, and instead refer to [3, Section 3] and the references therein.

Lemma 2.6. *Let G be a finite group and let M be a faithful, simple kG -module. Suppose that the Sylow p -subgroup U of G is cyclic and generated by u .*

- (i) *If $N_G(U)/U$ is abelian (for example, if $C_G(u) = Z(G) \cdot \langle u \rangle$) then u acts minimally actively on M if and only if $\dim(M) \leq o(u) + 1$.*
- (ii) *If $C_G(u)$ is abelian and M is minimally active then $\dim(M) < 2 \cdot o(u)$.*
- (iii) *If M is minimally active then $\dim(M) \leq o(u) + b$, where $b < |C_G(u)|$.*

Proof. By Lemma 2.5 we may assume that if M is minimally active, that it is u that acts minimally actively.

If $\dim(M) = a \cdot o(u)$ for some integer $a \geq 1$, then M is projective and u acts with a blocks of size $o(u)$; thus M is minimally active if and only if $\dim(M) = o(u)$. Hence we can suppose that M is not projective. Let V denote the Green correspondent of M in $N_G(U)$, so that $M \downarrow_{N_G(U)} = V \oplus X$, where V is an indecomposable $kN_G(U)$ -module and X is a relatively $\langle u^p \rangle$ -projective $kN_G(U)$ -module.

We next aim to understand the action of u on V . Note that V is indecomposable and all indecomposable modules for p -soluble groups are uniserial, and all composition factors of V lie in the same block of $N_G(U)$ and have the same dimension m . (This follows since the Brauer tree of a block of a p -soluble group is a star.) Thus the action of u is as m blocks of the same size r , where $\dim(V) = mr$.

Suppose that $X = 0$. Since u acts non-trivially on M , this means that $r > 1$, so that $m = 1$ if and only if u acts minimally actively on M . In particular, if M is minimally active then $\dim(M) < o(u)$. Since m

is the dimension of a simple $N_G(U)/U$ -module, if $N_G(U)/U$ is abelian then this means $m = 1$, so if $X = 0$ then u acts minimally actively on M . This proves all parts of the result when $X = 0$.

Since $X \neq 0$, and clearly u acts non-trivially on X , it must act trivially on V by Lemma 2.3(iii), so V is an indecomposable $kN_G(U)/U$ -module, hence a simple module as this is a p' -group. Since X is relatively $\langle u^p \rangle$ -projective, by Green's indecomposability criterion we see that all Jordan blocks of u on X have size a multiple of p , so there is exactly one, and $\dim(X) \leq o(u)$. This completes the proof of (i) since $\dim(V) = 1$ in this case.

Otherwise we need to bound the dimension of a simple $N_G(U)/U$ -module: since $|N_G(U)/C_G(U)|$ has order dividing $p - 1$, if $C_G(U)$ is abelian then any simple $N_G(U)$ -module has dimension at most $p - 1$: thus $\dim(M) \leq o(u) + (p - 1) < 2o(u)$, as needed. Finally, since $N_G(U)$ -modules are orbits of $C_G(U)$ -modules by Clifford's theorem, they have dimension at most $|C_G(U)| - 1$ (the '-1' is because the trivial is always in a separate orbit) proving the third part. \square

We move on to examining p -elements acting on direct sum and tensor product decompositions.

Lemma 2.7. *Let G be a finite group and let $u \in G$ be a p -element. Suppose that u acts minimally actively on a faithful, simple kG -module M . Suppose that H is a normal subgroup of G and that $G = H\langle u \rangle$. Let $1 < t = |G : H|$ and suppose that $M \downarrow_H$ is the sum of t non-isomorphic simple modules. The actions of u on M , and of u^t on each composition factor of $M \downarrow_H$, is as a single Jordan block, of size the dimension of the module.*

Conversely, if u^t acts on each composition factor of $M \downarrow_H$ as a single Jordan block, then u acts minimally actively on M with a single Jordan block.

Proof. We note at the start that t is a power of p . Since the restriction of M to H is the sum of t non-isomorphic modules, we have the decomposition

$$M \downarrow_H = M_1 \oplus M_2 \oplus \cdots \oplus M_t,$$

where $M_i \cdot u = M_{i+1}$. Suppose that $m = (m_1, \dots, m_t)$ is a fixed point of u , so that in particular each $m_i \in M_i$ is a fixed point of u^t . Note that, since $m \cdot u = m$, we must have that $m_i \cdot u = m_{i+1}$, whence there is a one-to-one correspondence between the u -fixed points of M and the u^t -fixed points of M_1 , and in particular their dimensions are equal.

Writing $d = \dim(M_1)$, so that $\dim(M) = dt$, if u has type $(a, 1^{dt-a})$, then $\dim(M)^{\langle u \rangle} = dt - a + 1$. This has to be equal to $\dim(M_1)^{\langle u^t \rangle}$, which is at most d . This yields

$$dt - a + 1 \leq d.$$

First suppose that $a \leq t$, so that $u^t = 1$. This yields $dt - t + 1 \leq dt - a + 1 = d$, i.e., $(d - 1)(t - 1) \leq 0$, yielding either $d = 1$ or $t = 1$, the latter of which is impossible.

Thus suppose that $u^t \neq 1$. Since u has type $(a, 1^{dt-a})$, we need to know how u^t acts: a block of size a , when raised to the t th power, has type $((\alpha + 1)^\beta, \alpha^{t-\beta})$, where $a = t\alpha + \beta$ and $0 \leq \beta < t$. Thus the action of u^t on M has type

$$((\alpha + 1)^\beta, \alpha^{t-\beta}, 1^{td-a}).$$

These must be distributed equally among the t distinct M_i , whence $t \mid \beta$ and this means that $\beta = 0$. This means that u^t acts on M with type $(t\alpha, 1^{t(d-\alpha)})$, which means that it acts on each M_i with type $(\alpha, 1^{d-\alpha})$.

Thus we now have that

$$1 + t(d - \alpha) = \dim(M)^{\langle u \rangle} = \dim(M_1)^{\langle u^t \rangle} = d - \alpha + 1,$$

so $t = 1$ (again, impossible) or $d = \alpha$, in other words, u^t acts with a single Jordan block, and therefore so does u , as claimed.

For the converse, since u^t fixes a unique 1-space on each M_i , any fixed point of u must lie inside this span. But u^t acts on this t -space as a transitive permutation module, hence fixes a unique 1-space. Thus u acts with a single Jordan block, as claimed. \square

Along with unipotent elements permuting direct sums, we need unipotent elements permuting tensor products.

Lemma 2.8. *Let G be a finite group and let u be a p -element. Let H be a normal subgroup of G such that $G = H\langle u \rangle$, and let M be a faithful, simple kG -module that is not isomorphic to a non-trivial tensor product of two modules, and whose restriction to H factors as a tensor product*

$$M_1 \otimes M_2 \otimes \cdots \otimes M_t,$$

of kH -modules, where $|G : H| = t > 1$ and $\dim(M_i) > 1$. If u acts minimally actively on M , then one of the following holds:

- (i) $p = t = 2$, $\dim(M_i) = 2, 3$, u^2 has type $(2, 2)$ or $(4, 4, 1)$ on M ;
- (ii) $p = t = 3$, $\dim(M_i) = 2$, u^3 has type $(3, 3, 2)$ on M .

Conversely, if $p, t, \dim(M)$ and the action of u^t is as above, then u acts minimally actively on M .

Proof. First suppose that $u^t = 1$. Notice that for any v_1 in M_1 , writing $v_{i+1} = v_i \cdot u$, we can arrange the M_i so that $v_i \in M_i$ and $v_1 \otimes \cdots \otimes v_t$ is fixed by u . The subspace spanned by all other monomials in the tensor product is also fixed by u , so $M \downarrow_{\langle u \rangle}$ is the sum of a trivial module of dimension $\dim(M_i)$ and a permutation module with basis the monomials in the tensor product. Since all other orbits than $v_1 \otimes \cdots \otimes v_t$ have length greater than 1, if u is minimally active then there is a single orbit on the monomials. However, this is clearly impossible, for example since the number of monomials is $\dim(M_i)^t - \dim(M_i) > t$, unless $p = t = \dim(M_i) = 2$.

We therefore may assume that $\langle u \rangle \cap H \neq 1$, so that u^t is a non-trivial p -element. If u acts minimally actively on M then u^t has at most t non-trivial blocks. We will prove that, for almost all possible Jordan normal forms of u^t , there must be more than t non-trivial blocks in its t -fold tensor power. Note that, if this is shown for a block of type $(\alpha_1, \dots, \alpha_r)$, then it is shown for any type $(\beta_1, \dots, \beta_s)$ with $s \geq r$ and $\alpha_i \leq \beta_i$ for all $1 \leq i \leq r$, because for the cyclic group of order a , the kZ_a -module with indecomposable summands of dimensions $\alpha_1, \dots, \alpha_r$ is a submodule of that with dimensions β_1, \dots, β_s , and hence the t -fold tensor power of the former is also a submodule of the latter.

Suppose that $p = 2$. If u^t is a single block of size 2, then the t -fold tensor power of the action of u^t is as 2^{t-1} blocks of size 2, and this is greater than t for $t \geq 3$. Thus if $t \geq 4$ (as t must be a power of 2) then u cannot act minimally actively at all. Thus we may assume that $t = 2$.

If u^t is a block of size 3 then the tensor square has type $(4, 4, 1)$, so is a candidate. If u^t is a block of size 4 then the tensor square has type $(4, 4, 4, 4)$, so we eliminate all blocks of size at least 4, leading to the result in the lemma.

We now check that these two cases occur. For $\dim(M_1) = 2$, we have that u has order 4 and its square acts as $(2, 2)$, so u must have a single block of size 4. For $\dim(M_1) = 3$, u has order 4 and its square acts

as $(4, 4, 1)$, so u must act as $(8, 1)$ (as blocks of sizes 5, 6 and 7 square to have types $(3, 2)$, $(3, 3)$ and $(4, 3)$ respectively). Thus u must be minimally active in these cases.

Now suppose that p is odd, and again we consider the t -fold tensor power of a block V of size 2. For the first few tensor powers, we describe them now. In this table we assume that $t < p$.

t	type
1	(2)
2	(3, 1)
3	(4, 2 ²)
4	(5, 3 ³ , 1 ²)
5	(6, 4 ⁴ , 2 ⁵)
6	(7, 5 ⁵ , 3 ⁹ , 1 ⁵)

These are easily generated as tensoring a block of size i by a block of size 2 yields two blocks, of size $i - 1$ and $i + 1$, at least when $i < p$. It is easy to see that the start of the t -fold product for arbitrary t is $(t + 1, (t - 1)^{t-1}, (t - 3)^{(t-1)(t-2)/2-1}, \dots)$, and so for $t \geq 5$ there are more than t non-trivial blocks of size less than t in the $(t - 1)$ -fold tensor power of a single block. This means that there are more than p non-trivial blocks in the p -fold power of a single block, and at least two of them have size p , for $p \geq 5$. (If $p = 3$, tensoring $(3, 1)$ by (2) yields $(3, 3, 2)$, so the second statement holds but not the first.)

For $t > p$, we write this as a single tensor product of $V^{\otimes p}$ and $V^{\otimes(t-p)}$. The first of these contains two blocks of size p , whose product with $V^{\otimes(t-p)}$ consists entirely of blocks of size p , and hence the product contains at least $2(t - p) > t$ blocks of size p when $t > p$ is a power of an odd p .

This proves that u is not minimally active if $p \geq 5$, or $p = 3$ and $t \geq 9$. If $p = t = 3$ and V is a block of size 3, then $V^{\otimes 3}$ is the sum of nine blocks of size 3, and if V is the sum of a 1- and 2-dimensional module, then $V^{\otimes 3}$ has type $(3^5, 2^2, 1^4)$. Thus if $\dim(M_i) \geq 3$ then we are also done.

We are left with V having dimension 2, in which case $V^{\otimes 3}$ has type $(3, 3, 2)$, as we saw above. If u has order 9 and u^3 acts as $(3, 3, 2)$, then we use the table below that displays the blocks of u^3 , given a block of u .

u	1	2	3	4	5	6	7	8
u^3	1	1 ²	1 ³	2, 1 ²	2 ² , 1	2 ³	3, 2 ²	3 ² , 2

We clearly see that u cannot have blocks of size other than 8, and hence u acts with a single block, as claimed. \square

The next two lemmas are needed in our analysis of minimally active modules for (central extensions of) symmetric groups.

Lemma 2.9. *Let $G = \text{Sym}_n$ for some n , and suppose that $u \in G$ is of order at least 4 and has no cycles of length 1 or 2. Then $\alpha(u) = 2$.*

Proof. Of course, $u = (1, 2, \dots, n)$ and $v = (1, 2)$ generate Sym_n , so $\langle u, u^v \rangle$ has index at most 2 in Sym_n , and we see that $\alpha(u) = 2$ when u is a single cycle.

Suppose that u has cycle type (m_1, \dots, m_r) , with all $m_i \geq 3$, and $m_r \geq 4$. Write $n_0 = 0$, $n_i = \sum_{j=1}^i m_j$, and for $1 \leq i \leq r - 1$, let $\sigma_i = (n_{i-1} - (i - 2), n_{i-1} - (i - 3), \dots, n_i - i, n - i)$, a cycle of length m_i . Finally,

let $\sigma_r = (n_{r-1} - (r - 2), n_{r-1} - (r - 3), \dots, n - r - 1, n, n - r)$, and let u be the product of the σ_i . The second generator is

$$v = (n_1, n - 1)(n_2 - 1, n - 2) \dots (n_{r-1} - (r - 2), n - (r - 1))(n - r, n).$$

Notice that uv is just the $(n - 1)$ -cycle $(1, \dots, n - 1)$, and that

$$[u, v] = (1, n_1 + 1)(n_1, n_2, n - 1)(n_2 - 1, n_3 - 1, n - 2) \dots (n_{r-2} - (r - 3), n_{r-1} - (r - 3), n - (r - 2)) \\ (n - r, n - (r - 1), n_{r-1} - (r - 2), n),$$

so that $[u, v]^6$ is a double transposition. Letting $H = \langle u, v \rangle$, we note that H is transitive and contains an $(n - 1)$ -cycle, hence 2-transitive and so primitive. Since it contains a double transposition, and by [4, Example 3.3.1] a primitive subgroup of Sym_n containing a double transposition contains Alt_n for $n \geq 9$, we get $\alpha(u) = 2$ in this case as well.

The remaining cases to check are for $n = 7, 8$ and u with cycle type $(4, 3)$, $(5, 3)$ and $(4, 4)$. In the first case, Sym_7 is generated by $(1, 2, 3, 4)(5, 6, 7)$ and $(1, 2, 3, 5)(4, 6, 7)$, and in the second and third, Alt_8 is generated by $(1, 2, 3, 4, 5)(6, 7, 8)$ and $(1, 2, 3, 4, 6)(5, 7, 8)$, and also by $(1, 2, 3, 4)(5, 6, 7, 8)$ and $(1, 2, 5, 6)(4, 3, 7, 8)$. \square

Lemma 2.10. *Let $G = \text{Sym}_n$ for some $n \geq 9$. If $u \in G$ has cycle type $(n - 2, 2)$, $(n - 4, 2, 2)$ or $(n - 6, 2, 2, 2)$, then $\alpha(u) = 2$. If $n = 10$ and u has cycle type $(4, 4, 2)$ then $\alpha(u) = 2$ also.*

Proof. In the first case, let $u = (1, n - 1)(2, 3, \dots, n - 3, n, n - 2)$ and $v = (2, n - 1)(n - 2, n)$. Again, $uv = (1, 2, \dots, n - 1)$ and the same proof applies as Lemma 2.9, as v is itself a double transposition.

In the second case, let $u = (1, n - 1)(2, n - 2)(3, \dots, n - 4, n, n - 3)$ and $v = (2, n - 1)(3, n - 2)(n - 3, n)$. Then uv is as in the previous case, but now we need to find an element of small support, and this is

$$[u, v] = (1, 3, n, n - 3, n - 2)(2, 4, n - 1),$$

and so $[u, v]^5$ is a 3-cycle, and we are done.

Finally, let $u = (1, n - 1)(2, n - 2)(3, n - 3)(4, \dots, n - 5, n, n - 4)$ and $v = (2, n - 1)(3, n - 2)(4, n - 3)(n - 4, n)$. Again, uv is as before, but if $n \geq 10$ then

$$[u, v] = (1, 3, 5, n - 2)(2, 4, n, n - 4, n - 3, n - 1),$$

and $[u, v]^6 = (1, 5)(3, n - 2)$, as needed. If $n = 9$ then u^2 is a 3-cycle, and we are again done.

Finally we simply give generators of Sym_{10} of the appropriate cycle types:

$$(1, 2, 3, 4)(5, 6, 7, 8)(9, 10) \quad \text{and} \quad (1, 5, 4, 7)(2, 3, 8, 10)(6, 9).$$

This completes the proof. \square

We also need to determine better bounds on $\alpha(u)$ for u a unipotent element in $\text{GL}_n(2)$ than $\alpha(u) \leq n$ given in [8]. While this bound is sharp for transvections, simply by considering the fixed-point subspace, we need elements close to regular elements. Indeed, by [6], with the exception of $\text{SL}_4(2)$, $\text{SL}_n(p)$ is generated by two regular unipotent elements for all primes p , and all $n \geq 3$. We give this in a lemma for reference.

Lemma 2.11 ([6]). *Let $G = \text{SL}_n(p)$ for some $n \geq 2$. If u is a regular unipotent element of G then $\alpha(u) = 2$.*

Proof. This is proved in [6] for all cases except for $\text{SL}_4(2) = \text{Alt}_8$, where the regular unipotent class is in bijection with the class containing $u = (1, 2, 3, 4)(5, 6)$. Letting $v = (1, 5, 7, 8)(4, 6)$, we note that $\langle u, v \rangle$ generates a primitive subgroup of Alt_n containing $(uv^2)^5 = (5, 8, 6)$. This completes the proof. \square

From this, we can get that if u is a 2-element of maximal order in $\mathrm{SL}_n(2)$, then $\alpha(u) \leq 4$; it is likely that this could be improved still further, but not without considerably more work.

Lemma 2.12. *Let $G = \mathrm{SL}_n(2)$, and let u be a unipotent element of maximal order in G . If n is even then $\alpha(u) \leq 3$ and if n is odd then $\alpha(u) \leq 4$.*

Proof. Write V for the natural module for G . Suppose that u has type $(a, 1^{n-a})$ on V for some $a \geq n/2 + 1$, so that in particular $C_V(u)$ has dimension less than half of $\dim(V)$. Let u_1 and u_2 be regular unipotent generators of $\mathrm{SL}_a(2)$, written as matrices in G , so with type $(a, 1^{n-a})$. The action of $H_1 = \langle u_1, u_2 \rangle$ on V has a single simple submodule W of dimension a , and all other simple submodules trivial.

Write v_1, \dots, v_a for a basis of W , and extend the basis to v_{a+1}, \dots, v_n on which H_1 acts trivially. Let u_3 act as follows:

$$v_i \cdot u_3 = \begin{cases} v_1 + v_n & i = 1, \\ v_i & 2 \leq i \leq n - a + 1, \\ v_i + v_{i-1} & n - a + 2 \leq i \leq n. \end{cases}$$

Of course, u_3 has the correct type. We claim that $H = \langle u_1, u_2, u_3 \rangle$ acts irreducibly on V , so let X denote an H -submodule of V . Since W is a simple H_1 -submodule, either $X \cap W = 0$ or $W \leq X$: if $W \leq X$ then $v_1 \in X$, so $v_n \in X$ and we see that each $v_i \in X$, so that $V = X$. Thus $X \cap W = 0$, and so H_1 acts trivially on X , yielding X is a subspace of $\langle v_{a+1}, \dots, v_n \rangle$. However, repeated application of u_3 to any element of this space eventually leaves it, as we must project onto v_a , so that $X = 0$. Thus H is irreducible on V , containing a copy of $\mathrm{SL}_a(2)$ acting on V in a non-self-dual way, hence $H \not\leq \mathrm{Sp}(V)$. Since H contains a transvection, we can apply [15, Theorem II]: either $H = G$, H is a classical group (all contained in $\mathrm{Sp}(V)$ as the characteristic is 2) or a symmetric group (again, contained in $\mathrm{Sp}(V)$ as the simple modules are self-dual), so since $H \not\leq \mathrm{Sp}(V)$ we have that $H = G$, as needed.

If $n - a + 1 = a$, i.e., $n = 2a + 1$, then the above argument fails: in this case, generate $\mathrm{SL}_{n-1}(2)$ with three elements, and use the fourth to get the full $\mathrm{SL}_n(2)$.

If the u_i have another type, with a single block of size a and various smaller blocks instead, then choose the u_i exactly as before: note that every subspace of V stabilized by u_i is also stabilized by the previous u_i , and so since we had an irreducible subgroup before we must have an irreducible subgroup again. Since it still contains a transposition, we still have $\mathrm{SL}_n(2)$, as needed. \square

We end this section by giving the notation used for almost quasisimple groups. Our groups G will have the property that $G = \langle F^*(G), u \rangle$ for some p -element u , that $G_0 = F^*(G)$ is quasisimple and that $Z(G) = Z(G_0)$. If M is a faithful simple kG -module then this yields an embedding of G into $\mathrm{GL}(M)$. Our conditions on G are equivalent to the image H of G in $\mathrm{PGL}(M)$ being almost simple, and H' being simple with H/H' generated by a p -element of H .

3 Reduction to almost simple groups

This section uses Aschbacher's classification of maximal subgroups of classical groups [1] (see also [17, 23, 31], and in particular [17], which modifies the classes of Aschbacher, and whose notation we will use here) to reduce to the case given at the end of the last section, where G is an almost quasisimple group. Thus we have eight classes $\mathcal{C}_1, \dots, \mathcal{C}_8$ of maximal subgroups, together with almost quasisimple groups \mathcal{S} . We will determine which elements of the \mathcal{C}_i contain minimally active elements.

We assume in this section that G is a subgroup of $\mathrm{GL}(M)$ for an n -dimensional k -vector space M , with $u \in G$ being a p -element acting minimally actively on M . As we are only concerned with irreducible modules, we stipulate that G acts irreducibly on M . In particular, G cannot lie in a parabolic subgroup, class \mathcal{C}_1 .

If G acts imprimitively on M then G stabilizes a direct sum decomposition

$$M = M_1 \oplus M_2 \oplus \cdots \oplus M_t$$

of M . Taking this decomposition to be as fine as possible, we see that G is a subgroup of $\mathrm{GL}_{n/t}(k) \wr \mathrm{Sym}_t$ (Aschbacher's class \mathcal{C}_2). The action of u on this group is in Lemma 2.7: u must act with a single non-trivial block, t is a power of p , and u^t acts with a single block on each M_i . Hence G lies in a wreath product $A \wr B$ where $u^t \in A$ acts on M_1 with a single Jordan block, and the t -cycle $(1, \dots, t)$ lies in B . Furthermore, given such a setup we always obtain an element u . Thus there are many groups acting imprimitively on M with minimally active elements.

If we extend the field and the module is no longer irreducible (G is contained in an extension field subgroup, class \mathcal{C}_3), then we can apply Lemma 2.3(iii) to see that G cannot have a minimally active element.

Suppose that G acts primitively and absolutely irreducibly on M , and preserves a tensor decomposition

$$M = M_1 \otimes M_2 \otimes \cdots \otimes M_t,$$

so that G is a subgroup of $\mathrm{GL}_m(k) \wr \mathrm{Sym}_t$ with $n = m^t$. Now we can apply Lemmas 2.3(v) and 2.8, which show that one of the following holds:

- (i) $n = 4$ and p is odd, with G a subgroup of $\mathrm{SL}_2(k) \times \mathrm{SL}_2(k)$ and $o(u) = p$ acting with type $(3, 1)$;
- (ii) $n = 4$ and $p = 2$, with $o(u) = 4$ acting with type (4) ;
- (iii) $n = 8$ and $p = 3$, with $o(u) = 9$ acting with type (8) ;
- (iv) $n = 9$ and $p = 2$, with $o(u) = 8$ acting with type $(8, 1)$.

We also showed in that lemma that these cases occur, and we will not comment further on this case. These are classes \mathcal{C}_4 and \mathcal{C}_7 .

If G is contained inside a subfield subgroup in \mathcal{C}_5 (of the form $N_{\mathrm{GL}_n(k)}(\mathrm{GL}_n(k_0))$), then since our property is independent of the field over which we take our module M , we replace k by k_0 and so this case can be ignored.

If G is contained inside another classical group (i.e., class \mathcal{C}_8), then we apply this classification of maximal subgroups to that group instead.

Thus we are left with \mathcal{C}_6 , extraspecial type subgroups, and \mathcal{S} , which is the focus of all subsequent sections of the paper.

Let $r \neq p$ be a prime, and let R denote an extraspecial group of order r^{1+2m} for some m . If $r = 2$, we allow R to be either an extraspecial group or the central product with Z_4 . Note that a faithful, irreducible representation of R in characteristic not r has dimension $n = r^m$, so we may embed R into $\mathrm{GL}_n(k)$ for k an algebraically closed field of characteristic p , and k any field of characteristic p except when $R = Z_4 \circ r^{1+2m}$ and k needs a fourth root of unity. Let G denote the normalizer in $\mathrm{GL}_n(k)$ of R , and let M be the natural module for G . More information about G can be found in [17, 23, 31].

We want to prove that if G contains a minimally active element then the parameters r, p, m are very tightly controlled. To do so, we need to know something about G/R , which is a classical group. In Section 7, particularly Proposition 7.3, we get information about the orders of p -elements of classical groups in characteristic different from p . Rather than deferring the proof of this result until then, we include it here, but use the definitions and notation from that section. The reader is recommended to skip the proof of this result until they have reached Section 7; the proof is similar to those contained in Section 8.

Proposition 3.1. *If $u \in G$ acts minimally actively on M , then $r = 2$ and one of the following holds:*

- (i) m is an odd prime, $p = 2^m - 1$ is a Mersenne prime, and u has order p , acting with type $(p, 1)$;
- (ii) m is a power of 2, $p = 2^m + 1$ is a Fermat prime, and u has order p , acting with type $(p - 1)$;
- (iii) $m = 2$, $p = 3$, $o(u) = 3$ acting with type $(3, 1)$ (not all elements of order 3 have this property);
- (iv) $m = 3$, $p = 3$, $o(u) = 9$, acting with type (8) .

Proof. If r is odd, then $O^{p'}(G) = R \rtimes \mathrm{Sp}_{2m}(r)$ is a split extension, and so if u is a p -element of G then we may assume that u lies in $H = \mathrm{Sp}_{2m}(r)$. By [7, Section 5], if p is odd then H acts on M as the direct sum of the two Weil modules, of dimensions $(r^m - 1)/2$ and $(r^m + 1)/2$, and so u cannot act minimally actively on M by Lemma 2.3(iii). If $p = 2$ then M is uniserial of length 3 [7, Lemma 5.2] with socle series W, k and W , where W is a Weil module, and since W is not minimally active for $p = 2$ by Theorem 8.1 we eliminate this case as well.

Thus $r = 2$ and hence p is odd. Here $G/R \cong H$ where $H = \mathrm{GO}_{2m}^{\pm}(2)$ or $\mathrm{Sp}_{2m}(2)$, and the orthogonal-type groups are contained in $R.\mathrm{Sp}_{2m}(2)$, so we will work solely with that group. Let d denote the order of 2 modulo p , so that $p \mid \Phi_d(2)$. We have that $\alpha(u) \leq m + 3$ in all cases, since p is odd.

Suppose that the image in H of u lies in a Levi subgroup of H , say $\mathrm{Sp}_{2m-2a}(2) \times \mathrm{Sp}_{2a}(2)$. Taking preimages in G yields a central product $G_1 \circ G_2$, where $G_1 = R_1.H_1 = (4 \circ 2^{2(n-a)+1}).\mathrm{Sp}_{2(m-a)}(2)$ and $G_2 = R_2.H_2 = (4 \circ 2^{2a+1}).\mathrm{Sp}_{2a}(2)$. The action of this group on M is a tensor product of actions of the G_i , and since the tensor product of two modules cannot be minimally active unless they both have dimension at most 2 by Lemma 2.3(v), we have that $m = 2$ and $p = 3$. This case will be considered later.

Thus we may assume that the Sylow p -subgroup of G , and hence H , does not lie in any proper Levi subgroup of H . This in particular means that d divides $2m$.

Suppose that d is either $2m$, or m is odd and $d = m$; in both cases d is regular, so $C_H(u)$ is abelian and of odd order since the image of u is semisimple in H and therefore $C_H(u)$ is reductive. Thus $C_G(u)$ splits as the direct product of $C_R(u)$ and a subgroup C isomorphic to $C_H(u)$. Thus $R = C_R(u) \circ [R, u]$, and since u does not lie in a Levi subgroup of H , $C_R(u) = Z(R) \leq Z(\mathrm{GL}(M))$. This shows that $C_G(u) = Z(R) \times C$, and in particular $C_G(u)$ is abelian, so that $2^m = \dim(M) \leq 2o(u)$ if u acts minimally actively on M by Lemma 2.6(ii). As $o(u) \mid \Phi_d(2) \mid (2^m \pm 1)$, if $(2^m \pm 1)$ is not a prime power then $2o(u) \leq 2(2^m \pm 1)/3 < 2^m$, and so $o(u) = 2^m \pm 1$. Thus $2^m \pm 1 = \Phi_d(2)$, so that m is either a power of 2 or is a prime, and $2^m \pm 1$ is a Fermat or Mersenne prime, or is $9 = 2^3 + 1$, but 2 has order 2 modulo 3, not 6.

In these cases, $C_H(u)$ is abelian and indeed is simply $\langle u \rangle$, so $C_G(u) = Z(R) \times \langle u \rangle$ is cyclic, and we may apply Lemma 2.6 again to show that u acts minimally actively on M , and we are done.

If $d = m$ for m even, then the Sylow p -subgroup of H has rank 2, and lies inside the Levi subgroup $\mathrm{Sp}_m(2) \times \mathrm{Sp}_m(2)$, so u can only act minimally actively if $m = 1$, i.e., $p = 3$, as we saw above.

Suppose that the Sylow p -subgroup of G is abelian, so that $o(u)$ is a divisor of $\Phi_d(2)$, for $d \mid 2m$ with $d \neq m, 2m$. If d is odd then $d \leq m/2$, and if d is even then $d \leq 2m/3$: in the first case, $o(u) \leq (2^{m/2} - 1)$ and in the second $o(u) \leq (2^{m/3} + 1)$. Since $\alpha(u) \leq m + 3$ and $\dim(M) = 2^m$, we get

$$(m + 3) \cdot (o(u) - 1) \leq 2^m,$$

which has solutions only for $m \leq 4$: noting that $d \neq 1$ and $d \mid 2m$, we get $d = 2$ for $m = 3, 4$. In this case $p \mid \Phi_2(2) = 3$ and the Sylow 3-subgroups of $\mathrm{Sp}_6(2)$ and $\mathrm{Sp}_8(2)$ are non-abelian, so there are no solutions.

If the Sylow p -subgroup of G is non-abelian then d divides m and also pd is at most m if d is odd and $2m$ if d is even. Suppose that $p \geq 5$: then $d \geq 3$, and if $d = 3$ then $p = 7$, if $d = 4$ then $p = 5$, and if $d \geq 5$ then $p \geq 11$. If $p = 5$ then $m \geq 10$, and if $p \geq 7$ then $m \geq 21$. We have $d \leq 2m/5$, and $o(u) \leq m(2^{2m/5} - 1)$, where the Weyl contribution is m and the toral contribution is of course $2^{2m/5} - 1$. Thus the inequality $\alpha(u) \cdot o(u) > \dim(M)$ for u to act minimally actively becomes

$$m(m + 3)(2^{2m/5} - 1) \geq 2^m$$

for $m \geq 10$, and the only solutions are for $m = 10, 11, 12$, so we must have $p = 5$. In this case $o(u) \leq 25$ so we get $25(m + 3) \leq 2^m$ for $m \geq 10$, and this obviously has no solutions.

We therefore have $p = 3$ so $d = 2$, and $o(u) \leq 3m$. This yields the inequality

$$(3m - 1)(m + 3) \geq 2^m,$$

which is satisfied for $m \leq 7$. The Sylow 3-subgroup of $\mathrm{Sp}_{2m}(2)$ lies inside the Levi subgroup $\mathrm{Sp}_6(2) \times \mathrm{Sp}_{2m-6}(2)$ for $4 \leq m \leq 7$, so these cases need not be considered.

We now collect together the cases we need to check, which are only $m = 2, 3$ for $p = 3$. For $(Z_4 \circ 2^{1+4}).\mathrm{Sp}_4(2)$, we have that u is contained in $(Z_4 \circ 2^{1+4}).(\mathrm{Sp}_2(2) \times \mathrm{Sp}_2(2))$, so the action of u on M can be factored as the tensor product of two matrices. If u lies in one of the factors then this would have type (2^2) , but if it is diagonal then it would act as $2 \otimes 2$, so type $(3, 1)$, minimally active.

For $(Z_4 \circ 2^{1+6}).\mathrm{Sp}_6(2) \leq \mathrm{SL}_8(9)$, if $o(u) = 3$ then $\alpha(u) \leq 4$ (and hence u cannot act minimally actively) unless u lies inside a Levi subgroup Sp_2 , which of course means that u is not minimally active (as u cannot lie inside such a Levi).

Thus $o(u) = 9$: inside $Z_3 \wr Z_3 \leq \mathrm{Sym}_9$, elements of order 9 square to the fixed-point-free class of elements of order 3, and so inside $\mathrm{Sp}_2(2) \wr \mathrm{Sym}_3 \leq \mathrm{Sp}_6(2)$ we see that elements of order 9 power to the class that lies diagonally across all three $\mathrm{Sp}_2(2)$ factors. Therefore the action of u^3 on M has type $(3, 3, 2)$, as this is the type of the third tensor power of a block of size 2. As blocks of size 5, 6, 7 and 8 power to have types $(2, 2, 1)$, $(2, 2, 2)$, $(3, 2, 2)$ and $(3, 3, 2)$ respectively, we see that u must have a single block of size 8, hence is minimally active, as needed. \square

Thus we have considered all of the \mathcal{C}_i , and so we may assume that G is a member of \mathcal{S} . Furthermore, this completes the proof of Theorem 1.1.

4 Alternating groups

In this section we use the notation at the very end of Section 2: let G_0 be a central extension of an alternating group Alt_n for $n \geq 5$, let u be a p -element of G such that $G = \langle G_0, u \rangle$. (Note that, since $\mathrm{Out}(G_0)$ is a 2-group, we have that $u \in G_0$ unless $p = 2$.) When $n = 5, 6, 7, 8$ we get very different answers to the general

case: for $n = 5$ this is because Alt_5 is isomorphic to $\text{SL}_2(4)$ and $\text{PSL}_2(5)$; for Alt_6 it is because of the extra outer automorphism, the exceptional triple cover, and the isomorphism with $\text{PSL}_2(9)$; for Alt_7 , it is the exceptional triple cover; and for Alt_8 it is the isomorphism with $\text{SL}_4(2)$.

Because of this, the first proposition deals with those four individual groups. Because of their small order, one can check all calculations easily on a computer, and we just say a few words about its proof. In this proposition, cases (ii) and (iii) are written as if they are general statements, which they are, even though there is only one instance of each in the range $5 \leq n \leq 8$.

Proposition 4.1. *Let G_0 be a central extension of Alt_n for $5 \leq n \leq 8$, and let u be a p -element of G such that $G = \langle G_0, u \rangle$. Let V denote the non-trivial composition factor of the permutation module for G . If u acts minimally actively on a non-trivial simple module M , then (up to outer automorphism in the case $n = 6$) one of the following holds:*

- (i) $G = \text{Alt}_n$ for p odd and $G = \text{Sym}_n$ for $p = 2$, u is a single cycle of length p^a for some $p^a \leq n$ and $M = V$;
- (ii) $G = \text{Alt}_n$ for $n = 2^a + 2$, $a \geq 2$, $p = 2$, u has cycle type $(2^a, 2)$ and $M = V$;
- (iii) $G = \text{Sym}_n$ for $n = 2^a + 4$ with $a \geq 2$, $p = 2$, u has cycle type $(2^a, 2, 2)$ and $M = V$, with u acting as $(2^a, 1, 1)$;
- (iv) $G = \text{Alt}_5$, $p = 2$, $o(u) = 2$ and $\dim(M) = 2$;
- (v) $G = \text{Sym}_5$, $p = 2$, $o(u) = 4$ and $\dim(M) = 4$;
- (vi) $G = \text{Alt}_5$, $p = 3$, $o(u) = 3$ and $\dim(M) = 3$;
- (vii) $G = 2 \cdot \text{Alt}_5$, $p = 3$, $o(u) = 3$ and $\dim(M) = 2$;
- (viii) $G = \text{Alt}_5$, $p = 5$, $o(u) = 5$ and $\dim(M) = 5$;
- (ix) $G = 2 \cdot \text{Alt}_5$, $p = 5$, $o(u) = 5$ and $\dim(M) = 2, 4$;
- (x) $G = \text{PGL}_2(9)$ or $G = M_{10}$, $p = 2$, $o(u) = 8$ and $\dim(M) = 8$ (three and one representation respectively);
- (xi) $G = 3 \cdot M_{10}$, $p = 2$, $o(u) = 8$ and $\dim(M) = 6, 9$;
- (xii) $G = 3 \cdot \text{Alt}_6$, $p = 2$, $o(u) = 2, 4$ and $\dim(M) = 3$ (four representations);
- (xiii) $G = \text{Alt}_6$, $p = 3$, $o(u) = 3$ and $\dim(M) = 3$;
- (xiv) $G = 2 \cdot \text{Alt}_6$, $p = 3$, $o(u) = 3$ and $\dim(M) = 2$ (two representations);
- (xv) $G = 2 \cdot \text{Alt}_6$, $p = 5$, $o(u) = 5$ and $\dim(M) = 4$ (two representations);
- (xvi) $G = 3 \cdot \text{Alt}_6$, $p = 5$, $o(u) = 5$ and $\dim(M) = 3, 6$ (two representations each);
- (xvii) $G = 6 \cdot \text{Alt}_6$, $p = 5$, $o(u) = 5$ and $\dim(M) = 6$ (four representations);
- (xviii) $G = \text{Alt}_7$, $p = 2$, $u = (1, 2, 3, 4)(5, 6)$ and $\dim(M) = 4$;
- (xix) $G = 2 \cdot \text{Alt}_7$, $p = 3$, u is the preimage of $(1, 2, 3)(4, 5, 6)$ and $\dim(M) = 4$;

- (xx) $G = 2 \cdot \text{Alt}_7$, $p = 5$, $o(u) = 5$ and $\dim(M) = 4$;
- (xxi) $G = 3 \cdot \text{Alt}_7$, $p = 5$, $o(u) = 5$ and $\dim(M) = 3, 6$ (two representations each);
- (xxii) $G = 6 \cdot \text{Alt}_7$, $p = 5$, $o(u) = 5$ and $\dim(M) = 6$ (four representations);
- (xxiii) $G = 2 \cdot \text{Alt}_7$, $p = 7$, $o(u) = 7$ and $\dim(M) = 4$;
- (xxiv) $G = 3 \cdot \text{Alt}_7$, $p = 7$, $o(u) = 7$ and $\dim(M) = 6$ (two representations);
- (xxv) $G = 6 \cdot \text{Alt}_7$, $p = 7$, $o(u) = 7$ and $\dim(M) = 6$ (four representations);
- (xxvi) $G = \text{Alt}_8$, $p = 2$, $u = (1, 2)(3, 4)(5, 6)(7, 8)$ and $\dim(M) = 4$;
- (xxvii) $G = \text{Alt}_8$, $p = 2$, $u = (1, 2, 3, 4)(5, 6, 7, 8)$ and $\dim(M) = 4$, with u acting as $(3, 1)$;
- (xxviii) $G = \text{Alt}_8$, $p = 2$, $u = (1, 2, 3, 4)(5, 6)$ and $\dim(M) = 4$;
- (xxix) $G = 2 \cdot \text{Alt}_8$, $p = 7$, $o(u) = 7$ and $\dim(M) = 8$.

In all cases, unless otherwise specified u acts on M with type $(\dim(M))$ if $\dim(M) \leq o(u)$, and as $(o(u), 1^{\dim(M)-o(u)})$ otherwise.

Proof. For $n = 5$, when p is odd we simply check all simple modules for $2 \cdot \text{Alt}_5$, and for $p = 2$ we check all simple modules for Alt_5 and Sym_5 .

Next, we deal with $n = 6$. For $p = 5$ we check all simple modules for $6 \cdot \text{Alt}_6$, and since $N_G(u)/\langle u \rangle$ is cyclic we only need to know which have dimension at most 6 by Lemma 2.6, so we get what is above.

For $p = 3$ we check all simple modules for $2 \cdot \text{Alt}_6 = \text{SL}_2(9)$, and the answer will be the same as for $\text{SL}_2(q)$ in defining characteristic.

For $p = 2$, here we have Alt_6 , the three extensions of Alt_6 by an outer automorphism, so Sym_6 , $\text{PSL}_2(9)$ and M_{10} , the central extension $3 \cdot \text{Alt}_6$, and the last group $3.M_{10}$, since the M_{10} outer automorphism is the only one preserving the centre of $3 \cdot \text{Alt}_6$. This group is not necessarily well defined, so we give more details now.

Let G be a group of the form $3 \cdot M_{10}$. By a quick computer calculation, G is generated by two conjugates of u for $o(u) = 4, 8$ and by three if $o(u) = 2$, by checking this is true for M_{10} .

The only faithful simple modules for G have dimensions 6 and 9, by [14], with the 6 restricting to $3 \cdot \text{Alt}_6$ as the sum of two non-isomorphic 3-dimensional simple modules. Thus here we are in the situation of Lemma 2.7, and u^2 must act with a single Jordan block of these 3s. This means that $o(u^2) = 4$ and so $o(u) = 8$. Furthermore, the only action of u that squares to Jordan block structure $(3, 3)$ is (6) , so that u does indeed act minimally actively on M of dimension 6.

For the module M of dimension 9, this restricts simply to $3 \cdot \text{Alt}_6$, and only u of order 8 could act minimally actively on M . This time, we are resigned to constructing the normalizer inside $\text{GL}_9(4)$ of $3 \cdot \text{Alt}_6$ and simply computing the action of these elements of order 8, and they do act as $(8, 1)$ on the two dual 9-dimensional simple modules.

For G_0 a central extension of Alt_7 , for $p = 5, 7$, $C_G(u) = \langle u \rangle Z(G_0)$ and so we only need $\dim(M) \leq p + 1$ by Lemma 2.6. For $p = 3$, we find the two simple modules for $2 \cdot \text{Alt}_7$ which are minimally active for the conjugacy class of elements of order 3 that have the smallest centralizer. For $p = 2$, the outer automorphism inverts the centre of $3 \cdot \text{Alt}_7$, so we only need concern ourselves with G one of Alt_7 , Sym_7 and $3 \cdot \text{Alt}_7$, all of which are easily constructible.

For Alt_8 , se simply need to consider Alt_8 and $2 \cdot \text{Alt}_8$ for p odd, and Alt_8 and Sym_8 for $p = 2$, which is easy to do directly. \square

From now on we let G_0 be a central extension of an alternating group Alt_n for some $n \geq 9$. We first need to decide which elements of Sym_n act minimally actively on the non-trivial composition factor of the permutation module. The next lemma does this.

Lemma 4.2. *Let $G = \text{Sym}_n$, and let V denote the non-trivial composition factor of the permutation module for G . If u acts minimally actively on V , then one of the following holds:*

- (i) u is a single cycle of length p^a for some $p^a \leq n$, acts on V with type $(p^a - 2)$ or $(o(u), 1^{n-o(u)})$;
- (ii) $n = 2^a + 2$, $a \geq 2$, $p = 2$, u has cycle type $(2^a, 2)$ and acts on V with type $(n - 2) = (2^a)$;
- (iii) $n = 2^a + 4$ with $a \geq 2$, $p = 2$, u has cycle type $(2^a, 2, 2)$ and acts on V with type $(n - 2, 1^2)$;
- (iv) $n = 3^a + 3$ with $a \geq 2$, $p = 3$, u has cycle type $(3^a, 3)$ and acts on V with type $(n - 1, 1)$.

Proof. Let M denote the permutation module for G . Note that V is obtained from M by removing either a single trivial summand if $p \nmid n$, or removing a trivial submodule and a trivial quotient if $p \mid n$.

Note that the action of u on M has type the cycle type of u , so if u has more than one cycle of length at least 4, more than two cycles of length at least 3, or more than three non-trivial cycles, then u cannot act minimally actively on V , as V is obtained from M by removing at most two trivials.

Thus we are left with checking the types $(m, 1^{n-m})$, $(m, 2, 1^{n-m-2})$, $(m, 2^2, 1^{n-m-4})$ and $(m, 3, 1^{n-m-3})$, with p any prime, 2, 2 and 3 respectively.

First suppose that u fixes a point, so that $u \in \text{Sym}_{n-1}$. The restriction of V to Sym_{n-1} is either simple if $p \nmid n$, or isomorphic to the permutation module on Sym_{n-1} if $p \mid n$. From this we can use induction to easily see that if u fixes a point and is minimally active then u acts like a single cycle, and the type of u on V is as above. Thus it remains to check cycles types of the form $(m, 2)$, $(m, 2, 2)$ and $(m, 3)$, for $m = 2^a$, 2^a and 3^a respectively.

If u has cycle type (p^a, p) for $p = 2, 3$ then u^p fixes a point, whence its action on V is known from the above working to have p blocks of size p^{a-1} and $p-2$ blocks of size 1. Since the action of u on the permutation module has Jordan blocks one of size p^a and one of size p , and the action of u on V is a subquotient of this, it must be that u acts with one block of size p^a and $p-2$ of size 1, as needed.

We now need to consider u of type $(2^a, 2, 2)$, which lies inside the subgroup $H = \text{Sym}_{n-4} \times \text{Sym}_4$. The permutation module for H is simply the direct sum of the permutation modules for Sym_{n-4} and Sym_4 , and has as a subquotient of codimension 4 a semisimple module obtained by removing all four trivials. The action of u on the 2-dimensional simple subquotient of the second summand is trivial, since $(1, 2)(3, 4)$ lies in the kernel of every simple module for Sym_4 . Thus u acts on this semisimple module with blocks $(m-2, 1, 1)$, where $m = 2^a$, with this semisimple module being itself a subquotient of V . As u^2 acts on V with blocks $(m/2, m/2, 1, 1, 1, 1)$, we see that the only possibility for the action of u consistent with both piece of information is that it has type $(m, 1, 1)$. \square

Proposition 4.3. *Let G_0 be a central extension of Alt_n for $n \geq 9$, and let u be a p -element of G with $G = \langle G_0, u \rangle$. Let V denote the non-trivial composition factor of the permutation module for G . If u acts minimally actively on a non-trivial simple module M , then (up to outer automorphism in the case $n = 6$) one of the following holds:*

- (i) $G = \text{Alt}_n$ for p odd and $G = \text{Sym}_n$ for $p = 2$, u is a single cycle of length p^a for some $p^a \leq n$ and $M = V$, and u acts with type (n) or $(o(u), 1^{n-o(u)})$;
- (ii) $G = \text{Alt}_n$ for $n = 2^a + 2$, $a \geq 2$, $p = 2$, u has cycle type $(2^a, 2)$ and $M = V$, acting with type (2^a) ;
- (iii) $G = \text{Sym}_n$ for $n = 2^a + 4$ with $a \geq 2$, $p = 2$, u has cycle type $(2^a, 2, 2)$ and $M = V$, acting with type $(n - 2, 1, 1)$;
- (iv) $G = \text{Alt}_n$ for $n = 3^a + 3$ with $a \geq 2$, $p = 3$, u has cycle type $(3^a, 3)$ and $M = V$, acting with type $(n - 1, 1)$;
- (v) $G = 2 \cdot \text{Alt}_9$, $p = 3$, $o(u) = 9$ and $\dim(M) = 8$, acting with type $(7, 1)$;
- (vi) $G = 2 \cdot \text{Alt}_9$, $p = 7$, $o(u) = 7$ and $\dim(M) = 8$, acting with type $(7, 1)$;

Proof. Case 1: p odd We first assume that $G_0 = \text{Alt}_n$. We start by checking that there are no simple modules for G_0 , other than the trivial module and V , that have dimension at most $2n - 2$. By [12, Theorem 7 and Table 1] for $n \geq 12$ we have that this holds, and the dimensions of simple modules are known for $n \leq 11$, so we can check that this holds.

We can do the same thing for $G_0 = 2 \cdot \text{Alt}_n$: by [18] for $n \geq 12$ we have that all faithful representations of G_0 are of dimension greater than $2n - 2$. Thus we need to check $9 \leq n \leq 11$: for all odd primes the minimal degrees are 8, 16 and 16, with all other faithful modules have dimension larger than $2n - 2$, unless $n = 10$ and $p = 5$, in which case the minimal degree is 8.

For $p = 11$ this cannot yield a minimally active module as $C_G(u) = \langle u \rangle \cdot Z(G)$ and so $\dim(M) \leq p + 1 = 12$ by Lemma 2.6.

For $p = 7$, since $\alpha(u) = 2$ we eliminate $n = 10, 11$, and for $n = 9$ we see that $C_G(u) = \langle u \rangle \cdot Z(G)$ and so u acts minimally actively on M if and only if $\dim(M) \leq p + 1 = 8$ by Lemma 2.6 again. Since $\dim(M) = 8$, this means they are minimally active.

For $p = 5$, we check that u acts on the 8-dimensional simple modules for $2 \cdot \text{Alt}_9$ as $(4, 4)$, so not minimally active, and for $2 \cdot \text{Alt}_{10}$ both classes act as $(4, 4)$, so again no minimally active faithful modules, hence none for $2 \cdot \text{Alt}_{11}$ either by restriction.

Finally, for $p = 3$ we have $o(u) = 9$. For Alt_9 the 8-dimensional module is minimally active, with action $(7, 1)$, but for $n = 10, 11$ the action on the 16-dimensionals has blocks $(9, 7)$, so not minimally active.

Thus if u is a p -element such that $\alpha(u) = 2$, the only simple modules on which u acts minimally actively are the trivial and V , unless $n = 9$ and $G_0 = 2 \cdot \text{Alt}_9$. Note that if $o(u) = 3$ then since $\alpha(G) \leq n/2$ by [8, Lemma 6.1], we have that $\dim(M) \leq 3n/2$ by Lemma 2.3(iv), so M is either trivial or V ; hence we will assume that $o(u) \geq 5$.

Let $G_0 = \text{Alt}_n$ or $G_0 = 2 \cdot \text{Alt}_n$ for some $n \geq 9$. By Lemma 2.9, if the image of u in Alt_n has no fixed points then $\alpha(u) = 2$ and we are done by the previous paragraph, so we may assume that the image of u in Alt_n lies in Alt_{n-1} ; we restrict a simple module M on which u acts minimally actively to $H = \text{Alt}_{n-1}$ or $H = 2 \cdot \text{Alt}_{n-1}$. Suppose that $H = \text{Alt}_{n-1}$ first. If a trivial submodule or quotient lies in this restriction then M is a composition factor of the permutation module on the cosets of H , so is either trivial or V . Moreover, since the composition factors of the restriction are minimally active, we know that the composition factors of $M \downarrow_H$ are either trivial or copies of V_H , the corresponding simple module for H . Since u acts on V_H with a block of size at least $o(u) - 2$, if there were more than one composition factor of $M \downarrow_H$ isomorphic with V_H then these two large Jordan blocks cannot form blocks of the form $o(u), 1^a$ unless $2(o(u) - 2) \leq o(u) + 1$,

i.e., $o(u) = 5$, and then $\dim(V_H)$ would need a block of size 3, only possible if $H = \text{Alt}_5$, but then $n = 6$, which is not allowed. Thus $M \downarrow_H$ has at most one copy of V_H , and can have no trivials as they would have to be submodules or quotients, so $\dim(M) \leq n - 1$, as needed.

If $H = 2 \cdot \text{Alt}_{n-1}$ then all composition factors of the restriction of M to H are non-trivial. If $n = 9$ then from Proposition 4.1 we see that $p = 7$. Since Alt_9 is generated by two 7-cycles, M is therefore a module that appears in the computations above. If $n = 10$, then again $p = 7$, this time by induction and using this proposition. Again, $\alpha(u) = 2$, and so M does not exist as it does not appear above. Finally, for $n \geq 11$ there can be no examples as there are no examples for $n = 10$. This completes the proof.

Case 2: $p = 2$ Here we do not need to consider $2 \cdot \text{Alt}_n$ but do need to consider $G = \text{Alt}_n$ and $G = \text{Sym}_n$. If $\alpha(u) = 2$ then again we need that $\dim(M) \leq 2n - 2$, in fact merely $\dim(M) \leq 2(o(u) - 1)$. From [12, Theorem 7 and Table 1] $\dim(M) > 2n - 2$ for $M \neq k, V$ for all $n \geq 15$, and for $n \leq 14$ we have that $o(u) \leq 8$, so we just need $\dim(M) \leq 14$, for which there are two modules for $G = \text{Alt}_9$ with this property, both of dimension 8 (but not isomorphic to V). However, Alt_9 does not contain elements of order 8, so these cannot be examples.

If $\alpha(u) = 2$, so that $M = V$, then by Lemma 4.2 we know that u acts minimally actively on V if and only if we are in cases (i)–(iii) of the proposition. Thus we may assume that $\alpha(u) > 2$, and in particular we cannot be in the situations given in Lemmas 2.9 and 2.10.

Suppose that u has at least three 2-cycles and does not have order 2: we can write $u = u_1 u_2$, where the supports of the u_i are disjoint, and where u_1 has cycle type $(2^a, 2, 2, 2)$ for some $a \geq 2$. Write H_i for the symmetric group on the support of u_i , $H = H_1 H_2 = H_1 \times H_2$, and note that the restriction $M \downarrow_H$ is minimally active. The simple kH -modules are tensor products of simple kH_i -modules, and by Lemma 2.3(v) a tensor product of two non-trivial simple modules is not minimally active (unless they both have dimension 2), we see that the composition factors of $M \downarrow_H$ are (minimally active) simple modules for one of the H_i . But H_1 has no non-trivial minimally active modules, so H_1 lies in the kernel of M , clearly nonsense as M is faithful.

Suppose that u has at least one 2-cycle and at least two cycles of length at least 4. Again, write $u = u_1 u_2$, this time with u the product of all of the cycles of length at least 4, unless there are exactly two of length exactly 4, in which case add another 2-cycle. (If u has cycle type $(4, 4, 2)$ then $\alpha(u) = 2$ by Lemma 2.10.) Defining the H_i and H as above, we again note that no non-trivial minimally active modules exist for H_1 , and so get the same contradiction. Since every fixed-point-free element has one of these properties, we have covered all fixed-point-free cases.

Thus u fixes a point and lies inside $H = \text{Sym}_{n-1}$. As for p odd, we restrict to H and note that the exact same proof works, as long as $n \geq 10$. In order to apply the argument for the case p odd to $p = 2$, we need to exclude the case $G_0 = \text{Alt}_9$, as for Alt_8 there are minimally active simple modules other than k and V . However, we already checked Alt_9 , so we may do this. Thus $M \downarrow_H$ has at most one copy of V and possibly trivial factors, and therefore M is a submodule of the permutation module on H , i.e., $M = k$ or $M = V$, as needed. \square

5 Lie type in defining characteristic

In the notation of the end of Section 2, this section considers G_0 a central extension of a simple group of Lie type in characteristic p .

Let \mathbf{G} be a simple, simply connected algebraic group defined over the field k , and let F be a Frobenius morphism on \mathbf{G} . With the exceptions of a few quasisimple groups (e.g., $2 \cdot \Omega_8^+(2)$, $3 \cdot \mathrm{PSL}_2(9)$, and so on) if G_0 is a quasisimple group of Lie type then for some choice of \mathbf{G} and F above, $G_0 = \mathbf{G}^F$. Moreover, from [5, Table 6.1.3] we see that if $Z(G_0)$ is a p' -group, which we require in order to have a faithful irreducible module, then G_0 is always \mathbf{G}^F (unless G_0 is an extension of $\mathrm{Sp}_4(2)' = \mathrm{Alt}_6$, which was examined in the previous section). Hence for this section we can always take G_0 to be the fixed points of \mathbf{G} under F .

Furthermore, every simple kG_0 -module is the restriction of a simple $k\mathbf{G}$ -module, and by Steinberg's tensor product theorem, every simple module for G_0 is a tensor product of Frobenius twists of p -restricted simple modules. Since tensor products of simple modules cannot be minimally active unless they have dimension 1 or 2 by Lemma 2.3(v), we thus consider p -restricted simple modules for \mathbf{G} .

As every unipotent class of \mathbf{G} appears in G_0 , when checking the action of a unipotent element on a given p -restricted simple module, we can use either \mathbf{G} or any of the quasisimple groups G_0 over the various ground fields. For explicit calculations, we of course will usually choose G_0 to be the smallest group, so over the field \mathbb{F}_p .

If $u \in G_0$ then the problem has been almost completely solved already, in [29] and [30], which dealt with all types apart from C and D in characteristic 2. Here we will finish that last case, and also consider the case where u induces an outer automorphism on G_0 . The next proposition completes the proof for $u \in G_0$. (In this proposition, (iv) and (v) can be viewed as subcases of (ii), using the isomorphisms $B_2 \cong C_2$ and $A_3 \cong D_3$, but we separate them out for clarity.) We use the notation $L(\lambda)$ for highest weight modules, using the same conventions as in for example [29] and [30] above. For weights of small rank we can write the weight in full, e.g., 110, but for larger rank it is more clear to write as a sum, so the previous example would be $\lambda_1 + \lambda_2$.

Proposition 5.1. *Let G_0 be a quasisimple group of Lie type, and let $u \in G_0$ be a non-trivial unipotent element. If M is a simple kG_0 -module on which u acts minimally actively, then up to outer automorphism of G_0 , one of the following holds:*

- (i) $G_0 = \mathrm{SL}_2(p^a)$, $M = L(i)$ for $0 \leq i \leq p-1$, or $M = L(1+p^j)$ of dimension 4 for some $1 \leq j \leq a/2$ and p odd, u has type $(i+1)$ and $(3,1)$ respectively;
- (ii) G_0 is of type A , 2A , B or C for all primes, D , 2D or 3D_4 for p odd, 2B_2 for $p=2$, M is the natural module;
- (iii) $p > 3$, $G_0 = \mathrm{SL}_3(p^a)$ or $\mathrm{SU}_3(p^a)$, u has type (3) on the natural module, $M = L(2\lambda_1)$ of dimension 6, u has type $(5,1)$;
- (iv) p is odd, $G_0 = \mathrm{SL}_4(p^a)$ or $\mathrm{SU}_4(p^a)$, u has type (4) or $(2,2)$ on the natural module, $M = L(\lambda_2)$ of dimension 6, u has type $(5,1)$ or $(3,1^3)$ respectively;
- (v) p is odd, $G_0 = \mathrm{Sp}_4(p^a)$, u has type (4) or $(2,2)$ on the natural module, $M = L(\lambda_2)$ of dimension 5, u has type (5) or $(3,1^2)$;
- (vi) p is odd, $G_0 = \mathrm{Spin}_7(p^a)$, u is regular unipotent, and $M = L(\lambda_3)$ of dimension 8, u has type $(7,1)$.
- (vii) $G_0 = G_2(p^a)$ or ${}^2G_2(3^{2a+1})$, u is regular unipotent, $M = L(\lambda_1)$ of dimension 7 (or 6 for $p=2$), u has type (7) (or (6) for $p=2$).

Proof. If $G_0 = \mathrm{SL}_2(p^a)$, then for $M = L(\lambda)$ p -restricted we see that u always acts with a single block, and for $L(\lambda)$ not p -restricted we use Lemma 2.3(v), which shows that p is odd and M is the tensor product of two 2-dimensional modules, i.e., $L(1 + p^j)$. By applying a field automorphism we may assume that $1 \leq j \leq a/2$.

For G_0 of type A , or types B , C and D and p odd, [29, Theorem 1.3] gives (ii)–(vi). (Note that the case of $p = 3$ for $G_0 = \mathrm{SL}_3(p^a)$ was erroneously included in [29], but can be excluded by Lemma 2.3(vii).) For G_0 of exceptional type [30] shows that only (vii) occurs, so we are left with types B/C and D in characteristic 2.

Let $G_0 = \mathrm{Sp}_{2n}(2^a)$ first. By [21, Chapter 4] every Jordan block of odd size appears an even number of times, so there can be no element of order 4 in G_0 that powers to a transvection. Thus if u is not itself a transvection, no power of u is one. By [8], $\alpha(u) \leq n + 3$ if u is not a transvection, and of course since G_0 acts on a $2n$ -dimensional space the order of u is at most $2a_n$, where a_n is the smallest power of 2 that is at least n . By Lemma 2.3(iv) we have $\dim(M) \leq \alpha(u) \cdot (o(u) - 1) \leq (n + 3)(2a_n - 1)$. (If u is a transvection then $\alpha(u) = 2n + 1$, and so $\dim(M) \leq 2n + 1$, so that M is the natural module.) By [22, Theorems 4.4 and 5.1], if $n \geq 8$ then M is one of the standard module, its symmetric square (does not occur for $p = 2$), or its exterior square with a trivial removed (with two trivials removed if n is even).

For $n \geq 3$, by Lemma 2.4 we see that this exterior square cannot be minimally active for u , and for $n = 2$ the simple modules for $\mathrm{Sp}_4(2)$ are the trivial, the natural and its image under the graph automorphism, and the Steinberg. Since $\mathrm{Sp}_4(2) = \mathrm{Sym}_6$, we use Proposition 4.1. Therefore we can assume that M is neither the natural $L(\lambda_1)$ nor the non-trivial factor of its exterior square $L(\lambda_2)$, and that $n \geq 3$.

For $3 \leq n \leq 7$, we use [22, Theorem 4.4] to get the following table of low-dimensional modules, with modules listed in order of increasing dimension.

n	Bound	Modules
3	42	100, 001, 010
4	49	1000, 0001, 0100, 0010
5	120	10000, 00001, 01000, 00100
6	135	$\lambda_1, \lambda_2, \lambda_6$
7	150	$\lambda_1, \lambda_2, \lambda_7$

As we have excluded $L(\lambda_1)$ and $L(\lambda_2)$, we need to consider the spin module $L(\lambda_n)$ for $\mathrm{Sp}_{2n}(2)$ for $3 \leq n \leq 7$, and the modules $L(0010)$ and $L(00100)$ for $\mathrm{Sp}_8(2)$ and $\mathrm{Sp}_{10}(2)$ respectively.

For $\mathrm{Sp}_8(2)$, $L(0010)$ is 48-dimensional, and since $\alpha(u) \leq 7$ for u not a transvection, we must have that $o(u) = 8$ if u acts minimally actively. If u has type $(6, 1^2)$, $(6, 2)$ or (8) on the natural module, then an easy computer calculation shows that u has type $(8^4, 6^2, 2^2)$, $(8^4, 6^2, 2^2)$ or (8^6) respectively on $L(0010)$, so cannot be minimally active.

For $\mathrm{Sp}_{10}(2)$, $L(00100)$ has dimension 100, and since $\alpha(u) \leq 8$ for u not a transvection, we must have that $o(u) = 16$ if u acts minimally actively, so u is the regular unipotent element. This element has type $(16^4, 14, 10, 6^2)$ on $L(00100)$, and so is not minimally active.

Finally, let $M = L(\lambda_n)$. Note that the restriction of M to the Sp_{2n-2} -parabolic has two composition factors, both isomorphic to $L(\lambda_{n-1})$. For $n = 3$, we compute the Jordan block structure of all unipotent elements on $L(\lambda_3)$ and get the following table.

Class	Action on $L(\lambda_1)$	Action on $L(\lambda_3)$
C_1	$2, 1^4$	2^4
A_1	$2^2, 1^2$	$2^2, 1^4$
$A_1^{(2)}$	$2^2, 1^2$	2^4
$A_1 + C_1$	2^3	2^4
C_2	$4, 1^2$	4^2
A_2	3^2	$3^2, 1^2$
$C_3(a_1)$	$4, 2$	4^2
C_3	6	$6, 2$

In particular, we see that there are no minimally active elements in $\mathrm{Sp}_6(2)$ on $L(001)$. Thus we proceed by induction on n . If u in $\mathrm{Sp}_{2n}(2)$ acts minimally actively on $L(\lambda_n)$, then place u inside an $\mathrm{Sp}_{2n-2}(2)$ -parabolic: it must act minimally actively on the indecomposable module for this group with socle $L(\lambda_{n-1})$, whence in particular it acts minimally actively on this submodule by Lemma 2.3(i). Since there are no non-trivial elements of $\mathrm{Sp}_{2n-2}(2)$ that act minimally actively on $L(\lambda_{n-1})$ by induction, u must act trivially on $L(\lambda_{n-1})$, whence $o(u) = 2$. One can see this either because the unipotent radical of the parabolic is elementary abelian, or because the action of u on $L(\lambda_n)$ must have blocks only of size 1 and 2, since there are two composition factors of the restriction of $L(\lambda_n)$ to the parabolic. At any rate, this is impossible since $\alpha(u) \leq 2n + 1$ and $\dim(M) = 2^n > 2n + 1 \geq \alpha(u) \cdot (o(u) - 1)$ by Lemma 2.3(iv).

We therefore need to consider groups of type D now. Let $G_0 = \Omega_{2n}^\pm(2)$, and note that $\alpha(u) \leq n + 3$ by [8, Theorem 4.4] (as transvections induce the graph automorphism on G_0 , so do not lie in G_0 itself). By placing G_0 inside $\mathrm{Sp}_{2n}(2)$, we see that no unipotent element can act minimally actively on $L(\lambda_2)$, since this is the restriction of the corresponding module for $\mathrm{Sp}_{2n}(2)$. (See for example [26, Table 1, MR₄].)

Using the bound $\dim(M) \leq \alpha(u) \cdot (o(u) - 1)$ from Lemma 2.3(iv), and [22, Theorems 4.4 and 5.1], we get the possible minimally active modules are the natural $L(\lambda_1)$, and $L(\lambda_2)$ (already eliminated above) for $n \geq 9$, and for $4 \leq n \leq 8$ we get the table below.

n	Bound	Modules
4	49	(1000, 0010, 0001), 0100, (0011, 1010, 1001)
5	56	$\lambda_1, \lambda_2, (\lambda_4, \lambda_5)$
6	135	$\lambda_1, \lambda_2, (\lambda_5, \lambda_6)$
7	150	$\lambda_1, \lambda_2, (\lambda_6, \lambda_7)$
8	165	$\lambda_1, \lambda_2, (\lambda_7, \lambda_8)$

The brackets indicate the groupings under the outer automorphism group. For $n \geq 5$ we need to check the two half-spin modules, but for $n = 4$ there are other modules to check, with the 48-dimensional modules $L(0011)$ and so on only occurring because our bound for $\alpha(u)$ is lax: checking by computer that $\alpha(u) = 2$ when $o(u) = 8$ inside $\Omega_8^+(2)$, we can therefore exclude them.

For $n = 5$, we note that D_5 lies inside E_6 acting as (up to automorphism) $L(0) \oplus L(\lambda_1) \oplus L(\lambda_5)$ on the minimal module for E_6 . As the dimension of $L(\lambda_5)$ is 16, we therefore need at least $17 - o(u)$ trivial Jordan blocks in the action of the corresponding unipotent class of E_6 on the minimal module: examining [20, Table 5], only the class A_1 of elements of order 2 have enough blocks of size 1, and of course if $o(u) = 2$ then $\dim(M) \leq 2n < 16$ (using [8, Theorem 4.4] and Lemma 2.3(iv)) which does not work either. For $n \geq 6$, we use induction, exactly the same as for type C . This completes the proof. \square

We now consider the case where u induces an outer automorphism on G_0 . We start with u inducing a graph automorphism on an untwisted group G_0 . In this case, $p = 2$ unless G_0 is type D_4 , and then $p = 2, 3$. We will then examine field and mixed field-graph automorphisms on untwisted groups, and finally how the automorphisms of the twisted groups compare with those of the untwisted groups.

In this proposition, (iii) may be thought of as the case $n = 3$ of (v), but we separate them for clarity.

Proposition 5.2. *Let G_0 be a quasisimple group of Lie type in characteristic p , and suppose that u lies in the coset of a graph automorphism on G_0 . If M is a minimally active, non-trivial simple module for G , then one of the following holds:*

- (i) $G = \mathrm{SL}_n(2^a).2$, u^2 is the regular unipotent element, and $M = L(\lambda_1) \oplus L(\lambda_{n-1})$ has dimension $2n$;
- (ii) $G = \mathrm{SL}_3(2^a).2$, u has order 8, and $M = L(11)$ has dimension 8;
- (iii) $G = \mathrm{SL}_4(2^a).2$, there are four possible classes for u , and $M = L(010)$ has dimension 6;
- (iv) $G = \mathrm{Sp}_4(2^a).2$ for a odd, u^2 is the regular unipotent element, and $M = L(10) \oplus L(01)$ has dimension 8;
- (v) $G = \mathrm{SO}_{2n}^+(2^a)$ for $n \geq 4$ and M is the natural module $L(\lambda_1)$.

Proof. Since u induces a graph automorphism on G_0 , we have that G_0 is untwisted by [5, Theorem 2.5.12(f)].

Suppose that u induces a graph automorphism on G_0 and that $p = 2$, so that G/G_0 has order 2. We go through each possibility in turn, of type A , type D , C_2 , F_4 and E_6 . (Note that G_2 only possesses the graph automorphism of order 2 when $p = 3$, so this case does not occur.) Notice that, if $M \downarrow_{G_0}$ is simple, then by Lemma 2.8 we may assume that it is 2-restricted or $G_0 = \mathrm{SL}_3(2^a)$ and $M \downarrow_{G_0}$ is the product of two 3-dimensional simple modules: but this is never graph stable, so we can ignore this case.

If $G_0 = E_6(2^a)$ then, since $o(u) \leq 32$ and $\alpha(u) \leq 9$, we have that $\dim(M) \leq 9 \cdot 31 = 279$ if M is minimally active, by Lemma 2.3(iv). The adjoint module $L(\lambda_2)$ is the only (non-trivial) graph-stable simple module with dimension at most 279, and the action of unipotent elements on this is given in [20]. For u to be minimally active, $u^2 \in G_0$ must have at most two non-trivial Jordan blocks, but this is not the case. If $M \downarrow_{G_0}$ is not simple then by Lemma 2.7 u^2 acts with a single Jordan block, which is not possible by Proposition 5.1.

For $G_0 = F_4(2^a)$, the exponent of the Sylow 2-subgroup of G_0 is 16, so $o(u) \leq 32$. Since $\alpha(u) \leq 8$ from [8, Theorem 5.1], this gives $\dim(M) \leq 248$. The dimensions of the simple modules are 1, 26, 26, 246, 246, and 676 and above. The (non-trivial) modules of dimension at most 246 are not graph-stable, and so as in the previous case we apply Lemma 2.7 and Proposition 5.1 to prove that no examples occur.

For $G_0 = \mathrm{Sp}_4(2^a)$ (we may assume that $a \geq 2$ since $a = 1$ yields Sym_6 , which has been considered already), there are only four 2-restricted modules: the trivial, the two 4-dimensional modules $L(10)$ and $L(01)$, swapped by the graph automorphism, and $L(11)$: the regular element acts with four blocks of size 4 on the $L(11)$ so this cannot extend to a minimally active module for G , and if u squares to the regular then $L(10) \oplus L(01)$ is minimally active by Lemma 2.7, since u^2 acts with a single Jordan block on $L(10)$. However, u can only induce a graph automorphism of order 2 if the Sylow 2-subgroup of $\mathrm{Out}(G_0)$ has order 2 by [5, Theorem 2.5.12(e)], with the graph automorphism squaring to a field automorphism in the other cases.

For G_0 of type D , note that $G_0.2 = \mathrm{SO}_{2n}^+(2^a)$ lies inside $\mathrm{Sp}_{2n}(2^a)$, with $\dim(M) \leq (n+3) \cdot (o(u) - 1)$ unless u is a transvection by [8, Theorem 4.4], and so we get the same bound as for the unipotent elements

for the simple group of type C in the proof of Proposition 5.1. Using the tables from [22], we see that for $n \geq 5$, every simple module for G that satisfies the bound on $\dim(M)$ is the restriction of a simple module for type C , and hence is minimally active only if the module is for type C . This yields only the natural module, which is of course minimally active. When $n = 4$, we get for G_0 the modules 0000 (trivial), 1000 (natural), $0010 \oplus 0001$ (sum of the two half-spins), 0100 (exterior square of natural), and 0011 of dimension 48. These are all also the restriction of a module for $\mathrm{Sp}_8(2^a)$, and so we are again done.

The last case for $p = 2$ is $G_0 = \mathrm{SL}_n(2^a)$. If $M \downarrow_{G_0}$ is not simple then by Lemma 2.7 we have that $v = u^2$ acts as a single Jordan block on each factor: thus v is the regular element and M is the sum of the natural and its dual.

Thus we may assume that $M \downarrow_{G_0}$ is simple, i.e., $M \downarrow_{G_0}$ is a graph-stable simple module, and 2-restricted by our discussion at the start of the proof. If v has maximal order in G_0 then $\alpha(u) \leq 4$ by Lemma 2.12, and writing a_n for the smallest power of 2 that is at least n , we have that $\dim(M) \leq 4 \cdot (2a_n - 1)$. We now use [22, Theorems 4.4 and 5.1], to get that $L(\lambda_1 + \lambda_{n-1})$ is the only graph-stable module of dimension at most this for $n \geq 7$. For $\mathrm{SL}_6(2^a)$ we have the module $L(\lambda_3)$, with $L(0110)$ for $\mathrm{SL}_5(2^a)$. For $n \leq 4$ there are several possibilities and we deal with them later.

If u does not have maximal order then $o(u) \leq a_n$ and $\alpha(u) \leq n$ by [8, Theorem 4.1], so that $\dim(M) \leq n(a_n - 1) < 2n^2$. We again use [22] to see which graph-stable modules we need to consider: doing so yields smaller bounds than the previous case, and so we need only consider $n = 3, 4$, and the specific modules for larger groups above.

We quickly show that $L(\lambda_1 + \lambda_{n-1})$ is not minimally active if $n \geq 5$: it is obtained from the tensor product of the natural and its dual by removing at most two trivial factors, and so if u inducing a graph on $L(\lambda_1 + \lambda_{n-1})$ is minimally active, then v acts with at most two non-trivial blocks on the module. As in the proof of Lemma 2.4, if v acts with at least two non-trivial blocks on $L(\lambda_1)$ then the action of v on the tensor square has at least eight non-trivial blocks, whence u cannot act minimally actively if one removes at most two trivials. If v acts with a trivial block and a block of size at least 3 then v acts on the tensor square with blocks at least of size $(4, 4, 3, 3, 1, 1)$, so u cannot be minimally active when removing two trivials. Similarly, if v contains a block of size at least 4 then v acts on the tensor square with at least four blocks of size at least 4, and if v acts on the natural with a block of size 2 and at least three trivial blocks, then v acts on the tensor square with at least five blocks of size at least 2, hence again u cannot be minimally active.

The actions of the classes of $\mathrm{SL}_6(2)$ on $L(\lambda_3)$ are as follows:

Type on $L(\lambda_1)$	Type on $L(\lambda_3)$
2, 1 ⁴	2 ⁶ , 1 ⁸
2 ² , 1 ²	2 ⁸ , 1 ⁴
2 ³	2 ¹⁰
3, 1 ³	3 ⁶ , 1 ²
3, 2, 1	4 ² , 3 ² , 2 ² , 1 ²
3 ²	4 ⁴ , 1 ⁴
4, 1 ²	4 ⁴ , 2 ²
4, 2	4 ⁴ , 2 ²
5, 1	7 ² , 3 ²
6	8 ² , 2 ²

None of these can be the square of a minimally active element, so we are done.

For $\mathrm{SL}_5(2)$, the dimension of $L(0110)$ is 74, so if $o(u^2) \leq 4$ then this module fails the bound $\alpha(u) \cdot (o(u) - 1)$, as $\alpha(u) \leq 5$. However, the only element v of $\mathrm{SL}_5(2)$ with order 8 is the regular unipotent element, and for this one $\alpha(v) = 2$ by Lemma 2.11, so this module cannot be minimally active.

We are thus left with $n = 3, 4$. For $n = 4$, we examine Proposition 4.1, which states that the 6-dimensional module $L(\lambda_2)$ is the only graph-stable minimally active module, and here there are several classes that work. For $n = 3$, we simply check the simple modules for $\mathrm{PGL}_2(7) = \mathrm{SL}_3(2).2$, and an element of order 8 acts with a single Jordan block on both $L(10) \oplus L(01)$, in line with the proposition, and also on the 8-dimensional module $L(11)$, as seen in [25, Theorem 1.2].

For $p = 3$, we only have the group $\Omega_8^+(3^a)$ to consider, and since the graph automorphism of order 3 permutes the central involutions of the Spin group regularly, we may assume that G_0 is simple. If M is minimally active and $M \downarrow_{G_0}$ is not simple then u^3 must act with a single Jordan block on each composition factor of $M \downarrow_{G_0}$ by Lemma 2.7, but this does not occur by Proposition 5.1. Thus $M \downarrow_{G_0}$ is simple: the smallest non-trivial, graph-stable simple module has dimension 28, and the next smallest has dimension 195 (see [22, Appendix A.41]), and since $\alpha(u) = 2$ if $o(u) = 9$ and $\alpha(u) \leq 4$ if $o(u) = 3$ for $G_0 = \mathrm{P}\Omega_8^+(3)$, we have that $\dim(M) = 28$ and $M = L(\lambda_2)$. Since u is minimally active, u^3 has at most three non-trivial Jordan blocks. However, of the 27 non-trivial unipotent classes of 3-elements in $\Omega_8^+(3)$, all have at least four non-trivial blocks on $L(\lambda_2)$. Thus there is no candidate for M . \square

Having dealt with the case where u induces a graph automorphism, we are left with the case where u induces either a field automorphism or a mixed field-graph automorphism on the untwisted group G_0 .

Proposition 5.3. *Let G_0 be a quasisimple group of Lie type in characteristic p , and suppose that u induces an outer automorphism on G_0 that is not a graph automorphism. If M is a minimally active simple module for G , then up to outer automorphism of G one of the following holds:*

- (i) $p = 2$, $G = \mathrm{SL}_2(2^{2a}).2$, $M = L(1 + 2^a)$ of dimension 4;
- (ii) $p = 3$, $G = \mathrm{SL}_2(3^{3a}).3$, $M = L(1 + 3^a + 3^{2a})$ of dimension 8;
- (iii) $p = 2$, $G = \mathrm{SL}_3(2^{2a}).2$, u induces either a field or the product of a field and graph automorphism on G_0 , $M = L(1 + 2^a, 0)$ or $M = (1, 2^a)$ respectively, both of dimension 9;
- (iv) $p = 2$, $G = \mathrm{SU}_3(2^{2a}).2$, u induces the unique outer automorphism of order 2 on G_0 , $\dim(M) = 9$ is such that $M \downarrow_{G_0}$ is simple;
- (v) $G = G_0.t$, $M \downarrow_{G_0} = M_1 \oplus \cdots \oplus M_t$ with $u^t \in G_0$ acting on each M_i with a single Jordan block, the possibilities for which are given in Proposition 5.1.

Proof. We begin with the case where G_0 is untwisted.

Suppose that M is minimally active and that $M \downarrow_{G_0}$ is the simple module $L(\lambda)$. By Lemma 2.8, either $L(\lambda)$ is (up to Frobenius twist) a p -restricted module, or $p = 2$ and $G_0 = \mathrm{SL}_3(2^a)$, or $p = 2, 3$ and $G_0 = \mathrm{SL}_2(p^a)$, with M a product of p twists of the natural module. If u induces a field automorphism then u replaces a highest weight $\lambda = a_1\lambda_1 + \cdots + a_n\lambda_n$ with $p^\alpha a_1\lambda_1 + \cdots + p^\alpha a_n\lambda_n$ for some $\alpha \geq 1$, and since graph automorphisms permute the a_i , for u to fix $L(\lambda)$ it cannot be p -restricted.

In the remaining cases of $\mathrm{SL}_3(2^a)$ and $\mathrm{SL}_2(p^a)$ for $p = 2, 3$, we have from Lemma 2.8 that u has order p and M is the tensor product of the modules in a single orbit under the action of u : for $G_0 = \mathrm{SL}_2(p^a)$, this orbit is clear, whereas for $\mathrm{SL}_3(2^a)$ if u is a pure field automorphism we get that $L(1, 0)$ and $L(2^a, 0)$ form

an orbit, and if u is the product of the field and the graph it is $L(1, 0)$ and $L(0, 2^a)$, yielding the modules in the statement of the proposition.

We therefore have that $M \downarrow_{G_0}$ is not simple, and is the sum of t simple factors M_1, \dots, M_t , each stabilized by u^t . By Lemma 2.7, u^t acts on each M_i with a single Jordan block, and since u^t cannot act as a pure graph automorphism on the M_i , we must therefore have that $u^t \in G_0$, and so the conclusion of the proposition holds here as well.

Now suppose that G_0 is twisted, so that u must be a field automorphism: if $M \downarrow_{G_0}$ is simple then it cannot be p -restricted, so as with the untwisted case we get that G_0 is of type A_2 and we proceed similarly to the case of $\mathrm{SL}_3(2^a)$. If $M \downarrow_{G_0}$ is not simple, then we get the same proof as for the untwisted case, yielding the result above. \square

There are many examples of (v) above, and some that may appear to be but are not. For example the group ${}^3D_4(3).3$ with M of dimension 24, is the sum of three 8-dimensional modules permuted by the graph automorphism. Each 8-dimensional module for ${}^3D_4(3)$ possesses minimally active elements by Proposition 5.1, but since the simple group of type D does not contain an element acting with a single block of size 8, there can be no minimally active u for M .

As another negative example, we consider $G = {}^2G_2(3)$, which has a module M of dimension 27 where the derived subgroup G' acts on M as the sum of three blocks, and G' contains an element v (of order 9) acting on each factor with a single Jordan block, but there is no element $u \in G$ such that $u^3 = v$ (as then the Sylow 3-subgroup would be cyclic).

We should enumerate which possibilities from Proposition 5.1 actually have an element acting with a single block, so that they can be inputs into case (v) in Proposition 5.3. For $G_0 = \mathrm{SL}_2(p^a)$, u acts as a single block on $L(i)$ for $i < p$, but acts as $(3, 1)$ on $L(1 + p^i)$. For G_0 of type A , 2A , B , 2B , and C , there is an element acting with a single Jordan block, but not for those of type D . We can include the 5-dimensional module for $\mathrm{Sp}_4(p^a)$ in those above via the isomorphism $B_2 \cong C_2$, and we also have G_2 with the minimal module.

Hence, if $G_0 = \mathrm{SU}_n(p^p)$ for some n, p for example, one may form the group $G = \mathrm{SU}_n(p^p).p$, and let u be an element of g whose p th power is the regular unipotent element of G_0 . In this case the hypotheses of (v) are satisfied, so this is an example.

6 Sporadic groups

In [24], all almost cyclic, and in particular minimally active elements were found for the case where G_0 is a central extension of a sporadic simple group, and where $G = G_0$. In this short section we deal with the case where u induces an outer automorphism of G_0 . Since $|\mathrm{Out}(G_0)| \leq 2$, we will always assume in this section that $p = 2$.

We can easily determine the outer classes of 2-elements from [2], and in Table 2 we give the largest order of such a 2-element, with the obvious intent to use the formula $\dim(M) \leq \alpha(u) \cdot (o(u) - 1)$ from Lemma 2.3(iv). For this we also need estimates for $\alpha(u)$ and $\dim(M)$: the latter appears in [13], and the former in [24, Theorem 3.1]. (This gives $\alpha(u^2)$, but of course $\alpha(u) \leq \alpha(u^2)$. If $o(u) = 2$ then we can use [8, Table 1].)

We get the following lemma.

Lemma 6.1. *Let G_0 be a sporadic simple group, and suppose that $u \in G \setminus G_0$ is a 2-element. If $o(u) \geq 8$ then $\alpha(u) = 2$. If $o(u) = 4$ then $\alpha(u) \leq 3$. If $o(u) = 2$ then $\alpha(u) \leq 8$, with $\alpha(u) \leq 4$ if $G_0 = J_2$.*

G_0	Largest 2-element of $G \setminus G_0$	G_0	Largest 2-element of $G \setminus G_0$
M_{12}	4	He	16
M_{22}	8	HN	8
HS	8	Fi_{22}	16
J_2	8	Fi'_{24}	16
McL	8	ON	8
Suz	16	J_3	8

Table 2: Exponents of Sylow 2-subgroups of $\text{Aut}(G_0/Z(G_0))$ for G_0 a sporadic quasisimple group with non-trivial outer automorphism group.

G_0	Minimal non-trivial simple module	G_0	Minimal non-trivial simple module
M_{12}	10	He	51
M_{22}	10	HN	132
HS	20	Fi_{22}	78
J_2	6	Fi'_{24}	3774
McL	22	ON	10944
Suz	110	J_3	78

Table 3: Minimal dimensions of non-trivial representations of G_0 .

Proof. From the tables in [2], we see that u cannot square to 4A when $G_0 = HS$, nor to 2A when $G_0 = Fi_{22}$. If $G_0 = J_2$ then the class 4B squares to 2A, so we need to check how many conjugates generate G in this case: a quick computer calculation shows that $\alpha(u) = 2$ for this class.

If $o(u) = 2$ then we use the bounds on $\alpha(G)$ given in [8]. □

Combining this information with Table 3, we see that the only possibilities for u acting minimally actively are that $G_0 = M_{22}, J_2$. For $G = M_{22}.2$, the only candidate simple module has dimension 10, and elements of order 8 act on this with type $(8, 2)$, so this is not an example. For $G = J_2.2$, the 6-dimensional simple modules of G_0 are swapped by the outer automorphism, so that $\dim(M) \geq 12$. In order for u to act minimally actively on this it must act as a single Jordan block by Lemma 2.7, but that is clearly impossible. The next smallest dimension is 28 (see [14, p.102]) so there are no examples here either.

This proves that $u \in G_0$ in all cases, so we get the following proposition, proved in [24].

Proposition 6.2. *Suppose that G_0 is a central extension of a sporadic simple group, and let u be a p -element of G such that $G = \langle G_0, u \rangle$. If u acts minimally actively on a non-trivial simple module M , then $G = G_0$ and one of the following holds:*

- (i) $G = M_{11}$, $p = 11$, $o(u) = 11$ and $\dim(M) = 9, 11$ (one representation each) or $\dim(M) = 10$ (two representations);
- (ii) $G = M_{12}$, $p = 11$, $o(u) = 11$ and $\dim(M) = 11$ (two representations);
- (iii) $G = 2 \cdot M_{12}$, $p = 11$, $o(u) = 11$ and $\dim(M) = 10$ (two representations) or $\dim(M) = 12$ (one representation);
- (iv) $G = 3 \cdot M_{22}$, $p = 2$, $o(u) = 8$ and $\dim(M) = 6$ (two representations);

- (v) $G = 2 \cdot M_{22}$, $p = 11$, $o(u) = 11$ and $\dim(M) = 10$ (two representations);
- (vi) $G = M_{23}$, $p = 23$, $o(u) = 23$ and $\dim(M) = 21$;
- (vii) $G = M_{24}$, $p = 23$, $o(u) = 23$ and $\dim(M) = 23$.
- (viii) $G = J_2$, $p = 2$, $o(u) = 8$ and $\dim(M) = 6$, (two representations) requires $\mathbb{F}_4 \subseteq k$;
- (ix) $G = 2 \cdot J_2$, $p = 7$, $o(u) = 7$ and $\dim(M) = 6$ (two representations) requires $\mathbb{F}_{49} \subseteq k$;
- (x) $G = 6 \cdot \text{Suz}$, $p = 11, 13$, $o(u) = p$ and $\dim(M) = 12$ (two representations);
- (xi) $G = Co_3$, $p = 23$, $o(u) = 23$, and $\dim(M) = 23$;
- (xii) $G = Co_2$, $p = 23$, $o(u) = 23$, and $\dim(M) = 23$;
- (xiii) $G = 2 \cdot Co_1$, $p = 23$, $o(u) = 23$, and $\dim(M) = 24$.
- (xiv) $G = J_1$, $p = 11$, $o(u) = 11$ and $\dim(M) = 7$;
- (xv) $G = 3 \cdot J_3$, $p = 2$, $o(u) = 8$ and $\dim(M) = 9$ (two representations);
- (xvi) $G = 3 \cdot J_3$, $p = 17, 19$, $o(u) = p$ and $\dim(M) = 18$ (four representations);
- (xvii) $G = 2 \cdot Ru$, $p = 29$, $o(u) = 29$ and $\dim(M) = 28$ (two representations).

Unless specified, the field of definition of these representations is the smallest k such that $|Z(G_0)| \mid |k^\times|$.

7 Groups of Lie type in cross characteristic: preliminaries

In this section we tackle groups of Lie type in characteristic r where $r \neq p$, so let q be a power of r . We specifically exclude the case where the group is also a group in characteristic p , so for $\text{SL}_3(2)$ for example, $p \neq 2, 7$. If G_0 is a classical group and M is a Weil representation, or if $G_0 = \text{PSL}_2(q)$, with u inducing an inner-diagonal automorphism on G_0 in both cases, then all minimally active u are classified in [25], but the case where u induces particularly a non-diagonal outer automorphism on M is missing. In this section we give some notation and introduce some previous results, particularly on minimal dimensions of irreducible representations.

Let G_0 be a quasisimple group of Lie type, defined over the field \mathbb{F}_q , let r be the prime dividing q , and let $p \neq r$ be another prime. With a few exceptions given in [5, Table 6.1.3], G_0 is (a quotient of) the fixed points of a Frobenius endomorphism of a simple, simply connected algebraic group in characteristic r .

Note that those groups of Lie type that are isomorphic to alternating groups have already been covered in Proposition 4.1, and we will ignore them from now on. We will also ignore cases where G_0 is isomorphic to a group in characteristic p , so under the isomorphisms:

- $\text{PSL}_2(4) \cong \text{PSL}_2(5) \cong A_5$;
- $\text{PSp}_4(2)' \cong \text{PSL}_2(9) \cong A_6$,
- $\text{PSL}_4(2) \cong A_8$;
- $\text{PSL}_3(2) \cong \text{PSL}_2(7)$,
- $\text{PSU}_3(3) \cong G_2(2)'$,

- $\mathrm{PSL}_2(8) = {}^2G_2(3)'$,
- $\mathrm{PSU}_4(2) \cong \mathrm{PSp}_4(3)$.

We find all simple, minimally active modules for classical G with G_0 possessing one of the central extensions given in [5, Table 6.1.3] now. For G_0 of exceptional type, see Proposition 9.1 except for ${}^2E_6(2)$, for which standard arguments work.

In the next result, if there is more than one central extension $n \cdot G$ (for example, $G = \mathrm{PSL}_3(4)$ has more than one 4-fold extension) we use the numbering of these from the Atlas [2], and we use [14] for information about the existence of certain simple modules for central extensions.

Proposition 7.1. *Let G_0 be a central extension of one of the following groups: $\mathrm{PSL}_3(2)$, $\mathrm{PSL}_3(4)$, $\mathrm{PSU}_4(2)$, $\mathrm{PSU}_4(3)$, $\mathrm{PSU}_6(2)$, $\mathrm{Sp}_6(2)$, $\Omega_7(3)$, $\Omega_8^+(2)$. Let u be a p -element of G such that $G = \langle G_0, u \rangle$. If u acts minimally actively on a non-trivial simple module M , then one of the following holds:*

- (i) $G = \mathrm{SL}_3(2)$, $p = 3$, $o(u) = 3$, $\dim(M) = 3$ (two representations) requires $\mathbb{F}_9 \subseteq k$;
- (ii) $G = 2 \cdot \mathrm{SL}_3(2)$, $p = 3$, $o(u) = 3$, $\dim(M) = 4$ (two representations) requires $\mathbb{F}_9 \subseteq k$;
- (iii) $G = 6 \cdot \mathrm{PSL}_3(4)$, $p = 5, 7$, $o(u) = p$ and $\dim(M) = 6$ (two representations), with u acting as $(5, 1)$ or (6) ;
- (iv) $G = 4_1 \cdot \mathrm{PSL}_3(4)$, $p = 7$, $o(u) = 7$, $\dim(M) = 8$ (four representations), with u acting as $(7, 1)$;
- (v) $G = \mathrm{PSU}_4(2)$, $p = 5$, $o(u) = 5$, $\dim(M) = 5$ (two representations) requires $\mathbb{F}_{25} \subseteq k$ or $\dim(M) = 6$ (one representation), with u acting as (5) or $(5, 1)$;
- (vi) $G = 2 \cdot \mathrm{PSU}_4(2)$, $p = 5$, $o(u) = 5$, $\dim(M) = 4$ (two representations) requires $\mathbb{F}_{25} \subseteq k$, with u acting as (4) ;
- (vii) $G = 6_1 \cdot \mathrm{PSU}_4(3)$, $p = 5, 7$, $o(u) = p$ and $\dim(M) = 6$ (two representations), with u acting as $(5, 1)$ or (6) ;
- (viii) $G = 3_1 \cdot \mathrm{PSU}_4(3)$, $p = 2$, $o(u) = 8$ and $\dim(M) = 6$ (two representations), with u acting as (6) ;
- (ix) $G = 3_1 \cdot \mathrm{PSU}_4(3).2_2$, $p = 2$, u can have order 2, 4, 8 (but not all elements of orders 2 or 4), and $\dim(M) = 6$ (two representations), with u acting as $(2, 1^4)$, $(4, 1^2)$ and (6) ;
- (x) $G = \mathrm{Sp}_6(2)$, $p = 3$, $o(u) = 3, 9$, $\dim(M) = 7$, with u acting as $(3, 1^4)$ or (7) (minimally active elements of order 3 are from the smallest conjugacy class);
- (xi) $G = \mathrm{Sp}_6(2)$, $p = 5, 7$, $o(u) = p$ and $\dim(M) = 7$, with u acting as $(5, 1^2)$ and (7) ;
- (xii) $G = 2 \cdot \mathrm{Sp}_6(2)$, $p = 3, 7$, $o(u) = 7, 9$ and $\dim(M) = 8$, with u acting as $(7, 1)$;
- (xiii) $G = 2 \cdot \Omega_8^+(2)$, $p = 3$, u has order 9, or u has order 3 with centralizer $Z_6 \times \mathrm{PSp}_4(3)$, and $\dim(M) = 8$;
- (xiv) $G = 2 \cdot \Omega_8^+(2)$, $p = 5$, u has order 5 with centralizer $Z_{10} \times \mathrm{PSL}_2(5)$, and $\dim(M) = 8$;
- (xv) $G = 2 \cdot \Omega_8^+(2)$, $p = 7$, $o(u) = 7$ and $\dim(M) = 8$.

Unless specified, the field of definition of these representations is the smallest k such that $|Z(G_0)| \mid |k^\times|$.

Proof. If G_0 has $\mathrm{PSL}_3(2)$ as a quotient then $p \neq 2, 7$, as these were considered in Section 5. Thus $p = 3$, $G = \mathrm{SL}_2(7)$, $o(u) = 3$, and $C_G(u) = \langle u \rangle \cdot Z(G)$, so that M is minimally active if and only if $\dim(M) \leq p+1 = 4$ by Lemma 2.6.

Now let G_0 be a central extension of $\mathrm{PSL}_3(4)$, and to begin let $p = 3$. The Sylow 3-subgroup of $\mathrm{PGL}_3(4)$ has exponent 3: $\mathrm{PSL}_3(4)$ has a unique class of elements u of order 3 and is generated by two conjugates of them, so if u acts minimally actively then $\dim(M) \leq 4$, but the minimal dimension for a simple module for $(4 \times 4) \cdot \mathrm{PSL}_3(4)$ is 6. (The Schur multiplier of $\mathrm{PSL}_3(4)$ is $4 \times 4 \times 3$.)

Alternatively, u could lie outside $\mathrm{PSL}_3(4)$ in G , and then G is generated by three conjugates of u , so $\dim(M) \leq 6$. However, we cannot form a central extension of $\mathrm{PGL}_3(4)$ by a 2-group as the outer automorphism acts transitively on the involutions in the $Z_4 \times Z_4$ Sylow 2-subgroup of the Schur multiplier, and the minimal dimension for $\mathrm{PGL}_3(4)$ is 19.

When $p = 5$, the normalizer of a Sylow 5-subgroup of $\mathrm{PSL}_3(4)$ is D_{10} , so we apply Lemma 2.6 to see that M is minimally active if and only if $\dim(M) \leq 6$. Similarly, the normalizer of a Sylow 7-subgroup of $\mathrm{PSL}_3(4)$ is $Z_7 \rtimes Z_3$, so again $\dim(M) \leq 8$ by Lemma 2.6. There are 6-dimensional representations of $6 \cdot \mathrm{PSL}_3(4)$ modulo 5 and 7, and 8-dimensional representations of $4_1 \cdot \mathrm{PSL}_3(4)$.

If G_0 is a central extension of $\mathrm{PSU}_4(2) = \mathrm{PSP}_4(3)$, then $p = 5$. For $p = 5$, the Sylow 5-subgroup has order 5, generated by u , and $C_G(u) = \langle u \rangle \cdot Z(G)$, so that M is minimally active if and only if $\dim(M) \leq p+1 = 6$. There are modules for $\mathrm{PSU}_4(2)$ of dimensions 5 and 6, and of $2 \cdot \mathrm{PSU}_4(2)$ of dimension 4, completing the proof.

Let G_0 be a central extension of $\mathrm{PSU}_4(3)$, so that $p = 2, 5, 7$. For $p = 5, 7$, we have that the Sylow p -subgroup is of order p , generated by u , and $C_G(u) = \langle u \rangle \cdot Z(G)$, so that $\dim(M) \leq p+1$ by Lemma 2.6. There is a module of dimension 6 for $6_1 \cdot \mathrm{PSU}_4(3)$, so this is minimally active for both primes. For $p = 2$, as $\mathrm{Out}(G_0)$ is D_8 and the Schur multiplier is $3 \times 3 \times 4$ there are many potential groups G .

If $G_0 = \mathrm{PSU}_4(3)$, then from [14] we see that $\dim(M) = 20$ or $\dim(M) \geq 34$. The order of $u \in G$ is 2, 4, 8 from [2], and for $u \in G_0$, $\alpha(u) = 2$ if $o(u) = 4, 8$, with $\alpha(u) = 3$ if $o(u) = 2$. As $\mathrm{Out}(G_0) = D_8$, $u^4 \in G_0$, so the only way that u can act minimally actively is if $o(u) = 8$, $u^2 \notin G_0$, and $\alpha(u) = 3$. However, by constructing $\mathrm{Aut}(G_0)$ in Magma, we check that $\alpha(u) = 2$ for all u of order 8, hence there are no non-trivial minimally active modules if $Z(G_0) = 1$. In $\mathrm{Aut}(G_0)$, we have that $\alpha(u) = 2$ if $o(u) = 8$, $\alpha(u) \leq 4$ if $o(u) = 4$, and $\alpha(u) \leq 6$ if $o(u) = 2$.

Thus G_0 is either $3_1 \cdot \mathrm{PSU}_4(3)$ or $3_2 \cdot \mathrm{PSU}_4(3)$. In the second case, $\dim(M) \geq 36$, so we again see that there are no minimally active modules by the above computations for $\alpha(u)$, as $\dim(M) \leq 14$ for M to be minimally active. Thus $G_0 = 3_1 \cdot \mathrm{PSU}_4(3)$, and from [2, p.53] we see that the only outer automorphism that centralizes G_0 is 2_2 , so we let G be either G_0 or $G_{0.2_2}$.

The only non-trivial simple module for G_0 of dimension at most 14 has dimension 6 (two up to duality), and this extends to $G_{0.2_2}$. Inside G_0 elements of order 8 act with type (6), and in $G_{0.2_2}$ (modulo a central involution, this is the complex reflection group G_{34} in Shephard–Todd notation) there are elements of orders 2, 4 and 8 (with the last one not in G_0) that act with types $(2, 1^4)$, $(4, 1^2)$ and (6) respectively.

Let G_0 be a central extension of $\mathrm{PSU}_6(2)$, so that $p = 3, 5, 7, 11$. If $p = 7, 11$ then $C_{G_0}(u) = \langle u \rangle \cdot Z(G_0)$, so that $\dim(M) \leq p+1$ if and only if M is minimally active by Lemma 2.6. However, $\dim(M) \geq 21$ for all odd p by [19], a contradiction. If $p = 5$ then $o(u) = 5$ and $\alpha(u) = 2$, so that $\dim(M) \leq 8$ for minimally active M , another contradiction. If $p = 3$ then there is an outer automorphism of order p , but in $\mathrm{PGU}_6(2)$ the exponent of the Sylow 3-subgroup is still 9, so if $\alpha(u) = 2$ when $o(u) = 9$ and $\alpha(u) \leq 10$ for $o(u) = 3$

then we are done. The former follows by a computer calculation, and the latter follows from [8, Theorem 4.1].

Let G_0 be a central extension of $\mathrm{Sp}_6(2)$, so that $p = 3, 5, 7$. The Sylow 7-subgroup of G_0 is cyclic, and if u is a generator for it then $C_{G_0}(u) = \langle u \rangle \cdot Z(G)$, so that $\dim(M) \leq p + 1 = 8$ if and only if M is minimally active. There is a 7-dimensional simple module for $\mathrm{Sp}_6(2)$ and an 8-dimensional simple module for $2 \cdot \mathrm{Sp}_6(2)$, so these are minimally active. If $p = 5$ then the Sylow 5-subgroup P has order 5, and $\alpha(u) = 2$ for u of order 5, yielding $\dim(M) \leq 8$ for minimally active modules. The module of dimension 7 is minimally active, with u of type $(5, 1^2)$, but on the module of dimension 8 the action has type (4^2) .

For $p = 3$, if $o(u) = 9$ then $\alpha(u) = 2$, and if $o(u) = 3$ then $\alpha(u) \leq 4$, so $\dim(M) \leq 16$ for M minimally active. There are three such non-trivial simple modules, of dimensions 7, 8 and 14. The 14-dimensional module is not minimally active, but elements of order 9 act with a single Jordan block on the other two modules. In addition, elements of order 3 from the smallest class, with centralizer of order 2160 in $\mathrm{Sp}_6(2)$, act on M of dimension 7 with type $(3, 1^4)$.

The next group is G_0 a central extension of $\Omega_7(3)$, with primes 2, 5, 7, 13. From [19], $\dim(M) \geq 27$ if M is non-trivial. Note that $o(u) \leq 13$, and $\alpha(u) = 2$ if $o(u) \geq 5$, $\alpha(u) \leq 3$ for $o(u) = 4$, $\alpha(u) \leq 4$ if $o(u) = 3$ and $\alpha(u) \leq 7$ if $o(u) = 2$. Thus there are no minimally active modules for G_0 . However, $\mathrm{Out}(G_0)$ has order 2, and from [2] we see that the exponent of the Sylow 2-subgroup of $\mathrm{Aut}(\Omega_7(3))$ has order 8; in this case, $\alpha(u)$ is as before if $o(u) = 2, 8$, and $\alpha(u) \leq 4$ if $o(u) = 4$. Again, there can be no non-trivial simple minimally active modules.

The final group on our list is $\Omega_8^+(2)$, where $p = 3, 5, 7$. For $G = 2 \cdot \Omega_8^+(2)$, $\alpha(u) = 2$ for $o(u) = 5, 7, 9$, and $\alpha(u) \leq 4$ for $o(u) = 3$. The 8-dimensional simple module for G is minimally active for $p = 7$, with type $(7, 1)$, and for $p = 5$ the three classes of elements of order 5 have type $(5, 1, 1, 1)$, and $(4, 4)$ twice, with the two classes of elements of order 5 having centralizer $Z_5 \times \mathrm{SL}_2(5)$ and the minimally active one having centralizer $Z_{10} \times \mathrm{PSL}_2(5)$.

For $p = 3$, we need to consider $G = 2 \cdot \Omega_8^+(2)$ and $\Omega_8^+(2).3$: the former case is easy, with elements of order 9 acting on the 8-dimensional module as $(7, 1)$, and one of the three classes of elements of order 3 with centralizer of order 155520 have type $(3, 1^5)$ and the other two having type (2^4) ; again these two have centralizer $Z_3 \times \mathrm{Sp}_4(3)$, and the one we want has centralizer $Z_6 \times \mathrm{PSp}_4(3)$. (There are two classes of elements of order 3 with smaller centralizer.)

To deal with $\Omega_8^+(2).3$, note that from [14] we get that $\dim(M) \geq 28$, and $o(u) = 3, 9$. Since $\alpha(u) \leq 4$ by [8, Theorem 4.4], elements of order 3 cannot work, and elements u of order 9 must cube to an element $v = u^3$ of order 3 in G_0 that acts on M with at most three blocks of size 3. Since $\dim(M) = 28$ or $\dim(M) \geq 48$, we have $\dim(M) = 28$, and the five conjugacy classes of elements of order 3 act on M as $(3^6, 1^{10})$, $(3^7, 2^2, 1^3)$ and $(3^9, 1)$, so it cannot be minimally active. This completes the proof for $\Omega_8^+(2)$. \square

Because of Proposition 7.1, if G is classical then we may take G_0 to be a quotient of one of the groups $\mathrm{SL}_n(q)$, $\mathrm{SU}_n(q)$, $\mathrm{Sp}_{2n}(q)$, $\mathrm{Spin}_{2n+1}(q)$ and $\mathrm{Spin}_{2n}^\pm(q)$. The order of $G_0(q)$ is given by a polynomial

$$q^N \prod_i \Phi_i(q)^{a_i},$$

where N and the a_i are integers, and Φ_i denotes the i th cyclotomic polynomial. If u is a p -element and $p \nmid q$ then p divides one of the $\Phi_i(q)$; let d denote the order of q modulo p , so that $p \mid \Phi_d(q)$ and $p \nmid \Phi_e(q)$ for all $1 \leq e < d$. (If $p = 2$, we let d be the order of q modulo 4.) This next well-known lemma tells us about the powers of p dividing various cyclotomic polynomials.

Lemma 7.2. *Let $p \neq r$ be a prime and suppose that q is a power of r .*

- (i) *Writing d for the order of q modulo p , $p \mid \Phi_e(q)$ if and only if $e = p^a d$ for some $a \geq 0$ (except if $p = 2$ and $d = 2$, where $2 \mid \Phi_1(q)$ as well);*
- (ii) *If e is not the order of q modulo p , then $p^2 \nmid \Phi_e(q)$;*
- (iii) *If $p \nmid d$, then $\Phi_d(q^p) = \Phi_d(q) \cdot \Phi_{pd}(q)$, and if $p \mid d$ then $\Phi_d(q^p) = \Phi_{pd}(q)$. Therefore for all d , the powers of p dividing $\Phi_d(q^{p^a})$ and $p^a \Phi_d(q)$ are the same.*

We now need information about the cross-characteristic Sylow structure of a group of Lie type, which is described in [5, Theorem 4.10.2]. We give a summary now, tailored to our needs.

Proposition 7.3. *Let $G_0 = G_0(q)$ denote a quasisimple group of Lie type, with $Z(G_0)$ a p' -group. Let d denote the order of q modulo p , and let p^a be the exact power of p dividing $\Phi_d(q)$. Let P be a Sylow p -subgroup of G_0 .*

There exists an abelian normal subgroup P_0 of G_0 , of exponent p^a , such that P/P_0 is isomorphic to a subgroup of the Weyl group of G_0 , unless one of the following holds:

- (i) $p = 3$, $G_0 = {}^3D_4(q)$, where P_0 has exponent p^{a+1} ;
- (ii) $p = 2$, $G_0 = {}^2G_2(q)$, where P is elementary abelian of order 8.

Furthermore, if G is an almost simple group containing G_0 as a normal subgroup, with G/G_0 consisting of diagonal automorphisms, then the same results hold.

This means that, in the notation of the proposition, if the exponent of the Sylow p -subgroup of the Weyl group of G_0 is p^b , then the exponent of P is at most p^{a+b} (except in the one case, where it is at most p^{a+b+1}).

To get a bound for the maximal order of u , we finally need to consider the outer automorphism group of G_0 , which is more or less completely described in [5, Theorem 2.5.12]. Thus the contribution to $o(u)$ comes from three sources: the *toral contribution*, the p -part of $\Phi_d(q)$ (except for 3D_4), the *Weyl contribution*, the exponent of the Sylow p -subgroup of the Weyl group, and the *outer contribution*, the exponent of the Sylow p -subgroup of the outer automorphism group of G_0 . This is usually far greater than the actual maximal order of u , and so we use this to reduce the possible options for G_0 , and then use more explicit techniques to get better bounds on $o(u)$ if required.

Notice that the Sylow p -subgroup of G_0 is abelian if and only if p divides exactly one of the $\Phi_d(q)$ that divide $|G_0|$, or in other words, if the Sylow p -subgroup of G_0 is non-abelian and the order of q modulo p is d , then both $\Phi_d(q)$ and $\Phi_{pd}(q)$ divide $|G_0(q)|$.

We also need to consider regular semisimple elements, in particular to know that their centralizer is abelian when u is a regular semisimple element in a cyclic Sylow subgroup. This result appears in [3, Proposition 9.1]. We remind the reader of the definition of a regular number. If W is a Coxeter group, a *regular element* is an element that acts regularly on the reflection representation, and a *regular number* is an number that is the order of a regular element. These are enumerated in [28, Section 5].

Lemma 7.4. *Let p be a prime and let q be a power of a prime $r \neq p$, and let $G(q)$ be a finite group of Lie type. Suppose that the order of q modulo p is a regular number, and that the Sylow p -subgroup of $G(q)$ is cyclic. If u is a generator for the Sylow p -subgroup of $G(q)$, then the centralizer $C_G(u)$ is abelian.*

8 Classical groups in cross characteristic

In this section we consider the case where G_0 is a central extension of a classical group in characteristic $r \neq p$.

In [25], Di Martino and Zalesski solve the problem of which elements of quasisimple classical groups act minimally actively on the Weil modules (in fact, they do all almost cyclic elements). However, they only allow u to induce an inner-diagonal outer automorphism on G_0 if it is linear or unitary, and only an inner automorphism if G_0 is symplectic. The theorem in [25], applied to minimally active modules only (i.e., where u is a p -element and the characteristic of the field is p), is as follows.

Theorem 8.1. *Let G_0 be one of $\mathrm{SL}_n(q)$ ($n \geq 3$), $\mathrm{SU}_n(q)$ or $\mathrm{Sp}_{2n}(q)$, and let u either be in G_0 or induce an inner-diagonal automorphism on G_0 if G_0 is not symplectic. Suppose that G_0 is not one of the groups considered in Proposition 7.1. If u acts minimally actively on a Weil module, then one of the following holds:*

- (i) $G = \mathrm{Sp}_{2n}(q)$, n is a power of 2, $p^a = (q^n + 1)/2$ for some $a \geq 1$, $o(u) = p^a$;
- (ii) $G = \mathrm{Sp}_{2n}(3)$, $n \neq p$ is an odd prime and $p^a = (3^n - 1)/2$ for some $a \geq 1$, with $o(u) = p^a$;
- (iii) $G_0 = \mathrm{SU}_n(q)$, $n \neq p$ is an odd prime at least 5, $p^a = (q^n + 1)/(q + 1)$ for some $a \geq 1$, and $o(u) = p^a$;
- (iv) $G = \mathrm{SU}_3(3)$, $p = 7$, $o(u) = 7$;
- (v) $G_0 = \mathrm{SL}_n(q)$, $n \neq p$ is an odd prime, $p^a = (q^n - 1)/(q - 1)$ for some $a \geq 1$, and $o(u) = p^a$.

We will add to this by proving the following result.

Proposition 8.2. *Let G_0 be a central extension of a simple special linear, unitary or symplectic group, but not one of the groups in Proposition 7.1. Let $u \in G$ be a p -element, and let M be a simple module on which u acts minimally actively. If $M \downarrow_{G_0}$ involves a Weil module, then u induces an inner-diagonal automorphism on G_0 .*

We begin by proving, for G_0 classical and not of type PSL_2 , that if M is not a Weil module then the possibilities for an element of G acting minimally actively are very limited, restricted mostly to cases of exceptional Schur multipliers given in Proposition 7.1.

Table 4 is a summary of what we will need about the dimensions of Weil modules, and lower bounds for the dimensions of non-Weil modules for classical groups, assuming that G_0 is not one of the groups in Proposition 7.1. (Let κ_n be 1 if p divides $(q^n - 1)/(q - 1)$ and 0 otherwise.)

8.1 $\mathrm{SL}_n(q)$, $n \geq 3$

For this section we let G_0 be a quotient of $\mathrm{SL}_n(q)$ for $n \geq 3$, and we exclude the cases of $\mathrm{PSL}_3(2) = \mathrm{PSL}_2(7)$, $\mathrm{PSL}_4(2) = \mathrm{Alt}_8$, and $\mathrm{PSL}_3(4)$ which are considered in Propositions 4.1 and 7.1. Suppose that M is a non-trivial simple module, but not a Weil module. From Table 4, the dimension of M is at least $(q^{n-1} - 1)((q^{n-2} - 1)/(q - 1) - 1)$ for $n \geq 5$.

Note that if u is a p -element of G then, as we saw in Proposition 7.3 and the discussion afterwards, the order of u is bounded by a product of numbers: the exponent of the Sylow p -subgroup of the outer automorphism group (the outer contribution); the exponent of the Sylow p -subgroup of the Weyl group (only if the Sylow p -subgroup of G_0 is non-abelian, the Weyl contribution); the p -part of $\Phi_d(q)$ (the toral contribution).

Group	Bound	Reference
$\mathrm{SL}_n(q)$ (Weil)	$(q^n - q)/(q - 1) - \kappa_n$	[9]
$\mathrm{SL}_n(q)$ (Weil)	$(q^n - 1)/(q - 1)$	[9]
$\mathrm{SL}_n(q)$, $n = 3, 4$ (non-Weil)	$(q - 1)(q^{n-1} - 1)/\mathrm{gcd}(n, q - 1)$	[9]
$\mathrm{SL}_n(q)$, $n \geq 5$ (non-Weil)	$(q^{n-1} - 1)((q^{n-2} - q)/(q - 1) - \kappa_{n-2})$	[9]
$\mathrm{SU}_n(q)$ (Weil)	$(q^n + q(-1)^n)/(q + 1)$	[10]
$\mathrm{SU}_n(q)$ (Weil)	$(q^n - (-1)^n)/(q + 1)$	[10]
$\mathrm{SU}_3(q)$ (non-Weil)	$(q - 1)(q^2 + 3q + 2)/6$	[10]
$\mathrm{SU}_4(q)$ (non-Weil)	$(q^2 + 1)(q^2 - q + 1)/2 - 1$	[10]
$\mathrm{SU}_n(q)$, $n \geq 5$ (non-Weil)	$q^{n-2}(q - 1)(q^{n-2} - q)/(q + 1)$	[10]
$\mathrm{Sp}_{2n}(q)$, q odd (Weil)	$(q^n \pm 1)/2$	[7]
$\mathrm{Sp}_{2n}(q)$, all q (non-Weil)	$q(q^n - 1)(q^{n-1} - 1)/2(q + 1)$	[7], [27]
$\Omega_{2n+1}(q)$	$q^{n-1}(q^{n-1} - 1)$	[11]
$\Omega_{2n}^+(q)$	$q^{n-2}(q^{n-1} - 1)$	[19]
$\Omega_{2n}^-(q)$	$(q^{n-1} + 1)(q^{n-2} - 1)$	[19]

Table 4: Minimal dimension of a non-trivial projective representation for simple classical groups

We will let G_0 be a group $G(q^t)$ and assume that u induces an automorphism on G_0 that projects onto a field automorphism of order t in $\mathrm{Out}(G_0)$.

Proposition 8.3. *Suppose that G_0 is a central extension of a special linear group $\mathrm{PSL}_n(q^t)$ for some $n \geq 3$, with $(n, q^t) \neq (3, 2), (3, 4), (4, 2)$, and let u be a p -element of G such that $G = \langle G_0, u \rangle$. If u acts minimally actively on a non-trivial simple module M that is not a Weil module, then $G = \mathrm{PSL}_3(3)$, $p = 13$, $o(u) = 13$, and $\dim(M) = 11, 13$.*

Proof. Let G_0 be a central extension of $\mathrm{PSL}_n(q^t)$ for some $n \geq 3$ and $t \geq 1$, with the exclusions given above of $\mathrm{PSL}_3(2)$, $\mathrm{PSL}_3(4)$ and $\mathrm{PSL}_4(2)$. First, let $n = 3, 4$, and note that if M is a non-Weil simple module then $\dim(M) \geq (q^t - 1)(q^{(n-1)t} - 1)/\mathrm{gcd}(n, q^t - 1)$. If the Sylow p -subgroup of $\mathrm{SL}_n(q^t)$ is abelian then, in the notation introduced after Proposition 7.3, the Weyl contribution is 1, the toral contribution is at most $\Phi_d(q^t)$, where $d = 1, \dots, n$, and the outer contribution is t , since diagonal automorphisms are not of concern by Proposition 7.3. Thus in all cases, $o(u) \leq \Phi_3(q^t) \cdot t$, in fact $o(u) \leq \Phi_3(q) \cdot t^2$ by Lemma 7.2. From [8, Theorem 4.1], $\alpha(u) \leq n$ for $p > 2$, and we have a lower bound for $\dim(M)$ from Table 4. The equation $\dim(M) \leq \alpha(u) \cdot (o(u) - 1)$ for minimally active M from Lemma 2.3(iv) now yields

$$(q^t - 1)(q^{(n-1)t} - 1)/\mathrm{gcd}(n, q^t - 1) \leq n((q^2 + q + 1)t^2 - 1);$$

since $p \mid t$, we can assume that t is odd, and all solutions are for $t = 1$, with $q = 2, 3, 4, 7$ for $n = 3$ and $q = 2, 3$ for $n = 4$. Removing those excluded from the start of this section, for $t = 1$ we need consider $(n, q) = (3, 3), (3, 7), (4, 3)$. For $\mathrm{PSL}_3(7)$, the only prime for which the Sylow p -subgroup is abelian is $p = 19$, $\dim(M) \geq 96$ (by Table 4) and $\alpha(u) \leq 3$, so it in fact fails the bound. For $\mathrm{PSL}_4(3)$, $\dim(M) \geq 26$ (again, table 4) and $p = 5, 13$, with $o(u) = p$. Since $\alpha(u) \leq 4$, this shows that $p = 5$ cannot yield a minimally active module, and for $p = 13$ we see that $C_{G_0}(u) = Z(G_0) \cdot \langle u \rangle$, so that $\dim(M) \leq p + 1$ for M to be minimally active by Lemma 2.6. Thus there are no examples here.

For $\mathrm{PSL}_3(3)$ we have $p = 13$, and again $C_{G_0}(u) = \langle u \rangle$ (there are no central extensions) so we may apply

Lemma 2.6. The simple modules from [14] have dimensions 1, 11, 13, 16 and 26, so those of dimensions 11 and 13 are minimally active.

Suppose that the Sylow p -subgroups are non-abelian, so that $p = 2, 3$. First assume $p = 3$: the exponent of the Sylow 3-subgroup of the Weyl group is 3, and the toral contribution of $o(u)$ is at most $q^t + 1$, in fact $t(q + 1)$ by Lemma 7.2. The outer contribution is at most t , so that $o(u) \leq 3t^2(q + 1)$. Thus our equation $\dim(M) < \alpha(u) \cdot o(u)$ becomes

$$(q^t - 1)(q^{(n-1)t} - 1) / \gcd(n, q^t - 1) < 3nt^2(q + 1);$$

which yields only $t = 1$ and $(n, q) = (3, 2), (3, 4), (4, 2)$, all of which are excluded.

For $p = 2$, we have a graph automorphism to consider as well. The toral contribution to $o(u)$ is at most $q + 1$, the Weyl contribution is at most n and the outer contribution if the lowest common multiple of 2 and t .

Hence $\alpha(u) \cdot o(u)$ is at most $nmt(q + 1) \cdot \text{lcm}(2, t)$, where $m = n$ for $n \geq 5$, $m = 4$ for $n = 3$ and $m = 6$ for $n = 4$. We will check both Weil and non-Weil modules simultaneously, and all n , so we need

$$nmt(q + 1) \cdot \text{lcm}(2, t) \geq \alpha(u) \cdot o(u) > \dim(M) \geq (q^{nt} - q^t) / (q^t - 1) - 1.$$

For $n \geq 5$ we only get $(n, q, t) = (5, 3, 1)$. For $n = 4$ we get $(n, q, t) = (4, 3, 1), (4, 5, 1)$, and for $n = 3$ we get $q^t \leq 23$.

For $\text{PSL}_5(3)$, the exponent of the Sylow 2-subgroup is 16, so that $o(u) \leq 32$, $\alpha(u) \leq 5$, and $\dim(M) \geq 120$. Thus we need $o(u) = 32$, and in this case $v = u^2$ is an element of $\text{PSL}_5(3)$ of order 16. However, a simple computer check confirms that $\alpha(v) = 2$ for these elements, so that there is no minimally active module.

For $\text{PSL}_4(3)$ and $\text{PSL}_4(5)$, we have $\alpha(u) \leq 4$ if $\langle u \rangle \cap G_0$, generated by v say, is non-trivial. Since the exponent of a Sylow 2-subgroup of both groups is 8, we get that $o(u) \leq 16$, and $\dim(M) \geq 26, 124$ respectively. This eliminates $\text{PSL}_4(5)$ as $\dim(M) > o(u) \cdot \alpha(u)$ (using Lemma 2.3(iv)). For $\text{PSL}_4(3)$, if $o(v) = 8$ then $\alpha(v) = 2$ and if $o(v) = 4$ then $\alpha(v) \leq 3$. Thus we can only get a minimal action of u on M if $\dim(M) = 26$ (the next smallest is dimension 38) and $o(u) = 16$, with u therefore inducing the graph automorphism on M . However, there is no element of $\text{Aut}(\text{PSL}_4(3))$ of order 16, as we see from [2, pp.68–69]. Thus we get no minimally active modules here either.

Finally, consider $n = 3$. If $t = 1$ then $\alpha(u) \leq 3$, and also the Weyl contribution to $o(u)$ is 2, not n which is 3. Thus in this case we get $o(u) \cdot \alpha(u)$ to be at most $12(q + 1)$, which yields $q \leq 13$ for there to be a minimally active module. If we replace $q + 1$ by the 2-part of $(q^2 - 1)/2$, which is the actual toral contribution, we obtain $q \leq 9$. For these groups we check in [2] that the exponents of the Sylow 2-subgroups of $\text{Aut}(\text{PSL}_3(q))$ are 8, 8, 16, 16, as $q = 3, 5, 7, 9$ respectively. Since $\dim(M) \geq 12, 30, 56, 90$ from the table above, and $\alpha(u) \leq 3$, we see that in fact $G_0 = \text{PSL}_3(3)$ is the only possibility. In this case, if $u \in G_{0.2}$ has order 4 or 8, we use a computer to check that $\alpha(u) = 2$, and if $u \in G_{0.2}$ has order 2 then $\alpha(u) \leq 4$ by [8, Theorem 4.1]. Since if M is minimally active then $\dim(M) \leq \alpha(u) \cdot (o(u) - 1)$, we see that $\dim(M) \leq 14$, so the 12-dimensional simple module is the only possibility, with $o(u) = 8$ (the simple modules for G_0 have dimensions 1, 12, 16 and 26). However, both classes of elements of order 8 in $G_{0.2}$ have type $(8, 4)$ on M of dimension 12, so it is not minimally active.

Thus let $n \geq 5$ and p be odd. If M is not a Weil module then $\dim(M) \geq (q^{(n-1)t} - 1) ((q^{(n-2)t} - q^t) / (q^t - 1) - 1)$, and for all u , $\alpha(u) \leq n$. If the Sylow p -subgroup of G_0 is abelian then $o(u)$ is at most $(q^{nt} - 1) / (q - 1) \cdot t$, and using the formula $\dim(M) < o(u) \cdot \alpha(u)$ yields only $\text{PSL}_5(2)$. As $\dim(M) \geq 75$ for non-Weil modules

from the formula above, and the order of prime-power elements of $\mathrm{PSL}_5(2)$ is at most 8 or 31, we see that $o(u) = 31 = \Phi_5(2)$. But in this case $C_{G_0}(u) = \langle u \rangle$, so that $\dim(M) \leq p + 1 = 32$, and there are therefore no examples.

Thus the Sylow p -subgroup of G_0 is non-abelian, and therefore p divides two separate Φ_d -tori: from Lemma 7.2, we see therefore that if q^t has order d modulo p , $dp \leq n$, and therefore the toral contribution to u is at most $q^{nt/p} - 1$, with the Weyl contribution at most n and the outer contribution t . We therefore have that $o(u) \leq nt(q^{\lfloor nt/3 \rfloor} - 1)$ (as $p \geq 3$), and using the formula yields

$$n^2 t (q^{\lfloor nt/3 \rfloor} - 1) > (q^{(n-1)t} - 1) \left((q^{(n-2)t} - q^t) / (q^t - 1) - 1 \right),$$

which has no solutions for $n \geq 5$. This completes the proof. \square

Having determined which non-Weil modules can be minimally active, we turn our attention to the Weil modules for odd primes p , where u induces a non-diagonal outer automorphism, which must involve a field automorphism of order at least 3.

As in the proof of the previous proposition, the order of an element of G_0 is at most either $t(q^n - 1)$ or $tn(q^{\lfloor n/3 \rfloor} - 1)$, and we multiply this by the outer contribution, which is t , and $\alpha(u)$, which is at most n , to get an estimate for $o(u) \cdot \alpha(u)$. We then compare that to $(q^{nt} - q^t)/(q^t - 1)$ for $t \geq 3$, and find only one possible solution: $\mathrm{PSL}_3(8).3$, of course with $p = 3$. In this case, $\dim(M) \geq 72$ by [14, p.187], and $o(u) \leq 9$ by [2, p.74], so this cannot work either.

Thus if $u \in G$ acts minimally actively on a Weil module, then it induces an inner-diagonal automorphism on G_0 , proving Proposition 8.2 for linear groups.

8.2 $\mathrm{SU}_n(q)$

This looks very similar to the linear case in the previous subsection. We start by dealing with non-Weil representations, with the cases $n = 3, 4$ having to be dealt with separately, and at the same time proving that there are no minimally active modules for $p = 2$, Weil or non-Weil. This then allows us to prove easily that u cannot act minimally actively on a simple module without inducing an inner-diagonal automorphism on G_0 , just as with the linear case.

Note that we exclude $\mathrm{PSU}_n(q)$ for $(n, q) = (3, 2), (4, 2), (4, 3), (6, 2)$. For $\mathrm{PSU}_3(3) = G_2(2)'$, we require $p \neq 2, 3$.

Proposition 8.4. *Suppose that G_0 is a central extension of a special unitary group $\mathrm{PSU}_n(q^t)$ for some $n \geq 3$, with $(n, q^t) \neq (3, 2), (4, 2), (4, 3), (6, 2)$, and let u be a p -element of G such that $G = \langle G_0, u \rangle$. If u acts minimally actively on a non-trivial simple module M , then M is a Weil module.*

Proof. Let G_0 be a central extension of $\mathrm{PSU}_n(q^t)$, excluding the groups listed above the proposition, and let M denote a non-Weil simple module. We first consider $n = 3$, where $\dim(M) \geq (q^t - 1)(q^{2t} + 3q^t + 1)/6$, and p is either 2 or 3, or divides one of $q^t - 1$, $q^t + 1$, or $q^{2t} - q^t + 1 = \Phi_6(q^t)$. We apply Lemma 7.2 to replace $\Phi_d(q^t)$ by $t\Phi_d(q)$, as in the case for $\mathrm{PSL}_n(q^t)$. Unless $q^t = 3$ and $p = 2$, from [8, Theorem 4.1] we have that $\alpha(u) \leq 3$.

If $p \neq 2, 3$, we have that $o(u) \leq t^2\Phi_d(q)$, with the Weyl contribution being trivial, the toral contribution being $t\Phi_d(q)$ for $d = 1, 2, 6$, and the outer contribution being t . (Note that diagonal automorphisms need not be considered, as in $\mathrm{PSL}_n(q^t)$, using Proposition 7.3.)

If $d = 1, 2$ then we only end up with $\mathrm{PSU}_3(3)$, and if $d = 6$ then since this is a regular number $C_{G_0}(u)$ is abelian by Lemma 7.4, and hence if M is minimally active then $\dim(M) \leq 2o(u)$ by Lemma 2.6. This forces

$t = 1$ and $q < 7$, so only $q = 3, 5$, since $q = 2, 4$ are excluded. For $q = 3, 5$, $q^2 - q + 1 = 7, 21$, and so $p = 7$ in both cases. However, this now excludes $q = 5$, leaving only $q = 3$. Here, if M is not a Weil module then $\dim(M) \geq 14$, and there are therefore no examples.

If $p = 3$ then the toral contribution is at most $q + 1$, the Weyl contribution is 3 and the outer contribution is t , so we place this in our formula to get only $(n, q, t) = (3, 2, 3), (3, 5, 1)$. The exponents of the Sylow 3-subgroups of $\text{Aut}(\text{PSU}_3(q^t))$ for $q^t = 5, 8$ are 3 and 9 respectively, whereas $\dim(M) \geq 20, 56$ respectively from [14]. Thus there are no minimally active modules for $p = 3$.

Suppose that $p = 2$. The toral contribution is at most $t(q + 1)$, the Weyl contribution is at most n , and the outer contribution is at most $2t$. Unless $n = 4$ or $G_0 = \text{PSU}_3(3)$, we have that $\alpha(u) \leq n$. Thus $\alpha(u) \cdot o(u) \leq 2t^2n^2(q + 1)$, and for this to be at least $\dim(M)$ (for any non-trivial M , not just non-Weil modules), for $n \geq 5$ we have $(n, q, t) = (5, 3, 1), (6, 3, 1)$. In these two cases, $\dim(M) \geq 60, 182$, whereas the exponent of the Sylow 2-subgroup of $\text{Aut}(\text{PSU}_5(3))$ is 16, and the exponent of the Sylow 2-subgroup of $\text{PSU}_6(3)$ is 16, so that of the automorphism group is at most 32. If $v \in \text{PSU}_n(3)$ for $n = 5, 6$ has order 8 or 16 then $\alpha(v) \leq 3$, so that M cannot be minimally active.

If $n = 4$ then $\alpha(u) \leq 6$, and this yields $q = 3, 5, 7$ for $t = 1$, and $q = 3$ for $t = 2$. For $q = 3, 5, 7, 9$, $\dim(M) \geq 20, 104, 300, 656$, with the exponents of the Sylow 2-subgroups of G_0 being 8, 8, 16, 16 respectively. Thus only $q = 3$ can yield a minimally active module, but $\text{PSU}_4(3)$ is excluded from consideration.

For $n = 3$ we need better bounds, because for $t = 1$ we get $q \leq 19$ satisfying the bound

$$2t^2n^2(q + 1) \geq (q^{nt} - q^t)/(q^t + 1),$$

and for $t = 2$ we get $q = 3$. The Weyl contribution (n in the above inequality) may be replaced by 2, and the toral contribution ($t(q + 1)$ above) may be replaced by the 2-part of $(q^{2t} - 1)/2$. Doing so yields $q^t \leq 9$, and replacing $o(u)$ with the correct exponents, which are 8, 8, 16, 16 for $q^t = 3, 5, 7, 9$ respectively, means that $q^t = 9$ can be excluded, as $\dim(M) \geq 6, 20, 42, 72$. If $\alpha(u) = 2$ for u of maximal order, then this will exclude $q^t = 5, 7$ as well: this can be checked and is indeed the case, yielding $G_0 = \text{PSU}_3(3) = G_2(2)'$, so already considered. This completes the proof for $p = 2$, all simple modules and all $n \geq 3$.

We now let $n = 4$, and now p is odd. If $p > 3$ then the Sylow p -subgroup is abelian, so $d = 1, 2, 4, 6$, and the toral contribution is at most $t(q^2 + 1)$; the Weyl contribution is 1; and the outer contribution is t . Since $\alpha(u) \leq n$ and $\dim(M) \geq (q^{2t} + 1)(q^{2t} - q^t + 1)/2 - 1$, plugged into $\alpha(u) \cdot o(u) > \dim(M)$ yields

$$nt^2(q^2 + 1) > (q^{2t} + 1)(q^{2t} - q^t + 1)/2 - 1,$$

which yields $t = 1$ and $q = 2, 3$, both of which are excluded from consideration.

If $p = 3$ then the Weyl contribution is 3 and the toral contribution is at most $t(q + 1)$, with t the outer contribution, yielding

$$3nt^2(q + 1) > (q^{2t} + 1)(q^{2t} - q^t + 1)/2 - 1.$$

Again, only $q^t = 2, 3$ satisfy this, which have been excluded.

Thus $n \geq 5$. First suppose that the Sylow p -subgroup is abelian. The toral contribution is at most $t(q^n - 1)/(q - 1)$ (as this is greater than $t(q^n + 1)/(q + 1)$, and the order d of q^t modulo p is either at most n or $2m$ for some odd $m \leq n$), the Weyl contribution is 1, and the outer contribution is at most t . For M a non-Weil module, the inequality $\alpha(u) \cdot o(u) < \dim(M)$ becomes

$$nt^2(q^n - 1)/(q - 1) > q^{(n-2)t}(q^t - 1)(q^{(n-2)t} - q^t)/(q^t + 1),$$

and this forces $t = 1$ and $(n, q) = (5, 2), (5, 3), (6, 2), (7, 2), (8, 2)$. To eliminate these, we first ignore the ts , and then produce better estimates for the toral contribution than $(q^n - 1)/(q - 1)$: for $n = 5, 7$ we get $(q^n + 1)/(q + 1)$, which eliminates $(5, 3)$ and $(7, 2)$, and for $n = 8$ we use $(q^7 + 1)/(q + 1)$, which eliminates $(8, 3)$. Since $(6, 2)$ is not being considered in this proposition, we are left with $\text{PSU}_5(2)$. Here $p = 5, 11$, $\dim(M) \geq 43$ for non-Weil modules from [14, pp.182–184], and it is easy to see that $\alpha(u) = 2$ for $o(u) = 5, 11$ by a computer check (alternatively we can use the fact that $C_G(u)$ is abelian and apply Lemma 2.6). Thus there are no minimally active non-Weil modules in this case.

If the Sylow p -subgroup is non-abelian then, as with the linear case, the toral contribution is at most $t(q^{\lfloor n/3 \rfloor} + 1)$ and the Weyl contribution is at most n , yielding

$$n^2 t^2 (q^{\lfloor n/3 \rfloor} + 1) > q^{(n-2)t} (q^t - 1) (q^{(n-2)t} - q^t) / (q^t + 1),$$

where we get $t = 1$ and $(n, q) = (5, 2), (6, 2)$, although $\text{PSU}_6(2)$ is excluded. For $G_0 = \text{PSU}_5(2)$, only the Sylow 3-subgroup (and the Sylow 2-subgroup of course) is non-abelian, and $\text{Out}(G_0)$ has order 2, we have $G_0 = G$ and so u has order at most 9, and $\alpha(u) = 2$ if $o(u) = 9$, and $\alpha(u) \leq 5$ if $o(u) = 3$, with $\dim(M) \geq 44$ from [14, p.181]. Thus there is no non-Weil simple minimally active module for this group. \square

As with linear groups, we now check that if u induces an automorphism that is not inner-diagonal on G_0 then u does not act minimally actively on a Weil module. From the previous proposition we may assume that p is odd, so that $t \geq 3$.

Suppose that the Sylow p -subgroup of G_0 is abelian: as in the proof of the proposition we see that $o(u) \leq t^2(q^n - 1)/(q - 1)$, and so we get

$$n t^2 (q^n - 1) / (q - 1) > (q^{nt} - q^t) / (q^t + 1),$$

yielding $(n, q) = (3, 2), (4, 2)$ for $t = 3$, and no solutions for $t \geq 5$. Thus $p = 3$, but the Sylow 3-subgroup of G_0 is definitely not abelian.

If the Sylow p -subgroup of G_0 is non-abelian then the toral contribution is at most $t(q^{\lfloor n/3 \rfloor} + 1)$, the Weyl contribution is at most n , and this time we get

$$n^2 t^2 (q^{\lfloor n/3 \rfloor} + 1) > (q^{nt} - q^t) / (q^t + 1),$$

and this yields $(n, q, t) = (3, 2, 3)$ as the only solution, so again $p = 3$. The Sylow 3-subgroup of $\text{Aut}(\text{PSU}_3(8))$ has exponent 9, and the dimension of a Weil module is 56, with $\alpha(u) \leq 3$ by [8, Theorem 4.1], so u cannot act minimally actively on a Weil module by Lemma 2.3(iv).

This completes the proof of Proposition 8.2 for unitary groups.

8.3 $\text{Sp}_{2n}(q)$

For this section we let G_0 be a quotient of $\text{Sp}_{2n}(q^t)$, and we exclude the cases of $\text{Sp}_4(2) = \text{Sym}_6$, $\text{PSp}_4(3) = \text{PSU}_4(2)$ and $\text{Sp}_6(2)$ (the last two appear in Proposition 7.1). Suppose that M is a non-trivial simple module, but not a Weil module, which exist only for odd q . From the table near the start of this section, the dimension of M is at least $q^t(q^{nt} - 1)(q^{(n-1)t} - 1)/2(q^t + 1)$.

Proposition 8.5. *Suppose that G_0 is a central extension of a symplectic group $\text{PSp}_{2n}(q^t)$ for some $n \geq 2$, with $(n, q^t) \neq (4, 2), (4, 3), (6, 2)$, and let u be a p -element of G such that $G = \langle G_0, u \rangle$. If u acts minimally actively on a non-trivial simple module M that is not a Weil module, then $G = \text{Sp}_4(4)$, $p = 17$, $o(u) = 17$, and $\dim(M) = 18$.*

Proof. Let $G_0 = \mathrm{Sp}_{2n}(q^t)$ for some $n \geq 2$, some prime power q and some $t \geq 1$. As in previous sections, u induces an automorphism that projects in $\mathrm{Out}(G_0)$ to a field automorphism of order t . Since M is not a Weil module, we have $\dim(M) \geq (q^{nt} - 1)(q^{nt} - q^t)/2(q^t + 1)$. Let d be the order of q^t modulo p .

If $d = 2n$ then u is regular, so $\dim(M) < 2o(u)$ by Lemma 7.4. We have that $o(u) \leq t^2(q^n + 1)$ by Lemma 7.2, since the outer contribution is t and the toral contribution is at most $t(q^n + 1)$, so we get

$$2t^2(q^n + 1) \geq (q^{nt} - 1)(q^{nt} - q^t)/2(q^t + 1).$$

If $t = 1$ then the solutions to this are $q \leq 5$ for $n = 2$, and $q = 2$ for $n = 3$. If $t > 1$ then we only get $(n, q, t) = (2, 2, 2)$, but this needs $p \mid t = 2$ and $p \nmid q = 4$, a contradiction. We of course exclude $(n, q) = (2, 2), (2, 3), (3, 2)$ for $t = 1$, as we stated above, so we are left with $(n, q) = (2, 4), (2, 5)$ for $t = 1$. Here $d = 4$, and $\Phi_d(q) = q^2 + 1$: $4^2 + 1 = 17$ and $5^2 + 1 = 26$. For $q = 5$ this means that $o(u) = 13$, and $\dim(M) \geq 40$, so this cannot work, but for $\mathrm{Sp}_4(4)$, the module of dimension 18 could be minimally active. Since $C_G(u) = \langle u \rangle$ in this case, it is minimally active by Lemma 2.6.

Suppose that the Sylow p -subgroup is abelian. If $d \neq n$, then we have that $o(u) \leq t^2(q^d + 1)$ with $d \leq n - 1$, and note that $n \geq 3$. We also have that $\alpha(u) \leq n + 3$ by [8, Theorem 4.3], so using Lemma 2.3(iv) we get

$$(n + 3) \cdot t^2(q^d + 1) \geq \alpha(u)o(u) \geq \dim(M) \geq (q^{nt} - 1)(q^{nt} - q^t)/2(q^t + 1),$$

and the only solutions are for $t = 1$, with $(n, q) = (3, 2), (4, 2)$. The first can be ignored using Proposition 7.1, and for the second we have $d = 1, 2, 3, 4, 6$, which yield $\Phi_d(2) = 1, 3, 7, 5, 3$. Since the Sylow 3-subgroup of $\mathrm{Sp}_8(2)$ is non-abelian, we only get the cases $d = 3, 4$, so $p = 7, 5$. For $p = 7$ it is easy to check with a computer that $\alpha(u) = 2$, $o(u) = 7$, and $\dim(M) \geq 35$ by the degree bound above. Thus there is no (non-trivial) minimally active simple module for this group. For $p = 5$ we have $o(u) = 5$ and $\alpha(u) \leq 3$, so that again there are no examples.

We also need to consider the case where $d = n$, so that $o(u) \leq t^2(q^n - 1)/(q - 1)$. Using $\alpha(u) \leq (n + 3)$ and Lemma 2.3(iv), we get that if M is minimally active then

$$(n + 3) \cdot t^2(q^n - 1)/(q - 1) \geq \alpha(u)o(u) > \dim(M) \geq (q^{nt} - 1)(q^{nt} - q^t)/2(q^t + 1),$$

which for $t = 1$ yields the solutions $(n, q) = (2, 2), (2, 3), (2, 4), (3, 2), (4, 2), (5, 2)$, and for $t \geq 2$ yields only the solution $(n, q) = (2, 2)$ for $t = 2$, which we noticed earlier is not an example because $p \mid t = 2$ and $p \nmid q = 4$. If n is odd then n is a regular number, so we may replace $n + 3$ by 2, as in the previous case, and this removes the case $(5, 2)$. The cases $(2, 2), (2, 3), (3, 2)$ are excluded from our analysis, and $(4, 2)$ has been dealt with, leaving only $\mathrm{Sp}_4(4)$, with $p \mid \Phi_2(4) = 5$. We have $\dim(M) \geq 18$, $o(u) = 5$ and $\alpha(u) \leq 3$ by a computer calculation, so there is no example here either.

We may therefore assume that the Sylow p -subgroup is non-abelian, and hence p divides the order of the Weyl group of type C , which is $Z_2 \wr \mathrm{Sym}_n$, and $p \leq n$ with p dividing two separate tori, so that $p \mid \Phi_d(q)$ and $p \mid \Phi_{pd}(q)$. If p is odd then in particular this means that $d \leq n/3$ or d is even and $d \leq 2n/3$, so in either case the toral contribution is at most $t(q^d + 1)$. The Weyl contribution is at most n , and the outer contribution is t , so $o(u) \leq nt^2(q^d + 1)$. This is of course maximized at $d = \lfloor n/3 \rfloor$. As $\alpha(u)$ is still at most $n + 3$, we get

$$(n + 3) \cdot nt^2(q^{\lfloor n/3 \rfloor} + 1) \geq \alpha(u)o(u) > \dim(M) \geq (q^{nt} - 1)(q^{nt} - q^t)/2(q^t + 1),$$

The only solution is $(n, q, t) = (3, 2, 1)$, which we have already excluded.

Thus $p = 2$. Here there is a diagonal automorphism, but we do not need to consider these by Proposition 7.3, so the outer contribution is t . The Weyl contribution is $2n$, and the toral contribution is at most $t(q + 1)$

by Lemma 7.2, so the order of a 2-element of G_0 is at most $2nt^2(q+1)$ by Proposition 7.3. Since $\alpha(u) \leq 2n$, this yields $\alpha(u) \cdot (o(u) - 1) \leq 2n(2nt^2(q+1) - 1)$. If M is minimally active then, by Lemma 2.3(iv)

$$2n(2nt^2(q+1) - 1) \geq \alpha(u)(o(u) - 1) \geq \dim(M) \geq (q^{nt} - 1)(q^{nt} - q^t)/2(q^t + 1),$$

which has no solutions for $t \neq 1$, and for $t = 1$ we get $(n, q) = (2, 2), (2, 3), (2, 5), (3, 2)$, with the first and last eliminated since q must be odd. Since $\mathrm{Sp}_4(3)$ is also excluded, this leaves $G_0 = \mathrm{Sp}_4(5)$: from [2, p.63] we see that $o(u) \leq 8$, and $\alpha(u) \leq 4$ with $\dim(M) \geq 40$, so M cannot be minimally active. This completes the proof. \square

We now have to complete the proof of Proposition 8.2 by checking that if u induces an outer automorphism on $G_0 = \mathrm{Sp}_{2n}(q^t)$ then u cannot act minimally actively on a Weil module. First suppose that p is odd, so that u induces a field automorphism and $t \geq 3$.

The dimension of M is $(q^{nt} \pm 1)/2$, (recall that q must be odd) and as we saw above, if the Sylow p -subgroup of G_0 is abelian then $o(u) \leq t^2(q^n + 1)$, and as $\alpha(u) \leq n + 3$, if u acts minimally actively then by Lemma 2.3(iv) we have

$$t^2(n+3)(q^n + 1) \geq (q^{nt} - 1)/2,$$

which yields $(n, q, t) = (2, 3, 3)$, but of course t , which is a power of p , cannot divide q , so we get no examples.

If the Sylow p -subgroup is non-abelian, then $\alpha(u)o(u) \leq (n+3)nt^2(q^{\lfloor n/3 \rfloor} + 1)$, as we saw in the proof of the previous proposition: thus we have

$$(n+3)nt^2(q^{\lfloor n/3 \rfloor} + 1) \geq (q^{nt} - 1)/2,$$

and this has no solutions.

We thus reduce to $p = 2$. In this case, from [7, Section 5], we see that there are two Weil modules, which have dimension $(q^n - 1)/2$ and are swapped by the diagonal automorphism. By Theorem 8.1, $v \in G_0$ cannot act on these Weil modules with a single Jordan block, and hence by Lemma 2.7 if u induces a diagonal automorphism on G_0 then it cannot act minimally actively on the sum of the two Weil modules.

Thus u acts as either a field automorphism or the product of a field and diagonal (whichever stabilizes the two Weil modules), but in either case $t \geq 2$.

We have already bounded $o(u)$ by $2nt^2(q+1)$, so with $\alpha(u) \leq 2n$ we get

$$2n(2nt^2(q+1) - 1) \geq \alpha(u) \cdot (o(u) - 1) \geq \dim(M) \geq (q^{nt} - 1)/2.$$

If $t \geq 4$ then there are no solutions, and for $t = 2$ we get solutions $(n, q) = (2, 3), (2, 5), (3, 3)$. The exponents of the Sylow 2-subgroups of $\mathrm{PSP}_4(9)$, $\mathrm{PSP}_4(25)$ and $\mathrm{PSP}_6(9)$ are 8, 8, 16 respectively, so $o(u) \leq 16, 16, 32$ respectively. The dimensions of the Weil modules are 40, 312, 364 respectively, and $\alpha(u) \leq 4, 4, 6$ respectively, so the formula $\dim(M) < \alpha(u) \cdot o(u)$ eliminates the second and third options from being minimally active. Finally, for $\mathrm{PSP}_4(9)$, if we can reduce $\alpha(u)$ for u of order 16 (hence $v = u^2 \in G_0$ of order 8) to 2 then we are done: this is the case by an easy computer calculation, and we complete the proof of Proposition 8.2.

8.4 $\Omega_{2n+1}(q)$ and $\Omega_{2n}^\pm(q)$

As we saw in Table 4, the minimal degree for $\mathrm{Spin}_{2n+1}(q)$ for $(n, q) \neq (3, 3)$ is $q^{n-1}(q^{n-1} - 1)$.

Recall that the polynomial order of $\mathrm{Spin}_{2n+1}(q)$ is

$$q^{n^2} \prod_{i=1}^n (q^{2i} - 1),$$

so that if $p \nmid q$ divides the order of $\text{Spin}_{2n+1}(q)$, p divides $q^d \pm 1$ for some $1 \leq d \leq n$.

Proposition 8.6. *Let G_0 be a central extension of one of the groups $\Omega_{2n+1}(q)$ for $(n, q) \neq (3, 3)$ and $n \geq 3$. Let u be a p -element of G such that $G = \langle G_0, u \rangle$. There are no non-trivial minimally active simple modules for G .*

Proof. Let $G_0 = \Omega_{2n+1}(q^t)$ for some $n \geq 3$, some prime power q , and some $t \geq 1$, and suppose that the Sylow p -subgroup of G_0 is abelian, so that p divides a single cyclotomic polynomial, and let d be the order of q^t modulo p . The order of u is at most $t \cdot t\Phi_d(q) \leq t^2 \cdot (q^n + 1)$. As $\alpha(u) \leq n + 3$ by [8, Theorem 4.4], we get using Lemma 2.3(iv)

$$(n + 3) \cdot t^2(q^n + 1) > \alpha(u) \cdot o(u) > q^{t(n-1)}(q^{t(n-1)} - 1)$$

if M is minimally active, and this forces $t = 1$ and $(n, q) = (3, 3), (3, 5)$. Omitting the t from now on, replacing the upper bound $q^n + 1$ for $\Phi_d(q)$ by each of $(q^n + 1)/(q + 1)$, $(q^n - 1)/(q - 1)$, $(q - 1)$ and $(q + 1)$ eliminates $(n, q) = (3, 5)$, and $(n, q) = (3, 3)$ is excluded already, so there are no solutions.

We now may assume that the Sylow p -subgroup is non-abelian, so that if p is odd then $o(u) \leq t \cdot n \cdot t(q^d + 1)$ (as the Weyl group of type B is the Weyl group of type C we can use the Weyl contribution from Proposition 8.5), but with both d and pd dividing $2n$. We thus get

$$(n + 3) \cdot n \cdot t^2(q^d + 1) > \alpha(u) \cdot o(u) > q^{t(n-1)}(q^{t(n-1)} - 1),$$

for $d \leq n/3$, which is obviously maximized at $d = \lfloor n/3 \rfloor$, still with no solutions.

If $p = 2$, then we get $o(u) \leq 2t \cdot 2n \cdot t(q + 1)$ using Proposition 7.3 and the fact that the order of an outer automorphism is at most $2t$, and so now we have

$$2n \cdot 4nt^2(q + 1) > \alpha(u) \cdot o(u) > q^{t(n-1)}(q^{t(n-1)} - 1),$$

(as $\alpha(u) \leq 2n$ this time) which again has no solutions for $(n, q, t) \neq (3, 3, 1)$. This completes the proof. \square

Having dispensed with the odd-dimensional orthogonal groups, we turn to the even-dimensional ones. For $\Omega_{2n}^+(q)$, the minimal degree is $q^{n-2}(q^{n-1} - 1)$ (unless $G_0 = \Omega_8^+(2)$) and for $\Omega_{2n}^-(q)$ the minimal degree is $(q^{n-2} - 1)(q^{n-1} + 1)$, so in both cases $\dim(M) > (q^{n-1} - 1)(q^{n-2} - 1)$. If we use this bound then we can deal with both cases simultaneously. The polynomial order of $\text{Spin}_{2n}^\pm(q)$ is

$$q^{n(n-1)}(q^n \mp 1) \prod_{i=1}^{n-1} (q^{2i} - 1).$$

We already found minimally active modules for $2 \cdot \Omega_8^+(2)$ in Proposition 7.1, and the next proposition says that there are no more.

Proposition 8.7. *Let G_0 be a central extension of one of the groups $\Omega_{2n}^\pm(q)$ other than $\Omega_8^+(2)$, for $n \geq 4$. Let u be a p -element of G such that $G = \langle G_0, u \rangle$. There are no non-trivial minimally active simple modules for G .*

Proof. Our proof works the same as Proposition 8.6. If the Sylow p -subgroup is abelian then p is odd and $o(u) \leq t^2\Phi_d(q) \leq t^2(q^n + 1)$. Placing this in our standard formula from Lemma 2.3(iv), using $\alpha(u) \leq n + 3$ from [8, Theorem 4.4] gives

$$(n + 3)t^2(q^n + 1) \geq \dim(M) > (q^{t(n-1)} - 1)(q^{t(n-2)} - 1).$$

This bound yields no solutions for $t \geq 3$, and for $t = 1$ the solutions

$$(n, q) = (4, 2), (4, 3), (4, 4), (4, 5), (4, 7), (5, 2), (6, 2).$$

If q is odd then we may replace $(q^n + 1)$ by $(q^n + 1)/2$, removing $(4, 5)$ and $(4, 7)$ from the list.

If $d = n$ or $d = 2n$ then u is regular, so $C_{G_0}(u)$ is abelian by Lemma 7.4, and so in this case we may replace $\alpha(u) \cdot (o(u) - 1)$ by $2o(u)$ via Lemma 2.6, and so (removing the t , which is equal to 1 anyway)

$$2(q^n + 1) > (q^{n-1} - 1)(q^{n-2} - 1),$$

which only has a solution for $(n, q) = (4, 2)$. Thus we may assume that $d \neq n, 2n$, in which case we may replace $o(u)$ by $t^2(q^{n-1} + 1)$. Using this we reduce our possibilities to $(4, 2)$ and $(5, 2)$.

As $\Omega_8^+(2)$ is excluded, we just consider $G_0 = \Omega_8^-(2)$: from [14] we see that $\dim(M) \geq 33$, and $o(u) = 3, 5, 7, 9, 17$. Furthermore, $\alpha(u) = 2$ for $o(u) > 3$, and $\alpha(u) \leq 4$ for $o(u) = 3$, so there are no examples using the formula $\dim(M) \leq \alpha(u) \cdot (o(u) - 1)$.

For G_0 a central extension of $\Omega_{10}^\pm(2)$, $\dim(M)$ is at least the smallest of $2^{5-2}(2^{5-1} - 1) = 120$ and $(2^{5-1} + 1)(2^{5-2} - 1) = 119$, so $\dim(M) \geq 119$. For $\Omega_{10}^+(2)$, $o(u) \in \{3, 5, 7, 9, 17, 31\}$, and for $o(u) \geq 7$ we have $\alpha(u) = 2$, with $\alpha(u) \leq 3$ for $o(u) = 5$ and $\alpha(u) \leq 5$ for $o(u) = 3$, which shows that u cannot act minimally actively on a non-trivial M . For $\Omega_{10}^-(2)$, $o(u) \in \{3, 5, 7, 9, 11, 17\}$, and the same statements hold for $\alpha(u)$, so again u cannot act minimally actively on a non-trivial M .

Suppose that p is still odd, but that the Sylow p -subgroup is non-abelian. Thus p divides both $\Phi_d(q^t)$ and $\Phi_{dp}(q^t)$, and $d \leq n/3$. Since the Weyl group of type D is a subgroup of the Weyl group of type B , we see that the exponent of the Sylow p -subgroup of the Weyl group is at most n . Thus $o(u) \leq t^2 \cdot n \cdot (q^{\lfloor n/3 \rfloor} + 1)$ and $\alpha(u) \leq n + 3$, and thus we need to check

$$t^2 n(n + 3)(q^{\lfloor n/3 \rfloor} + 1) \geq (q^{t(n-1)} - 1)(q^{t(n-2)} - 1),$$

which only has solutions for $t = 1$, and then $(n, q) = (4, 2), (5, 2)$, which we have already checked. This completes the proof for p odd.

Suppose that $p = 2$, so that the order of u is at most $4t^2 \cdot 2n \cdot (q + 1)$: $\text{Out}(G_0)$ has exponent at most $4t$, the Weyl contribution is at most $2n$, and the toral contribution is at most $t(q + 1)$. (To see that the exponent of Sylow 2-subgroup of $\text{Out}(G_0)$ is at most $4t$ and not $8t$, note that if n is even then the diagonal automorphisms form $Z_2 \times Z_2$, so we are done, and if n is odd then the diagonal automorphisms form either Z_2 or Z_4 , with the graph automorphism inverting this group [5, Theorem 2.5.12(i)].) Since $\alpha(u) \leq 2n$, we get

$$16t^2 n^2 (q + 1) \geq (q^{t(n-1)} - 1)(q^{t(n-2)} - 1),$$

which has a solution only for $(n, q, t) = (4, 3, 1)$. However, although information on the Sylow 2-subgroups of $\text{Aut}(\Omega_8^\pm(3))$ is not available in [2], there are constructions of them on the online Atlas, and hence a computer algebra package immediately tells you that the exponent is 8, not 32 as suggested by the formula above. This proves that there are no minimally active modules for $p = 2$. \square

8.5 $\text{SL}_2(q)$

This short subsection deals with $G_0 = \text{SL}_2(q)$, where $p \nmid q$, $q \geq 4$ and $q \neq 4, 5, 7, 9$ (as these are alternating groups or are given in Proposition 7.1). In [25, Theorem 1.2], if $\text{SL}_2(q) \leq G \leq \text{GL}_2(q)$ then all possibilities for u acting minimally actively are determined, and given by the following lemma.

Lemma 8.8. *Let G_0 be a central extension of $\mathrm{PSL}_2(q)$ for $q \neq 4, 5, 7, 9$. Suppose that u induces an inner-diagonal automorphism on G_0 . If u acts minimally actively on M then one of the following holds:*

- (i) $G = \mathrm{SL}_2(q)$ for $q = 2^a$, $p = 2^a \pm 1$ is a Fermat or Mersenne prime, $o(u) = p$, M is any simple module;
- (ii) $G = \mathrm{SL}_2(q)$ for q odd, p is odd, $(q \pm 1)/2 = p^a$, $o(u) = p^a$, $\dim(M) \leq o(u) + 1$;
- (iii) $G = \mathrm{PSL}_2(q)$ or $\mathrm{PGL}_2(q)$, q is a Fermat or Mersenne prime, $p = 2$, and $\dim(M) \leq o(u) + 1$;

Thus we can assume that $u \in G$ acts as a field or product of a field and diagonal automorphism. Let G_0 be a central extension of $\mathrm{PSL}_2(q^t)$ for some prime power q and some $t \geq 2$ a power of p , with $G = \langle G_0, u \rangle$ and $|G : G_0| = t$. Note that $(q^t - 1)/2$ is the smallest dimension of a non-trivial simple module for G_0 if q is odd, and $q^t - 1$ is the smallest dimension if q is even.

Suppose that $\langle u \rangle \cap G_0 = 1$. If $o(u)$ is even then $\alpha(u) \leq 4$ by [8, Lemma 3.1], so $\dim(M) \leq 4(t - 1)$ by Lemma 2.3(iv). This yields $(q^t - 1)/2 \leq 4(t - 1)$, so $t = 2$ and $q^t \leq 9$. Having excluded 4, 5, 7, 9, and since 8 need not be considered, there are no solutions.

If $o(u)$ is odd then as $\alpha(u) = 2$ by [8, Lemma 3.1], we see that $\dim(M) \leq 2(t - 1)$, and $\dim(M) \geq (q^t - 1)/2$. As $t \geq 3$, we only get $q^t = 8$, but then the minimal degree is $q^t - 1$, not $(q^t - 1)/2$, and so there are no solutions here either.

Thus $u^t \neq 1$, and so $p \mid (q \pm 1)$. If $p = 2$ then the order of u is at most $t^2(q + 1)$, and $\alpha(u) = 3$ by [8, Lemma 3.1], so if u acts minimally actively on M then $\dim(M) < 3t^2(q + 1)/2$, whereas $\dim(M) \geq (q^t - 1)/2$. For $t = 2$ this yields $q \leq 11$, for $t = 4$ this yields $q = 3$, and there are no solutions for $t \geq 8$.

Replacing $t(q + 1)$ with the 2-part of $(q^{2t} - 1)/4$ (which is the exponent of the Sylow 2-subgroup of G_0) yields $q \leq 9$. Finally, in the remaining cases, one may check that the exponent of the Sylow 2-subgroup of $\mathrm{Aut}(\mathrm{PSL}_2(q^t))$ is 8, 16 and 16, for $q^t = 25, 49, 81$ respectively, and $\dim(M) \geq 12, 24, 40$ respectively. This eliminates the case where $q^t = 81$. Finally, in the remaining cases if $\dim(M) \geq q^t - 1$ then it cannot be minimally active, so it is only the two modules of dimension $(q^t - 1)/2$ that are important: for these, a computer calculation shows that any element v of G_0 acts on them with only blocks of size $o(v)$, whence u cannot act minimally actively.

Thus p is odd. First assume that q is even. The order of p is at most $t^2(q + 1)/2$, and $\alpha(u) = 2$ by [8, Lemma 3.1], so that if u acts minimally actively then $\dim(M) \leq t^2(q + 1)$, whereas $\dim(M) \geq q^t - 1$. The only solution to this is $q = 2$ and $t = p = 3$, which is the small Ree group ${}^2G_2(3)$, hence will not be considered as it is defining characteristic.

Hence we may assume that q is odd, in which case $\dim(M) \geq (q^t - 1)/2$. We still have that $\alpha(u) = 2$ and $o(u) \leq t^2(q + 1)/2$. This yields only one solution again, namely $q = 3$ and $t = 3$, but then this is defining characteristic and not in consideration. Thus there are no solutions when u does not induce an inner-diagonal automorphism.

9 Exceptional groups in cross characteristic

In this section we deal with G_0 a central extension of an exceptional group of Lie type. We start by dealing with a few small groups, which feature because they have exceptional Schur multipliers and so can have unusually small minimal faithful degrees, the analogue of Proposition 7.1.

Proposition 9.1. *Let G_0 be a central extension of one of the following simple groups: $G_2(3)$, $G_2(4)$, $F_4(2)$, ${}^2B_2(8)$. Let u be a p -element of G such that $G = \langle G_0, u \rangle$, and let M be a non-trivial simple module on which u acts minimally actively. One of the following holds:*

- (i) $G = G_2(3)$, $p = 13$, $o(u) = 13$ and $\dim(M) = 14$;
- (ii) $G = 2 \cdot G_2(4)$, $p = 13$, $o(u) = 13$ and $\dim(M) = 12$;
- (iii) $G = 2 \cdot {}^2B_2(8)$, $p = 13$, $o(u) = 13$ and $\dim(M) = 14$ (two representations).

The field of definition is always \mathbb{F}_{13} .

Proof. $G_2(3)$ has outer automorphism group of order 2 and Schur multiplier of order 3, with the outer automorphism of $3 \cdot G_2(3)$ inverting the centre, so we only have to consider the groups $G = G_2(3)$, $G = 3 \cdot G_2(3)$ and $G = G_2(3).2$. We simply check these one by one for $p = 2, 7, 13$, and get the single example above.

The group $G_2(4)$ has outer automorphism group of order 2 and Schur multiplier of order 2, so we need to consider $G_2(4)$ and $2 \cdot G_2(4)$ for p odd (so $p = 3, 5, 7, 13$), where the minimal degree is 12. The order of u must be p , and for $p = 5, 7, 13$ two conjugates of u generate G , so we can exclude $p = 3, 5$ by Lemma 2.3(iv) and focus on the 12-dimensional simple module for $2 \cdot G_2(4)$ for $p = 7, 13$. As the centralizer of an element of order 7 in $G_2(4)$ has order 21 we get $\dim(M) \leq 10$ by Lemma 2.6 and so can exclude this as well, leaving just $p = 13$, where for $G_2(4)$ the centralizer has order exactly 13, so an application of the same lemma shows that the simple module of dimension 12 is minimally active for u .

For $G = F_4(2)$, there is an exceptional Schur multiplier of order 2: for $G = 2 \cdot F_4(2)$, the character degrees are known for $p = 5, 7, 13, 17$, but the full set of character degrees is not known for $p = 3$. For $p \geq 5$, the minimal faithful degree is 52, and for $p = 3$ [27] states that it is at least 44. A computer calculation shows that G is generated by two conjugates of u for $o(u) = 5, 7, 9, 13, 17$, and so there are no minimally active modules for these elements. For $o(u) = 3$, three conjugates suffice, and so there are no minimally active modules here either.

Now let G_0 be a central extension of ${}^2B_2(8)$, where the exceptional Schur multiplier is a Klein four group, but all extensions $2 \cdot {}^2B_2(8)$ are isomorphic because the outer automorphism of order 3 permutes them. Here $p = 5, 7, 13$, as we can discount $p = 3$, since $o(u) = 3$ and $\alpha(u) \leq 3$ by [8, Proposition 5.8], and $\dim(M) \geq 14$. Thus we just check all simple modules for $G = 2 \cdot {}^2B_2(8)$ and $o(u) = 5, 7, 13$, noting that M is minimally active if and only if $\dim(M) \leq p + 1$ because $C_{G_0}(u) = Z(G_0) \cdot \langle u \rangle$ via Lemma 2.6. We find just the one example for $p = 13$. \square

In [8, Theorem 5.1], it is shown that $\alpha(G) \leq \ell + 3$ if G is of exceptional type, where ℓ is the untwisted rank of G (with one exception of G of type F_4 and $p = 2$, where $\alpha(G) \leq 8$).

We also need a bound on the orders of p -elements in an exceptional group of Lie type. Broadly speaking, by Proposition 7.3 if q has order d modulo p , then the order of a p -element is at most $q^d - 1$ multiplied by the exponent of the Weyl group. We give a general bound now.

Proposition 9.2. *Let G_0 be a central extension of one of $G_2(q)$, $F_4(q)$, $E_6(q)$, ${}^2E_6(q)$, $E_7(q)$ and $E_8(q)$ other than $G_2(2)$, with $p \nmid q$, and let u be a p -element of G . The order of u is at most $q^{\ell+1} - 1$, where ℓ is the untwisted rank of G .*

Proof. Let u be a p -element, and let d be the order of q modulo p . If p does not divide the order of the Weyl group, then $p \geq 5$ ($p \geq 7$ for $E_8(q)$) and the Sylow p -subgroup of the socle of G is abelian and has exponent at most $\Phi_d(q)$ by Proposition 7.3. The only outer automorphism of G that u can induce is a field automorphism, as diagonal and graph automorphisms have order 2 or 3, and field automorphisms have order less than $q - 1$.

Thus $o(u) \leq \Phi_d(q) \cdot (q-1)$. All we therefore need is an upper estimate for all $\Phi_d(q)$ where d divides one of the reflection degrees for G .

For $G = G_2$, the largest is $\Phi_3(q) = q^2 + q + 1$, so $o(u) \leq (q^3 - 1)$. For F_4 , the largest is $\Phi_8(q) = q^4 + 1$, so $o(u) \leq (q^4 + 1)(q-1) \leq q^5 - 1$. For E_6 the largest is $\Phi_9(q) = q^6 + q^3 + 1$, so $o(u) \leq (q^6 + q^3 + 1)(q-1) \leq (q^7 - 1)$. For ${}^2E_6(q)$ the largest is $\Phi_{18}(q) = q^6 - q^3 + 1$, so again we have $o(u) \leq q^7 - 1$. For E_7 the largest is $\Phi_7(q) = q^6 + q^5 + q^4 + q^3 + q^2 + q + 1$, so $o(u) \leq q^7 - 1$ again. Finally, for E_8 the largest is $\Phi_{30}(q) = q^8 + q^7 - q^5 - q^4 - q^3 + q + 1$, so that $o(u) \leq \Phi_{30}(q)(q-1) \leq q^9 - 1$. Thus in all cases, $o(u) \leq q^{\ell+1} - 1$.

Thus now p divides the order of the Weyl group, so $p \leq 7$, and we use Proposition 7.3 again. The exponents of the Sylow p -subgroups of the Weyl groups of exceptional groups are given below.

Group	Exponents
G_2	2, 3
F_4	8, 3
E_6	8, 9, 5
E_7	8, 9, 5, 7
E_8	8, 9, 5, 7

For $p = 7$, $G = E_7, E_8$ and either the Sylow p -subgroup is abelian, hence we are done by above, or $p \mid (q \pm 1)$, whence the order of u is at most $7(q \pm 1)(q-1) \leq 7(q^2 - 1)$, where the Weyl contribution is 7, the toral contribution is $(q \pm 1)$, and the outer contribution is at most $(q-1)$. The product of these is clearly less than $q^8 - 1$.

For $p = 5$, we have that the order of u is at most $5(q^2 + 1)(q-1) \leq 5(q^3 - 1)$, using the same argument, as $d = 1, 2, 4$. Of course, this is still smaller than $q^7 - 1$, which is the required bound as $G = E_6, E_7, E_8$.

For $p = 3$, the order of u depends on which group we are in. The toral contribution is at most $q + 1$, the Weyl contribution is 3 for G_2, F_4 and 9 otherwise, and the outer contribution is at most $3(q-1)$ (diagonal for ${}^\epsilon E_6$ and field automorphisms). For $G = G_2, F_4$ we get $3(q^2 - 1)$, which is at most $(q^3 - 1)$ for $G_2(q)$ (as $q \neq 2, 3$), and at most $q^5 - 1$ for all $q \geq 2$, so we get at most $q^{\ell+1} - 1$. For $G = E_6, E_7, E_8$, we get $27(q^2 - 1)$, which is less than $(q^7 - 1)$ for all $q \geq 2$, so in all cases again $q^{\ell+1} - 1$ will do.

For $p = 2$, we get that the toral contribution is at most $q + 1$, the Weyl contribution is 2 for G_2 and 8 for all other groups, and the outer contribution is at most $(q-1)$ for G_2, F_4 and E_8 , $2(q+1)$ for ${}^\epsilon E_6$, and $(q-1)$ for E_7 , by [5, Theorem 2.5.12], yielding at most $2(q^2 - 1)$ for G_2 , which is less than $q^3 - 1$ for $q \geq 3$.

For F_4 we get that $o(u)$ is at most $8(q+1) \leq (q^5 - 1)$, for ${}^\epsilon E_6$ we have $16(q^2 - 1)(q+1) \leq (q^7 - 1)$ for all $q \geq 3$, and for E_7, E_8 we have $o(u) \leq 16(q+1)(q-1) \leq (q^8 - 1)$, as needed. \square

Now we know that every p -element in G has order at most $q^{\ell+1} - 1$, then we get that $\dim(M) < \alpha(G) \cdot (q^{\ell+1} - 1)$, and we can apply the Landazuri–Seitz–Zalesskii bounds from [19] and [27].

Group	Landazuri–Seitz	LS evaluated	$\alpha(G)$	$\alpha(G) \cdot (q^{\ell+1} - 1)$ evaluated
F_4 ($q \geq 4$ even)	$q^7(q^3 - 1)(q-1)/2$	448 ($q = 4$)	≤ 8	≤ 248 ($q = 4$)
F_4 (q odd)	$q^6(q^2 - 1)$	5832 ($q = 3$)	≤ 7	≤ 1694 ($q = 3$)
${}^\epsilon E_6$	$q^9(q^2 - 1)$	1536 ($q = 2$)	≤ 9	≤ 1143 ($q = 2$)
E_7	$q^{15}(q^2 - 1)$	98304 ($q = 2$)	≤ 10	≤ 2550 ($q = 2$)
E_8	$q^{27}(q^2 - 1)$	402653184 ($q = 2$)	≤ 11	≤ 5621 ($q = 2$)

This proves that these groups have no non-trivial minimally active modules, but slightly better bounds are needed for the other groups, as the minimal faithful degrees are closer to q^ℓ .

Proposition 9.3. *Let G_0 be the simple group $G_2(q^t)$ for $q^t \geq 5$, and let u be a p -element of G such that $G = \langle G_0, u \rangle$. There are no non-trivial, minimally active modules for G .*

Proof. Let G_0 be a central extension of $G_2(q^t)$ for some prime power q and some $t \geq 1$, and let G be obtained by adding on a field automorphism of order t to G_0 . Since $G_2(2)' = \text{PSU}_3(3)$ we have already dealt with it, and we dealt with $G_2(3)$ and $G_2(4)$ in Proposition 9.1, we may assume that $q^t \geq 5$.

First suppose that $p \geq 5$, so that $t = 1$ or $t \geq 5$. From [8] we have that $\alpha(G) \leq 5$, and the Landazuri–Seitz bound [19] for G is $q^t(q^{2t} - 1)$. As $p \geq 5$, the order of u is at most one of $t\Phi_d(q)$ for $d = 1, 2, 3, 6$, with $d = 3$ maximizing this, so we get

$$5t^2(q^2 + q + 1) > \alpha(G) \cdot (o(u) - 1) \geq \dim(M) \geq q^t(q^{2t} - 1),$$

with $t = 1$ and $q = 5$ as solutions, and no solutions for $t \geq 5$. For $G = G_0 = G_2(5)$, the primes other than 2, 3, 5 dividing $|G|$ are 7 and 31, each dividing it exactly once, whence we need a simple module of dimension at most $4 \cdot (31 - 1) = 120$, but 124 is the minimal degree.

Suppose that $p = 3$, or $p = 2$ and q is not a power of 3. The toral contribution is at most $t(q + 1)$, the Weyl contribution is at most 3, and the outer contribution is t . Since $\alpha(u) \leq 5$, we get

$$15t^2(q + 1) > \alpha(G) \cdot (o(u) - 1) \geq \dim(M) \geq q^t(q^{2t} - 1),$$

which has no solutions for $q^t \geq 5$.

If $p = 2$ and q is a power of 3, then we get the toral contribution to be $t(q + 1)$, the Weyl contribution to be 2, and the outer contribution to be $2t$, so similar to the above expression, and we get

$$20t^2(q + 1) > q^t(q^{2t} - 1),$$

which has no solutions for $q^t \geq 9$. This completes the proof. \square

Proposition 9.4. *Let G_0 be the simple group ${}^3D_4(q^t)$ for some q and t , and let u be a p -element of G such that $G = \langle G_0, u \rangle$. There are no non-trivial, minimally active modules for G .*

Proof. Let $G_0 = {}^3D_4(q^t)$ for some prime power q and some $t \geq 1$ (there are no central extensions), and let G be obtained by adding on a field automorphism of order dividing $3t$ to G_0 (see [5, Theorem 2.5.12]). Note that $\alpha(G) \leq 7$ by [8, Proposition 5.7].

If $p \geq 5$ then the Sylow p -subgroup of G_0 is abelian, so let d be the order of q^t modulo p , so that $d = 1, 2, 3, 6, 12$. If $p \mid \Phi_{12}(q^t)$ then from the list of maximal subgroups in [16], we see that $C_{G_0}(u)$ is abelian, so $\dim(M) \leq 2o(u)$ if M is minimally active, by Lemma 2.6. Furthermore, $o(u) \leq t^2\Phi_{12}(q) = t^2(q^4 - q^2 + 1)$, and since $\dim(M) \geq q^{3t}(q^{2t} - 1)$ from [19], we get

$$2t^2(q^4 - q^2 - 1) \geq \dim(M) \geq q^{3t}(q^{2t} - 1),$$

which has only the solution $q^t = 2$, where $p = 13$. Here $\dim(M) \geq 26 = 2o(u)$ by [14, p.253] so there are no non-trivial minimally active modules here.

If $d = 1, 2, 3, 6$ then the toral contribution is at most $t(q^2 + q + 1)$, and the outer contribution is t ; since $\alpha(u) \leq 7$ we get that if M is minimally active then

$$7t^2(q^2 + q + 1) \geq q^{3t}(q^{2t} - 1),$$

where $q^t = 2$ is again the only solution, this time with $p = 7$. A quick computer check shows that for $p = 7$ we actually have $\alpha(u) = 2$, so that there are no non-trivial minimally active modules here either.

Thus $p = 2, 3$ remain. If $p = 3$ then the order of u is at most $9t^2(q+1)$ by Proposition 7.3, using the fact that the exponent of the Weyl group of type D_4 is 12. Since $\alpha(u) \leq 7$ we have that

$$7 \cdot 27t^2(q+1) \geq \dim(M) \geq q^{3t}(q^{2t} - 1),$$

which is satisfied only for $t = 1, q = 2, 3$, with $q = 3$ not allowed as $p \nmid q$. If $p = 2$, then u has order at most $4t^2(q+1)$, and again $\alpha(u) \leq 7$ so that

$$7 \cdot 4t^2(q+1) \geq \dim(M) \geq q^{3t}(q^{2t} - 1),$$

which only has a solution for $t = 1$ and $q = 2$, not of interest as $p = 2$. Thus we need to consider $p = 3$, $G_0 = {}^3D_4(2)$.

It is easy to check by computer that for any 3-element in G_0 , two conjugates of it generate G_0 , and $o(u) \leq 9$, so if $u \in G_0$ then $\dim(M) \leq 2 \cdot 8 = 16$, smaller than the minimal dimension of 25 [14, p.251]. If G/G_0 has order 3, then the Sylow 3-subgroup still has exponent 9, so either $o(u) = 9$, in which case two conjugates of u generate G and $\dim(M) \leq 16$, or $o(u) = 3$ and $\langle u \rangle$ lies outside G_0 , and then $\alpha(G)(o(u) - 1) \leq 14$, less than 25. This completes the proof. \square

Proposition 9.5. *Let G_0 be a central extension of a Ree or Suzuki group other than ${}^2B_2(8)$. There are no non-trivial, minimally active modules for G .*

Proof. Let G_0 be a central extension of a Suzuki group ${}^2B_2(2^{2n+1})$ for some $n \geq 2$, so that p is odd. The minimal faithful degree for G_0 is $2^n(2^{2n+1} - 1)$ from [19], and from [8, Proposition 5.8] we have that $\alpha(G) \leq 3$. Note also that there is no Weyl contribution as p is odd: the toral contribution is a divisor of one of $2^{2n+1} - 1$, $2^{2n+1} + 2^{n+1} + 1$ and $2^{2n+1} - 2^{n+1} + 1$, hence at most $2^{2n+1} + 2^{n+1} + 1$, and the outer contribution is $t \mid (2n + 1)$.

This yields

$$3t(2^{2n+1} + 2^{n+1} + 1) > \alpha(u) \cdot (o(u) - 1) \geq \dim(M) \geq 2^n(2^{2n+1} - 1),$$

which has no solutions for $t = 1$, and for $t = 2n + 1$ only works for $n \leq 5$. If $t > 1$ then $p \mid (2n + 1)$: if $p \mid (2^{2n+1} \pm 2^{n+1} + 1)$ and $p \mid (2n + 1)$ then $p \mid (2 \pm 2 + 1)$, so $p = 5$. The other alternative is that $p \mid (2^{2n+1} - 1)$, in which case $p = 1$, which is not allowed. Thus $p = 5$ always, so we need to consider $n = 2$ only, as this is the only case where 5 divides $2n + 1$.

Here we just need to be more precise, noting that the Sylow 5-subgroup of ${}^2B_2(32).5$ has order 125 but exponent 25, so actually $\dim(M) \leq 72$, less than the minimal degree of 124.

We perform a similar analysis for the Ree groups $G_0 = {}^2G_2(3^{2n+1})$ for $n \geq 1$, where the Landazuri–Seitz bound is $3^{2n+1}(3^{2n+1} - 1)$, and from [8, Proposition 5.8] we have that $\alpha(G) \leq 3$. If $p = 2$ then $o(u) = 2$ by Proposition 7.3, so $\dim(M) \leq 3$ if M is minimally active, absurd; thus p is odd.

The order of any semisimple element of G_0 is a divisor of one of $3^{2n+1} - 1$, $3^{2n+1} + 1$, $3^{2n+1} + 3^{n+1} + 1$ and $3^{2n+1} - 3^{n+1} + 1$, and the outer contribution is at most $2n + 1$: whence for $u \in G$,

$$3 \cdot (2n + 1) \cdot (3^{2n+1} + 3^{n+1} + 1) \geq 3 \cdot (o(u) - 1) \geq \dim(M) \geq 3^{2n+1}(3^{2n+1} - 1),$$

if u acts minimally actively, which fails for all $n \geq 1$.

We end with $G_0 = {}^2F_4(2^{2n+1})$. Here the Landazuri–Seitz bound is $2^{9n+4}(2^{2n+1} - 1)$, and $\alpha(G) \leq 7$. The toral contribution is at most one of

$$2^{2n+1} \pm 1, \quad 2^{2n+1} \pm 2^{n+1} + 1, \quad 2^{4n+2} \pm 2^{3n+2} + 2^{2n+1} \pm 2^{n+1} + 1,$$

the Weyl contribution is 3, and the outer contribution divides $2n + 1$. Thus from the formula $\dim(M) \leq \alpha(G)(o(u) - 1)$ for minimally active M , we get

$$21(2n + 1)(2^{4n+2} + 2^{3n+2} + 2^{2n+1} + 2^{n+1} + 1) > \alpha(G)(o(u) - 1) \geq \dim(M) \geq 2^{9n+4}(2^{2n+1} - 1).$$

The only solution to this is $n = 0$, i.e., G_0 is the Tits group. Here it is easy to check that G_0 is generated by two conjugates of any element of order at least 3, that u has order at most 13, and that $\dim(M) \geq 26$, thus there is no example here. \square

10 Proof of Theorem 1.2

In this section we need to check that all of the minimally active modules that we have found satisfy Theorem 1.2.

Proposition 10.1. *If $G_0/Z(G_0)$ is an alternating group then Theorem 1.2 is satisfied.*

Proof. We check the twenty-nine examples from Proposition 4.1 first, noting that the symmetric group in its natural module is a complex reflection group; this checks (i)–(iii). Cases (iv) and (v) are $G_0 = \mathrm{SL}_2(4)$, so these are fine. Most of the other cases are where $G/Z(G)$ has a self-centralizing cyclic Sylow p -subgroup, where $\dim(M) \leq p + 1$: all of the cases where $p = 5, 7$ have this property, so we are left with (x)–(xiv), (xviii)–(xix) and (xxvi)–(xxviii), with these last three cases being the natural module for $\mathrm{SL}_4(2)$, so covered.

For (x)–(xiv) we have $G_0/Z(G_0)$ being Alt_6 . Case (xii) is the complex reflection group G_{27} , and Cases (xiii) and (xiv) are $\mathrm{SL}_2(9)$, leaving (x) and (xi). In (x) the module is imprimitive, as is the 6-dimensional module in (xi), so we are left with the 9-dimensional module for $3 \cdot M_{10}$, which is on our list of exceptions.

Finally, Case (xviii) is the restriction of the same module for Alt_8 , and Case (xix) is again on our list of exceptions.

The remaining alternating groups are from Proposition 4.3, where we either have the permutation module or $G = 2 \cdot \mathrm{Alt}_9$ and $\dim(M) = 8$, and this is the restriction of the reflection representation of the Weyl group of E_8, G_{37} . \square

Proposition 10.2. *If $G_0/Z(G_0)$ is a sporadic group then Theorem 1.2 is satisfied.*

Proof. These are given in Proposition 6.2. We first remove those cases where there is a self-centralizing, cyclic Sylow p -subgroup, which are (i)–(iii), (v)–(vii), (ix)–(xiv), (xvi) and (xvii). The remaining three cases are all for $p = 2$, and are on our list of exceptions. \square

Proposition 10.3. *If $G_0/Z(G_0)$ is a group of Lie type in cross characteristic then Theorem 1.2 is satisfied.*

Proof. We start with the groups and modules in Proposition 7.1. Cases (vi)–(vii) are G_{34} , Cases (ix) and (x) are the Weyl group of E_7, G_{36} , and (xii)–(xiv) are the Weyl group of type E_8, G_{37} .

Cases (i)–(v) have cyclic Sylow p -subgroups that are self centralizing, while (ii) is also contained in G_{34} and (iv) with the 6-dimensional module if G_{35} . The remaining case is (xi), which is contained in the Weyl group of type E_8 .

The companion proposition to Proposition 7.1 for exceptional groups is Proposition 9.1, and all the groups in this have self-centralizing, cyclic Sylow 13-subgroups, and $\mathrm{Sp}_4(4)$ has a self-centralizing, cyclic Sylow 17-subgroup, as in Proposition 8.5. The remaining non-Weil module is from Proposition 8.3, and $\mathrm{PSL}_3(3)$ has a self-centralizing Sylow 13-subgroup of order 13.

If G_0 is a central extension of $\mathrm{PSL}_2(q)$ then in all cases the Sylow p -subgroup is cyclic and self centralizing, so this case is covered.

If M is a Weil module then u is a Singer cycle and so the Sylow p -subgroup is cyclic and self-centralizing, as is the Sylow 7-subgroup of $\mathrm{SU}_3(3)$, so these are also covered by the theorem. \square

Proposition 10.4. *If $G_0/Z(G_0)$ is a group of Lie type in defining characteristic then Theorem 1.2 is satisfied.*

Proof. Each of the modules listed in Proposition 5.1 appears on our list, so u induces an outer automorphism, and appears in Propositions 5.2 and 5.3.

For the groups in Proposition 5.2, (i) and (iv) are imprimitive, (v) is the natural module, (iii) is the exterior square of the natural, and (ii) is mentioned explicitly. The groups in Proposition 5.3 are either imprimitive in case (v), or stabilize a tensor product in cases (i)–(iv). \square

Acknowledgements The author would like to thank Donna Testerman and Alex Zalesski for giving him access to unpublished manuscripts of theirs.

The author is financially supported by the Royal Society, being awarded a University Research Fellowship.

References

- [1] Michael Aschbacher, *On the maximal subgroups of the finite classical groups*, Invent. Math. **76** (1984), 469–514.
- [2] John Conway, Robert Curtis, Simon Norton, Richard Parker, and Robert Wilson, *Atlas of finite groups*, Oxford University Press, Eynsham, 1985.
- [3] David A. Craven, Jason Semeraro, and Bob Oliver, *Reduced fusion systems over p -groups with abelian subgroup of index p : II*, Adv. Math. **322** (2017), 201–268.
- [4] John Dixon and Brian Mortimer, *Permutation groups*, Springer–Verlag, 1996.
- [5] Daniel Gorenstein, Richard Lyons, and Ronald Solomon, *The classification of the finite simple groups, Number 3. Part I. Chapter A.*, American Mathematical Society, Providence, RI, 1998.
- [6] Roderick Gow and M. Chiara Tamburini, *Generation of $\mathrm{SL}(n, Z)$ by a Jordan unipotent matrix and its transpose*, Lin. Algebra Appl. **181** (1993), 63–71.
- [7] Robert Guralnick, Kay Magaard, Jan Saxl, and Pham Huu Tiep, *Cross characteristic representations of symplectic and unitary groups*, J. Algebra **257** (2002), 291–347.
- [8] Robert Guralnick and Jan Saxl, *Generation of almost simple groups by conjugates*, J. Algebra **268** (2003), 519–571.
- [9] Robert Guralnick and Pham Huu Tiep, *Low-dimensional representations of special linear groups in cross characteristics*, Proc. London Math. Soc. **78** (1999), 116–138.
- [10] Gerhard Hiss and Gunter Malle, *Low-dimensional representations of special unitary groups*, J. Algebra **236** (2001), 745–767.

- [11] Corneliu Hoffman, *Cross characteristic projective representations for some classical groups*, J. Algebra **229** (2000), 666–677.
- [12] Gordon James, *On the minimal dimensions of irreducible representations of symmetric groups*, Math. Proc. Camb. Phil. Soc. **94** (1983), 417–424.
- [13] Christoph Jansen, *The minimal degrees of faithful representations of the sporadic simple groups and their covering groups*, LMS J. Comput. Math. **8** (2005), 122–144.
- [14] Christoph Jansen, Klaus Lux, Richard Parker, and Robert Wilson, *An atlas of Brauer characters*, Oxford University Press, New York, 1995.
- [15] William Kantor, *Subgroups of classical groups generated by long root elements*, Trans. Amer. Math. Soc. **248** (1979), 347–379.
- [16] Peter Kleidman, *The maximal subgroups of the Steinberg triality groups ${}^3D_4(q)$ and of their automorphism groups*, J. Algebra **115** (1988), 182–199.
- [17] Peter Kleidman and Martin Liebeck, *The subgroup structure of the finite classical groups*, London Mathematical Society Lecture Note Series, vol. 129, Cambridge University Press, Cambridge, 1990.
- [18] Alexander Kleshchev and Pham Huu Tiep, *Small-dimensional projective representations of symmetric and alternating groups*, Algebra Number Th. **6** (2012), 1773–1816.
- [19] Vicente Landazuri and Gary Seitz, *On the minimal degrees of projective representations of the finite Chevalley groups*, J. Algebra **32** (1974), 418–443.
- [20] Ross Lawther, *Jordan block sizes of unipotent elements in exceptional algebraic groups*, Comm. Algebra **23** (1995), 4125–4156.
- [21] Martin Liebeck and Gary Seitz, *Unipotent and nilpotent classes in simple algebraic groups and Lie algebras*, Mathematical Surveys and Monographs, vol. 180, American Mathematical Society, Providence, RI, 2012.
- [22] Frank Lübeck, *Small degree representations of finite Chevalley groups in defining characteristic*, LMS J. Comput. Math. **4** (2001), 135–169.
- [23] Gunter Malle and Donna Testerman, *Linear algebraic groups and finite groups of lie type*, Cambridge University Press, 2011.
- [24] Lino Di Martino, M.A. Pellegrini, and Alexandre Zalesski, *On generators and representations of the sporadic simple groups*, Comm. Alg. **42** (2014), 880–908.
- [25] Lino Di Martino and Alexandre Zalesski, *Almost cyclic elements in Weil representations of finite classical groups*, Comm. Alg., to appear.
- [26] Gary Seitz, *The maximal subgroups of classical algebraic groups*, Mem. Amer. Math. Soc. **67** (1987), no. 365, iv+286.
- [27] Gary Seitz and Alexandre Zalesskii, *On the minimal degrees of projective representations of the finite Chevalley groups*, J. Algebra **158** (1993), 233–243.

- [28] Tonny Springer, *Regular elements of finite reflection groups*, Invent. Math. **25** (1974), 159–198.
- [29] I.D. Suprunenko, *Unipotent elements of nonprime order in representations of the classical algebraic groups: two big Jordan blocks*, J. Math. Sci. **199** (2014), 350–374.
- [30] Donna Testerman and Alexandre Zalesski, *Irreducible representations of simple algebraic groups in which a unipotent element is represented by a matrix with single non-trivial Jordan block*, preprint, 2017.
- [31] Robert A. Wilson, *The finite simple groups*, Graduate Texts in Mathematics, vol. 251, Springer-Verlag London Ltd., London, 2009.
- [32] Thilo Zieschang, *Primitive permutation groups containing a p -cycle*, Arch. Math. **64** (1995), 471–474.