

Recasting Human V δ 1 Lymphocytes in an Adaptive Role

Davey, Martin; Willcox, Carrie; Baker, Alfie; Hunter, Stuart; Willcox, Benjamin

DOI:

[10.1016/j.it.2018.03.003](https://doi.org/10.1016/j.it.2018.03.003)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Davey, M, Willcox, C, Baker, A, Hunter, S & Willcox, B 2018, 'Recasting Human V δ 1 Lymphocytes in an Adaptive Role', *Trends in Immunology*, vol. 39, no. 6, pp. 446-459. <https://doi.org/10.1016/j.it.2018.03.003>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Published in *Trend in Immunology* on 18/04/2018

DOI: 10.1016/j.it.2018.03.003

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Opinion

Recasting Human V δ 1 Lymphocytes in an Adaptive RoleMartin S. Davey,^{1,3} Carrie R. Willcox,^{1,3} Alfie T. Baker,¹ Stuart Hunter,^{1,2} and Benjamin E. Willcox^{1,*}

$\gamma\delta$ T cells are unconventional lymphocytes commonly described as ‘innate-like’ in function, which can respond in both a T cell receptor (TCR)-independent and also major histocompatibility complex (MHC)-unrestricted TCR-dependent manner. While the relative importance of TCR recognition had remained unclear, recent studies revealed that human V δ 1 T cells display unexpected parallels with adaptive $\alpha\beta$ T cells. V δ 1 T cells undergo profound and highly focussed clonal expansion from an initially diverse and private TCR repertoire, most likely in response to specific immune challenges. Concomitantly, they differentiate from a V δ 1 T cell naïve (T_{naïve}) to a V δ 1 T cell effector (T_{effector}) phenotype, marked by the downregulation of lymphoid homing receptors and upregulation of peripheral homing receptors and effector markers. This suggests that an adaptive paradigm applies to V δ 1 T cells, likely involving TCR-dependent but MHC-unrestricted responses to microbial and non-microbial challenges.

 $\gamma\delta$ T Cells and the Lymphoid Stress Surveillance Hypothesis

$\gamma\delta$ T cells have coevolved alongside $\alpha\beta$ T cells and B cells for at least the past ~450 million years of vertebrate evolution [1,2], each distinguished by related but distinct somatically recombined antigen receptors. However, our understanding of these different lineages is strikingly imbalanced. Critical to our understanding of $\alpha\beta$ T cell and B cells is the classical adaptive paradigm (Box 1). Within this, seminal discoveries have established the core function of the $\alpha\beta$ T cell lineage: to enable immune responses to target cells based on the presence on their surface of antigenic peptide in the context of MHC molecules; similarly, we understand that B cells, which underpin humoral immunity, enable the production of soluble antibodies capable of recognising a diverse range of antigenic targets in native, 3D conformation. In keeping with Burnet’s suggestion that ‘receptor occupation’ is key in driving the activation and clonal selection of adaptive lymphocytes [3], structural studies have confirmed both the involvement of clonotypically unique hypervariable loops in $\alpha\beta$ TCR/peptide-MHC and B cell receptor (BCR)/antigen engagement, and the significance of such interactions in regulating multiple facets of their immunobiology (Box 1).

Originally identified serendipitously during studies defining $\alpha\beta$ TCR genes [4,5] $\gamma\delta$ T cells have by contrast remained somewhat mysterious both in terms of the immunological niche they occupy and the key reason(s) for their evolutionary preservation as a third lymphocyte lineage within vertebrate immunity. Moreover, although $\gamma\delta$ T cells are implicated in a range of immune settings, including antimicrobial immunity, antitumour immunity, and tissue homeostasis (reviewed in [6]), the central paradigms that govern their development and antigen recognition functions are unresolved. Finally, despite remaining a focus of ongoing interest, the closely related issue of the importance and exact role of $\gamma\delta$ TCR occupation in $\gamma\delta$ T cell biology remains a central question.

Highlights

V δ 1 T cells are the predominant tissue-associated $\gamma\delta$ T cell subset in humans, and can recognise signs of cellular dysregulation, including viral infection and transformation. They are often assumed to be innate-like effectors.

The V δ 1 TCR repertoire is initially diverse, yet a few clonotypes typically expand heavily over time. Such expanded clonotypes are diverse in sequence both within and between individuals.

Clonal expansion occurs concurrently with differentiation from V δ 1 T_{naïve} to V δ 1 T_{effector} status, alongside a switch from lymphoid to peripheral homing receptors, and upregulation of cytotoxic pathways.

This concurrent expansion and differentiation suggests an adaptive-like response, likely driven by diverse stimuli, including microbial pathogen infection.

¹Cancer Immunology and Immunotherapy Centre, Institute of Immunology and Immunotherapy, School of Medicine and Dental Sciences, University of Birmingham, Birmingham, B15 2TT, UK

²Centre for Liver Research and NIHR Biomedical Research Unit in Liver Disease, Institute of Immunology and Immunotherapy, School of Medicine and Dental Sciences, University of Birmingham, Birmingham, B15 2TT, UK

³These authors contributed equally

*Correspondence: b.willcox@bham.ac.uk (B.E. Willcox).

Box 1. Hallmarks of Classical Adaptive Immunity

Notably, $\alpha\beta$ T cells and B cells share key hallmarks of classical adaptive immunity.

Generation of a Diverse Antigen Receptor Repertoire and Tolerance Mechanisms

Both $\alpha\beta$ T cell and B cell lineages feature somatically recombined TCRs and BCRs, with repertoires featuring high diversity in their hypervariable complementarity-determining region loops, particularly CDR3. For both lineages, selection events during lymphocyte development are critical for immune tolerance. $\alpha\beta$ T cells undergo positive and negative selection in the thymus; B cells, in the bone marrow, undergo both antigen-independent positive selection, based on tonic BCR signalling, and processes that eliminate or mitigate autoreactive specificities, including negative selection and anergy induction.

Clonal Expansion from a Diverse Immune Receptor Repertoire

The selection of individual clonotypes from within the diverse naïve immune receptor repertoire allows expansion of specific $\alpha\beta$ T cell and B cell clonotypes bearing receptors that critically enable amplified responses to specific immune challenges, such as pathogen infection.

Differentiation into Long-Lived Effectors

Concurrent with clonal expansion, both $\alpha\beta$ T cell and B cell lineages not only undergo differentiation to effectors, but also permit the maintenance of long-lived clonotypically expanded populations, enabling immunological memory, whereby faster and more potent immune responses are induced in response to secondary antigenic challenge.

Critical Importance of Antigen Receptor–Ligand Interactions

Diverse studies highlight the central role for TCR–pMHC and BCR–ligand interactions in directing $\alpha\beta$ T cell and B cell development, maintenance, clonal amplification and activation, and memory formation, emphatically validating the concept that ‘receptor occupancy’ is a central driver of adaptive lymphocyte biology.

One concept emerging from mouse studies of $\gamma\delta$ T cells is that certain $\gamma\delta$ T cell subsets, instead of functioning via conventional adaptive paradigms, may instead act as ‘innate-like’ lymphocytes. Notably, murine $\gamma\delta$ T cells express distinct TCR γ and TCR δ combinations at different anatomical sites, and often display semi-invariant TCR repertoires, in some cases featuring highly restricted CDR3 regions [7–9]. They can be preprogrammed during thymic development to differentiate into discrete effector populations producing either interleukin-17 (IL-17) or interferon-gamma (IFN- γ) [10,11]. More recently, intra-epithelial lymphocyte populations have been shown to be selected in tissues after birth, dependent on the expression of particular butyrophilin-like molecules (BTNLS) [12]. Such populations of ‘activated-but-resting’ unconventional lymphocytes are thought to be capable of reacting directly to dysregulated target cells without the need for clonal expansion and differentiation. These data align with the idea such subsets may recognise a limited range of host-encoded stress ligands [13], and suggest that their TCRs act like surrogate pattern recognition receptors (PRRs) for molecular signals of microbial/non-microbial stress. In humans, the $\gamma\delta$ T cell subset that aligns most clearly to this biology is characterised by a V γ 9/V δ 2 chain pairing, and represents the predominant peripheral blood subset (1–10% of T cells) [14]. Based on their restricted TCR V γ and V δ gene segment usage and CDR3 lengths, presence of common CDR3 motifs, foetal generation, polyclonal production of IFN γ and Tumour Necrosis Factor- α (TNF α) following exposure to pyrophosphate antigens (P-Ags), and strong dependency of TCR-mediated recognition on the BTN3A1 Ig-like protein, this subset arguably conforms to such an innate-like functional paradigm, although the exact mechanisms underlying its recognition of target cells remain unclear. To some extent, the features of these $\gamma\delta$ T cell subsets mirror those of unconventional $\alpha\beta$ T cell subsets [e.g., mucosal-associated invariant T cells (MAITs) and invariant natural killer cells

(iNKTs)], which also feature highly restricted TCR repertoires [15,16], and have been shown to recognise relatively nonpolymorphic ligands [MHC class I-related gene protein (MR1) and Cluster of differentiation 1 (CD1), respectively].

These observations led to the development of the lymphoid stress surveillance hypothesis [17], which postulates that such effectors, by circumventing the requirement for clonal selection and differentiation, may provide protection from microbial or non-microbial stress challenges during the initial phase of the response, before adaptive immune responses have been generated.

A key finding in this area has been that such subsets can be activated not only via their TCR, but also independently by TCR-extrinsic signals. For example, mouse dendritic epidermal T cells (DETC, a subset of $\gamma\delta$ T cells present in murine skin) can be activated directly via NK receptor (NKR)-mediated recognition of stress ligands expressed on stressed epithelium, independently of the TCR [18]. Moreover, while human V γ 9/V δ 2 T cells exhibit potent TCR-dependent recognition of P-Ag-exposed target cells, they can also be activated by NKG2D–ligand interactions and are responsive to cytokines such as IL-12/IL-18 [19–21]. In addition, recent studies on mouse skin and gut $\gamma\delta$ T cell subsets suggest that TCR signals, potentially mediated via interactions with BTNLs, are required for the development, homing, and establishment of their effector program [11,12]; however, these cells can become hyporesponsive to TCR signals and function in an TCR-independent manner to respond to signs of cellular stress [22].

In addition to V γ 9/V δ 2 T cells, a second human $\gamma\delta$ T cell compartment exists, bearing V δ 2-negative TCRs, of which the V δ 1 component is dominant. V δ 1 T cells are the most prevalent subtype of $\gamma\delta$ T cells at birth [23], and the dominant $\gamma\delta$ T cell subtype in peripheral tissues in adults, such as the gut [24,25] and skin [26]. V δ 1 T cells have remained very much an enigma in terms of the fundamental paradigms underlying their biology. Based on their predominant effector phenotype, their potent cytotoxicity/cytokine production, combined with their ability to recognise both virally infected and also cancerous cells, and their expression of NKRs [27–29], they have often been assumed to act in an innate-like fashion, similar to NK cells, potentially enabling recognition of diverse cellular stress signals in target cells. However, here we review recent data that have revised this picture, and suggest instead that V δ 1 T cells exhibit a radical new adaptive immunobiology. These data highlight some of the most significant questions in $\gamma\delta$ T cell biology, including the importance and exact role of the $\gamma\delta$ TCR that defines the lineage, but ironically, is so poorly understood.

TCR Repertoire Analyses Reveal V δ 1 T Cell Clonal Amplification

The advent of next-generation sequencing (NGS) approaches has allowed in-depth analyses of the TCR/BCR repertoire within the $\alpha\beta$ and B cell lineage, respectively [30,31]. Furthermore, application of these technologies to human peripheral blood $\gamma\delta$ T cells has provided valuable information on clonal evolution within the $\gamma\delta$ T cell lineage. The approaches used include either DNA-based or RNA-based methods, with the former highlighting the requirement to purify $\gamma\delta$ T cells from $\alpha\beta$ subsets to avoid contamination of TCR γ sequences recombined in mature $\alpha\beta$ T cells [32]. One inherent challenge of these methods is the potential for polymerase chain reaction (PCR) and sequence errors [33]. In the absence of *in silico* error correction and appropriate data handling, this can result in retention of erroneous sequences, often resulting from single-base errors of other higher frequency clonotypes. Therefore, caution should be applied when interpreting highly similar TCR base sequences. The availability of public software packages dedicated to the interpretation of NGS TCR sequencing data (MiTCR [34], MiXCR [35], and TcR [36]), which provide robust error correction, should alleviate such challenges. Moreover, since RNA-based approaches are potentially vulnerable to bias, based on the overall

level of RNA differing between cells in different activation and/or differentiation states, ideally parallel techniques, such as single-cell PCR-based TCR analysis, should be considered to validate NGS-based TCR frequencies correlate with cell number on a specific platform [37–39].

Such studies have not only shed substantial light on V δ 1 T cell immunobiology, but equally importantly, have also highlighted features of the V δ 1 repertoire that contrast markedly with that of both peripheral blood V γ 9/V δ 2 T cells and thymically programmed mouse $\gamma\delta$ T cells. First, based on analyses of cord blood samples, the V δ 1 TCR repertoire at the start of life, which features a variety of V γ chain pairings, is clonotypically diverse, featuring a range of CDR3 lengths apparently unrelated in sequence. Importantly, these neonatal cord blood V δ 1⁺ repertoires are essentially entirely unfocussed: in other words, it appears that ‘all clonotypes are created equal’, with no single sequence exceeding 1–2% of the repertoire (Figure 1). This contrasts with the V γ 9/V δ 2 population, which, even in foetal and cord blood, includes relatively prevalent V γ 9-J γ P clonotypes that are public (i.e., shared at either the nucleotide or amino acid level between individuals) and are present throughout life, consistent with selection of a semi-invariant repertoire preprogrammed in development for polyclonal P-Ag recognition [32,40–42].

A second important finding from such studies is that, in comparison to cord blood repertoires, adult V δ 1 repertoires are in general substantially more focussed, typically resulting from the presence of a relatively small number (e.g., <5) of heavily expanded clonotypes, and which often account for a large proportion of the total adult V δ 1 repertoire. Of note, parallel single-cell TCR analysis was used by Davey *et al.* [38] and Ravens *et al.* [39] to confirm that these represented genuine numerical clonal expansions, and were not biased by RNA abundance. Strikingly, both adult and cord blood V δ 1⁺ TCR repertoires were overwhelmingly private (i.e., unique to an individual at both a nucleotide and amino acid level; even more so than TCR β [38]), with the Complementarity determining region 3 (CDR3) lengths of expanded clonotypes highly diverse, and their sequences apparently unrelated, both within and between individuals [38]. These features appeared to stem from the addition of high levels of nontemplated (N)





Figure 1. Clonal Expansion in the V δ 1 Repertoire. At birth, neonatal V δ 1 T cell populations comprise a broad set of private clonotypes. During the progression to adulthood, most human V δ 1 T cell repertoires undergo clonotypic focussing towards a limited set of private clonally expanded T cell receptors (TCRs). Despite this, in some individuals, clonal expansion and focussing is not evident and their V δ 1 T cell repertoires remain diverse. While human cytomegalovirus (CMV) infection is directly implicated in driving clonal expansion in V δ 1 T cell repertoires, other immunological stimuli are clearly capable of driving TCR-specific responses. The TCR δ tree plots depict representative V δ 1 T cell repertoires at each stage of life. Each coloured block represents a single unique Complementarity determining region (CDR)-3 δ . Each repertoire was private and clonotypes did not overlap between individuals. Percentages are from Davey *et al.* [38], and intended as a guide only.

nucleotides (introduced by terminal deoxynucleotidyl transferase) or occasionally palindromic (P) nucleotides (a mean of 19 N/P nucleotides for V δ 1 CDR3) during variable (diversity) joining [V(D)J] recombination. Both the private nature of the V δ 1 repertoire and the presence of such dominant clonal expansions contrast markedly with the V γ 9/V δ 2 TCR repertoire, which displays less pronounced focussing and contains several public V γ 9 sequences in cord blood and adults [32,38,41]. Moreover, compared with V δ 1 T cells, the V γ 9/V δ 2 T cell repertoire features restricted CDR3 γ and CDR3 δ lengths, with CDR3 δ 2 sequences generally considerably shorter than CDR3 δ 1 sequences. Similarly, TCR $\alpha\beta$ repertoires also display CDR3 length restriction, which is likely imposed by the structural constraints of peptide–MHC recognition [43].

$\gamma\delta$ TCR repertoire analyses have also highlighted factors driving such clonal expansions. In particular, Ravens *et al.* showed that acute cytomegalovirus (CMV) infection following stem cell transplantation (SCT) can drive expansion of V δ 2^{neg} (predominantly V δ 1) TCR clonotypes [39]; V δ repertoires were also noted to be private. These findings build on numerous studies highlighting the importance of V δ 2-negative T cells in responses to CMV infection following kidney transplantation and in healthy donors [29,44,45], including spectratyping data consistent with a degree of clonotypic focussing in CMV^{pos} healthy donors [29]. Despite this, the link to CMV infection is not necessarily straightforward, because both Ravens *et al.* [39] and Davey *et al.* [38] noted that the V δ 1 repertoire of some CMV^{pos} individuals lacked clonal expansions, indicating that V δ 1⁺ TCR clonal expansion was not an inevitable consequence of CMV infection (Figure 1). The reason for this is unclear, but could conceivably reflect different routes of infection, greater/lesser dependency of the anti-CMV response on V δ 2^{neg} V δ 1^{neg} $\gamma\delta$ T cells, and/or the presence of ‘holes’ in the V δ 1 TCR repertoire in some individuals. Further studies are required to address these possibilities. Moreover, several CMV^{neg} individuals also had heavily expanded V δ 1 clonotypes, indicating that CMV infection is not the sole immune challenge that stimulates V δ 1 T cell responses (Figure 1). Consistent with diverse stimuli for the V δ 1 subset, expansion of V δ 1 T cells has also been noted in response to HIV [46] and in synovial fluid in Lyme disease [47], while two case reports describe V δ 1 clonal focussing after EBV infection in SCT [48,49]. Nevertheless, infectious stimuli underlying clonotypic focussing in healthy donors, other than CMV, remain to be identified. Finally, irrespective of the stimuli inducing such responses, both Davey *et al.* [38] and Ravens *et al.* [39] provide evidence that such expanded clonotypes can be long-lived, and persist for at least 2 years, consistent with long-term contributions to immunosurveillance.

Clonal Selection in V δ 1 T Cells Induces Adaptive Changes in Phenotype and Function

Davey *et al.* [38] combined NGS TCR sequencing, single-cell TCR analyses, and a flow cytometric immunophenotyping approach to delineate different V δ 1 subsets in adult peripheral blood and cord blood samples. Notably, a naïve-like CD27^{hi} V δ 1 subset was identified expressing highly diverse TCRs and multiple markers common to naïve T cells, including IL-7R, CD28, CD62L, and CCR7 (Figure 2); we hereafter apply the term ‘V δ 1 T_{naïve}’ to this subset. Importantly, although such T_{naïve} cells were typically a minor fraction of adult peripheral blood V δ 1 T cells, essentially the entire V δ 1 T cell subset in cord blood was clonotypically unfocussed. By contrast, clonotypically expanded V δ 1 TCRs present in adults invariably resided within a differentiated effector CD27^{lo/neg} compartment largely absent in cord blood, which was detected to different extents within adult peripheral blood V δ 1 T cells across a 20-person cohort. This compartment shared several phenotypic features with conventional T_{effector} populations, including expression of granzymes, perforin, and CX3C chemokine receptor 1 (CX₃CR1; Figure 2); we hereafter apply the term ‘V δ 1 T_{effector}’ to this subset. Importantly, the observation that the V δ 1 compartment of the minority of adult donors who



Trends in Immunology

Figure 2. Phenotypic Changes in Vδ1 T Cells upon Adaptive Expansion. Vδ1 T cells displaying a diverse T cell receptor (TCR) repertoire expressed high levels of Cluster of differentiation 27 (CD27). Conversely, clonally focussed TCR repertoires either displayed reduced expression or had completely downregulated CD27. These CD27^{hi} and CD27^{lo/neg} Vδ1 T cells also displayed markers and functional responses consistent with naïve and effector T cells, respectively. Vδ1 T_{naïve} cells, alongside a broad γδ TCR repertoire, expressed the co-stimulatory receptor CD28, lymphoid tissue homing receptor CCR7, tissue access molecule CD62L, and mounted a proliferative response to the lymphoid tissue-associated homeostatic cytokine interleukin 7 (IL-7). By contrast, Vδ1 T_{effector} cells had downregulated CD28, CCR7, CD62L, and IL7Rα and upregulated cytotoxic granzymes, perforin, and endothelial homing receptor CX₃C chemokine receptor 1 (CX₃CR1), and proliferated in response to the peripheral tissue-associated cytokine IL-15. Both populations were unresponsive to innate stimuli (IL-12 and IL-18) but retained TCR responsiveness (anti-CD3 stimulation), with Vδ1 T_{effector} cells becoming rapidly activated, whereas Vδ1 T_{naïve} cells responded over a longer period of time.

retained relatively diverse Vδ1 TCR repertoires was dominated by Vδ1 T_{naïve} cells confirmed the validity of this phenotypic distinction, and highlighted that clonal expansion and differentiation were not inevitable consequences of Vδ1 T cell maturation.


These phenotypic features of Vδ1 T_{naïve} and T_{effector} subsets point towards an adaptive biology. First, the transition from Vδ1 T_{naïve} to T_{effector} subset is accompanied by a reprogramming of homing receptor expression. Vδ1 T_{naïve} cells uniformly express high levels of central lymphoid homing markers. By contrast, Vδ1 T_{effector} cells exhibit strong downregulation of CCR7 and CD62L, but increased CX₃CR1, which binds to fractalkine, an endothelial homing chemokine. The respective expression profiles of these markers on Vδ1 T_{naïve} and T_{effector} subsets closely mirrored expression on CD8 naïve and T_{EMRA} cell populations, respectively. Correspondingly, and as suggested for CX₃CR1^{hi} CD8 memory T_{effector} cells [50], CX₃CR1^{hi} Vδ1 T_{effector} cells may be involved in endothelial immunosurveillance. In terms of function, whereas Vδ1 T_{naïve} cells, such as naïve CD8 T cells, were devoid of cytotoxic effector markers (e.g., perforin, and granzymes A and B), these were heavily upregulated in Vδ1 T_{effector} cells (equivalent to CD8 T_{EMRA} populations). Moreover, Vδ1 T_{effector} cells retained a rapid proliferative capacity and TCR

sensitivity, and were preferentially sensitive to IL-15 relative to $V\delta 1 T_{naive}$ cells, which conversely (and similar to naive CD8 T cells) were preferentially responsive to IL-7 and exhibited slower proliferation following TCR stimulation. These findings strongly suggest that the TCR-diverse $CD27^{hi}$ and highly TCR focussed $CD27^{lo/neg}$ populations represent *bona fide* naive and effector $V\delta 1$ subsets, respectively (Figures 2 and 3).

These findings demonstrate that, far from representing a preformed effector subset from birth, $V\delta 1$ T cells are initially highly naive in phenotype and feature an entirely unfocussed TCR repertoire. They also suggest that differentiation to an effector phenotype is not an inevitable developmental process, but is inextricably linked to clonal amplification, and drastically affects both cytotoxic capability and homing receptor expression (Figure 3).

Adaptive MHC-Unrestricted $\gamma\delta$ T Cell Stress Surveillance: A Paradigm for $V\delta 1$ T Cells

The observations outlined above highlight surprising parallels between $V\delta 1$ T cells and classical adaptive T cell subsets, particularly CD8 T cells; conversely, they emphasise key distinctions



Trends in Immunology

Figure 3. Adaptive Stress Surveillance Paradigm. Both $V\delta 1 T_{naive}$ and $T_{effector}$ cells circulate in the peripheral blood. $V\delta 1 T_{naive}$ populations (expressing CCR7 and CD62L) are likely to migrate to secondary lymphoid tissue, via CCL19 and CCL21 chemokine gradients. Access to secondary lymphoid tissue permits encounter of homeostatic interleukin 7 (IL-7), maintaining $V\delta 1 T_{naive}$ cells and allowing their persistence throughout adulthood. $V\delta 1 T_{naive}$ cells may also encounter cognate antigen either in the lymphoid tissues, akin to $\alpha\beta$ T cells, or elsewhere, and give rise to $V\delta 1 T_{effector}$ cells. Circulating $V\delta 1 T_{effector}$ populations may enter peripheral tissues, accessing homeostatic IL-15 concentrations. Access to peripheral tissues may indicate a stress surveillance role and antimicrobial function, through T cell receptor (TCR)–ligand engagement.

Box 2. Key Principles of Adaptive Stress Surveillance

We propose that three key tenets of an adaptive immunobiology apply to V δ 1 T cells. We suggest that:

- (i) specific stress stimuli, including microbial infection and/or colonisation, are capable of triggering the clonal expansion and differentiation of V δ 1 T_{effector} clonotypes from a highly diverse, unfocussed V δ 1 T_{naïve} pool;
- (ii) as for $\alpha\beta$ T cells and B cells, 'receptor occupancy', in this context the ability of the V δ 1 $\gamma\delta$ TCR to engage cognate ligands, drives clonal expansion and initiates differentiation to the effector state; and
- (iii) clonally expanded V δ 1 T_{effector} populations, which are relatively long-lived, provide enhanced protection against recurrent immune challenges.

between V δ 1 T cells and the V γ 9/V δ 2 T cell subset. They suggest that the $\gamma\delta$ TCR is central to the biology of V δ 1 T cells, and lead us to suggest a previously unrecognised mode of MHC-unrestricted adaptive immunobiology applies to the V δ 1 T cell subset (Box 2). This paradigm and the evidence that underpins it, has several implications worthy of consideration that will likely frame future investigations.

Tolerance Induction

The V δ 1 and V γ 9/V δ 2 T cell subsets develop at different stages, consistent with a distinct underlying immunobiology and TCR repertoire for the two compartments. V γ 9/V δ 2 T cells are generated during development in the foetal liver, and later the foetal thymus, and are only present in small numbers in the postnatal thymus [51,52]. Development may require positive selection for BTN3A1 reactivity and/or self phosphoantigens, such as isopentenyl pyrophosphate (IPP), and subsequent expansion after microbial exposure during early childhood [53]. By contrast, V δ 1 T cells are the dominant $\gamma\delta$ T cell population in the postnatal thymus. It is unclear whether human $\gamma\delta$ T cells undergo thymic selection. The diversity of CDR3 lengths and sequences for V δ 1 and associated V γ TCR chains would suggest a strong potential for autoreactivity; however, there is little evidence of negative selection in $\gamma\delta$ T cells [32]. Alternatively, peripheral tolerance mechanisms may be involved in deleting or inducing the anergy of strongly autoreactive cells, or other mechanisms similar to NK licensing or arming (reviewed in [54]) may be used to set thresholds for V δ 1 reactivity. V δ 2^{neg} $\gamma\delta$ T cells express inhibitory NKRs, such as LILRB1/ILT2, which may be involved in this process [28]. Additionally, TCR self-reactivity may involve low-affinity TCR–ligand interactions and may require additional co-stimulatory signals or adhesion molecules to lead to productive TCR signalling [55].

T Cell Priming and Migration

Although the cellular mechanisms underlying initiation of V δ 1 clonal expansions are unclear, the immune phenotype of V δ 1 T_{naïve} cells suggests preferential access to secondary lymphoid organs, as opposed to clonally expanded V δ 1 T_{effector} subsets. This implies that V δ 1 T_{naïve} cells most likely recirculate between blood and lymph tissues (Figure 3), and suggests that clonal amplification requires lymphoid tissue-derived factors and/or is initiated during a priming step in secondary (or possibly tertiary) lymphoid organs. Future studies assessing the differentiation status and localisation of V δ 1 T cells in secondary and/or tertiary lymphoid organs will be required to address these suggestions, and may shed light on whether specific antigen-presenting cell types are involved in priming. Conversely, the increased expression of CX₃CR1 on V δ 1 T_{effector} populations and their enhanced sensitivity to IL-15 suggests that they preferentially home to solid tissues, and contribute to a more clonotypically focussed repertoire at such sites. Further studies comparing the phenotypic features and TCR diversity of peripheral blood V δ 1 T cells with those in solid tissues will address these issues.

Table 1. $\gamma\delta$ TCR Ligands in the Context of Adaptive Stress Surveillance^a

Candidate $\gamma\delta$ TCR ligand	Chain usage of T cells	Origin of T cells	Frequency of response	Memory phenotype of T cells	Direct ligand binding/affinity	CDR3 involvement	Comments/potential physiological significance	Refs
EPCR	V γ 4/V δ 5	PBMC from immunosuppressed lung transplant patient with acute CMV	25% of total CD3 ⁺ T cells in one CMV ⁺ individual	CD45RO ^{neg} CD28 ^{neg}	~90 μ M (BIAcore)	Yes; CDR3 γ , CDR3 δ ^b	Single private clonotype Expanded clonotype, likely $\gamma\delta$ T _{effector} status Potential 'restriction factor' for endothelial cells during CMV infection Allows detection of 'multimolecular stress signature' on CMV-infected cells EPCR upregulated on various cancer cells	[27,28,55,65,66]
PE	V δ 1 ⁺ , various V γ chains	PE staining of healthy donor blood	0.025% of total CD3 ⁺ T cells	ND	Mouse TCR-PE 2.7 μ M (BIAcore) Human TCR – ND	Yes (mouse $\gamma\delta$ TCR)	Various clonotypes involved Collectively low frequency in V δ 1 T cells; therefore, unlikely to represent dominant $\gamma\delta$ T _{effector} expansions PE derived from marine blue-green algae; therefore, may reflect potential of V δ 1 ⁺ T cells to recognise foreign antigens Immunisation of mice upregulated activation markers on PE ⁺ $\gamma\delta$ T cells and led to IL-17 production	[62]
CD1d	V δ 1 ⁺ , various V γ chains	CD1d (unloaded or with various lipids) tetramer staining of healthy donor PBMC	<0.05% of total T cells	ND	16 μ M α -GalCer/CD1d 35 μ M unloaded CD1d (BIAcore); 33 μ M sulfatide/CD1d, 240 μ M unloaded CD1d (BLI)	CDR3 δ required	Various clonotypes involved Collectively low frequency in V δ 1 T cells; therefore, unlikely to represent dominant $\gamma\delta$ T _{effector} expansions May reflect $\gamma\delta$ T _{naive} population CD1d-reactive $\gamma\delta$ T cells could expand upon recognition of CD1d-restricted stress and/or infection-linked lipids	[61,63,64]
CD1c	V δ 1 ⁺ , various V γ chains	CD1c (loaded with Mtb or self lipids) tetramer staining of healthy donor PBMC	0.16% of total T cells	ND	23–30 μ M foreign lipids; 28–150 μ M self lipids LPA, LPC, sulfatide (BLI)	Yes: chain swap	Various clonotypes involved Collectively low frequency in V δ 1 T cells; therefore, unlikely to represent dominant $\gamma\delta$ T _{effector} expansions May reflect $\gamma\delta$ T _{naive} population CD1c-reactive $\gamma\delta$ T cells could expand upon recognition of CD1d-restricted stress and/or infection-linked lipids	[60]
Annexin A2	V δ 3 ⁺ clone	Healthy donor PBMC cultured with Raji + IL-2	ND	ND	3 μ M (BIAcore)	ND	Various clonotypes involved Frequency of clone unclear in original donor; may reflect $\gamma\delta$ T _{naive} population Annexin A2 upregulated in cellular stress (tumourigenesis, oxidative stress) May permit V δ 2 ^{neg} $\gamma\delta$ T cells to recognise tumour cells or metabolically stressed cells	[59]

^aAbbreviations: CCR7, C-C chemokine receptor type 7; EBV, Epstein–Barr virus; EPCR, endothelial protein C receptor; ICAM-1, intercellular adhesion molecule 1; LILRB1/LT2, leukocyte immunoglobulin-like receptor B1/immunoglobulin-like transcript2; ND, not determined; NKG2D, natural killer group 2 member D; PBMC, peripheral blood mononuclear cell; Skint, selection and upkeep of intraepithelial T cells.

^bCarrie Willcox, unpublished data.

Mechanisms and Targets Underpinning V δ 1 $\gamma\delta$ TCR Ligand Recognition

Defining the ligands recognised by the $\gamma\delta$ TCR remains a major goal and one that may help unlock the molecular basis by which $\gamma\delta$ T cells recognise abnormal target cells (Table 1).

Most $\gamma\delta$ TCR ligand investigations have focussed on peripheral blood populations. Importantly, our understanding of the V δ 1 TCR repertoire in solid tissues is limited. However, a recent study highlighted a V γ 4V δ 1 subpopulation, present at variable frequencies in colorectal tissue in different individuals, which underwent TCR-dependent activation in response to BTNL3/8-positive target cells [12]. A priority for future investigations is to understand whether BTNL3/8 act as direct ligands for this V γ 4V δ 1 subpopulation, and whether the TCR repertoire and phenotype of such cells reflect a semi-invariant innate-like paradigm or an adaptive immunobiology. In addition, a more comprehensive understanding of the V δ 1 TCR repertoire in different solid tissues is required.

Current data on peripheral blood V δ 1 T cells indicate a highly diverse TCR repertoire both in terms of V γ chain pairing and V δ 1/V γ CDR3 regions. Notably, there are no obvious similarities in the CDR3 regions of clonally amplified V δ 1 TCRs, either in terms of lengths or sequences. Moreover, these features may apply to other V δ 2-negative TCRs [38]. The high degree of CDR3 diversity within the V δ 1 TCR repertoire, including use of diverse V γ gene segments, contrasts markedly with other unconventional T cell populations thought to recognise single germline-encoded ligands (e.g., iNKTs [15], MAITs [16], and V γ 9/V δ 2 T cells [38,39]), which feature semi-invariant TCR repertoires, including prominent public CDR3 clonotypes; by contrast, the V δ 1 TCR repertoire is both diverse and essentially private [38,39].

Importantly, the high diversity of the peripheral blood V δ 1 TCR repertoire, including in expanded clonotypes, may not necessarily exclude recognition of a limited range of physiologically relevant ligands shared between individuals. Of note, there is precedent for degeneracy in $\alpha\beta$ TCR recognition, whereby the same peptide–MHC complex can be recognised by TCRs of diverse sequence [56]. Similarly, antibodies featuring diverse CDR loops can recognise the same protein antigen, either via distinct or similar surface epitopes [57,58]. However, an alternative, and possibly more likely scenario, is that the high V δ 1 TCR diversity reflects a diversity of ligands recognised. This is arguably supported by the diverse array of ligands proposed for V δ 2-negative TCRs [55,59–62] (Table 1). A major challenge is to make sense of this seemingly unrelated group of ligands and decipher key underlying principles; however, the basis for making such judgements has hitherto remained entirely unclear. Based on the central assumption that TCR–ligand binding (‘receptor occupation’) drives clonal expansion, the adaptive paradigm we outline suggests a set of criteria (based on clonotype frequency, phenotype, CDR3 involvement, and ligand expression pattern; Table 1) by which the position of candidate ligands within this adaptive framework could be assessed.

Re-evaluation of current $\gamma\delta$ T cell ligands in light of these criteria prompts several observations. Since reactivities to CD1c [60] and CD1d [61,63,64] described to date reflected extremely low percentages of the V δ 1 T cell repertoire (contributing <0.05% and <0.16% of total T cells, respectively; Table 1), they are unlikely to represent *in vivo* expanded V δ 1 T_{effector} clonotypes. However, they could derive from V δ 1 T_{naïve} subsets, and conceivably changes in lipid cargo in different physiological settings may drive TCR-mediated V δ 1 clonal expansions. Similarly, phycoerythrin (PE) [62], a model BCR antigen derived from a marine alga, was also convincingly demonstrated to be a direct $\gamma\delta$ TCR ligand in mice, and for an extremely low proportion of the human V δ 1 T cell repertoire (<0.025% of CD3⁺ T cells, Table 1), excluding recognition by

expanded V δ 1 T_{effector} clonotypes. Despite this, PE recognition may reflect the potential of V δ 1 TCRs to recognise foreign antigens in intact form.

One human $\gamma\delta$ TCR reactivity, to Endothelial Protein C Receptor (EPCR) [55], arguably fulfils the criteria for recognition by an expanded V δ 1 T_{effector} clonotype, albeit involving a V δ 5 TCR (Table 1). In addition, the LES TCR that recognised EPCR was a private TCR sequence, in keeping with the properties of V δ 1 TCRs and the finding that recognition of EPCR was restricted to a single patient. Despite this, it enabled $\gamma\delta$ TCR and EPCR-mediated recognition of CMV-infected fibroblast and/or endothelial cells. Interestingly, although EPCR expression was not enhanced by CMV infection, LES $\gamma\delta$ T cell activation was dependent on a TCR-extrinsic 'multimolecular stress signature', which included induction of increased expression of intercellular adhesion molecule 1 (ICAM-1) on target cells after CMV infection [55]. Moreover, while similar changes were evident in some tumour lines, it is also clear that overexpression of EPCR itself is linked to genetic changes during tumourigenesis [65], and has also been linked to chemoresistance [66]. Conceivably, the dependence of effector function on TCR-extrinsic changes in addition to the presence of TCR ligands could be an important factor in the maintenance of $\gamma\delta$ T cell tolerance in the absence of relevant microbial and/or non-microbial stress stimuli. If, as previously suggested, the LES–EPCR reactivity proves to be 'unique but paradigmatic', then other private ligands may map onto other private expanded clonotypes. A repertoire-based ligand identification strategy, ideally focussing on multiple $\gamma\delta$ TCR specificities expanded in response to a single immune challenge, should confirm this, and may reveal commonalities between ligands (e.g., expression on a particular tissue relevant to the specific pathogen infection). In addition, other modes of stress stimulus-induced $\gamma\delta$ TCR–ligand-mediated activation could be envisaged, for example involving increased expression of the ligand, altered post-translational modification of the ligand, or decreased levels of target cell-expressed ligands for $\gamma\delta$ T cell inhibitory receptors. Clearly, future studies in this area, guided by the criteria outlined above, may resolve some of these key questions.

Nature of Immune Stimuli for V δ 1 Responses

Although CMV appears to trigger clonal expansion of V δ 1 T cells, the fact that CMV-seronegative adults still frequently harbour major V δ 1 T_{effector} populations indicates that other immune stimuli, possibly other infectious challenges, must trigger specific V δ 1 responses. Consistent with this, several pathogens have been linked with increased numbers or clonality of V δ 2-negative T cells, including HIV [46], Epstein–Barr virus (EBV) [48,49], and other microbial infections [1]. Despite this, the link between a given pathogen infection, specific V δ 1 clonal expansion, and an augmented recall response to that challenge warrants further study. Several reasons suggest that the transition from V δ 1 T_{naïve} to T_{effector} status accompanying such clonal expansions would increase the speed and potency of effector responses: notably, V δ 1 T_{effector} cells express perforin and granzymes, whereas V δ 1 T_{naïve} cells do not; moreover, V δ 1 T_{effector} cells exhibit enhanced and more rapid production of cytokines and proliferation relative to V δ 1 T_{naïve} cells following CD3/CD28 stimulation [38]. In addition, previous studies have highlighted V δ 2-negative T_{effector} responses to CMV-infected target cells following CMV infection [27], alongside an increased proportion of V δ 1 T cells bearing an effector phenotype relative to CMV-seronegative individuals [29]. However, an increased understanding of how such alterations link with clonotypic changes is needed. Given the diverse stimuli that could underpin the generation of clonotypic V δ 1 T_{effector} responses, this will require analysis of human samples before and after relevant infections, and comparison with individuals who either remained uninfected or did not exhibit postinfection clonal expansions. Such analyses will also allow the relative kinetics of the phenotypic transitions and clonal expansion to be assessed.

Evolutionary Advantage of Adaptive MHC-Unrestricted $\gamma\delta$ T Cell Stress Surveillance

The universal presence of $\gamma\delta$ T cells in vertebrates suggests that compelling reasons must exist for their evolutionary conservation [1]. In addition to providing semi-invariant $\gamma\delta$ T cell populations that have evolved recognition modes highly distinct from $\alpha\beta$ T cell subsets, potentially involving germline-encoded targets, such as BTN3A1 [67,68] and BTNL/Selection and upkeep of intraepithelial T cells (Skint) family members [12,69], it is interesting to consider what evolutionarily advantageous immune functions adaptive $\gamma\delta$ T cell subsets might alternatively provide. With regards to CMV infection, the only pathogenic challenge currently confirmed as driving the clonotypic expansion of human V δ 1 T cells, much evidence supports a role for NK cells [70] and CD8 T cells [71] in anti-CMV immunity. However, CMV has also evolved numerous immune evasion strategies targeted at disrupting essential components of CD8/NK immunosurveillance, including the MHC presentation pathway [72], and sequestration of ligands for conserved activatory ligands for germline-encoded NKRs, such as natural killer group 2 member D (NKG2D) [70]. Evolution of a stochastically recombined immune receptor repertoire allowing MHC-unrestricted recognition of 'altered/stressed self' via diverse and potentially private $\gamma\delta$ TCR reactivities to intact cell surface antigens, would likely complement such strategies and may prove more challenging for pathogens to evade. Conceivably, such subsets could also enable direct recognition of foreign pathogen proteins on the surface of infected host cells. Recent data from mouse models suggesting that $\gamma\delta$ T cells provide as potent protection against CMV as the CD8 T cell compartment [73,74] are consistent with these ideas; however, further studies are required to understand the immunobiology of the $\gamma\delta$ subsets involved and the molecular basis of the recognition events in which they are involved.

Concluding Remarks

Recent studies have substantially revised our understanding of V δ 1 T cells. The coincident clonal expansion and differentiation of V δ 1 T cells not only represent a 'smoking gun' that the $\gamma\delta$ TCR is likely to be central to their immunobiology, but also lead us to propose that V δ 1 function is underpinned by a novel MHC-unrestricted adaptive paradigm. This contrasts with V γ 9/V δ 2 T cell immunobiology, which appears to be predominantly innate-like, highlighting the coexistence of adaptive and innate-like paradigms within the human $\gamma\delta$ T cell lineage.

This adaptive perspective for V δ 1 T cells prompts a reassessment of previous studies, and also provides an intellectual framework around which future investigations can be designed, which should help answer the many unresolved questions (see Outstanding Questions). Of these, the question of what represent *bona fide* physiological ligands for V δ 1 $\gamma\delta$ TCRs in the context of this adaptive paradigm is central, but importantly the repertoire and immunophenotype-based observations upon which the paradigm is built suggest objective criteria by which to assess such reactivities and plan future studies. Moreover, there is an expanding evidence base for the importance of Skint/BTN/BTNL molecules in mouse $\gamma\delta$ T cell development and biology [12,69,75]. Understanding the full significance of this family in human $\gamma\delta$ T cell biology in the context of parallel innate-like and adaptive-like paradigms is another aim.

An improved understanding of human $\gamma\delta$ T cell biology will hopefully accelerate therapeutic exploitation of their function. Despite limited understanding of $\gamma\delta$ TCR ligand recognition, there is already substantial therapeutic interest in $\gamma\delta$ T cells, particularly in the cancer immunotherapy arena, not least due to their MHC-unrestricted recognition of target cells and potent cytotoxic function. In addition, the ability to immunophenotypically delineate V δ 1 T_{naïve} from T_{effector} subpopulations should provide a means to probe V δ 1 T cell responses against diverse microbial and non-microbial immunological challenges at different life stages, which would

Outstanding Questions

What thymic processes shape the V δ 1 repertoire?

How is V δ 1 clonal amplification initiated? Does this occur in lymphoid organs or in peripheral tissues and, if so, which factors and/or antigen-presenting cells are involved?

When do adaptive V δ 1 responses occur throughout life? Do they compensate for when $\alpha\beta$ T cell responses are suppressed, such as in neonates or after transplantation?

Aside from CMV, which stress stimuli trigger clonal selection? What is the range of pathogens involved, and which non-microbial stimuli can drive V δ 1 responses *in vivo*?

What is special about clonally expanded V δ 1 TCRs? Are they capable of recognising ligands that relate to physiologically relevant stress stimuli and, if so, what are these? Are the ligands private to each individual or public, and how do they relate to each other?

Why are the V δ 1 repertoires of some individuals more clonotypically focussed than others? Does this reflect differences in pathogen exposure, or are there 'holes' in the V δ 1 repertoires of some individuals with respect to specific stress challenges?

How do the repertoire and immunobiology of V δ 1 T cells in solid tissues relate to peripheral blood V δ 1 T cells? Do they overlap with adaptive V δ 1 subsets in peripheral blood, or are they distinct, and might they also contain semi-invariant populations, as defined in mouse tissue-associated $\gamma\delta$ T cells?

Is there a role for BTNL molecules (e.g., BTNL3/8) in the development, selection, or antigenic stimulation of the V δ 1 T cell compartment? If so, does this involve innate-like or adaptive V δ 1 populations?

How phenotypically plastic are V δ 1 T cells? Can they be exploited therapeutically, such as in vaccination or adoptive transfer approaches?

represent an important early step along the pathway towards the successful therapeutic exploitation of $\gamma\delta$ T cell function.

Acknowledgements

We thank Adrian Hayday, Anton van der Merwe, and Simon Davis for helpful discussions. This work was supported by Wellcome Trust Investigator award funding to B.E.W., supporting M.S.D. and C.R.W. (Grant code: 099266/Z/12/Z).

References

- Hayday, A.C. (2000) $\gamma\delta$ cells: a right time and a right place for a conserved third way of protection. *Annu. Rev. Immunol.* 18, 975–1026
- Hirano, M. *et al.* (2013) Evolutionary implications of a third lymphocyte lineage in lampreys. *Nature* 501, 435–438
- Burnet, F.M. (1957) A modification of Jerne's theory of antibody production using the concept of clonal selection. *Aust. J. Sci.* 20, 67–69
- Chien, Y.H. *et al.* (1987) A new T-cell receptor gene located within the alpha locus and expressed early in T-cell differentiation. *Nature* 327, 677–682
- Saito, H. *et al.* (1984) A third rearranged and expressed gene in a clone of cytotoxic T lymphocytes. *Nature* 312, 36–40
- Vantourout, P. and Hayday, A. (2013) Six-of-the-best: unique contributions of gammadelta T cells to immunology. *Nat. Rev. Immunol.* 13, 88–100
- Asanow, D.M. *et al.* (1989) Distinct antigen receptor repertoires of two classes of murine epithelium-associated T cells. *Nature* 341, 60–62
- Asanow, D.M. *et al.* (1988) Limited diversity of gamma delta antigen receptor genes of Thy-1+ dendritic epidermal cells. *Cell* 55, 837–847
- Itoharu, S. *et al.* (1990) Homing of a gamma delta thymocyte subset with homogeneous T-cell receptors to mucosal epithelia. *Nature* 343, 754–757
- Jensen, K.D. *et al.* (2008) Thymic selection determines gamma-delta T cell effector fate: antigen-naïve cells make interleukin-17 and antigen-experienced cells make interferon gamma. *Immunity* 29, 90–100
- Turchinovich, G. and Hayday, A.C. (2011) Skint-1 identifies a common molecular mechanism for the development of interferon-gamma-secreting versus interleukin-17-secreting gamma-delta T cells. *Immunity* 35, 59–68
- Di Marco Barros, R. *et al.* (2016) Epithelia use butyrophilin-like molecules to shape organ-specific gammadelta T cell compartments. *Cell* 167, 203–218 e217
- Janeway, C.A., Jr. *et al.* (1988) Specificity and function of T cells bearing gamma delta receptors. *Immunol. Today* 9, 73–76
- De Libero, G. *et al.* (1991) Selection by two powerful antigens may account for the presence of the major population of human peripheral gamma/delta T cells. *J. Exp. Med.* 173, 1311–1322
- Dellabona, P. *et al.* (1994) An invariant V alpha 24-J alpha Q/V beta 11 T cell receptor is expressed in all individuals by clonally expanded CD4⁺ T cells. *J. Exp. Med.* 180, 1171–1176
- Tilloy, F. *et al.* (1999) An invariant T cell receptor alpha chain defines a novel TAP-independent major histocompatibility complex class Ib-restricted alpha/beta T cell subpopulation in mammals. *J. Exp. Med.* 189, 1907–1921
- Hayday, A.C. (2009) Gammadelta T cells and the lymphoid stress-surveillance response. *Immunity* 31, 184–196
- Strid, J. *et al.* (2008) Acute upregulation of an NKG2D ligand promotes rapid reorganization of a local immune compartment with pleiotropic effects on carcinogenesis. *Nat. Immunol.* 9, 146–154
- Garcia, V.E. *et al.* (1998) IL-15 enhances the response of human gamma delta T cells to nonpeptide [correction of nonpeptide] microbial antigens. *J. Immunol.* 160, 4322–4329
- Garcia, V.E. *et al.* (1999) IL-18 promotes type 1 cytokine production from NK cells and T cells in human intracellular infection. *J. Immunol.* 162, 6114–6121
- Rincon-Orozco, B. *et al.* (2005) Activation of V gamma 9V delta 2 T cells by NKG2D. *J. Immunol.* 175, 2144–2151
- Wencker, M. *et al.* (2014) Innate-like T cells straddle innate and adaptive immunity by altering antigen-receptor responsiveness. *Nat. Immunol.* 15, 80–87
- Morita, C.T. *et al.* (1994) TCR usage and functional capabilities of human gamma delta T cells at birth. *J. Immunol.* 153, 3979–3988
- Deusch, K. *et al.* (1991) A major fraction of human intraepithelial lymphocytes simultaneously expresses the gamma/delta T cell receptor, the CD8 accessory molecule and preferentially uses the V delta 1 gene segment. *Eur. J. Immunol.* 21, 1053–1059
- Halstensen, T.S. *et al.* (1989) Intraepithelial T cells of the TCR gamma/delta+ CD8- and V delta 1/J delta 1+ phenotypes are increased in coeliac disease. *Scand. J. Immunol.* 30, 665–672
- Bos, J.D. *et al.* (1990) T-cell receptor gamma delta bearing cells in normal human skin. *J. Invest. Dermatol.* 94, 37–42
- Halary, F. *et al.* (2005) Shared reactivity of V(delta)2(neg) (gamma){delta} T cells against cytomegalovirus-infected cells and tumor intestinal epithelial cells. *J. Exp. Med.* 201, 1567–1578
- Lafarge, X. *et al.* (2005) Expression of MHC class I receptors confers functional intraclonal heterogeneity to a reactive expansion of gammadelta T cells. *Eur. J. Immunol.* 35, 1896–1905
- Pitard, V. *et al.* (2008) Long-term expansion of effector/memory Vdelta2-gammadelta T cells is a specific blood signature of CMV infection. *Blood* 112, 1317–1324
- Woodsworth, D.J. *et al.* (2013) Sequence analysis of T-cell repertoires in health and disease. *Genome Med.* 5, 98
- Yaari, G. and Kleinstein, S.H. (2015) Practical guidelines for B-cell receptor repertoire sequencing analysis. *Genome Med.* 7, 121
- Sherwood, A.M. *et al.* (2011) Deep sequencing of the human TCRgamma and TCRbeta repertoires suggests that TCRbeta rearranges after alphabeta and gammadelta T cell commitment. *Sci. Transl. Med.* 3, 90ra61
- Heather, J.M. *et al.* (2017) High-throughput sequencing of the T-cell receptor repertoire: pitfalls and opportunities. *Brief. Bioinform.* 2017, bbw138
- Bolotin, D.A. *et al.* (2013) MITCR: software for T-cell receptor sequencing data analysis. *Nat. Methods* 10, 813–814
- Bolotin, D.A. *et al.* (2015) MIXCR: software for comprehensive adaptive immunity profiling. *Nat. Methods* 12, 380–381
- Nazarov, V.I. *et al.* (2015) tcR: an R package for T cell receptor repertoire advanced data analysis. *BMC Bioinform.* 16, 175
- Guo, X.Z. *et al.* (2016) Rapid cloning, expression, and functional characterization of paired alphabeta and gammadelta T-cell receptor chains from single-cell analysis. *Mol. Ther. Methods Clin. Dev.* 3, 15054
- Davey, M.S. *et al.* (2017) Clonal selection in the human Vdelta1 T cell repertoire indicates gammadelta TCR-dependent adaptive immune surveillance. *Nat. Commun.* 8, 14760
- Ravens, S. *et al.* (2017) Human gammadelta T cells are quickly reconstituted after stem-cell transplantation and show adaptive clonal expansion in response to viral infection. *Nat. Immunol.* 18, 393–401
- Cairo, C. *et al.* (2010) Impact of age, gender, and race on circulating gammadelta T cells. *Hum. Immunol.* 71, 968–975
- Cairo, C. *et al.* (2013) Human cord blood gammadelta T cells expressing public Vgamma2 chains dominate the response to bisphosphonate plus interleukin-15. *Immunology* 138, 346–360

42. Dimova, T. *et al.* (2015) Effector Vgamma9Vdelta2 T cells dominate the human fetal gammadelta T-cell repertoire. *Proc. Natl. Acad. Sci. U. S. A.* 112, E556–565
43. Rock, E.P. *et al.* (1994) CDR3 length in antigen-specific immune receptors. *J. Exp. Med.* 179, 323–328
44. Dechanet, J. *et al.* (1999) Implication of gammadelta T cells in the human immune response to cytomegalovirus. *J. Clin. Invest.* 103, 1437–1449
45. Lafarge, X. *et al.* (2001) Cytomegalovirus infection in transplant recipients resolves when circulating gammadelta T lymphocytes expand, suggesting a protective antiviral role. *J. Infect. Dis.* 184, 533–541
46. Autran, B. *et al.* (1989) T cell receptor gamma/delta+ lymphocyte subsets during HIV infection. *Clin. Exp. Immunol.* 75, 206–210
47. Vincent, M.S. *et al.* (1996) Apoptosis of Fas-high CD4+ synovial T cells by borrelia-reactive Fas-ligand(high) gamma delta T cells in Lyme arthritis. *J. Exp. Med.* 184, 2109–2117
48. Farnault, L. *et al.* (2013) Clinical evidence implicating gammadelta T cells in EBV control following cord blood transplantation. *Bone Marrow Transplant.* 48, 1478–1479
49. Fujishima, N. *et al.* (2007) Skewed T cell receptor repertoire of Vdelta1(+) gammadelta T lymphocytes after human allogeneic haematopoietic stem cell transplantation and the potential role for Epstein-Barr virus-infected B cells in clonal restriction. *Clin. Exp. Immunol.* 149, 70–79
50. Gerlach, C. *et al.* (2016) The chemokine receptor CX3CR1 defines three antigen-experienced CD8 T Cell subsets with distinct roles in immune surveillance and homeostasis. *Immunity* 45, 1270–1284
51. McVay, L.D. and Carding, S.R. (1996) Extrathymic origin of human gamma delta T cells during fetal development. *J. Immunol.* 157, 2873–2882
52. McVay, L.D. *et al.* (1991) Regulated expression and structure of T cell receptor gamma/delta transcripts in human thymic ontogeny. *EMBO J.* 10, 83–91
53. Parker, C.M. *et al.* (1990) Evidence for extrathymic changes in the T cell receptor gamma/delta repertoire. *J. Exp. Med.* 171, 1597–1612
54. Bessoles, S. *et al.* (2014) Adaptations of natural killer cells to self-MHC Class I. *Front. Immunol.* 5, 349
55. Willcox, C.R. *et al.* (2012) Cytomegalovirus and tumor stress surveillance by binding of a human gammadelta T cell antigen receptor to endothelial protein C receptor. *Nat. Immunol.* 13, 872–879
56. Ding, Y.H. *et al.* (1998) Two human T cell receptors bind in a similar diagonal mode to the HLA-A2/Tax peptide complex using different TCR amino acids. *Immunity* 8, 403–411
57. Goshorn, S.C. *et al.* (1991) Common structural features among monoclonal antibodies binding the same antigenic region of cytochrome c. *J. Biol. Chem.* 266, 2134–2142
58. Padlan, E.A. *et al.* (1989) Structure of an antibody-antigen complex: crystal structure of the HyHEL-10 Fab-lysozyme complex. *Proc. Natl. Acad. Sci. U. S. A.* 86, 5938–5942
59. Marlin, R. *et al.* (2017) Sensing of cell stress by human gamma-delta TCR-dependent recognition of annexin A2. *Proc. Natl. Acad. Sci. U. S. A.* 114, 3163–3168
60. Roy, S. *et al.* (2016) Molecular analysis of lipid-reactive Vdelta1 gammadelta T cells identified by CD1c tetramers. *J. Immunol.* 196, 1933–1942
61. Uldrich, A.P. *et al.* (2013) CD1d-lipid antigen recognition by the gammadelta TCR. *Nat. Immunol.* 14, 1137–1145
62. Zeng, X. *et al.* (2012) Gammadelta T cells recognize a microbial encoded B cell antigen to initiate a rapid antigen-specific interleukin-17 response. *Immunity* 37, 524–534
63. Bai, L. *et al.* (2012) The majority of CD1d-sulfatide-specific T cells in human blood use a semiinvariant Vdelta1 TCR. *Eur. J. Immunol.* 42, 2505–2510
64. Luoma, A.M. *et al.* (2013) Crystal structure of Vdelta1 T cell receptor in complex with CD1d-sulfatide shows MHC-like recognition of a self-lipid by human gammadelta T cells. *Immunity* 39, 1032–1042
65. Lal, N. *et al.* (2017) Endothelial protein C receptor is overexpressed in colorectal cancer as a result of amplification and hypomethylation of chromosome 20q. *J. Pathol. Clin. Res.* 3, 155–170
66. Scheffer, G.L. *et al.* (2002) Expression of the vascular endothelial cell protein C receptor in epithelial tumour cells. *Eur. J. Cancer* 38, 1535–1542
67. Harly, C. *et al.* (2012) Key implication of CD277/butyrophilin-3 (BTN3A) in cellular stress sensing by a major human gammadelta T-cell subset. *Blood* 120, 2269–2279
68. Vavassori, S. *et al.* (2013) Butyrophilin 3A1 binds phosphorylated antigens and stimulates human gammadelta T cells. *Nat. Immunol.* 14, 908–916
69. Barbee, S.D. *et al.* (2011) Skint-1 is a highly specific, unique selecting component for epidermal T cells. *Proc. Natl. Acad. Sci. U. S. A.* 108, 3330–3335
70. Goodier, M.R. *et al.* (2018) CMV and natural killer cells: shaping the response to vaccination. *Eur. J. Immunol.* 48, 50–65
71. Klenerman, P. and Oxenius, A. (2016) T cell responses to cytomegalovirus. *Nat. Rev. Immunol.* 16, 367–377
72. Wiertz, E. *et al.* (1997) Cytomegaloviruses use multiple mechanisms to elude the host immune response. *Immunol. Lett.* 57, 213–216
73. Khairallah, C. *et al.* (2015) Gammadelta T cells confer protection against murine cytomegalovirus (MCMV). *PLoS Pathog.* 11, e1004702
74. Sell, S. *et al.* (2015) Control of murine cytomegalovirus infection by gammadelta T cells. *PLoS Pathog.* 11, e1004481
75. Rhodes, D.A. *et al.* (2016) Regulation of immunity by butyrophilins. *Annu. Rev. Immunol.* 34, 151–172