

## A novel cross-species inhibitor to study the function of CatSper Ca<sup>2+</sup> channels in sperm

Rennhack, Andreas; Schiffer, Christian; Brenker, Christoph; Fridman, Dmitry ; Nitao, Elis; Cheng, Yi-Min; Tamburrino, Lars; Balbach, Melanie ; Stölting, Gabriel; Berger, Thomas K.; Kierzek, Michalina; Alvarez, Luis; Wachten, Dagmar; Zeng, Xu-Hui; Baldi, Elisabetta; Publicover, Stephen; Kaupp, U. Benjamin; Strünker, Timo

DOI:  
[10.1111/bph.14355](https://doi.org/10.1111/bph.14355).

License:  
None: All rights reserved

*Document Version*  
Peer reviewed version

*Citation for published version (Harvard):*  
Rennhack, A, Schiffer, C, Brenker, C, Fridman, D, Nitao, E, Cheng, Y-M, Tamburrino, L, Balbach, M, Stölting, G, Berger, TK, Kierzek, M, Alvarez, L, Wachten, D, Zeng, X-H, Baldi, E, Publicover, S, Kaupp, UB & Strünker, T 2018, 'A novel cross-species inhibitor to study the function of CatSper Ca<sup>2+</sup> channels in sperm', *British Journal of Pharmacology*. <https://doi.org/10.1111/bph.14355>.

[Link to publication on Research at Birmingham portal](#)

### **Publisher Rights Statement:**

This is the peer reviewed version of the following article: Rennhack, A., Schiffer, C., Brenker, C., Fridman, D., Nitao, E. T., Cheng, Y.-M., Tamburrino, L., Balbach, M., Stölting, G., Berger, T. K., Kierzek, M., Alvarez, L., Wachten, D., Zeng, X.-H., Baldi, E., Publicover, S., Kaupp, U. B., and Strünker, T. (2018) A novel cross-species inhibitor to study the function of CatSper Ca<sup>2+</sup> channels in sperm. *British Journal of Pharmacology*, doi: 10.1111/bph.14355., which has been published in final form at [Link to final article using the 10.1111/bph.14355]. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.

### **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

### **Take down policy**

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

# 1 **A novel cross-species inhibitor to study the function of CatSper** 2 **Ca<sup>2+</sup> channels in sperm**

3 Andreas Rennhack<sup>1#</sup>, Christian Schiffer<sup>2</sup>, Christoph Brenker<sup>2</sup>, Dmitry Fridman<sup>1</sup>, Elis T.  
4 Nitao<sup>3</sup>, Yi-Min Cheng<sup>4</sup>, Lara Tamburrino<sup>5</sup>, Melanie Balbach<sup>1</sup>, Gabriel Stölting<sup>6</sup>, Thomas K.  
5 Berger<sup>1</sup>, Michelina Kierzek<sup>2</sup>, Luis Alvarez<sup>1</sup>, Dagmar Wachten<sup>8,9</sup>, Xu-Hui Zeng<sup>4</sup>, Elisabetta  
6 Baldi<sup>5</sup>, Stephen Publicover<sup>3</sup>, U. Benjamin Kaupp<sup>1</sup>, Timo Strünker<sup>2</sup>

7  
8 <sup>1</sup>Center of Advanced European Studies and Research (caesar), Department Molecular Sensory Systems, Bonn,  
9 Germany. <sup>2</sup>University Hospital Münster, Centre of Reproductive Medicine and Andrology, Münster, Germany.  
10 <sup>3</sup>School of Biosciences, University of Birmingham, Edgbaston, Birmingham, UK. <sup>4</sup>Institute of Life Science and  
11 School of Life Science, Nanchang University, Nanchang, Jiangxi, PR China. <sup>5</sup>Department of Experimental and  
12 Clinical Medicine, Center of Excellence DeNothe, University of Florence, Firenze, Italy. <sup>6</sup>Institute of Complex  
13 Systems – Zelluläre Biophysik 4, Forschungszentrum Jülich, Jülich, Germany. <sup>8</sup>Max-Planck Research Group  
14 Molecular Physiology, Center of Advanced European Studies and Research, Bonn, Germany. <sup>9</sup>Institute of Innate  
15 Immunity, University Hospital, University of Bonn, Germany. #present address: Concept Life Science,  
16 Sandwich, UK  
17

## 18 **Background and Purpose**

19 Sperm from many species share the sperm-specific Ca<sup>2+</sup> channel [CatSper](#) (cation channel of  
20 sperm) that controls the intracellular Ca<sup>2+</sup> concentration and, thereby, the swimming  
21 behaviour. A growing body of evidence suggests that the mechanisms controlling CatSper  
22 activity and the role of the channel during fertilization differ among species. However, a lack  
23 of suitable pharmacological tools has hampered the elucidation of the function of CatSper.  
24 Known CatSper inhibitors exhibit considerable side effects and inhibit also [Slo3](#), the K<sup>+</sup>  
25 channel in mammalian sperm.

## 26 **Experimental Approach**

27 The drug RU1968 was reported to suppress Ca<sup>2+</sup> signaling in human sperm by an unknown  
28 mechanism. We resynthesized the drug and revisited its mechanism of action in sperm from  
29 humans, mice, and sea urchins.

## 30 **Key Results**

31 We show by Ca<sup>2+</sup> fluorimetry, single-cell Ca<sup>2+</sup> imaging, electrophysiology, opto-chemistry,  
32 and motility analysis that RU1968 inhibits CatSper in sperm from invertebrates and  
33 mammals. The drug lacks toxic side effects in human sperm, does not affect mouse Slo3, and  
34 inhibits human Slo3 with about 15-fold lower potency than CatSper. Moreover, in human  
35 sperm, the inhibitor mimics CatSper dysfunction and suppresses motility responses evoked by  
36 progesterone, an oviductal steroid that activates CatSper. Finally, we show that the drug  
37 abolishes CatSper-mediated chemotactic navigation in sea urchin sperm.

## 38 **Conclusion and Implications**

39 We propose RU1968 as a novel tool to elucidate the function of CatSper in sperm across  
40 species.

41 **Non-standard abbreviations**

|    |               |  |
|----|---------------|--|
| 42 | 2-AG          | 2-arachidonoylglycerol (2-AG)                    |
| 43 | ABHD2         | alpha/beta hydrolase domain-containing protein 2 |
| 44 | ASW           | artificial sea water                             |
| 45 | BSA           | bovine serum albumin                             |
| 46 | $[Ca^{2+}]_i$ | intracellular $Ca^{2+}$ concentration            |
| 47 | CASA          | computer-assisted sperm analysis                 |
| 48 | CatSper       | cation channel of sperm                          |
| 49 | CI            | confidence interval                              |
| 50 | F             | fluorescence                                     |
| 51 | HC            | HC-056456  |
| 52 | HSA           | human serum albumin                              |
| 53 | HTF           | human tubal fluid                                |
| 54 | LED           | light-emitting diode                             |
| 55 | MDL           | MDL 12330A                                       |
| 56 | NNC           | NNC 0936   |
| 57 | PGE1          | prostaglandin E1                                 |
| 58 | $pH_i$        | intracellular pH                                 |
| 59 | RT            | room temperature                                 |
| 60 | sEBSS         | supplemented Earle's balanced salt solution      |
| 61 | Slo3          | slowpoke channel isoform 3                       |
| 62 | TYH           | Toyoda, Yokoyama and Hosi's medium               |
| 63 | UV            | ultraviolet light                                |
| 64 | VAP           | velocity average path                            |
| 65 | $V_m$         | membrane potential                               |

66

67

68

## 69 Introduction

70 The intracellular  $\text{Ca}^{2+}$  concentration ( $[\text{Ca}^{2+}]_i$ ) modulates the beat of the sperm  
71 flagellum and, thereby, the swimming behavior (Alvarez et al., 2014, Publicover et al., 2008).  
72 In many but not all species,  $[\text{Ca}^{2+}]_i$  is controlled by the sperm-specific CatSper  $\text{Ca}^{2+}$  channel  
73 (Lishko et al., 2010, Ren et al., 2001, Kirichok et al., 2006, Quill et al., 2001, Seifert et al.,  
74 2015, Loux et al., 2013). CatSper appeared early in evolution, before the branching of  
75 eukaryotes into unikonts and bikonts (Cai and Clapham, 2008, Chung et al., 2017, Cai et al.,  
76 2014). So far, most of our knowledge about CatSper originates from physiological studies of  
77 native channels in mammalian and sea urchin sperm. In general, CatSper is activated by  
78 depolarization of the membrane potential ( $V_m$ ) and by alkalization of the intracellular pH  
79 ( $\text{pH}_i$ ) (Lishko et al., 2011, Lishko et al., 2010, Kirichok et al., 2006, Strünker et al., 2011,  
80 Seifert et al., 2015). In sea urchin sperm, the egg's chemoattractant evokes rapid changes in  
81  $V_m$  and  $\text{pH}_i$  and, thereby, stimulates  $\text{Ca}^{2+}$  influx via CatSper (Seifert et al., 2015, Espinal-  
82 Enriquez et al., 2017); the chemoattractant-evoked  $\text{Ca}^{2+}$  influx controls chemotactic steering.  
83 Targeted ablation of genes encoding CatSper subunits provided insight into the function of  
84 CatSper in mouse sperm (Chung et al., 2017, Chung et al., 2011, Wang et al., 2009, Liu et al.,  
85 2007, Carlson et al., 2003, Ren et al., 2001, Quill et al., 2001, Carlson et al., 2005, Qi et al.,  
86 2007, Zeng et al., 2013): mouse  $\text{CatSper}^{-/-}$  sperm suffer from impaired motility (Qi et al.,  
87 2007, Ren et al., 2001, Miki and Clapham, 2013), fail to traverse the oviduct (Ho et al., 2009,  
88 Chung et al., 2014, Miki and Clapham, 2013), and are unable to penetrate the egg coat (Ren et  
89 al., 2001), resulting in male infertility (Qi et al., 2007, Quill et al., 2001, Ren et al., 2001).  
90 CatSper is essential for fertilization also in humans: mutations in *CATSPER* genes (Avenarius  
91 et al., 2009, Hildebrand et al., 2010) and CatSper dysfunction (Williams et al., 2015) are  
92 associated with male infertility. However, mouse and human CatSper have distinct properties,  
93 indicating that the channel might serve various different functions (Alvarez, 2017, Kaupp and  
94 Strünker, 2016). For example, in human but not in mouse sperm, CatSper serves as a  
95 polymodal sensor that integrates diverse chemical cues (Brenker et al., 2018, Schiffer et al.,  
96 2014, Brenker et al., 2012): human CatSper is activated by progesterone and prostaglandins  
97 (Brenker et al., 2012, Lishko et al., 2011, Strünker et al., 2011), two hormones present in the  
98 oviductal fluid (Schuetz and Dubin, 1981). The ensuing  $\text{Ca}^{2+}$  influx controls the swimming  
99 behaviour and promotes the penetration of the egg coat (Schaefer et al., 1998, Harper et al.,  
100 2003, Publicover et al., 2008, Oren-Benaroya et al., 2008, Baldi et al., 2009, Tamburrino et  
101 al., 2015, Tamburrino et al., 2014, Alasmari et al., 2013a, Kilic et al., 2009, Schiffer et al.,  
102 2014). Moreover, progesterone facilitates the migration of human sperm in viscous medium

103 encountered by the sperm during their voyage across the female genital tract (Alasmari et al.,  
104 2013b). However, in humans, neither the role of CatSper nor that of progesterone and  
105 prostaglandins during fertilization has been fully established. The function of CatSper in  
106 species other than sea urchin, mouse, and human is largely unknown. To address these  
107 questions, we rely on pharmacological tools that allow manipulating CatSper function.

108 Several drugs have been identified that suppress CatSper activity, for example [NNC-](#)  
109 [0396](#) (NNC) (Lishko et al., 2011, Strünker et al., 2011), [Mibefradil](#) (Strünker et al., 2011),  
110 MDL12330A (MDL) (Brenker et al., 2012), and [HC-056456](#) (HC) (Carlson et al., 2009). In  
111 patch-clamp experiments, NNC, Mibefradil, and MDL abolish CatSper currents (Brenker et  
112 al., 2012, Lishko et al., 2011, Strünker et al., 2011); HC attenuates CatSper currents (Carlson  
113 et al., 2009), but it is unknown whether the drug inhibits the channel completely. Of note,  
114 none of these drugs is selective for CatSper: the drugs also inhibit the sperm-specific K<sup>+</sup>  
115 channel Slo3 (Carlson et al., 2009, Brenker et al., 2014, Navarro et al., 2007, Mansell et al.,  
116 2014) - the principal K<sup>+</sup> channel in mouse (Zeng et al., 2011, Santi et al., 2010) and human  
117 sperm (Brenker et al., 2014). Notably, each drug inhibits CatSper and Slo3 with similar  
118 potency. Moreover, NNC, Mibefradil, and MDL exhibit serious adverse actions in human  
119 sperm: at high micromolar concentrations required to abolish Ca<sup>2+</sup> influx via CatSper, NNC  
120 and Mibefradil evoke a sizeable and sustained increase of [Ca<sup>2+</sup>]<sub>i</sub> and pH<sub>i</sub> (Strünker et al.,  
121 2011, Brenker et al., 2012, Chavez et al., 2017) and stimulate acrosomal exocytosis (Chavez  
122 et al., 2017) (Figure S3). Similarly, MDL at high micromolar concentrations also evokes a  
123 sustained [Ca<sup>2+</sup>]<sub>i</sub> increase in human sperm (Brenker et al., 2012). Finally, the drugs affect the  
124 vitality and overall motility of sperm (Tamburrino et al., 2014) (Figure S3). HC has not been  
125 further characterized in human sperm, because it is not commercially available. In conclusion,  
126 novel potent and selective CatSper inhibitors without toxic side effects are required.

127 Before the discovery of CatSper, the steroidal sigma-receptor ligand RU1968 was  
128 reported to suppress progesterone- and prostaglandin-induced Ca<sup>2+</sup> signals in human sperm  
129 (Schaefer et al., 2000). The mechanism of RU1968 action in sperm has remained unclear,  
130 except that it does not involve the activation of sigma receptors (Schaefer et al., 2000). We  
131 wondered whether RU1968 might inhibit CatSper and revisited the drug's action in sperm.  
132 We show that RU1968 potently abolishes CatSper-mediated Ca<sup>2+</sup> signals in mouse, human,  
133 and sea urchin sperm. Patch-clamp recordings from mouse and human sperm corroborated  
134 that RU1968 inhibits CatSper. The drug does not affect mouse Slo3 and inhibits human Slo3  
135 with about 15-fold lower potency than human CatSper. When present during the capacitation  
136 process, RU1968 suppresses hyperactivation in human sperm. The drug also inhibits

137 progesterone-evoked motility responses, showing that these involve  $\text{Ca}^{2+}$  influx via CatSper.  
138 Finally, we demonstrate that RU1968 abolishes chemotaxis of sea urchin sperm. In summary,  
139 RU1968 is a potent cross-species CatSper inhibitor that is selective for CatSper over Slo3.  
140 The drug seems well-suited to study CatSper function in sperm from invertebrates to  
141 mammals.

142

## 143 **Material and Methods**

144

### 145 **Sperm preparation**

146 The studies involving human sperm were performed in agreement with the standards  
147 set by the Declaration of Helsinki. Samples of human semen were obtained from volunteers  
148 with their prior written consent. Approval of the institutional ethics committees of the medical  
149 association Westfalen-Lippe and the Medical Faculty of the University of Münster: 4INie;  
150 approval of the ethical committee of the University of Birmingham Life and Health Sciences:  
151 ERN12-0570R. For  $\text{Ca}^{2+}$  fluorimetry in sperm populations, single-cell  $\text{Ca}^{2+}$  imaging, patch-  
152 clamp recordings, single-cell motility studies, and assays for acrosomal exocytosis and  
153 viability, sperm were purified by the swim-up procedure in human tubal fluid (HTF) medium  
154 containing (in mM): 93.8 NaCl, 4.69 KCl, 0.2  $\text{MgSO}_4$ , 0.37  $\text{KH}_2\text{PO}_4$ , 2.04  $\text{CaCl}_2$ , 0.33 Na-  
155 pyruvate, 21.4 lactic acid, 2.78 glucose, 21 HEPES, and 4  $\text{NaHCO}_3$ , pH 7.35 (adjusted with  
156 NaOH). Sperm were washed and re-suspended in HTF containing 3 mg/ml human serum  
157 albumin (HSA, Irvine Scientific, Santa Ana, CA, USA). For Kremer test, sperm were purified  
158 by the swim-up procedure in Supplemented Earle's Balanced Salt Solution (sEBSS),  
159 containing (in mM): 98.5 NaCl, 5.4 KCl, 1.8  $\text{CaCl}_2$ , 1  $\text{MgCl}_2$ , 5.5 glucose, 25  $\text{NaHCO}_3$ , 2.5  
160 Na-pyruvate, 19 Na-lactate, 0.81  $\text{MgSO}_4$ , 15 HEPES, and 0.3% bovine serum albumin, pH  
161 7.4 (adjusted with NaOH). Before experiments, sperm were incubated for at least 300 min at  
162  $37^\circ\text{C}$  and 5%  $\text{CO}_2$  atmosphere. For computer-assisted sperm analysis (CASA), sperm were  
163 purified by the swim-up procedure in HTF lacking  $\text{NaHCO}_3$  and HSA. Sperm were washed  
164 and re-suspended in this medium (non-capacitating conditions) or in HTF fortified with 25  
165 mM  $\text{NaHCO}_3$  and 3 mg/ml HSA (capacitating conditions). Before experiments, sperm were  
166 incubated for at least 300 minutes at  $37^\circ\text{C}$  and 5%  $\text{CO}_2$  atmosphere.

167 C57BL/6N wildtype and C57BL/6N CatSper1<sup>-/-</sup> mice were kept specific pathogen-free  
168 in ventilated cages (Greenline, Tecniplast). Maximally five mice were housed per cage and  
169 handled and sacrificed in accordance with the guideline set by the Animal Center of  
170 Nanchang University (Approval: SYXK2010-0002) and in accordance with the German

171 Animal Welfare Act and the district veterinary office under approval by the LANUV  
172 (AZ.02.05.50.16.011 and AZ.84-02.04.2012.A192). Mouse epididymides were obtained from  
173 at least 15 weeks old male mice that were anaesthetized with CO<sub>2</sub> or isoflurane (Abbvie  
174 Deutschland, Ludwigshafen, Germany) and sacrificed by cervical dislocation. For patch-  
175 clamp recordings, sperm were isolated from the cauda epididymis by swim-out in HS solution  
176 containing (in mM): 135 NaCl, 5 KCl, 1 MgSO<sub>4</sub>, 2 CaCl<sub>2</sub>, 20 HEPES, 5 glucose, 10 lactic  
177 acid, 1 Na-pyruvate, pH 7.4 (adjusted with NaOH). After 20 min swim-out at 37 °C and 10%  
178 CO<sub>2</sub>, the supernatant was collected. Sperm were washed twice and re-suspended in HS  
179 solution. For Ca<sup>2+</sup> fluorimetry, sperm were isolated by swim-out in TYH-medium containing  
180 (in mM): 138 NaCl, 4.8 KCl, 2 CaCl<sub>2</sub>, 1.2 KH<sub>2</sub>PO<sub>4</sub>, 1 MgSO<sub>4</sub>, 5.6 glucose, 0.5 Na-pyruvate,  
181 10 Na-lactate, 10 HEPES, pH 7.4 (adjusted with NaOH). After 15 min swim-out at 37 °C and  
182 5% CO<sub>2</sub>, sperm were counted and capacitated in TYH-medium supplemented with 25 mM  
183 NaHCO<sub>3</sub> and 3 mg/ml BSA.

184 Sperm from the sea urchin *Arbacia punctulata* were obtained by injecting 0.5 M KCl  
185 into the body cavity or electrical stimulation of the animal. The ejaculate (“dry sperm”) was  
186 diluted in artificial sea water (ASW) containing (in mM): 423 NaCl, 9.27 CaCl<sub>2</sub>, 9 KCl, 22.94  
187 MgCl<sub>2</sub>, 25.5 MgSO<sub>4</sub>, 0.1 EDTA, 10 HEPES, pH 7.8 (adjusted with NaOH).

188

### 189 **Measurement of changes in intracellular Ca<sup>2+</sup>**

190 In human sperm populations, changes in [Ca<sup>2+</sup>]<sub>i</sub> and pH<sub>i</sub> were measured with the  
191 fluorescent Ca<sup>2+</sup> indicator Fluo4 and BCECF (Thermo Fisher, Waltham, MA, USA),  
192 respectively, in 384 multi-well plates in a fluorescence plate reader (Fluostar Omega, BMG  
193 Labtech, Ortenberg, Germany) at 29°C, or in a rapid-mixing device in the stopped-flow mode  
194 (SFM400, Bio-Logic, Grenoble, France) at 37°C. Sperm were loaded with Fluo4-AM (10  
195 μM) in the presence of Pluronic F-127 (0.05% w/v) at 37°C for 45 min or with BCECF-AM  
196 (10 μM) at 37°C for 15 min. After incubation, excess dye was removed by centrifugation (700  
197 x g, 10 min, room temperature (RT)). Sperm concentration was adjusted to 5 x 10<sup>6</sup> cells /ml  
198 in HTF and equilibrated for 5 min at 29°C (Fluostar) or 37°C (stopped-flow).

199 In plate-reader experiments, wells were filled with 50 μl of the sperm suspension; the  
200 fluorescence was excited at 480 nm (Fluo-4) or 440 nm and 480 nm (dual excitation, BCECF)  
201 and fluorescence emission was recorded at 520 nm. Fluorescence was monitored before and  
202 after injection of 25 μl (1:3 dilution) RU1968F1, followed after 5 min by injection of stimuli  
203 (1:10 dilution). The solutions were injected into the wells with an electronic multichannel  
204 pipette. Changes in Fluo-4 fluorescence are depicted as ΔF/F (%), i.e. the change in

205 fluorescence ( $\Delta F$ ) relative to the mean basal fluorescence ( $F$ ) before application of buffer or  
206 stimuli, to correct for intra- and inter-experimental variations in basal fluorescence among  
207 individual wells. Changes in BCECF-fluorescence ratio ( $R$ , 480/440 nm) are depicted as  
208  $\Delta R/R$  (%), i.e. the change in ratio ( $\Delta R$ ) relative to the mean basal ratio ( $R$ ) before application  
209 of buffer or stimuli, to correct for intra- and inter-experimental variations in the basal  
210 fluorescence ratio among individual wells. In stopped-flow experiments, the sperm  
211 suspension was rapidly mixed (1:1; flow rate = 1 ml/s) with HTF containing RU1968 and  
212 other stimuli, or with K8.6-, KCl-, or pH<sub>0</sub>8.6-HTF containing RU1968. Fluorescence was  
213 excited with a SpectraX Light Engine modulated at 10 kHz (Lumencor, Beaverton OR, USA)  
214 and passed through a 494/20 nm excitation filter (Semrock, Buffalo NY, USA). Emission was  
215 passed through a 536/40 nm filter (Semrock) and recorded with a photomultiplier (H9656-20;  
216 Hamamatsu Photonics, Hamamatsu, Japan). Signals were amplified with a lock-in amplifier  
217 (7230 DSP, Signal Recovery, Oak Ridge TN, USA) and recorded with a data acquisition pad  
218 (PCI-6221; National Instruments, Germany) and BioKine software v. 4.49 (Bio-Logic). K8.6-  
219 HTF (in mM): 98.5 KCl, 0.2 MgSO<sub>4</sub>, 0.37 KH<sub>2</sub>PO<sub>4</sub>, 2.04 CaCl<sub>2</sub>, 0.33 Na-pyruvate, 21.4 lactic  
220 acid, 2.78 glucose, 21 TAPS, and 4 KHCO<sub>3</sub>, pH 8.6 (adjusted with KOH). KCl-HTF (in mM):  
221 98.5 KCl, 0.2 MgSO<sub>4</sub>, 0.37 KH<sub>2</sub>PO<sub>4</sub>, 2.04 CaCl<sub>2</sub>, 0.33 Na-pyruvate, 21.4 lactic acid, 2.78  
222 glucose, 21 HEPES, and 4 KHCO<sub>3</sub>, pH 7.35 (adjusted with KOH). pH<sub>0</sub>8.6-HTF (in mM):  
223 93.8 NaCl, 4.69 KCl, 0.2 MgSO<sub>4</sub>, 0.37 KH<sub>2</sub>PO<sub>4</sub>, 2.04 CaCl<sub>2</sub>, 0.33 Na-pyruvate, 21.4 lactic  
224 acid, 2.78 glucose, 21 TAPS, and 4 NaHCO<sub>3</sub>, pH 8.6 (adjusted with NaOH). Changes in Fluo-  
225 4 fluorescence are depicted as  $\Delta F/F$  (%), i.e. the change in fluorescence ( $\Delta F$ ) relative to the  
226 fluorescence ( $F$ ) right after the mixing, to correct for intra- and inter-experimental variations  
227 in basal fluorescence. For single-cell Ca<sup>2+</sup> imaging, sperm were incubated in the wells of  
228 PLL-coated Greiner Cellview glass slides with Fluo-4-AM (5  $\mu$ M) for 30 min at 37°C,  
229 followed by another 15 min at room temperature to allow settling of sperm on the glass  
230 surface. Afterwards, the buffer was replaced twice with 90  $\mu$ l of “fresh” HTF to remove  
231 excess extracellular dye. Progesterone and RU1968F1 were injected in a 1:10 dilution (10  $\mu$ l)  
232 into the well and the ensuing changes in [Ca<sup>2+</sup>]<sub>i</sub> were observed under an Olympus IX73  
233 inverted microscope, equipped with a 20x/0.75 objective (U Plan S Apo, Olympus, Germany),  
234 coupled to an Andor Zyla 4.2 sCMOS camera (Andor Technology, Belfast, UK). Images were  
235 captured at 1 Hz. Changes in [Ca<sup>2+</sup>]<sub>i</sub> were determined from a region of interest around the  
236 head and neck of single sperm. Signals are displayed as  $F-F_0/F_{\max}-F_0$ ;  $F_0$  is the mean  
237 fluorescence of  $\geq 5$  images before injection of RU1968F1 or progesterone, whereas  $F_{\max}$  is the  
238 peak fluorescence signal evoked by a subsequent injection of ionomycin. This procedure



239 corrects for intra- and inter-experimental variations in resting  $[Ca^{2+}]_i$  and dye loading among  
240 individual sperm.

241 In mouse sperm populations, changes in  $[Ca^{2+}]_i$  were measured in sperm loaded with  
242 Cal520-AM (5  $\mu$ M) (ATT Bioquest, USA) in the presence of Pluronic F-127 (0.02% w/v) for  
243 45 min at 37°C in TYH buffer. After loading, excess dye was removed by three  
244 centrifugations (700 x g, 7 min, RT). Recordings were performed using the stopped-flow  
245 apparatus as described above, but with mixing at a flow rate of 0.5 ml/s. K8.6-TYH (in mM):  
246 4.8 NaCl, 138 KCl, 2 CaCl<sub>2</sub>, 1.2 KH<sub>2</sub>PO<sub>4</sub>, 1 MgSO<sub>4</sub>, 5.6 glucose, 0.5 Na-pyruvate, 10 lactic  
247 acid, 10 TAPS, pH 8.6 (adjusted with KOH).

248 In sea urchin sperm populations, changes in  $[Ca^{2+}]_i$  were recorded in Fluo4-loaded  
249 sperm. To this end, dry sperm (diluted 1:6 (v/v)) were loaded with Fluo4-AM (10  $\mu$ M) in the  
250 presence of Pluronic F-127 (0.02% w/v) for 45 min at 18°C in ASW. After loading, sperm  
251 were diluted 1:20 (v/v) in ASW and allowed to equilibrate for 5 min. Recordings were  
252 performed using the stopped-flow apparatus with a flow rate of 1 ml/s. Fluorescence was  
253 excited, recorded, and processed as described above.

254 KCl-ASW (in mM): 216 KCl, 216 NaCl, 9.27 CaCl<sub>2</sub>, 22.94 MgCl<sub>2</sub>, 25.5 MgSO<sub>4</sub>, 0.1 EDTA,  
255 10 HEPES, pH 7.8, (adjusted with NaOH).

256

### 257 **Patch-clamp recordings**

258 Patch-clamp recordings from human sperm were performed in the whole-cell  
259 configuration, as previously described (Strünker et al., 2011). Seals between pipette and  
260 sperm were formed either at the cytoplasmic droplet or the neck region in standard  
261 extracellular solution (HS) containing (in mM): 135 NaCl, 5 KCl, 1 MgSO<sub>4</sub>, 2 CaCl<sub>2</sub>, 5  
262 glucose, 1 Na-pyruvate, 10 lactic acid, and 20 HEPES, pH 7.4 (adjusted with NaOH). CatSper  
263 currents were recorded in divalent-free solutions containing (in mM): 140 CsCl, 40 HEPES, 1  
264 EGTA, pH 7.4 (adjusted with CsOH); the pipette solution contained (in mM): 130 Cs-  
265 aspartate, 50 HEPES, 5 EGTA, 5 CsCl, pH 7.3 (adjusted with CsOH). Slo3 currents were  
266 recorded in HS with a pipette solution containing (in mM): 140 K-aspartate, 50 HEPES, 10  
267 NaCl, 5 KCl, 0.5 CaCl<sub>2</sub>, pH 7.3 (adjusted with KOH). Hv1 currents were recorded in a bath  
268 and pipette solution containing (in mM): 120 NMDG, 100 MES, 5 TEA-Cl, 2 EGTA, pH 6  
269 (adjusted with methanesulfonic acid). To depict mean changes in CatSper currents, CatSper  
270 current amplitudes were normalized to that of the monovalent currents in the absence of any  
271 progesterone, NH<sub>4</sub>Cl, or RU1068F1. This procedure corrects for variations in amplitudes  
272 among individual sperm to ease and to improve the clarity of the graphical illustration. Patch-

273 clamp recordings from mouse sperm were performed in the whole-cell configuration, as  
274 previously described (Kirichok et al., 2006, Zeng et al., 2011). Seals between pipette and  
275 sperm were formed at the cytoplasmic droplet. For Slo3 recordings, the extracellular solution  
276 contained (in mM): 160 KOH, 10 HEPES, 150 MES, and 2 Ca(MES)<sub>2</sub>, adjusted to pH 7.4  
277 with MES; the pipette solution contained (in mM): 155 KOH, 5 KCl, 10 BAPTA, 20 HEPES,  
278 115 MES, pH 8.0 (adjusted with KOH). CatSper currents were recorded in divalent-free  
279 solutions containing (in mM): 150 NaCl, 20 HEPES, 5 EDTA, pH 7.4 (adjusted with NaOH);  
280 and with a pipette solution containing (in mM) 135 Cs-MES, 10 HEPES, 10 EGTA, and 5  
281 CsCl, pH 7.2 (adjusted with CsOH). Current amplitudes were normalized to that of the  
282 monovalent currents in the absence of NH<sub>4</sub>Cl or RU1968F1 (control) to correct for variations  
283 in amplitudes among individual sperm.

284 Human T-type (Ca<sub>v</sub>3.2) and L-type (Ca<sub>v</sub>1.2+β2b+α2δ1) Ca<sup>2+</sup> channels were studied in  
285 HEK293T cells (The European Collection of Cell Cultures, Porton Down, UK) that were  
286 cultured according to the supplier's protocol in the presence of penicillin G (100 U/ml) and  
287 streptomycin (10 mg/ml). Cells were transfected at 40% confluency with pcDNA3.1-  
288 CACNA1C, pcDNA3.1-CaVb2b, and pIRES-dsRed-CaVa2d1 in a ratio of 2:1:1 μg, or with  
289 2 μg of pCMV-Entry-CACNA1H, using the calcium-phosphate precipitation method. Patch-  
290 clamp recordings from HEK293T cells were performed in the whole-cell configuration, using  
291 a HEKA EPC 10 amplifier with PatchMaster software (both HEKA Elektronik, Lambrecht,  
292 Germany). The extracellular solution contained (in mM): 125 TEA-Cl, 15 glucose, 10  
293 HEPES, 5 CaCl<sub>2</sub>, pH 7.4, (adjusted with CsOH); the pipette solution contained (in mM): 100  
294 CsCl, 10 EGTA, 10 HEPES, 5 TEA-Cl, 5 MgATP, 0.2 NaGTP, pH 7.4 (adjusted with CsOH).  
295 RU1968F1 was applied via a gravity-driven perfusion system.

296

### 297 **Analysis of sperm motility**

298 To evaluate the acute action of RU1968F1 on motility parameters, sperm from a  
299 particular sample were incubated side-by-side in HTF lacking NaHCO<sub>3</sub> and HSA (non-  
300 capacitating medium) and capacitating medium (25 mM NaHCO<sub>3</sub> / 3 mg/ml HSA). After 3  
301 hours of incubation at 37°C, sperm kinematic parameters were analyzed by a CASA system  
302 (CEROS, Hamilton Thorn Research, Beverly, MA, USA) before and after application of  
303 RU1968F1. Sperm were bathed in RU1968F1 for 5 min prior to the experiment. To evaluate  
304 the long-term action of RU1968F1 on sperm motility parameters, sperm were re-suspended in  
305 capacitating medium (HTF containing 25 mM NaHCO<sub>3</sub> and 3 mg/ml HSA) with or without  
306 RU1968F1. After 3 hours of incubation at 37°C, sperm kinematic parameters were analyzed

307 by CASA. To evaluate the action of RU1968F1 on progesterone-induced hyperactivation,  
308 capacitated sperm were incubated for 5 min in the absence (control) and presence of  
309 progesterone, RU1968F1, and progesterone plus RU1968F1, and the motility was analyzed by  
310 CASA. The following parameters were determined by CASA: curvilinear velocity (VCL,  
311  $\mu\text{m/s}$ ), amplitude of lateral head displacement (ALH,  $\mu\text{m}$ ), linearity of progression (LIN, %),  
312 percentage of total, progressive, and rapid motility as well as percentage of motile,  
313 hyperactivated sperm. The threshold values for hyperactivation were manually set (VCL >150  
314  $\text{mm/s}$ , ALH >7  $\text{mm}$ , LIN < 50% (Mortimer et al., 1998; Tamburrino et al, 2014). A minimum  
315 of 100 cells and 5 fields of view were analysed for each aliquot. The experiments were  
316 performed at 37°C.

317 Motility in human sperm evoked by uncaging of progesterone were studied in an  
318 observation chamber (100  $\mu\text{m}$  depth) under an Olympus IX71 inverted microscope (Olympus,  
319 Tokyo, Japan), equipped with a 4x microscope objective (0.13 NA, UPLFLN-PH, Olympus)  
320 under dark-field illumination (red LED, M660L3-C1, Thorlabs). Movies were recorded at a  
321 total magnification of 6.4x with a high-speed CMOS camera (Dimax HD, PCO, Kelheim,  
322 Germany) at 150 Hz. Photolysis of caged progesterone (1  $\mu\text{M}$ ) (Kilic et al., 2009) was  
323 achieved using a 200 ms light flash delivered by a 365 nm LED (M365L2-C, Thorlabs,  
324 Munich, Germany). Movies were processed and analyzed using a customized CASA  
325 (computer-assisted sperm analysis) plugin for ImageJ. Changes in the average path velocity  
326 (VAP) are depicted as VAP (%), i.e. the change in VAP relative to the VAP right before the  
327 UV flash, to correct for intra- and inter-experimental variations and for the different resting  
328 VAP in the absence and presence of the inhibitor. This procedure eases and improves the  
329 clarity of the graphical illustration.

330 Kremer penetration assays were performed in sEBSS supplemented with  
331 methylcellulose (1% w/v) and 0.3% BSA, equilibrated overnight at 4°C (penetration  
332 medium). The penetration medium, with or without RU1968F1, was filled into flattened glass  
333 capillary tubes (dimensions: 1.2 x 4.8 x 50 mm, 400  $\mu\text{m}$  depth; CM scientific, UK); one end  
334 of the tubes was sealed with CristaSeal wax (Hawksley, UK). The open ends of the tubes  
335 were submersed in a sperm suspension ( $3 \times 10^6$  /ml) with or without stimuli and/or  
336 RU1968F1. Penetration was assessed after 1h (37°C, 5.5%  $\text{CO}_2$ ) by counting sperm at 2 cm  
337 using a phase contrast microscope at a 200x magnification. 3 fields of view were chosen and  
338 in each field, three focal planes were counted, yielding 9 fields altogether.

339 Human sperm viability and motility as shown in Figure S3 was tested following  
340 incubation of sperm for 5 min at room temperature with RU1968F1, NNC-55-0396,

341 Mibefradil, or the vehicle (DMSO). The fraction of immotile and dead sperm was assessed by  
342 counting and by an eosin vitality test, respectively, at 200x magnification under a phase-  
343 contrast microscope (Axiostar, Carl Zeiss), in accordance with the WHO guidelines for semen  
344 analysis (WHO, 2010). For the eosin staining, 5  $\mu$ l of the sperm suspension was mixed with 5  
345  $\mu$ l of eosin staining solution (0.5 % (w/v) eosin Y dissolved in a 0.9 % NaCl solution) on a  
346 microscope slide, covered with a 22 x 22 mm coverslip, and incubated for 30 s at room  
347 temperature. Eosin-positive (dead) vs. eosin-negative (live) sperm were counted. To  
348 determine the fraction of immotile and viable sperm, a total number of 400 sperm was  
349 assessed.

350 Sea urchin sperm chemotaxis was studied as described (Seifert et al., 2015). In brief,  
351 sperm ( $\sim 10^8$  cells/ml) were observed in a recording chamber (150  $\mu$ m depth) under an IX71  
352 microscope (Olympus), equipped with a 10x microscope objective (UPlanSApo; NA 0.4;  
353 Olympus), with stroboscopic (500 Hz) dark-field illumination (white LED; K2 star; Luxeon).  
354 Movies were recorded with an EMCCD camera (DU-897D; Andor) at 20 Hz through a  
355 bandpass filter (HQ520/40; Chroma). Photolysis of caged resact was achieved using a 200 ms  
356 pulse from a 365 nm LED (M365L2-C, Thorlabs). The relative dispersion was calculated as  
357 described before (Seifert et al., 2015).

358

### 359 **Acrosomal exocytosis**

360 Human sperm, capacitated for at least 300 min, were incubated with either 0.1%  
361 DMSO (vehicle control), RU1968F1 (10  $\mu$ M), progesterone (10  $\mu$ M), or a mixture of both (10  
362  $\mu$ M each) for 1 h at 37°C. Afterwards, sperm were washed by centrifugation and re-suspended  
363 in 0.5 ml of hypo-osmotic swelling medium (WHO, 2010). After 1 h at 37°C, sperm were  
364 washed again and fixed in 50  $\mu$ l ice-cold methanol. The sperm were layered on a slide, air-  
365 dried, and stored at -20°C. For acrosome staining, sperm were incubated for 20 min in the  
366 dark with 1 mg/ml FITC-labeled *Arachis hypogaea* (peanut) lectin (PNA-FITC, Sigma  
367 Aldrich) in PBS. Slides were analyzed using an Axiolab A1 FL microscope (Carl Zeiss, Jena,  
368 Germany). For each condition, 200 curled-tail (viable) cells were analyzed for their acrosomal  
369 status, as previously described (Tamburrino et al., 2014).

370

### 371 **Data analysis and statistical evaluation**

372 The data analysis complies with the recommendations on experimental design and  
373 analysis in pharmacology (Curtis et al, 2015). All data are presented as mean  $\pm$  standard  
374 deviation. Statistical analysis and fitting of dose-response relations were performed using

375 GraphPad Prism 5 (Prism, La Jolla, USA). Half-maximal inhibitory concentrations (IC<sub>50</sub>)  
376 were derived by nonlinear regression analysis, using a four parameter fit:

$$377 \quad Y = \text{bottom} + \frac{(\text{top} - \text{bottom})}{(1 + 10^{(\log IC_{50} - x)n})}$$

378 Y = signal amplitude; bottom and top = plateaus in the units of Y; x = log(concentration of  
379 inhibitor); IC<sub>50</sub> = concentration of agonist that gives the response half way between bottom  
380 and top; n = Hill coefficient.

381 Most of the experiments were performed in a randomized block design, i.e. for each  
382 experimental replicate, sperm prepared from one particular semen sample were subjected in  
383 parallel to all treatment conditions. If the experiment involved two conditions (control and  
384 treatment), we used the paired t-test. If the experiment involved  $\geq 3$  conditions, we used one-  
385 way randomized block ANOVA, assuming sphericity. When ANOVA's F-test and the test for  
386 matching efficacy achieved  $P < 0.05$ , means were compared to the control's mean by  
387 Dunnett's multiple comparisons post-hoc test, unless otherwise indicated. If experiments were  
388 not performed in a randomized block design, we used unpaired t-test or one-way ANOVA;  
389 when ANOVA's F-test achieved  $P < 0.05$  and Bartlett's test yielded no significant variance  
390 inhomogeneity, means were compared to the control's mean or to each other by Dunnett's or  
391 Bonferron's multiple comparisons post-hoc test, respectively. In Figure 6E, J, and Figure 8I,  
392 for the ease of illustration and for clarity, we show data normalized to the control. Yet, we  
393 normalized the data only after the statistical analysis using one-way ANOVA, because  
394 normalization makes any data set violate the ANOVA.

395

### 396 **Randomization and blinding**

397 Experiments and data analysis were performed without randomization and blinding,  
398 except for the Eosin test and the manual counting of motile/immotile sperm. Otherwise, non-  
399 treated and treated conditions were measured and analyzed side-by-side by the same  
400 experimenter, using objective measures and analysis methods.

401

### 402 **Nomenclature of targets and ligands**

403 Key protein targets and ligands in this article are hyperlinked to corresponding entries  
404 in <http://www.guidetopharmacology.org>, the common portal for data from the IUPHAR/BPS  
405 Guide to PHARMACOLOGY (Harding et al., 2018), and are permanently archived in the  
406 Concise Guide to PHARMACOLOGY 2017/18 (Alexander et al., 2017a, b).

407

### 408 **Results**

## 409 **Synthesis of RU1968**

410 We synthesized RU1968 from ( $\pm$ ) estrone methyl ether (Figure 1A, SI) that is readily  
411 accessible via the Torgov route (Ananchenko et al., 1962, Ananchenko and Torgov, 1963).  
412 Van Leusen reaction yielded the nitrile (1) (Van Leusen and Van Leusen, 2004), followed by  
413 addition of methyllithium to yield the ketone (2). Reductive amination with *N,N*-  
414 dimethylethylenediamine established the aza side chain, yielding a mixture of four  
415 diastereomers (3) (at C-17 and C-20; Figure 1A, 3). We separated two diastereomers (Figure  
416 1A, compounds 3a and 3b), each a mixture of a *cis* and a *trans* isomer. Relative  
417 configurations (C-18 (CH<sub>3</sub>) and C-20 (CH)) were assigned by NMR spectroscopy (H,H-  
418 COSY and NOE, Figure S1 and S2); the *cis* isomers were the dominant species (*cis/trans*  
419 ratio for compound 3a was 3:1; and for 3b it was 4:1). Finally, cleavage of the phenolic  
420 methyl ether yielded a diastereomeric mixture of RU1968. The diastereomers eluted from a  
421 preparative HPLC in four fractions called RU1968F1-4; RU1968F1 and 2 are derived from 3a  
422 *cis* and *trans*, respectively, whereas RU1968F3 and 4 are derived from 3b *trans* and *cis*,  
423 respectively (Figure 1C). Because the actions of RU1968F2-4 in sperm were similar to that of  
424 RU1968F1 (see Figure 2, S3, and S6), we chose RU1968F1 to characterize its action in  
425 sperm. We first examined the action of RU1968F1 in populations of human sperm loaded  
426 with a fluorescent pH<sub>i</sub> or Ca<sup>2+</sup> indicator (Figure S3A-D). The drug evoked negligible changes  
427 in pH<sub>i</sub>. At concentrations  $\leq 7.5$   $\mu$ M, RU1968F1 caused a small, transient Ca<sup>2+</sup> increase; [Ca<sup>2+</sup>]<sub>i</sub>  
428 peaked and returned to basal levels within about 250 s. At concentrations  $> 7.5$   $\mu$ M,  
429 RU1968F1 evoked a slow decrease of [Ca<sup>2+</sup>]<sub>i</sub>. The mechanism(s) underlying the drug-evoked  
430 changes in [Ca<sup>2+</sup>]<sub>i</sub> are unclear; yet, in the absence of extracellular Ca<sup>2+</sup>, the drug did not  
431 change [Ca<sup>2+</sup>]<sub>i</sub>, indicating that Ca<sup>2+</sup> release from internal stores is not involved (Figure S3C).  
432 Most importantly, even at high micromolar concentrations (30  $\mu$ M), the drug did not affect  
433 overall human sperm motility and viability and did not evoke acrosomal exocytosis (Figure  
434 S3E-G). Thus, RU1968F1 lacks the toxic and adverse actions of NNC, Mibefradil, and MDL  
435 in human sperm.

## 436 **RU1968F1 is a potent cross-species CatSper inhibitor**

437 To investigate whether RU1968F1 inhibits human CatSper, we studied progesterone-  
438 and PGE1-evoked Ca<sup>2+</sup> signals in human sperm bathed in the drug. RU1968F1 slowed down  
439 and completely suppressed the Ca<sup>2+</sup> signals in a dose-dependent fashion; the IC<sub>50</sub> values of  $4$   
440  $\pm 2$   $\mu$ M (progesterone) and  $3.8 \pm 0.5$   $\mu$ M (PGE1) ( $n = 7$ , mean  $\pm$  SD) were similar to those  
441 reported previously (Schaefer et al., 2000) (Figure 2A-D). The drug was effective within a

442 range of extracellular pH ( $pH_o$ ) values: at  $pH_o$  6.8 and 7.8, the  $IC_{50}$  value of the progesterone-  
443 evoked  $Ca^{2+}$  signal was  $4.4 \pm 1.4 \mu M$  and  $2.2 \pm 0.6 \mu M$ , respectively ( $n = 6$ ) (Figure 2B). The  
444 actions of RU1968F2-4 alone and on progesterone-induced  $Ca^{2+}$  signals were similar to those  
445 of RU1968F1 (see Figure 2, S3, and S6).

446 Furthermore, we studied whether RU1968F1 also inhibits  $Ca^{2+}$  signals evoked by  
447 intracellular alkalization via weak bases, e.g.  $NH_4Cl$  (Figure 2E-G). The drug slowed down  
448 and almost completely suppressed  $Ca^{2+}$  signals evoked by  $NH_4Cl \leq 3$  mM (Figure 2E, F); the  
449  $IC_{50}$  value for 3 mM  $NH_4Cl$  was  $4.0 \pm 2.8 \mu M$  ( $n = 5$ ) (Figure 2F).  $Ca^{2+}$  signals evoked by 10  
450 mM  $NH_4Cl$  were only slightly attenuated, whereas for 30 mM  $NH_4Cl$ , the signal was rather  
451 similar in the absence and presence of RU1968F1. We conclude that RU1968F1 inhibits also  
452  $\Delta pH_i$ -evoked  $Ca^{2+}$  responses; its potency seems to decrease with increasing  $\Delta pH_i$ .  
453 Alternatively, the presence of  $NH_4^+$  or  $NH_3$  might impair binding of RU1968F1 to its blocking  
454 site. Arguing against that notion, the drug readily suppressed progesterone responses in sperm  
455 that were bathed for about 20 min in  $NH_4Cl$  (30 mM) (Figure S5;  $IC_{50} = 3.5 \pm 1.4 \mu M$ ,  $n = 3$ );  
456 with time, the  $pH_i$  slowly recovers from the  $NH_4Cl$ -evoked alkalization (Strünker et al. 2011).

457 Influx of  $Ca^{2+}$  via CatSper can also be evoked by simultaneous extracellular  
458 alkalization (which increases  $pH_i$ ) and depolarization by  $K^+$  (K8.6 buffer) (e.g. Carlson et al.,  
459 2003). RU1968F1 suppressed  $Ca^{2+}$  signals evoked by simultaneous  
460 alkalization/depolarization (Figure 2K, L;  $IC_{50} = 1.2 \pm 0.6 \mu M$ ,  $n = 3$ ) and by alkalization or  
461 depolarization alone (Figure S4). Finally, when sperm were mixed simultaneously with  
462 progesterone and RU1968F1 in a stopped-flow apparatus, the inhibition of  $Ca^{2+}$  responses was  
463 similar to that under pre-incubation conditions (Figure 2I, J) ( $IC_{50} = 3.0 \pm 1.1 \mu M$ ;  $n = 4$ ).  
464 This result suggests that RU1968F1 rapidly reaches its blocking site.

465 Next, we studied the action of RU1968F1 in single human sperm by  $Ca^{2+}$  imaging. At  
466 concentrations  $\geq 10 \mu M$ , RU1969F1 evoked a slow decrease of  $[Ca^{2+}]_i$  (Figure 3A); For low  
467 micromolar RU1969F1 concentrations, we did not observe  $Ca^{2+}$  transients, which might  
468 reflect differences in sensitivity of population *versus* single-cell fluorimetry. In the presence  
469 of RU1968F1, progesterone-induced  $Ca^{2+}$  responses were suppressed in a dose-dependent  
470 fashion (Figure 3B-D) with an  $IC_{50}$  of  $4.8 \pm 1.2 \mu M$  (standard error of the fit). Altogether, the  
471 action of the RU1868F1 itself on  $[Ca^{2+}]_i$  and on progesterone-evoked  $Ca^{2+}$  responses is  
472 similar when investigated in sperm populations and in single sperm.

473 We further tested whether the drug also inhibits  $Ca^{2+}$  influx via CatSper in mouse  
474 sperm. Mouse CatSper is insensitive to progesterone and prostaglandins (Lishko et al., 2011).  
475 Therefore, we activated CatSper via simultaneous alkalization/depolarization or via 8-Br-

476 cAMP, which activates mouse (Ren et al., 2001) and human (Brenker et al., 2012) CatSper at  
477 high concentrations. RU1968F1 suppressed  $\text{Ca}^{2+}$  responses evoked by  
478 alkalization/depolarization or 8-Br-cAMP with an  $\text{IC}_{50}$  of  $0.83 \pm 0.07 \mu\text{M}$  and  $0.84 \pm 0.03$   
479  $\mu\text{M}$ , respectively ( $n = 3$ ) (Figure 4A-D).

480 Finally, we investigated the action of RU1968F1 on CatSper in sperm of the sea  
481 urchin *Arbacia punctulata*. To this end, we studied CatSper-mediated  $\text{Ca}^{2+}$  responses evoked  
482 either by the chemoattractant resact, depolarization of  $V_m$ , or by  $\text{NH}_4\text{Cl}$ . Irrespective of the  
483 stimulus, RU1968F1 suppressed the  $\text{Ca}^{2+}$  responses with  $\text{IC}_{50}$  values of  $1.3 \pm 0.1$ ,  $1.1 \pm 0.4$ ,  
484 and  $4 \pm 2$ , respectively ( $n = 3$ ) (Figure 5A-F). Altogether, these results suggest that  
485 RU1968F1 is a potent cross-species CatSper inhibitor.

486 To scrutinize this conclusion by an independent technique, we recorded by whole-cell  
487 patch-clamping CatSper currents in human and mouse sperm. In human sperm, monovalent  
488 CatSper currents were evoked by stepping the membrane voltage from  $-100 \text{ mV}$  to  $+150 \text{ mV}$   
489 in increments of  $10 \text{ mV}$  from a holding potential of  $0 \text{ mV}$ . RU1968F1 completely suppressed  
490 the currents with an  $\text{IC}_{50}$  of  $0.4 \pm 0.3 \mu\text{M}$  ( $n = 5$ ) (Figure 6A, B). Superfusion with  
491 progesterone or  $\text{NH}_4\text{Cl}$  enhanced the current amplitudes (Fig, 6C, D). The progesterone- and  
492  $\text{NH}_4\text{Cl}$ -evoked currents were either completely suppressed or strongly attenuated by  
493 RU1968F1 (Figure 6C-E). In mouse sperm, monovalent CatSper currents were evoked by  
494 ramping the membrane voltage between  $-100$  and  $+100 \text{ mV}$  from a holding potential of  $0 \text{ mV}$ .  
495 Superfusion with RU1968F1 completely suppressed the currents with an  $\text{IC}_{50}$  of  $10 \pm 1 \mu\text{M}$   
496 (Figure 6F, G). CatSper currents evoked at  $\text{pH}_i 8$  and by  $\text{NH}_4\text{Cl}$  were strongly attenuated by  
497 the drug (Figure 6H-J). Thus, RU1968F1 inhibits human and mouse CatSper at rest and upon  
498 activation by ligands and  $\Delta\text{pH}_i$ . Similar to the results obtained by  $\text{Ca}^{2+}$  fluorimetry, the  
499 potency of the drug seems to decrease with increasing  $\text{pH}_i$ . We did not test whether higher  
500 RU1968F1 concentrations completely suppress the currents evoked by  $\text{NH}_4\text{Cl}$  and at  $\text{pH}_i 8$ .

501

### 502 **RU1968F1 inhibits human but not mouse Slo3**

503 We studied the interaction of RU1968F1 with sperm ion channels other than CatSper.  
504 In mouse sperm, currents carried by the  $\text{K}^+$  channel Slo3 were similar in the absence and  
505 presence of RU1968F1 (Figure 7A, B). By contrast, in human sperm, the Slo3 current was  
506 inhibited with an  $\text{IC}_{50}$  of  $7 \pm 6 \mu\text{M}$  ( $n = 4$ ) (Figure 7C, D). Thus, although not perfectly  
507 selective for CatSper, about 15-fold higher RU1968F1 concentrations are required to block  
508 human Slo3 channels. At concentrations up to  $10 \mu\text{M}$ , the drug does not inhibit the voltage-  
509 gated proton channel Hv1 and the ATP-gated  $\text{P}_2\text{X}$  channel (Figure S7), which are expressed



510 in human and mouse sperm, respectively (Navarro et al., 2011, Lishko et al., 2010). We  
511 conclude that in mouse sperm, RU1968F1 acts rather selectively on CatSper. In human sperm,  
512 the drug inhibits also Slo3, yet, with about 15-fold lower potency. Finally, RU1968F1  
513 inhibited heterologously expressed L- and T-type Ca<sup>2+</sup> channels with IC<sub>50</sub> values of about 20  
514 and 10 μM, respectively (Figs. S9, S10), indicating that the drug acts with lower potency also  
515 on classic voltage-gated Ca<sup>2+</sup> channels of somatic cells.

516

### 517 **RU1968F1 suppresses progesterone-evoked motility responses in human sperm**

518 Next, we tested the action of RU1968F1 on the motility of human sperm using  
519 classical computer-assisted sperm analysis (CASA). A brief incubation (5 min) of non-  
520 capacitated or capacitated sperm with the drug did not impair overall motility (Figure 8A, B,  
521 black), whereas the fraction of progressively motile sperm decreased about twofold with  
522 increasing RU1968F1 concentrations (Figure 8A, B, blue). Whether this is due to the  
523 inhibition of CatSper or represents an adverse action of the drug is unclear.

524 Furthermore, the penetration of the egg coat requires hyperactivated motility, which is  
525 characterized by an asymmetric flagellar beat, lower beating frequency, wiggly swimming  
526 trajectory, and lower average path velocity (VAP) (Suarez, 2008). In mouse sperm, CatSper is  
527 required for hyperactivation (Ren et al., 2001), whereas the control of hyperactivation by  
528 CatSper in human sperm is debated (Tamburrino et al., 2014, Alasmari et al., 2013b). We  
529 studied whether RU1968F1 affects hyperactivation in human sperm. A brief incubation (5  
530 min) of capacitated sperm with RU1968F1 did not suppress spontaneous hyperactivated  
531 swimming. In fact, RU1968F1 concentrations < 10 μM seem to slightly enhance  
532 hyperactivation, whereas higher drug concentrations had no effect (Figure 8B, green).  
533 Spontaneous hyperactivation develops during the capacitation process (compare Figure 8A  
534 and B, green). In sperm that were capacitated in the presence of RU1968F1 (10 μM), i.e.  
535 incubated for some hours under capacitating conditions, the drug suppressed spontaneous  
536 hyperactivation (Figure 8C); the fraction of progressively motile sperm or overall motility  
537 was not affected (Figure 8D, E). This result suggests that, in human sperm, CatSper is  
538 involved in the ability to undergo hyperactivation; though, the partial inhibition of Slo3 might  
539 contribute to this action of RU1968F1.

540 Finally, we studied the action of RU1968F1 on progesterone-induced changes in  
541 swimming behavior. Incubation of capacitated sperm with progesterone seemingly promoted  
542 hyperactivation, which was inhibited by RU1968F1 (Figure 8F); the effect of progesterone  
543 was, however, not statistically significant. Therefore, we studied the motility of human sperm

544 before and after rapid activation of CatSper using caged progesterone (Kilic et al., 2009).  
545 Uncaging of progesterone by a brief (200 ms) UV flash instantaneously evoked a wiggly  
546 swimming trajectory (Figure 8G) and a decrease of VAP (Figure 8G, I), reminiscent of  
547 hyperactivated motility. The VAP reached its minimum ~5 s after uncaging of progesterone  
548 and did not recover within the recording time of 10 s. In the presence of RU1968F1, the  
549 swimming trajectory and the swimming pattern and VAP remained unchanged upon uncaging  
550 of progesterone (Figure 8H, I). We conclude that progesterone-evoked hyperactivation  
551 requires  $\text{Ca}^{2+}$  influx via CatSper.

552 Furthermore, it is well established that progesterone facilitates the migration of human  
553 sperm into viscous medium (Alasmari et al., 2013b). In sperm from an infertile man lacking  
554 functional CatSper channels, this facilitation was abolished (Williams et al., 2015). Using a  
555 modified Kremer's sperm-mucus penetration test, we investigated whether CatSper inhibition  
556 by RU1968F1 recapitulates this phenotype. To this end, an open glass capillary, which  
557 contained medium fortified with methylcellulose, was submersed in a sperm suspension. The  
558 number of sperm at a penetration distance of 2 cm (Figure 9A-C) was determined; data for  
559 shorter or longer penetration distances are presented in Figure S8. Consistent with previous  
560 results (Alasmari et al., 2013b), bathing sperm in progesterone enhanced the number of sperm  
561 penetrating the viscous medium (Figure 9A, C; S8A, C). The progesterone action was  
562 abolished by 1  $\mu\text{M}$  RU1968F1 (Figure 9A, S8A). Of note, at this concentration, the drug itself  
563 did not affect the number of penetrating cells (Figure 9B, S8B). The progesterone action was  
564 also abolished when RU1968F1 was added to the capillary medium instead of to the sperm  
565 suspension (Figure 9C, S8C). These results support the notion that progesterone acts via  
566 CatSper to promote swimming in high-viscosity media and shows that RU1968F1 mimics the  
567 lack of functional CatSper channels. Of note, at concentrations  $> 1 \mu\text{M}$ , RU1968F1 in a dose-  
568 dependent fashion lowered the number of penetrating sperm both in the absence and presence  
569 of progesterone (Figure 9A, B; S8A, B); this probably reflects the drug-related decrease of the  
570 fraction of progressively motile sperm.

571 Incubation of human sperm in high micromolar concentrations of progesterone evokes  
572 acrosomal exocytosis (Baldi et al., 2009) (Figure 9D), i.e. the release of proteolytic enzymes  
573 from a secretory vesicle in the sperm head. In the presence of RU1968F1, the progesterone  
574 action was attenuated (Figure 9D), whereas RU1968F1 itself did not evoke acrosomal  
575 exocytosis (Figure 9D, S3). These results suggest that the progesterone-induced acrosome  
576 reaction involves  $\text{Ca}^{2+}$  influx via CatSper and that RU1968F1 might allow unraveling the role  
577 of CatSper in this process in more detail. Altogether, we conclude that RU1968F1 can provide

578 important insight on the role of progesterone action on CatSper to control various sperm  
579 functions.

### 580 **RU19681F1 inhibits chemotaxis of sea urchin sperm**

581 Finally, we tested whether RU1968F1 affects CatSper-mediated chemotactic steering  
582 of sea urchin sperm. In a shallow observation chamber under a dark-field microscope, sperm  
583 were bathed in a caged derivative of the chemoattractant resact (Alvarez et al., 2012, Böhmer  
584 et al., 2005, Kaupp et al., 2003). A chemoattractant gradient was established by photolysis of  
585 caged resact via a UV flash in the center of the recording chamber (Figure 10A). After the  
586 flash, sperm accumulated in the irradiated area, indicated by a decrease of sperm dispersion in  
587 the field of view (Figure 10B); the accumulation was abolished by RU1968F1 (Figure 10), but  
588 the drug did not affect the overall motility of the sperm.

589

### 590 **Discussion**

591 Here, we introduce RU1968F1 as a new pharmacological tool to elucidate the  
592 presence and role of CatSper in sperm. RU1968F1 is superior to hitherto known inhibitors:  
593 the drug is rather selective for CatSper and lacks toxic side effects in human sperm. Yet,  
594 because the action of the drug is complex, we cannot exclude adverse actions in other sperm  
595 species. This cautious note has to be considered in future studies using RU1968F1.

596 What is the molecular mechanism underlying CatSper inhibition by RU1968F1? It has  
597 been proposed that progesterone acts via an endocannabinoid-signaling pathway, involving  
598 the receptor alpha/beta hydrolase domain-containing protein 2 (ABHD2) (Miller et al., 2016):  
599 at rest, CatSper is inhibited by the endocannabinoid 2-arachidonoylglycerol (2-AG) in the  
600 flagellar membrane. Upon progesterone binding, ABHD2 degrades 2-AG and, thereby,  
601 relieves CatSper from inhibition (Miller et al., 2016). Considering that RU1968F1 is a steroid,  
602 the drug might act as an antagonist at the steroid-binding site on ABHD2. However, CatSper  
603 activation by prostaglandins does not involve ABHD2 (Miller et al., 2016), and in mouse and  
604 sea urchin sperm, CatSper is not activated by progesterone or prostaglandins (Lishko et al.,  
605 2011, Seifert et al., 2015). Moreover, CatSper activation by alkaline pH<sub>i</sub> and depolarization  
606 does probably not involve a ligand-binding site. Therefore, we suspect that RU1968F1 binds  
607 to residues in the pore region and, thereby, directly block ion flux. The drug's inhibitory  
608 action on classical voltage-gated Ca<sup>2+</sup> channels also supports this conclusion. Of note, the  
609 potency of RU1968F1 to inhibit activation of CatSper by alkalization seems to decrease with

610 increasing amplitude of  $\Delta\text{pH}_i$ . This might reflect a pH sensitivity of the blocking mechanism  
611 or pH-dependent distribution of the drug across membranes. The latter is rather unlikely: upon  
612 rapid mixing, the drug blocks  $\text{Ca}^{2+}$  influx via CatSper without a measurable latency,  
613 suggesting that the drug acts rather from the outside. However, the mechanism of CatSper  
614 inhibition will be difficult to elucidate rigorously by structure-function analysis or site-  
615 directed mutagenesis, because CatSper resists functional expression.

616         What is the nature of the blocking site in Slo3 in human sperm? Human, but not  
617 mouse Slo3, is inhibited by micromolar concentrations of progesterone (Brenker et al., 2014).  
618 This inhibition results from binding of progesterone either to a site on the channel itself or on  
619 its accessory subunit LRRC52 (Brenker et al., 2014). RU1968F1 might act via this steroid-  
620 binding site on human Slo3. To improve the inhibitor's selectivity, a structure-activity  
621 analysis is required to identify RU1968F1 derivatives that do not act on human Slo3, but  
622 display a similar or even enhanced potency to inhibit CatSper. The fact that human Slo3 can  
623 be functionally expressed in cultured cells (Brenker et al., 2014) facilitates this endeavor.

624         Although the make-up of  $\text{Ca}^{2+}$ -signaling pathways in sperm is quite diverse (Kaupp  
625 and Strünker, 2016, Alvarez, 2017), the CatSper channel is a common component in many,  
626 but not all species (Cai et al., 2014). Our finding that RU1968F1 inhibits CatSper across  
627 species opens the possibility to use the drug in diverse experimental settings. First, teleost fish  
628 seem to lack CatSper genes (Cai and Clapham, 2008), yet, the swimming behavior of  
629 zebrafish is controlled by  $\text{Ca}^{2+}$  (Fechner et al., 2015). However, the absence of CatSper in fish  
630 has been contested (Yanagimachi et al., 2017). RU1968F1 might help to solve this  
631 controversy. Second, the genome of many marine species, including the saprophytic fungus  
632 *Allomyces macrogynus*, the tunicate *Ciona intestinalis*, and the seastar *Asterias amurensis*  
633 harbor CatSper genes (Cai and Clapham, 2008, Cai et al., 2014). Sperm from these species  
634 also undergo chemotaxis (Matsumoto et al., 2003, Miller, 1975, Pommerville, 1978, Yoshida  
635 et al., 2002). RU1968F1 might reveal whether chemotaxis involves CatSper. Third, the drug  
636 might help to define the diverse CatSper functions among mammalian sperm. For example,  
637 mouse sperm undergo rotational motion that governs rheotaxis in gradients of flow velocities  
638 (Miki and Clapham, 2013). By contrast, CatSper<sup>-/-</sup> mouse sperm do not rotate and fail to  
639 undergo rheotaxis, suggesting that  $\text{Ca}^{2+}$  influx via CatSper is required. However, another  
640 study describes rheotaxis as a passive process that does not require  $\text{Ca}^{2+}$  influx (Zhang et al.,  
641 2016). CatSper recruits several proteins into  $\text{Ca}^{2+}$ -signaling domains that form a quadrilateral  
642 arrangement along the flagellar membrane (Chung et al., 2014, Chung et al., 2017). Targeted  
643 deletion of CatSper subunits disrupts these signaling domains (Chung et al., 2014, Chung et

644 al., 2017). Therefore, the motility defects of CatSper<sup>-/-</sup> mouse sperm might be caused by the  
645 lack of Ca<sup>2+</sup> influx via CatSper, by disruption of the supramolecular flagellar ultrastructure, or  
646 by a combination of both. Fourth, in human sperm, neither the role of oviductal CatSper  
647 ligands, nor the role of CatSper during fertilization has been fully established. This is due to  
648 the demanding challenge to mimic the complex chemical, hydrodynamic, and topographical  
649 environment of the oviduct *in vitro* (Xiao et al., 2017). We envision the use of RU1968F1 as a  
650 tool to study the role of CatSper and its ligands in human sperm navigating across artificial or  
651 explanted oviducts.

652 Finally, mutations in *CATSPER* genes (Avenarius et al., 2009, Hildebrand et al., 2010)  
653 and the lack of functional CatSper channels (Williams et al., 2015) are associated with male  
654 infertility. In human sperm, at least *in vitro*, RU1968F1 mimics the lack of CatSper,  
655 indicating that inhibition of CatSper *in vivo* might prevent fertilization. Thus, RU1968F1  
656 could serve as a lead structure to develop new non-hormonal contraceptives. Drugs that  
657 specifically target CatSper should exhibit no side effects, because the expression of the  
658 channel is confined to sperm.

659

## 660 **Author contributions**

661 All persons designated as authors qualify for authorship, and all those who qualify for  
662 authorship are listed. AR and TS conceived and designed the study, coordinated the  
663 experiments, and wrote the manuscript. AR, CS, CB, DF, TEN, YMC, LT, MB, GS, TKB,  
664 MK, LA, DW, XHZ, EB, SP, UBK, and TS acquired, analyzed, and/or interpreted data and  
665 revised the manuscript critically for important intellectual content. All authors approved the  
666 manuscript.

## 667 **Acknowledgments**

668 This work was supported by the German Research Foundation (SFB645 to TS, UBK, and  
669 DW; SFB 1089 to UBK; SPP1926, 1726, and TRR83 to DW; and CRU326 to TS); the Cells-  
670 in-Motion (CiM) Cluster of Excellence, Münster, (FF-2016-17 to TS); the ImmunoSensation  
671 Cluster of Excellence, Bonn (to UBK and DW); the Fritz-Thyssen-Foundation and the  
672 Boehringer Ingelheim Foundation (to DW), the fund “Innovative Medical Research (IMF) of  
673 the University of Münster Medical School, Münster, (I-BR121507 to CB); and the CAPES  
674 Foundation (Ministry of Education, Brazil, to ETN). We thank David Clapham (Janelia  
675 Research Campus, USA) for kindly providing the CatSper1-KO mice.

676 **Conflict of interest**

677 The authors declare that they have no conflict of interest.

678 **References**

679 Alasmari W, Barratt CLR, Publicover SJ, Whalley KM, Foster E, Kay V, *et al.* (2013a). The  
680 clinical significance of calcium-signalling pathways mediating human sperm hyperactivation.  
681 *Hum Reprod* 28: 866-876.

682 Alasmari W, Costello S, Correia J, Oxenham SK, Morris J, Fernandes L, *et al.* (2013b). Ca<sup>2+</sup>  
683 signals generated by CatSper and Ca<sup>2+</sup> stores regulate different behaviors in human sperm. *J*  
684 *Biol Chem* 288: 6248-6258.

685 Alvarez L (2017). The tailored sperm cell. *J Plant Res* 130: 455-464.

686 Alvarez L, Dai L, Friedrich BM, Kashikar ND, Gregor I, Pascal R, *et al.* (2012). The rate of  
687 change in Ca<sup>2+</sup> concentration controls sperm chemotaxis. *J Cell Biol* 196: 653-663.

688 Alvarez L, Friedrich BM, Gompper G, Kaupp UB (2014). The computational sperm cell.  
689 *Trends Cell Biol* 24: 198-207.

690 Alexander SPH, Kelly E, Marrion NV, Peters JA, Faccenda E, Harding SD, *et al.* (2017) The  
691 Concise Guide to PHARMACOLOGY 2017/18. *Br J Pharmacol* 174 (Suppl 1): S1-S446.

692 Alexander SP, Striessnig J, Kelly E, Marrion NV, Peters JA, Faccenda E, *et al.* (2017) The  
693 Concise Guide to PHARMACOLOGY 2017/18: Voltage-gated ion channels. *Br J Pharmacol*  
694 174 Suppl 1:S160-S194. doi: 10.1111/bph.13884.

695 Ananchenko SN, Limanov VY, Leonov VN, Rzhiznikov VN, Torgov IV (1962). Syntheses  
696 of derivatives of oestrane and 19-norsteroids from 6-methoxytetralone and 6-hydroxytetralone.  
697 *Tetrahedron* 18: 1355-1367.

698 Ananchenko SN, Torgov IV (1963). New Syntheses of estrone, d,1-8-iso-estrone and d,1-19-  
699 noresterone. *Tetrahedron Lett* 4: 1553-1558.

700 Avenarius MR, Hildebrand MS, Zhang Y, Meyer NC, Smith LLH, Kahrizi K, *et al.* (2009).  
701 Human male infertility caused by mutations in the CATSPER1 channel protein. *Am J Hum*  
702 *Genet* 84: 505-510.

703 Baldi E, Luconi M, Muratori M, Marchiani S, Tamburrino L, Forti G (2009). Nongenomic  
704 activation of spermatozoa by steroid hormones: facts and fiction. *Mol Cell Endocrinol* 308:  
705 39-46.

706 Böhmer M, Van Q, Weyand I, Hagen V, Beyermann M, Matsomoto M, *et al.* (2005). Ca<sup>2+</sup>  
707 spikes in the flagellum control chemotactic behavior of sperm. *EMBO J* 24: 2741-2752.

708 Brenker C, Goodwin N, Weyand I, Kashikar ND, Naruse M, Krähling M, *et al.* (2012). The  
709 CatSper channel: a polymodal chemosensor in human sperm. *EMBO J* 31: 1654-1665.

710 Brenker C, Zhou Y, Müller A, Echeverry FA, Trötschel C, Poetsch A, *et al.* (2014). The Ca<sup>2+</sup>  
711 -activated K<sup>+</sup> current of human sperm is mediated by Slo3. *Elife* 3: e01438.

712 Brenker C, Schiffer C, Wagner IV, Tüttelmann F, Röpke A, Rennhack A, *et al.* (2018).  
713 Action of steroids and plant triterpenoids on CatSper Ca<sup>2+</sup> channels in human sperm. *Proc*  
714 *Natl Acad Sci USA* 115(3):E344-E346.

715 Cai X, Clapham DE (2008). Evolutionary genomics reveals lineage-specific gene loss and  
716 rapid evolution of a sperm-specific ion channel complex: CatSper and CasSper $\beta$ . *PLoS One*  
717 3: e3569.

718 Cai X, Wang X, Clapham DE (2014). Early evolution of the eukaryotic Ca<sup>2+</sup> signaling  
719 machinery: conservation of the CatSper channel complex. *Mol Biol Evol* 31: 2735-2740.

720 Carlson AE, Burnett LA, del Camino D, Quill TA, Hille B, Chong JA, *et al.* (2009).  
721 Pharmacological targeting of native CatSper channels reveals a required role in maintenance  
722 of sperm hyperactivation. *PLoS One* 4: e6844.

723 Carlson AE, Quill TA, Westenbroek RE, Schuh SM, Hille B, Babcock DF (2005). Identical  
724 phenotypes of CatSper1 and CatSper2 null sperm. *J Biol Chem* 280: 32238-32244.

725 Carlson AE, Westenbroek RE, Quill TA, Ren D, Clapham DE, Hille B, *et al.* (2003).  
726 CatSper1 required for evoked Ca<sup>2+</sup> entry and control of flagellar function in sperm. *Proc Natl*  
727 *Acad Sci USA* 100: 14864-14868.

728 Chávez JC, De la Vega-Beltrán JL, José O, Torres P, Nishigaki T, Treviño CL, *et al.* (2017).  
729 Acrosomal alkalization triggers Ca<sup>2+</sup> release and acrosome reaction in mammalian  
730 spermatozoa. *J Cell Physiol*. doi: 10.1002/jcp.26262.

731

732 Chung JJ, Miki M, Kim D, Shim SH, Shi HF, Hwang JY, *et al.* (2017). CatSper $\zeta$  regulates the  
733 structural continuity of sperm Ca<sup>2+</sup> signaling domains and is required for normal fertility.  
734 *Elife* 6: e23082.

735 Chung JJ, Navarro B, Krapivinsky G, Krapivinsky L, Clapham DE (2011). A novel gene  
736 required for male fertility and functional CatSper channel formation in spermatozoa. *Nat*  
737 *Commun* 2: 153.

738 Chung JJ, Shim SH, Everley RA, Gygi SP, Zhuang X, Clapham DE (2014). Structurally  
739 distinct Ca<sup>2+</sup> signaling domains of sperm flagella orchestrate tyrosine phosphorylation and  
740 motility. *Cell* 157: 808-822.

741 Curtis MJ, Bond RA, Spina D, Ahluwalia A, Alexander SP, Giembycz MA, (2015) *et al.*  
742 Experimental design and analysis and their reporting: new guidance for publication in *BJP*. *Br*  
743 *J Pharmacol* 172(14):3461-71

744 Espinal-Enríquez J, Priego-Espinosa DA, Darszon A, Beltrán C, Martínez-Mekler G (2017).  
745 Network model predicts that CatSper is the main Ca<sup>2+</sup> channel in the regulation of sea urchin  
746 sperm motility. *Sci Rep* 7(1):4236.

747 Fechner S, Alvarez L, Bönigk W, Müller A, Berger TK, Pascal R, *et al.* (2015). A K<sup>+</sup>-  
748 selective CNG channel orchestrates Ca<sup>2+</sup> signaling in zebrafish sperm. *Elife* 4: e07624.

749 Harding SD, Sharman JL, Faccenda E, Southan C, Pawson AJ, Ireland S *et al.* (2018). The  
750 IUPHAR/BPS Guide to PHARMACOLOGY in 2018: updates and expansion to encompass  
751 the new guide to IMMUNOPHARMACOLOGY. *Nucl Acids Res* 46: D1091-D1106.

752 Harper CV, Kirkman-Brown JC, Barratt CLR, Publicover SJ (2003). Encoding of  
753 progesterone stimulus intensity by intracellular [Ca<sup>2+</sup>] ([Ca<sup>2+</sup>]<sub>i</sub>) in human spermatozoa.  
754 *Biochem J* 372: 407-417.

755 Hildebrand MS, Avenarius MR, Fellous M, Zhang Y, Meyer NC, Auer J, *et al.* (2010).  
756 Genetic male infertility and mutation of CatSper ion channels. *Eur J Hum Genet* 18: 1178-  
757 1184.

758 Ho K, Wolff CA, Suarez SS (2009). CatSper-null mutant spermatozoa are unable to ascend  
759 beyond the oviductal reservoir. *Reprod Feril Dev* 21: 345-350.

760 Kaupp UB, Solzin J, Hildebrand E, Brown JE, Helbig A, Hagen V, *et al.* (2003). The signal  
761 flow and motor response controlling chemotaxis of sea urchin sperm. *Nat Cell Biol* 5: 109-  
762 117.

763 Kaupp UB, Strünker T (2016). Signaling in sperm: more different than similar. *Trends Cell*  
764 *Biol* 27: 101-109.

765 Kilic F, Kashikar ND, Schmidt R, Alvarez L, Dai L, Weyand I, *et al.* (2009). Caged  
766 progesterone: a new tool for studying rapid nongenomic actions of progesterone. *J Am Chem*  
767 *Soc* 131: 4027-4030.

768 Kirichok Y, Navarro B, Clapham DE (2006). Whole-cell patch-clamp measurements of  
769 spermatozoa reveal an alkaline-activated Ca<sup>2+</sup> channel. *Nature* 439: 737-740.

770 Lishko PV, Botchkina IL, Federenko A, Kirichok Y (2010). Acid extrusion from human  
771 spermatozoa is mediated by flagellar voltage-gated proton channel. *Cell* 140: 327-337.

772 Lishko PV, Botchkina IL, Kirichok Y (2011). Progesterone activates the principal Ca<sup>2+</sup>  
773 channel of human sperm. *Nature* 471: 387-391.

774 Liu J, Xia J, Cho KH, Clapham DE, Ren D (2007). CatSperβ, a novel transmembrane protein  
775 in the CatSper channel complex. *J Biol Chem* 282: 18945-18952.

776 Loux SC, Crawford KR, Ing NH, Gonzales-Fernandez L, Macias-Garcia B, Love C, *et al.*  
777 (2013). CatSper and the relationship of hyperactivated motility to intracellular calcium and  
778 pH kinetics in equine sperm. *Biol Reprod* 89: 123.

779 Mansell SA, Publicover SJ, Barratt CLR, Wilson SM (2014). Patch clamp studies of human  
780 sperm under physiological ionic conditions reveal three functionally and pharmacologically  
781 distinct cation channels. *Mol Hum Reprod* 20: 392-408.



782 Matsumoto M, Solzin J, Helbig A, Hagen V, Ueno SI, Kawase O, *et al.* (2003). A sperm-  
783 activating peptide controls a cGMP-signaling pathway in starfish sperm. *Dev Biol* 260: 314-  
784 324.

785 Miki M, Clapham DE (2013). Rheotaxis guides mammalian sperm. *Curr Biol* 23: 443-452.

786 Miller MR, Mannowetz N, Iavarone AT, Safavi R, Gracheva EO, Smith JF, *et al.* (2016).  
787 Unconventional endocannabinoid signaling governs sperm activation via the sex hormone  
788 progesterone. *Science* 352: 555-559.

789 Miller RL (1975) Chemotaxis of the spermatozoa of *Ciona intestinalis*. *Nature* 254: 244-245.

790 Mortimer ST, Swan MA, Mortimer D (1998). Effect of seminal plasma on capacitation and  
791 hyperactivation in human spermatozoa. *Hum Reprod* 13(8):2139-46.

792 Navarro B, Kirichok Y, Clapham DE (2007). KSper, a pH-sensitive K<sup>+</sup> current that controls  
793 sperm membrane potential. *Proc Natl Acad Sci USA* 104: 7688-7692.

794 Navarro B, Miki K, Clapham DE (2011). ATP-activated P2X2 current in mouse spermatozoa.  
795 *Proc Natl Acad Sci USA* 108: 14342-14347.

796 Oren-Benaroya R, Orvieto R, Gakamsky A, Pinchasov M, Eisenbach M (2008). The sperm  
797 chemoattractant secreted from human cumulus cells is progesterone. *Hum Reprod* 23: 2339-  
798 2345.

799 Pommerville J (1978). Analysis of gamete and zygote motility in *allomyces*. *Exp Cell Res*  
800 113: 161-172.

801 Publicover SJ, Giojalas LC, Teves ME, Mendes Machado de Oliveira GS, Morales Garcia  
802 AA, Barratt CLR, *et al.* (2008). Ca<sup>2+</sup> signaling in the control of motility and guidance in  
803 mammalian sperm. *Front Biosci* 13: 5623-5637.

804 Qi H, Moran MM, Navarro B, Chong JA, Krapivinsky G, Krapivinsky L, *et al.* (2007). All  
805 four CatSper ion channel proteins are required for male fertility and sperm cell hyperactivated  
806 motility. *Proc Natl Acad Sci USA* 104: 1219-1223.

807 Quill TA, Ren D, Clapham DE, Garbers DL (2001). A sperm ion channel required for sperm  
808 motility and male fertility. *Nature* 413: 603-609.

809 Santi CM, Martinez-Lopez P, De la Vega-Beltran JL, Butler A, Alisio A, Darszon A, *et al.*  
810 (2010). The Slo3 sperm-specific potassium channel plays a vital role in male fertility. *FEBS*  
811 *Lett* 584: 1041-1046.

812 Schaefer M, Habenicht UF, Bräutigam M, Gudermann T (2000). Steroidal sigma receptor  
813 ligands affect signaling pathways in human spermatozoa. *Biol Reprod* 63: 57-63.

814 Schaefer M, Hofmann T, Schultz G, Gudermann T (1998). A new prostaglandin E receptor  
815 mediates calcium influx and acrosome reaction in human spermatozoa. *Proc Natl Acad Sci*  
816 *USA* 95: 3008-3013.

817 Schiffer C, Müller A, Egeberg DL, Alvarez L, Brenker C, Rehfeld A, *et al.* (2014). Direct  
818 action of endocrine disrupting chemicals on human sperm. *EMBO Rep* 15: 758-765.

819 Schuetz AW, Dubin NH (1981). Progesterone and prostaglandin secretion by ovulated rat  
820 cumulus cell-oocyte complexes. *Endocrinology* 108: 457-463.

821 Seifert R, Flick M, Bönigk W, Alvarez L, Trötschel C, Poetsch A, *et al.* (2015). The CatSper  
822 channel controls chemosensation in sea urchin sperm. *EMBO J* 34: 379-392.

823 Strünker T, Goodwin N, Brenker C, Kashikar ND, Weyand I, Seifert R, *et al.* (2011). The  
824 CatSper channel mediates progesterone induced  $Ca^{2+}$  influx in human sperm. *Nature* 471:  
825 382-386.

826 Suarez SS (2008). Control of hyperactivation in sperm. *Hum Reprod Update* 14: 647-657.

827 Tamburrino L, Marchiani S, Minetti F, Forti G, Muratori M, Baldi E (2014). The CatSper  
828 calcium channel in human sperm: relation with motility and involvement in progesterone-  
829 induced acrosome reaction. *Hum Reprod* 29: 418-428.

830 Tamburrino L, Marchiani, S, Vicini E, Muciaccia B, Cambi M, Pellegrini S, *et al.* (2015).  
831 Quantification of CatSper1 expression in human spermatozoa and relation to functional  
832 parameters. *Hum Reprod* 30: 1532-1544.

833 Van Leusen D, Van Leusen AM (2004). Synthetic uses of tosylmethyl isocyanide (TosMic).  
834 *Organic Reactions* 57: 417-666.

835 Wang H, Liu J, Cho KH, Ren D (2009). A novel, single, transmembrane protein CatSperγ is  
836 associated with CatSper1 channel protein. *Biol Reprod* 81: 539-544.

837 Williams HL, Mansell S, Alasmari W, Brown SG, Wilson SM, Sutton KA, *et al.* (2015).  
838 Specific loss of CatSper function is sufficient to compromise fertilization capacity of human  
839 spermatozoa. *Hum Reprod* 30: 2737-2746.

840 Yanagimachi R, Harumi T, Matsubara H, Yan W, Yuan S, Hirohashi N, *et al.* (2017).  
841 Chemical and physical guidance of fish spermatozoa into the egg through the micropyle. *Biol*  
842 *Reprod* 1;96(4):780-799.

843 Xiao S, Copetta JR, Rogers HB, Isenberg BC, Zhu J, Olalekan SA, *et al.* (2017). A  
844 microfluidic culture model of the human reproductive tract and 28-day menstrual cycle. *Nat*  
845 *Commun* 8: 14584.

846 Yoshida M, Murata M, Inaba K, Morisawa M (2002). A chemoattractant for ascidian  
847 spermatozoa is a sulfated steroid. *Proc Natl Acad Sci USA* 99: 14831-14836.

848 Zhang Z, Liu J, Meriano J, Ru C, Xie S, Luo J, Sun Y (2016). Human sperm rheotaxis: a  
849 passive physical process. *Sci Rep* 23;6:23553.

850 Zeng XH, Navarro B, Xia XM, Clapham DE, Lingle CJ (2013). Simultaneous knockout of  
851 Slo3 and CatSper1 abolishes all alkalization- and voltage-activated currents in mouse  
852 spermatozoa. *J Gen Physiol* 142: 305-313.

853 Zeng XH, Yang C, Kim ST, Lingle CJ, Xia XM (2011). Deletion of the Slo3 gene abolishes  
854 alkalization-activated K<sup>+</sup> current in mouse spermatozoa. Proc Natl Acad Sci USA 108: 5879-  
855 5884.

856

857

858 **Figure legends**

859

860 **Figure 1: Synthesis of RU1968F1-F4** (A) Synthesis of RU1968. Carbon atoms referred to in the text are  
861 marked with circles. (B) Structure of RU1968. (C) HPLC-elution profile of the four diastereomers. Isomers are  
862 named according to their order of elution.

863

864 **Figure 2: Action of RU1968F1 on CatSper-mediated Ca<sup>2+</sup> signals in human sperm populations.** (A)  
865 Progesterone-induced Ca<sup>2+</sup> signals in human sperm in the absence and presence of RU1968F1.  $\Delta F/F$  (%)  
866 indicates the percentage change in fluorescence ( $\Delta F$ ) with respect to the mean basal fluorescence ( $F$ ) before  
867 application of progesterone (500 nM). (B) Dose-response relation for the maximal signal amplitudes of the data  
868 from (A) (pH 7.35) ( $IC_{50} = 5.5 \mu M$ ) and of progesterone responses studied at an extracellular pH of 6.8 ( $IC_{50} =$   
869  $4.2 \mu M$ ) or 7.8 ( $IC_{50} = 2.6 \mu M$ ). (C) PGE1-induced Ca<sup>2+</sup> signals in human sperm in the absence and presence of  
870 RU1968F1; PGE1 = 500 nM. (D) Dose-response relation for the maximal signal amplitudes of the data from (C)  
871 ( $IC_{50} = 3.1 \mu M$ ). (E) NH<sub>4</sub>Cl-induced Ca<sup>2+</sup> signals in the absence and presence of RU1968F1, NH<sub>4</sub>Cl = 3 mM. (F)  
872 Dose-response relation for the maximal signal amplitude of the data from (E) ( $IC_{50} = 1.8 \mu M$ ) (G) Ca<sup>2+</sup> signals  
873 evoked by various NH<sub>4</sub>Cl concentrations in the absence (control) and presence of RU1968F1 (30  $\mu M$ ). (H) Mean  
874 relative amplitude of Ca<sup>2+</sup> signals evoked by various NH<sub>4</sub>Cl concentrations in the presence of RU1968F1  
875 (30  $\mu M$ ) (n = 5); amplitude evoked in the absence of RU1968F1 = 1 (control). Error bars indicate SD. \*P < 0.05  
876 versus control. (I) Ca<sup>2+</sup> signals evoked by simultaneous mixing of sperm with progesterone (500 nM) and  
877 RU1968F1 in a stopped-flow apparatus.  $\Delta F/F$  (%) indicates the percentage change in fluorescence ( $\Delta F$ ) with  
878 respect to the fluorescence ( $F$ ) immediately after mixing. (J) Dose-response relation of the data from (I) ( $IC_{50} =$   
879  $2.4 \mu M$ ). (K) Ca<sup>2+</sup> signals evoked by mixing of sperm with K8.6-HTF and RU1968F1. The final K<sup>+</sup>  
880 concentration and pH after mixing was 51.25 mM and 8.1, respectively. (L) Dose-response relation of the data  
881 from (K) ( $IC_{50} = 1.7 \mu M$ ).  
882

883

884 **Figure 3: Action of RU1968F1 on progesterone-evoked Ca<sup>2+</sup> signals in single human sperm.** (A) Changes in  
885 [Ca<sup>2+</sup>]<sub>i</sub> evoked by RU1968F1 in immobilized sperm. Sperm were challenged with RU1968F1 at t = 0. Traces  
886 represent averages of 122 (0 RU1968F1), 344 (1), 42, (3), 165, (10), and 109 (30) sperm from 3 donors. Signals  
887 are displayed as  $F - F_0 / F_{max} - F_0$ ;  $F_0$  is the mean fluorescence of  $\geq 5$  images before application of RU1968F1;  $F_{max}$   
888 is the peak fluorescence signal evoked by ionomycin (not shown) to gauge the maximal response amplitude. (B)  
889 Progesterone-evoked Ca<sup>2+</sup> responses (2  $\mu M$ ) in the absence and presence of RU1968F1 (30  $\mu M$ ); averages of 50  
890 (control) and 109 (RU1968F1) sperm from 3 donors. Progesterone and RU1968F1 were applied at t = 0;  
891 following application of RU1968F1, progesterone was applied at the time point indicated by the arrow. (C)  
892 Amplitude of progesterone-evoked Ca<sup>2+</sup> responses in the absence and presence of different RU1968F1  
893 concentrations; averages of 122 (0  $\mu M$  RU1968F1), 344 (1), 42 (3), 165 (10), 109 (30) sperm from 3 donors. (D)  
894 Mean amplitude of progesterone-evoked Ca<sup>2+</sup> signals (2  $\mu M$ ) in the presence of RU1968F1; number of sperm:  
895 310 (0  $\mu M$  RU1968F1), 552 (1), 181 (3) 302 (10), 222 (30). Error bars indicate SD. Fitting of a dose-response  
896 curve to the data yielded an  $IC_{50}$  of  $4.8 \pm 1.2 \mu M$  (standard error of the fit).

897

897 **Figure 4: Action of RU1968F1 on CatSper-mediated Ca<sup>2+</sup> signals in mouse sperm populations.** (A) Ca<sup>2+</sup>  
898 signals evoked by simultaneous mixing of mouse sperm with K8.6-TYH and RU1968F1 in a stopped-flow  
899 apparatus. After mixing, the final K<sup>+</sup> concentration and pH was 69 mM and 8.1, respectively (B) Dose-response  
900 relation of the data from (A) ( $IC_{50} = 0.90 \mu M$ ). (C) Ca<sup>2+</sup> signals evoked by simultaneous mixing of mouse sperm  
901 with 8-Br-cAMP (20 mM) and RU1968F1. (D) Dose-response relation of the data from (C) ( $IC_{50} = 0.84 \mu M$ ).

902

902 **Figure 5: Action of RU1968F1 on CatSper-mediated Ca<sup>2+</sup> responses in sea urchin sperm.** (A) Resact-  
903 induced Ca<sup>2+</sup> signals in sea urchin sperm evoked by simultaneous mixing of sperm with resact (20 pM) and  
904 RU1968F1 in a stopped-flow apparatus. (B) Dose-response relation of the data from (A) ( $IC_{50} = 0.7 \mu M$ ). (C)  
905 Depolarization-induced Ca<sup>2+</sup> signals evoked by mixing of sperm with KCl-ASW and RU1968F1. Final K<sup>+</sup>  
906 concentration after mixing was 108 mM. (D) Dose-response relation of the data from (C) ( $IC_{50} = 1.0 \mu M$ ). (E)  
907 Alkaline-evoked Ca<sup>2+</sup> signals in the presence of RU1968F1; the final NH<sub>4</sub>Cl concentration after mixing was 30  
908 mM. (F) Dose-response relation of the data from (E) ( $IC_{50} = 4.6 \mu M$ ).  
909  
910  
911  
912  
913  
914  
915

916 **Figure 6: RU1968F1 inhibits monovalent CatSper currents in human and mouse sperm.** (A) Representative  
917 current-voltage relationship of CatSper currents recorded from a human sperm cell in divalent-free extracellular  
918 and intracellular solution (pH 7.4) in the absence and presence of increasing RU1968F1 concentrations. Voltage  
919 was stepped from -100 mV to +150 mV in increments of 10 mV. Inset: Voltage protocol. (B) Dose-response  
920 relation for the inhibition of human CatSper currents by RU1968F1 at +100 mV ( $IC_{50} = 0.4 \pm 0.3 \mu M$ ;  $n = 5$ ).  
921 (C) Representative monovalent CatSper currents recorded from a human sperm cell before (control) and after  
922 perfusion with progesterone (2 $\mu M$ ) and progesterone plus RU1968F1 (3 $\mu M$ ), evoked by the voltage protocol  
923 shown in (A). The dotted red line indicates the current at 0 mV. (D) Monovalent CatSper currents recorded from  
924 a human sperm cell before (control) and after perfusion with  $NH_4Cl$  (10 mM) and  $NH_4Cl$  plus RU1968F1  
925 (3 $\mu M$ ), evoked by the voltage protocol shown in (A). The dotted red line indicates the current at 0 mV. (E) Mean  
926 amplitudes of monovalent currents at +100 mV recorded in the presence of RU1968F1, progesterone (2  $\mu M$ ),  
927 progesterone plus RU1968F1,  $NH_4Cl$  (10 mM), and  $NH_4Cl$  plus RU1968F1. Amplitudes were normalized to that  
928 evoked in the absence of any drug (control, dashed line). Error bars indicate SD ( $n = 5$ ). \* $P < 0.05$  versus  
929 control. Data were normalized only after performing the statistical analysis using one-way ANOVA (see  
930 Methods for details and explanations). (F) Representative CatSper currents recorded from a mouse sperm cell in  
931 divalent-free extracellular and intracellular solution (pH 7.2) in the absence and presence of increasing  
932 RU1968F1 concentrations. Voltage was ramped between -100 and +100 mV from a holding potential of 0 mV.  
933 Inset: Voltage protocol. (G) Dose-response relation for the inhibition of mouse CatSper currents at +100 mV  
934 ( $IC_{50} = 10 \pm 1 \mu M$ ,  $n = 3$ ). Error bars indicate SD. (H) Currents in the presence of extracellular divalent ions  
935 (HS) and monovalent currents in divalent-free conditions (control) recorded from a mouse sperm cell evoked at a  
936  $pH_i$  8 before (control) and after perfusion with RU1968F1, using the voltage protocol shown in (F). (I) Currents  
937 recorded from a mouse sperm cell at  $pH_i$  7.2 before (control) and after perfusion with  $NH_4Cl$  (30 mM) and  
938  $NH_4Cl$  plus RU1968F1, using the voltage protocol shown in (F). (J) Mean amplitudes of monovalent currents at  
939 +100 mV recorded from mouse sperm in the presence of RU1968F1,  $NH_4Cl$  plus RU1968F1, and at  $pH_i$  8 in the  
940 presence of RU1968F1, using the voltage protocol shown in (F). Amplitudes were normalized to the monovalent  
941 currents evoked in the absence of any drug (control, dashed line). Error bars indicate SD. ( $n = 5$ ). \* $P < 0.05$   
942 versus control. Data were normalized only after performing the statistical analysis using one-way ANOVA (see  
943 Methods for details and explanations).  
944

945 **Figure 7: RU1968F1 inhibits human but not mouse Slo3.** (A) Representative Slo3 currents in a mouse sperm  
946 cell, recorded in the presence of extracellular divalent ions at symmetric intra- and extracellular  $K^+$   
947 concentrations in the absence and presence of 50  $\mu M$  RU1968F1. Inset: Voltage protocol. (B) Mean Slo3  
948 currents in mouse sperm at +100 mV in the absence and presence of 50  $\mu M$  RU1968F1 ( $n = 5$ ). (C)  
949 Representative Slo3 currents recorded from a human sperm cell in the absence and presence of RU1968F1. Inset:  
950 Voltage protocol. (D) Dose-response relation for the inhibition of human Slo3 currents by RU1968F1 at  
951 +100 mV ( $IC_{50} = 7 \pm 6 \mu M$ ,  $n = 4$ ). Error bars indicate SD.  
952

953 **Figure 8: RU1968F1 interferes with hyperactivation and abolishes progesterone-induced motility  
954 responses in human sperm.** (A, B) Motility parameters of non-capacitated (A) and capacitated (B) human  
955 sperm in the absence and presence of RU1968F1 ( $n = 8$ ); sperm were bathed in the drug for 300 s. Error bars  
956 indicate SD. \* $P < 0.05$  versus control (absence of RU1968F1). (C-E) Fraction of hyperactivated (C), motile (D),  
957 and progressively swimming (E) sperm after capacitation in the absence and presence of RU1968F1 ( $n = 11$ ).  
958 Error bars indicate SD. \* $P < 0.05$  versus control (absence of RU1968F1). (F) Hyperactivation evoked by bathing  
959 sperm for 300 s in RU1968F1, progesterone, or progesterone plus RU1968F1 ( $n = 11$ ). Error bars indicate SD. (G)  
960 Track of a single sperm cell recorded before (3 s, black), during (0.2 s, red), and after (2.8 s, blue) uncaging of  
961 progesterone. The arrow indicates the direction of movement. Inset: time course of the average path velocity  
962 (VAP); the red bar indicates the uncaging of progesterone. (H) Track of a single sperm cell recorded before (3 s,  
963 black), during (0.2 s, flash, red) and after (2.8 s, blue) uncaging of progesterone in the presence of RU1968F1  
964 (30  $\mu M$ ). Inset: time course of VAP; the red bar indicates the uncaging of progesterone. (I) Mean relative  
965 changes in VAP averaged over 20-30 sperm in the field of view after uncaging of progesterone ( $n = 11$ ). Error  
966 bars indicate SD. \* $P < 0.05$  versus control (before flash, 0 s). Data were normalized only after performing the  
967 statistical analysis using one-way ANOVA (see Methods for details and explanations).  
968

969 **Figure 9: RU1968F1 suppresses penetration of sperm into viscous media.** (A) Number of sperm at a  
970 penetration distance of 2 cm in a modified Kremer's sperm-mucus penetration test. The sperm were incubated in  
971 buffer (control), progesterone, or progesterone plus RU1968F1 ( $n = 21$ ). Error bars indicate SD. \* $P < 0.05$  versus  
972 control; # $P < 0.05$  versus progesterone without RU1968F1. (B) Number of sperm after incubation in buffer  
973 (control) or RU1968F1 ( $n = 21$ ). Error bars indicate SD. \* $P < 0.05$  versus control ( $n = 21$ ). (C) Number of sperm  
974 when the sperm were bathed in buffer (control) or progesterone, in the absence (0) or presence of RU1968F1 in  
975 the capillary ( $n = 6$ ). Error bars indicate SD. \* $P < 0.05$  versus control; # $P < 0.05$  versus progesterone without  
976 RU1968F1. (D) Acrosome reaction evoked by RU1968F1, progesterone, or progesterone and RU1968F1 ( $n$   
977 =10). Error bars indicate SD. \* $P < 0.05$  versus control, # $P < 0.05$  versus progesterone.

978  
979  
980  
981  
982  
983  
984  
985  
986  
987

**Figure 10: RU1968F1 abolishes chemotaxis of sea urchin sperm.** (A) Dark-field microscopy images of a sperm suspension before (top) and after (bottom) establishing a resact gradient by photolysis of caged resact (middle) in the absence (control, left panel) or presence of RU1968F1 (30  $\mu$ M, right panel). RU1968F1 abolishes resact-induced sperm accumulation. (B) Relative change of the sperm dispersion in the field of view evoked by uncaging of resact ( $t = 0$ , flash) in the absence (control, red) or presence of RU1968F1 (black); a decrease of dispersion indicates sperm accumulation in the irradiated area (control,  $n = 5$ , RU1968F1,  $n = 6$ ). Error bars (grey) indicate SD.