

# Dorsal and ventral cortices are coupled by cross-frequency interactions during working memory

Popov, Tzvetan; Jensen, Ole; Schoffelen, Jan-Mathijs

DOI:

[10.1016/j.neuroimage.2018.05.054](https://doi.org/10.1016/j.neuroimage.2018.05.054)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

*Document Version*

Peer reviewed version

*Citation for published version (Harvard):*

Popov, T, Jensen, O & Schoffelen, J-M 2018, 'Dorsal and ventral cortices are coupled by cross-frequency interactions during working memory', *NeuroImage*, pp. 277-286.  
<https://doi.org/10.1016/j.neuroimage.2018.05.054>

[Link to publication on Research at Birmingham portal](#)

**Publisher Rights Statement:**

Checked for eligibility 30/05/2018

<https://doi.org/10.1016/j.neuroimage.2018.05.054>

**General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

**Take down policy**

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

1 **Dorsal and ventral cortices are coupled by cross-frequency interactions during**  
2 **working memory**

3  
4  
5  
6

7 Tzvetan Popov<sup>1</sup>, Ole Jensen<sup>2,3</sup>, Jan-Mathijs Schoffelen<sup>2</sup>

8 Author's affiliations:

9 1- Department of Psychology, Universität Konstanz, Germany

10 2- Radboud University Nijmegen, Donders Institute for Brain, Cognition and Behaviour,

11 Center for Cognitive Neuroimaging, The Netherlands

12 3- School of Psychology, University of Birmingham, Hills Building, Birmingham B15 2TT

13

14

15 Corresponding author: Jan-Mathijs Schoffelen

16 e-mail: jan.schoffelen@donders.ru.nl

17 Corresponding address: Kapitelweg 29, 6525 EN Nijmegen, P.O. Box 91101, 6500 HB

18 Nijmegen, The Netherlands

19 Tel. +31 (0)24 3614793

20

21 Keywords: N-back task, working memory, alpha oscillations, gamma oscillations,

22 magnetoencephalography

23

24

25

26

27

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16

**Abstract**

Oscillatory activity in the alpha and gamma bands is considered key in shaping functional brain architecture. Power increases in the high-frequency gamma band are typically reported in parallel to decreases in the low-frequency alpha band. However, their functional significance and in particular their interactions are not well understood. The present study shows that, in the context of an N-back working memory task, alpha power decreases in the dorsal visual stream were related to gamma power increase in early visual areas. Granger causality analysis revealed directed interregional interactions from dorsal to ventral stream areas, in accordance with task demands. Present results reveal a robust, behaviorally relevant and architectonically decisive power-to-power relationship between alpha and gamma activity. This relationship suggests that anatomically distant power fluctuations in oscillatory activity can link cerebral network dynamics on trial-by-trial basis during cognitive operations such as working memory.

1 **Introduction**

2  
3 Rhythmic neuronal activity is a ubiquitous phenomenon that underlies the spectral  
4 components that can be readily observed in electrophysiological recordings. [In recent](#)  
5 decades a large body of empirical evidence has been collected [that provides](#) a link between  
6 oscillatory activity in specific frequency bands and their functional role in cognition. For  
7 instance, amplitude fluctuations [in](#) alpha oscillatory activity (8-14Hz) have been found to be  
8 related to cognitive processes such as perception, attention, and memory (Gevins et al.,  
9 1997; Cooper et al., 2003; Klimesch et al., 2007; Jensen and Mazaheri, 2010; Saalman et al.,  
10 2012; van Kerkoerle et al., 2014; Bastos et al., 2015; Michalareas et al., 2016; Popov et al.,  
11 2017). Specifically, decreases in alpha amplitude have been proposed to index engagement  
12 of a cortical area, whereas increases mark reduced processing capabilities. This has been  
13 demonstrated in the visual (Adrian and Matthews, 1934; Foxe et al., 1998), sensorimotor  
14 (Haegens et al., 2010; Haegens et al., 2011; van Ede et al., 2014), and auditory (Weisz et al.,  
15 2011; Mazaheri et al., 2014) domains, suggesting an overarching principle for effective  
16 neuronal resource allocation in [a regionally](#) specific manner.

17  
18 [Faster](#) rhythms (gamma oscillations,  $>30\text{Hz}$ ), on the other hand, have been frequently linked  
19 to coherent stimulus processing (Tallon-Baudry and Bertrand, 1999; Fries, 2005), attention  
20 (Bauer et al., 2014; Marshall et al., 2015a; Marshall et al., 2015b), and working memory  
21 (Tallon-Baudry et al., 1998; Roux et al., 2012). It is well known that fast and slow rhythms  
22 frequently co-occur and that they are often co-modulated (or anti-modulated) as a  
23 consequence of an experimental manipulation. This has led to the idea that dynamic  
24 interactions between slow and fast oscillatory activity might be a key mechanism [shaping](#)  
25 functional interactions in cortical networks (Buzsaki and Draguhn, 2004; Canolty and Knight,  
26 2010; Lisman and Jensen, 2013).

27  
28 One possible mechanism [by](#) which fast rhythms could interact with slower rhythms is by  
29 means of phase-amplitude coupling (PAC) (Jensen and Colgin, 2007; Canolty and Knight,  
30 2010), where slow oscillations phasically modulate the amplitude of faster rhythms. PAC has  
31 been identified in a variety of species, including rodents (Tort et al., 2009; Tort et al., 2010),  
32 nonhuman primates (Whittingstall and Logothetis, 2009; Spaak et al., 2012), and humans  
33 (Canolty and Knight, 2010). Moreover, PAC has been demonstrated during cognitive

1 operations such as item-context binding (Tort et al., 2009), spatial navigation and decision  
2 making (Tort et al., 2008), working (Axmacher et al., 2010; Leszczynski et al., 2015) and  
3 episodic memory (Staudigl and Hanslmayr, 2013; Park et al., 2016), sleep (Staresina et al.,  
4 2015), and resting conditions (Florin and Baillet, 2015) and in a variety of clinical conditions  
5 such as Parkinson's disease (van Wijk et al., 2016), schizophrenia (Allen et al., 2011; Kirihara  
6 et al., 2012; Popov and Popova, 2015), autism (Berman et al., 2015), and affective disorders  
7 (Miskovic et al., 2011).

8

9 Another view of the relationship between slow and fast oscillatory activity comes from  
10 recent work investigating the network properties of directed oscillatory coupling [e.g.  
11 Granger causality (GC)] between brain regions in humans and non-human primates. Long-  
12 range inter-regional gamma synchronization has been shown to reflect feed-forward  
13 interactions within the visual cortical hierarchy, whereas slower alpha-beta synchronization  
14 reflects feedback interactions (van Kerkoerle et al., 2014; Bastos et al., 2015; Michalareas et  
15 al., 2016). These findings are in line with generic anatomical connection profiles between  
16 cortical layers, both within and between cortical areas, and with the cortical-layer-specific  
17 distribution of different neuronal rhythms.

18

19 A third type of interaction could involve a relationship between the oscillations' amplitudes.  
20 In contrast to the PAC and GC measures, this type of functional interaction does not require  
21 a strict relationship of the oscillations' phase between brain areas. In fact, a number of  
22 studies have reported stimulus-induced increases in gamma band activity with a  
23 concomitant decrease in alpha-beta power [e.g. (Schoffelen et al., 2005; Hoogenboom et al.,  
24 2006; Swettenham et al., 2009; Hoogenboom et al., 2010; Muthukumaraswamy and Singh,  
25 2013; Perry et al., 2013; Bauer et al., 2014; Kujala et al., 2015; Michalareas et al., 2016)].  
26 These empirical observations suggest a negative correlation between the task-induced  
27 amplitude modulations of slow versus fast rhythms.

28

29 Yet a functionally relevant mechanistic relationship between low- and high-frequency  
30 oscillatory activity, where a temporary increase in low-frequency power leads to a  
31 temporary reduction in high-frequency power, would suggest a negative correlation within  
32 trials. Reports of this nature are far less frequent in the literature (de Lange et al., 2008; Park

1 | et al., 2011; Popov et al., 2017; Wang et al., 2017), which may be a consequence of the fact  
2 | that single-trial estimates of oscillatory amplitudes typically have low signal-to-noise ratio, in  
3 | particular when estimated from non-invasively recorded data.

4 |  
5 | **The present study** leveraged the availability of a large number of subjects (n = 83), analyzing  
6 | MEG data from a publicly available dataset. Participants performed a visual working memory  
7 | (WM) N-back task, which reliably induces modulations in both alpha and gamma activity  
8 | (Roux and Uhlhaas, 2014). Thus, these data are well suited to investigate the relationship  
9 | between WM-load-dependent local changes in low- and high-frequency oscillatory activity,  
10 | both across time within trials and across subjects. We hypothesized that WM demands  
11 | would manifest as power-power interactions, where reductions in alpha activity would be  
12 | associated with a spatially specific power increase in gamma activity. Second, these power  
13 | fluctuations should be related to behavioral performance both within and between trials.  
14 | Third, “top-down” influence within key network nodes should be reflected in low-frequency  
15 | activity, whereas “bottom-up” communication should be evident in high-frequency activity.

## 17 **Material and methods**

### 18 *Participants and experimental procedures*

19 Publicly available data provided by the human connectome consortium  
20 ([www.humanconnectome.org](http://www.humanconnectome.org)) were analyzed. Eighty-three subjects (37 female and 46 male,  
21 mean age 28.5 years, range 22-35) participated in the experiment. Most were right-handed  
22 as measured with the Edinburgh Handedness Inventory with an mean lateralization quotient  
23 of 65% and SD = 44% (Oldfield, 1971). **Participants gave written informed consent for  
24 participation in accordance with the Declaration of Helsinki. In two runs, participants  
25 performed an N-back WM paradigm, memorizing pictures of faces and tools. Subjects were  
26 presented with 16 blocks per run of 0-back and 2-back trials and were asked to indicate  
27 match and no-match responses via button press with their right index and right middle  
28 finger, respectively (Figure 1). An initial cue presented for 2500 ms signaled the beginning of  
29 either a 0- or a 2-back block. After this, a serial presentation of face or tool stimuli displayed  
30 for 2000 ms was intermixed with an inter-stimulus intervals (ISI) of 500 ms. Participants  
31 were asked to respond as fast as possible via button press after the presentation of each  
32 stimulus. Responses had to take place within the stimulus presentation window. Following a**

1 | fixation interval of 500 ms, the next trial was presented.

2

3 | *Data acquisition and analysis*

4 | Data were recorded using a whole-head 248-channel magnetometer system (MAGNES 3600  
5 | WH, 4D Neuroimaging, San Diego, CA) with subjects supine. Data were continuously  
6 | recorded with a sampling rate of 2034.5101 Hz and a bandwidth of DC-400Hz. Digitization of  
7 | each head shape and of the locations of the fiducial coils was accomplished with a Polhemus  
8 | 3Space Fasttrak system. The WM task was a part of a 3-hour session, where both task and  
9 | resting-state MEG data were collected. Participants performed a sequence of tasks,  
10 | described in detail in the reference manual provided by the human connectome consortium  
11 | (<http://www.humanconnectome.org/documentation/S500/index.html>). Just prior to the N-  
12 | back paradigm the participant underwent three runs of approximately 6 minutes of resting-  
13 | state MEG recording.

14

15 | The analysis described below was performed on the ‘minimally preprocessed’ data that were  
16 | downloaded from the human connectome database. In brief, epochs lasting from 1.5 s  
17 | before to 2.5 s after each picture’s onset had been extracted from the continuous recording.  
18 | Epochs containing superconducting quantum interference device (SQUID) jumps, bad  
19 | sensors, or bad segments, defined as excessive signal amplitude changes  $> \sim 10^{-12} T_z$  were  
20 | excluded from further processing. Eye-movement-related signals and cardiac signals had  
21 | been identified with independent component analysis (ICA) (Jung et al., 2001) and projected  
22 | out of the data. All data was analysed using custom scripts in MATLAB and the FieldTrip  
23 | toolbox (Oostenveld et al., 2011).

24

25 | *Spectral analysis*

26 | For each epoch, spectral analysis was performed using a windowed Fast Fourier Transform  
27 | (FFT). For frequencies below 40 Hz, we used a 500 ms sliding window (sliding in steps of 50  
28 | ms) multiplied by a Hanning taper, achieving an effective frequency resolution of  $\sim 2$  Hz. For  
29 | high frequency activity ( $>40$  Hz), we used multitapers (Mitra and Pesaran, 1999). Eleven  
30 | orthogonal Slepian tapers were used, resulting in a spectral smoothing of  $\pm 10$ Hz.

31

32 | *Source level analysis*

1 Source reconstruction of oscillatory activity was performed using the dynamic imaging of  
2 coherent sources (DICS) algorithm (Gross et al., 2001). This algorithm uses the sensor-level  
3 cross-spectral density matrix and a set of location-specific forward models to construct a set  
4 of spatial filters optimized for a given frequency at the specific locations. The data of both  
5 conditions were used for the spatial filter computation, which were subsequently used to  
6 compute the pattern of oscillatory activity for each condition separately. A realistic, single-  
7 shell brain model (Nolte, 2003) was constructed based on the individual anatomical MRI as  
8 implemented in FieldTrip. Forward models were computed using the anatomical information  
9 provided in the database, consisting of a subject-specific, realistically shaped single shell  
10 volume conduction model and a 3-dimensional grid of dipole locations with equidistant  
11 positions in normalized MNI-space (spacing of 8 mm).

12

13 For a set of predefined locations, we reconstructed time courses of neuronal activity using a  
14 linearly constrained minimum variance (LCMV) beamformer (Van Veen et al., 1997). These  
15 locations were identified by contrasting source-reconstructed power estimates of alpha and  
16 gamma activity for the 2- vs. 0-back conditions resulting in differential effects in primary  
17 visual areas (V1), intra-parietal sulcus (IPS), and fusiform gyrus (FF) (see Figure 1). [Local](#)  
18 [maxima were identified in right FF \[MNI coordinates: 55 -58 -12\], right intraparietal sulcus](#)  
19 [IPS \[MNI coordinates: 34 -76 28\], and primary visual cortex V1 \[MNI coordinates: 4 -80 8\].](#)  
20 [Spatial filters were estimated based](#) on the unfiltered data covariance matrix [for all trials](#) and  
21 forward models of the locations of interest. Subsequently, these filters were multiplied [by](#)  
22 the data in order to obtain 'virtual channel' time series at the respective locations.

23

#### 24 *Correlation analyses*

##### 25 *Brain-behavior relationships*

26 The relationship between behavioral measures (average reaction times, [RT](#), or [RT](#)  
27 differences) and trial-averaged neural data were computed [as](#) Spearman's rank correlation  
28 across participants. This relationship was computed for each sensor, time, and frequency  
29 bin, and statistical inference was done using cluster-based permutation tests (see below).

30 [Within-subject Spearman](#) correlations across trials, using single trial RTs and neural data at  
31 sensor level, [were also computed.](#)

32



## 1 *Cross-frequency power-power correlations*

2 In order to assess cross-frequency power-power correlations, [Spearman correlations](#)  
3 [characterized cross-frequency](#) alpha-gamma power-[power relationships](#). At the sensor level,  
4 stimulus-induced gamma activity (60-80Hz) in visual areas was correlated [with](#) the estimated  
5 power at each sensor, time [point](#), and frequency bin (up to 40 Hz). Gamma power was  
6 quantified as a single value per participant, computed as the mean over occipito-parietal  
7 sensors, frequencies, and time points. The inclusion criteria for sensors, [time points, and](#)  
8 frequencies were based on the condition differences (2-back vs. 0-back) in gamma activity  
9 (see Figure [1A](#)). In addition, trial-by-trial relationships were evaluated at source level. First,  
10 generators of alpha and gamma oscillatory signals were determined by a spatial filtering  
11 algorithm (see below) and time courses of neuronal activity at these locations were  
12 extracted. Next, within participants, two sets of high and low alpha trials per location were  
13 identified on the basis of a median split [on alpha power](#). Finally, the null hypothesis of no  
14 group differences in gamma amplitudes in high and low alpha trials was evaluated using the  
15 cluster-based permutation framework.

16

## 17 *Granger causality analysis [and parcellation of source-reconstructed activity](#)*

18

19 Spectrally resolved [Granger causality \(GC\)](#) analysis (Granger, 1969; Ding et al., 2006)  
20 was used to dissociate potential directionality in inter-nodal communication. Briefly, GC  
21 represents the result of a model comparison where the predictive strength of past values of  
22 time series  $x$  [predicting the](#) future behavior of time series  $x$  (a univariate auto-regressive  
23 model) is evaluated against the past values of time series  $x$  and another time series  $y$  (a  
24 bivariate auto-regressive model) [predicting the](#) future behavior of time series  $x$ . Initially  
25 formulated in the time domain, GC can also be estimated in the frequency domain (Geweke,  
26 1982; Kaminski et al., 2001; Chen et al., 2006) [see for detailed review (Bastos and  
27 Schoffelen, 2015) and (Ding et al., 2006)], which requires the estimation of the spectral  
28 transfer matrix between a set of signals.

29 A whole-cortex all-to-all GC analysis was performed. In a first step, the dimensionality  
30 of the data was reduced by applying an atlas-based parcellation scheme (Glasser et al.,  
31 2016). This atlas discretizes the neocortex into 180 parcels per hemisphere, which can [be](#)  
32 further grouped into 22 modality-specific areas, e.g. auditory, somatosensory, visual, task-

1 positive, and task-negative. Following procedures [similar to those of](#) (Schoffelen et al.,  
2 2017), single-dipole-specific spatial filters were concatenated across vertices comprising a  
3 parcel, yielding a set of 22 multivariate time courses of activity per hemisphere. For each  
4 parcel, the spatial components explaining 95% of the variance within the corresponding  
5 parcel were selected. [Out of these preselected components, only the first \(largest\)](#)  
6 [component was considered for further analysis. This step is motivated by the fact that](#)  
7 [differences in local dipole orientations preclude averaging over a set of components within a](#)  
8 [parcel, potentially leading to cancelation of effects.](#) For comparability purposes, source  
9 power analysis of alpha and gamma activity was re-computed as described above followed  
10 by whole-brain (all to all parcels) conditional GC analysis. [Parcel time series were re-](#)  
11 [segmented to include 1 to 2 seconds post-stimulus-onset \(the post-stimulus task window\).](#)  
12 [For each trial, Fourier coefficients were computed for the entire spectrum up to the Nyquist](#)  
13 [frequency. Non-parametric matrix factorization of the cross-spectral density matrix \(Wilson,](#)  
14 [1972; Dhamala et al., 2008\) estimated the spectral transfer matrix. For a given pair of](#)  
15 [locations, GC was computed conditional on the rest of locations,](#) similar to the procedures  
16 applied in (Wen et al., 2013) and (Bastos et al., 2015). [In addition, GC influences were](#)  
17 [computed twice: once on the original and once on the time-reversed time series.](#) The latter  
18 strategy proposed by (Haufe et al., 2013; Winkler et al., 2015) accounts for the presence of  
19 so-called weak asymmetries as a possible interpretational confound, contributing to an  
20 apparent dominant directional drive between two areas. Briefly, as opposed to strong  
21 asymmetries, which are caused by actual time-lagged relations between signals, weak  
22 asymmetries are the consequence of differences in univariate signal properties (for instance,  
23 [the local signal-to-noise ratio\).](#) [Time reversal of the signals does not affect these univariate](#)  
24 [signal properties, whereas it should reverse the dominant direction of interaction. In other](#)  
25 [words, if similar patterns of GC are observed after the time reversal, these are most likely](#)  
26 [artificial, whereas a reversal of the direction of the asymmetry implies true time-lagged, thus](#)  
27 [directed relationships. Only relationships surviving the contrast \[forward minus backward vs.](#)  
28 [forward<sub>flip</sub> minus backward<sub>flip</sub>\] are presented and evaluated in this report.](#)

## 29 *Statistical analysis*

30 [Inferential statistical](#) evaluation was carried out by non-parametric permutation tests  
31 (Maris and Oostenveld, 2007). [A cluster-based approach, clustering across space, time](#)  
32 [points,](#) and frequency (wherever appropriate) [controlled](#) for multiple comparisons, [using](#)

1 | [1000 permutations and an alpha threshold of 0.025.](#)  
2

|

10

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24

## Results

The present study evaluated publicly available neuromagnetic data from human participants performing an alternating WM task. In a block design, participants performed either 0 or 2-back task. Each block began with a cue indicating the WM load condition for 2.5 s followed by presentation of face or tool stimuli. Stimulus duration was 2 s separated by a 0.5 s interstimulus interval (offset to onset) during which participants were asked to provide a response (Figure 1).




Figure 1: Experimental task. A- At the beginning of the 0-back block, participants are presented with a target face X. On subsequent trials different faces are presented. After each face, participants indicate via button press whether the presented stimulus matches the target stimulus. Responses had to occur within 500 ms after the stimulus offset. B- 2-back blocks were signaled by a presentation of "2-back" for 2500 ms. Participants indicated whether the presented stimulus matched the stimulus two trials earlier. Responses had to occur in 500 ms after stimulus offset.

## Alpha\_beta and gamma activity in object-selective areas scale with working memory load

As a first step, we evaluated the effect of WM load on the time-frequency response of the brain. Compared to the 0-back condition, the 2-back condition was associated with an amplitude decrease in alpha\_beta power (8-15Hz) and an increase in gamma power (60-80 Hz) ( $p < 0.01$ , non-parametric randomization)(Figure 2A). Sensor-level effects were widespread, with the largest difference at occipital sensors (Figure 2B). Source reconstruction of these effects, however, suggests different cortical generators of the observed topographies (Figure 2C). Alpha\_beta condition differences

Unknown  
Formatiert: Schriftart:(Standard) Calibri,  
Schriftfarbe: Automatisch

1 were strongest in bilateral fusiform gyri and right intra parietal sulcus (IPS). In contrast,  
 2 gamma effects were strongest in early visual areas around the primary visual cortex  
 3 (V1). The chosen threshold of 80 % of maximum is arbitrary, and source extent  
 4 be interpreted with caution. The largest observed power difference occur on average  
 5 at peak minimum of 800 ms for alpha and peak maximum of 650 ms for gamma  
 6 activity. Button responses following stimulus presentation occurred on average at 631  
 7 ms for 0-back (min/max = 394/1053 ms, SD = 128 ms) and 827 ms for 2-back (min/max  
 8 = 511/1161 ms, SD = 131 ms).  
 9



10 **Figure 2: High- and low-frequency activity scales with WM load.** A- Time-frequency representations of memory-load-  
 11 dependent differences in power for low (left) and high (right) frequencies. The color bar indicates difference in oscillatory  
 12 power (2-back vs. 0-back), with dashed rectangles approximating the time-frequency range contributing to the largest  
 13 extent of the cluster providing the basis for the rejection of the null-hypothesis, at a p-value of 0.01. Dots highlight the spatial  
 14 extent of the cluster providing the basis for the rejection of the null-hypothesis, at a p-value of 0.01. C- Source-level  
 15 representation of condition differences highlighting the strongest (80% of minimum) effects with respect to alpha activity  
 16 (left) in bilateral fusiform (FF) gyri as well as right intra-parietal sulcus (IPS). In contrast, gamma modulation by WM load  
 17 was strongest (80% of maximum) in early visual (V1) areas (right graphs). MNI coordinates: right FF [55 -58 -12], right IPS  
 18 [34 -76 28], V1 [4 -80 8]. Note: alpha activity shows two different loci, whereas the maxima of gamma band activity is  
 19 predominantly localized in the primary visual cortex.  
 20

21  
 22

23 **Alpha-beta and gamma oscillations are inversely related to behavioral performance**  
 24 **between and within participants**

25  
 26

On average, an increase in WM load resulted in increased RT to the target stimulus.


1 The difference in RT ( $\Delta RT = RT_{2\text{-back}} - RT_{0\text{-back}}$ ) was on average 131 ms (range 175 to 432 ms;  $p <$   
2  $0.0001$ ,  $t_{82} = 10.05$ , paired  $t$ -test). Figure 3 shows the analysis of the correlation, across  
3 subjects, between the behavioral effect and the effect on the time-frequency response. The  
4 left panel of Figure 3A illustrates the association between posterior alpha-beta power  
5 modulations and RT differences. This indicates that, across participants, stronger alpha  
6 power modulations (larger memory-load-dependent decreases) were associated with a  
7 larger behavioral penalty ( $\Delta RT$ ), reflecting the increase in WM demands. Participants  
8 characterized by stronger alpha-beta decrease with load were also those with a larger RT  
9 increase with load. The opposite was observed for gamma activity (Figure 3A right panels). A  
10 strong WM-load induced increase in gamma band power was related to an increase in RT  
11 differences. Next, we investigated the correlations across trials. The left panels in Figure 3B  
12 illustrate the trial-to-trial relationship between alpha-beta oscillations and RT for 0- and 2-  
13 back conditions separately. Each line indicates the correlation across trials as a function of  
14 time per participant, sorted along the mean reaction time of the respective participant. Prior  
15 to the button press there was a positive relationship between alpha-beta power modulation  
16 and RT and negative one after response indication. Thus, WM-induced alpha/beta  
17 modulations were predictive for individual performance on a trial-by-trial basis. Moreover,  
18 this brain-behavior relationship was modulated by WM load (Figure 2B, bottom left panel,  $p$   
19  $< 0.01$  cluster-permutation approach). Similar association differentiating between memory  
20 loads was also evident for gamma band activity and RT (Figure 3B, right panels,  $p < 0.01$ ).

21

22 In summary, WM load dependent modulations of alpha-beta and gamma oscillations were  
23 found related to behavioral performance both between and within participants. Condition  
24 specific differences in alpha-beta and gamma oscillations were characterized by different  
25 activation patterns: alpha-beta activity was most pronounced in IPS, whereas maxima of  
26 condition differences in gamma band activity were more confined to early visual areas. The  
27 correlation between the WM-load dependent gamma response and alpha-beta modulation  
28 observed at the sensor-level, combined with the neural sources of the observed effects  
29 suggests a cross-frequency interaction between “early” (visual) and “late” (ventral and  
30 dorsal) hierarchical levels, which we explored next.

31

32



**Figure 3: Relationship between neural data and overt performance, and between alpha and gamma power. A) Between-subject relationships.** Time-frequency representation of the correlation between RT differences and differences in low- (left) and high- (middle) frequency activity (alpha and RT  $p < 0.001$ ; gamma and RT  $p < 0.03$ ; gamma and alpha  $p < 0.001$ ; cluster-permutation approach). The scatterplots in A are an additional illustration of the respective effects, with each dot corresponding to an individual participant. Time-frequency spectrograms are averaged over occipito-posterior sensors with the following labels: A75, A105, A106, A107, A138, A139, A140, A166, A167, A189, A136, A137, A138, A139, A164, A165. **B- Within-subject relationships.** Time by participant graphs illustrating the variation of the association between RT and alpha-beta power (left) and gamma power (right). Each line represents a single subject. The color bar indicates correlation coefficients. Participants are sorted according to mean RT time (indicated by the thick black lines). Bottom graphs provide mean correlation over participants with shading indicating  $\pm 1$  SEM.

### Alpha\_beta oscillations in IPS regulate gamma amplitudes in early visual cortex

The results so far suggest an inverse relationship between memory-load-dependent changes in alpha-beta activity on the one hand and the changes in gamma activity on the other. We reconstructed the broadband neural activity at the source locations identified and illustrated in Figure 2. This yielded three virtual sensors per subject, in right IPS, fusiform (FF), and visual (V1) areas. For each participant and brain location, we computed single-trial alpha power averaged between 0.5-2 s and 8-15 Hz. For each of the sets of single-trial alpha power estimates, we performed a median split, dividing the trials into region-specific high


1 | and low alpha power trials for each participant. Subsequently, we computed time-frequency  
2 | representations of power in the gamma range (40-120 Hz) for subsets of trials with low and  
3 | high alpha power.

4 |         The outcome of this analysis is illustrated in Figure 4A. Alpha power decreases in IPS  
5 | were associated with a sustained relative increase of gamma band activity in V1. That is, on  
6 | trials with less alpha activity in IPS, gamma amplitude in V1 was higher. This finding was  
7 | observed only for the IPS-V1 relationships and not for the other possible pairs illustrated in  
8 | Figure 4A. A control analysis confirmed a robust cross-participant relationship between  
9 | alpha\_beta and gamma activity (60-80Hz) (Figure 4B,  $p < 0.02$  cluster-permutation  
10 | approach). Whole\_brain analysis confirmed that this effect was most pronounced in early  
11 | visual areas in addition to activation of left motor areas, potentially reflecting the  
12 | preparation for right-hand button press. Specifically, gamma band activity was reconstructed  
13 | for high and low IPS alpha trials and subsequently contrasted, resulting in a difference in  
14 | gamma activity per voxel and per load. Figure 3C illustrates this contrast for 0\_ and 2-back,  
15 | respectively as well as their interaction (IPS alpha<sub>[high, low]</sub> × WM Load<sub>[0-back, 2-back]</sub>). Finally,  
16 | Figure 4D illustrates this trial-to-trial relationship as a function of time for 0\_ and 2-back  
17 | conditions. Each line indicates the correlation across trials as a function of time per  
18 | participant, sorted along the mean RT of the respective participant. In line with the source\_  
19 | space results, there was significant condition difference between the alpha/gamma  
20 | correlations in 0\_ vs. 2-back (Figure 4D,  $p < 0.02$  cluster-permutation approach).

21 |         This cross-frequency power relationship is correlative in nature and does not permit  
22 | conclusions about the potential directionality of effects. These findings suggest a cross-  
23 | frequency anti-correlation between alpha oscillations in IPS and gamma oscillations in early  
24 | visual cortex. The extent to which this effect is mediated by top-down (e.g., IPS to V1) or  
25 | bottom\_up (e.g., V1 to IPS) interactions can not be addressed with the analyses reported  
26 | above, so Granger causality (GC) between the regions of interest derived from that analyses  
27 | was undertaken.

28






**Figure 4: Inter-regional power-power correlation of alpha and gamma activity.** **A-** Time-frequency representation of power differences in high-frequency activity between high and low alpha trials as a function of median split location collapsed across conditions. Color bar illustrates the strength of the observed differences, and dashed rectangle approximates the time-frequency cluster of significant condition differences ( $p < 0.05$ , cluster-based permutation approach). **B-** Time-frequency representation of correlation strength between alpha-beta (??-?? Hz) and gamma activity (60-80 Hz) predominantly over occipital sensors ( $p < 0.01$ , cluster-based permutation approach). **C-** Whole-brain analysis of gamma activity as a function of high vs. low IPS alpha. Effects were located primarily in early visual areas with spectro-temporal characteristics similar to those illustrated by the spectrogram on the left. WM load and gamma in high vs. low alpha trials interacted ( $p < 0.01$ , uncorrected). The data used in the time-frequency spectrogram are derived from V1 (MNI 4 -80 8). **D-** Time by participant graphs illustrating the variation of the association between alpha-beta and gamma power based on sensor level data. In the upper panels, time is depicted on the abscissa and participant number on the ordinate. Each row represents a single subject. The color bar indicates correlation coefficients. Participants are sorted according to mean RT (indicated by the thick black lines). Bottom panel depicts mean correlation over participants in 0-back (red) and 2-back (blue) with shading indicating SEM.

## Dorsal stream exerts top-down control over ventral stream

Conditional GC differentiates direct and mediated influences (Chen et al., 2006; Wen et al., 2013) using a late post-stimulus time window (1 to 2 s) in order to avoid interference from stimulus- and response-locked components. Directed connectivity during WM was probed by a parcellation-based GC analysis (see Granger causality analysis method section above). Figure 5A confirms the primary condition difference in alpha-beta power in posterior parietal cortex (Figure 5A, left), whereas differences in gamma activity with load were confined mainly to early visual areas (Figure 5A, right). Directed interactions between

1 cortical structures that survived the time-reversal test are shown in Figure 5B. Dorsal-stream  
 2 structures such as dorsal visual cortex exhibited directed interactions with ventral-stream  
 3 structures, such as ventral visual cortex (vVis), medial temporal gyrus (MTG), and area MT as  
 4 well as visual areas V1, V2/V3 (Figure 5B). This connectivity architecture was evident during  
 5 both WM conditions. Moreover, this directed communication was predominantly  
 6 established in the alpha-beta bands (Figure 5B). No GC gamma effects (40+ Hz) were  
 7 observed. Overall, GC results complement the power-power findings reported above,  
 8 strengthening the evidence that the dorsal stream exerts top-down control of visual and  
 9 ventral stream regions via alpha-beta oscillations.



10  
11  
12  
13  
14  
15  
16  
17  
18  
19

**Figure 5: Whole-cortex GC analysis.** A- Source-level representation of cortical parcels illustrating the strongest condition differences with respect to alpha (left) and gamma (right). B- **Left panel:** Directed connectivity architecture collapsed across WM load. Only connections surviving the time-reversal procedure (see methods) are shown. **Right panels:** GC spectra (original data) collapsed across WM load averaged over participants per area pair. Red indicates "bottom-up" and black "top-down" directions. Shading depicts SEM. 0-back and 2-back conditions analyzed separately (not shown) showed similar GC spectral profiles.

Unknown  
 Formatiert: Schriftart:(Standard) Calibri,  
 Schriftfarbe: Automatisch

## 1 Discussion

2 |       The present [study](#) examined the relationship between low- and high-frequency  
3 | oscillatory activity during a working memory task. Participants performed an N-back working  
4 | memory task under low and high load conditions (0-back and 2-back), which allowed us to  
5 | compare WM-load-dependent stimulus-related changes in oscillatory activity. A large  
6 | sample of 83 human participants [showed](#) modulations of alpha-beta and gamma oscillatory  
7 | activity [that 1\)](#) were related to each other across participants, 2) [were related to each other](#)  
8 | across trials [within participants](#), and 3) functionally [connected](#) distant brain areas by means  
9 | of power-power interactions. Moreover, [Granger causality analyses showed](#) directed  
10 | interactions from dorsal-stream parietal areas to ventral-stream areas.

### 11 [Within- and between-subject relationships between power modulations and reaction time](#) 12 [index effortful processing](#)

13 |       A key finding of the present report is a robust relationship between the neural and  
14 | [overt performance](#) data, both across subjects and across [time within](#) trials (Figure 2B).  
15 | [Cognitive](#) effort was quantified as the increase in [RT](#) due to the number of items to be held  
16 | in memory. [This](#) increase in cognitive demand was associated with [more](#) decrease [in](#) alpha-  
17 | beta power in IPS and a concurrent increase in gamma power in V1 (Figure 2A). This [pattern](#)  
18 | fits well with the interpretation that decreases in alpha-beta power and increases in gamma  
19 | power indicate the involvement and overall activation level of task-relevant cortical areas.

20 |       Yet, across participants, larger differences in WM-load-dependent power were  
21 | related to a larger behavioral penalty, i.e. slower RTs with increasing WM load. [There was a](#)  
22 | [stronger positive alpha to RT correlation prior to the actual response in the 2-back as](#)  
23 | [compared to the 0-back conditions \(Figure 3B\). It is conceivable that the parameters of the](#)  
24 | [response-related rebound event-related synchronization \(ERS\) at alpha-beta frequencies in](#)  
25 | [response to stimulus N-1 correlate with the RT to stimulus N. The magnitude of the ERS](#)  
26 | [indexes the overall readiness of the motor system for a fast response, where higher](#)  
27 | [amplitude is associated with slow responses. Consequently, the observed positive](#)  
28 | [correlation in the 2-back condition at early latencies is increased. Due to the overall slower](#)  
29 | [responses in the 2-back condition, this effect may persist until after onset of the next](#)  
30 | [stimulus.](#) This finding challenges the often tacitly assumed expectation of a positive  
31 | relationship between oscillatory power modulations and cognitive performance. A number

1 of studies have demonstrated that the amount of [alpha-beta](#) power reduction relates to  
2 faster [RT](#) (Gonzalez Andino et al., 2005; Schoffelen et al., 2005; van Ede et al., 2011; van Ede  
3 et al., 2012; Buchholz et al., 2014). A common feature of these studies is the explicit  
4 manipulation of attention prior to the behavioral response. Anticipatory attention and/or  
5 informative sensory cues might [facilitate](#) an anticipatory state, [including](#) a decrease in alpha-  
6 beta activity. This anticipatory state will result in faster [RT](#) to an upcoming target stimulus.  
7 One important difference between the [present](#) experimental paradigm and the referenced  
8 literature is the distinction between stimulus-induced changes in power (induced by task-  
9 relevant stimuli that may require a response) and power modulations [in](#) oscillatory activity in  
10 anticipation [of](#) a behaviorally relevant stimulus. Thus, in the [results](#) reported here, the  
11 increase in gamma/decrease in alpha-beta may reflect increased allocation of local  
12 computational resources to process the incoming stimulus and to plan the appropriate  
13 response, while having to keep and continuously update [two](#) stimuli in memory. Rather  
14 than [being](#) directly linked to the speed of behavioral responses, oscillatory power  
15 modulations [may](#) reflect the increased level of simultaneous item maintenance [efforts](#). The  
16 reported decrease [in](#) alpha activity [may](#) reflect the engagement of the visual system in active  
17 information processing, with the predominant interplay between ventral and dorsal brain  
18 areas potentially reflecting the visual imagery involved in the context of the present task.

19 [Alternatively, instead of neural processes supporting WM that correlate with RT,](#)  
20 [present findings may reflect motor response processes. In fact, processes related to motor](#)  
21 [response and WM update processes can co-occur and are not readily dissociable in the](#)  
22 [present task. However, the strongest condition-specific effects were observed at latencies](#)  
23 [within the range of the RTs \(0.4-1.2 s\), thus surrounding the actual motor response.](#)  
24 [Furthermore, the clear occipito-parietal topography of the correlations and the well-known](#)  
25 [involvement of these regions during WM tasks suggest that the findings reflect engagement](#)  
26 [of cortical areas that subserve the more core WM-related processes.](#)

### 27 ***Parietal alpha-beta power fluctuations are related to gamma activity in early visual areas***

28 Extending previous reports showing concurrent stimulus-induced power modulations  
29 of alpha and gamma band activity, [present](#) findings support a mechanistic view [of](#)  
30 [relationships](#) between neuronal oscillations in different frequency bands and between  
31 neuronal oscillations and cognitive performance. Specifically, participants with a stronger

1 WM-load-dependent decrease in alpha-beta power were also characterized by a stronger  
2 increase in gamma power. Whereas neuronal generators of gamma activity were confined  
3 mainly to early visual areas, the strongest alpha-beta power reduction was predominantly in  
4 higher-order parietal areas (Figure 2C). These power-power relationships between low- and  
5 high-frequency activity were anatomically specific and are in line with previous reports,  
6 highlighting one possible role of parietal brain regions, exerting a top-down control over  
7 task-relevant visual cortex (Liu et al., 2014; Michalareas et al., 2016). Furthermore, previous  
8 MEG work also considering interactions between gamma and alpha band activity during  
9 working memory operations identified the superior temporal gyrus (STG) as a hub region  
10 coordinating network activity during maintenance (Park et al., 2011). We observed little  
11 engagement of the STG. However, there were several differences in the tasks used. The  
12 executive component of the present task involves IPS-V1 interaction, where WM  
13 maintenance is typically reflected in frontal to parietal communication (Christophel et al.,  
14 2017). Importantly, on a trial-by-trial basis, more alpha power decrease in parietal cortex  
15 was associated with a more increase in gamma power in early visual areas (Figure 4),  
16 suggesting a direct functional relationship between spatially distant brain areas. Observing a  
17 negative correlation in both conditions suggests an involvement of and coordinated  
18 interaction between parietal and early visual areas, in line with task requirements. This  
19 correlation was stronger for the 0-back than the 2-back condition (Figure 4C). However, the  
20 interpretability of this difference is not straightforward. This reflects estimates of both mean  
21 power and variance, and on the other hand there was an overall increase in gamma power  
22 during the 2-back conditions. Thus, a ceiling effect on gamma power potentially resulted in a  
23 smaller difference between low and high alpha power trials. The extent to which this reflects  
24 a behaviourally relevant functional architecture or rather more stimulus-related perceptual  
25 processes remains to be demonstrated. On the basis of this functional power to power  
26 relationship between IPS alpha and V1 gamma, we next examined the effective  
27 communication between these nodes.

28

### 29 ***Dorsal-stream-to-ventral-stream interactions at alpha-beta frequencies***

30 GC analysis indicated a unidirectional relationship from dorsal visual to ventral visual  
31 areas relying on alpha-beta frequencies (Figure 5B), with no clear GC spectral peak at  
32 gamma frequencies. Early visual areas were receiving input from dorsal areas primarily at

1 | alpha-beta frequencies (Figure 4A). These results are partially in line with reports from  
2 | human and animal research, demonstrating that feedback interactions operate in the  
3 | intermediate alpha-beta range (Saalman et al., 2012; van Kerkoerle et al., 2014; Bastos et  
4 | al., 2015; Michalareas et al., 2016; Popov et al., 2017). [Previous literature suggests that, for](#)  
5 | [perceptual tasks](#) strongly engaging the dorsal stream, dorsal regions affect mid-level ventral-  
6 | stream regions [more](#) than early visual areas (Michalareas [et al., 2016](#)). Present findings are  
7 | consistent with this, demonstrating that, during high WM load, [dorsal alpha activity is a](#)  
8 | [mechanism of](#) top-down [control](#) over ventral regions, despite an experimental arrangement  
9 | strongly engaging the ventral stream (e.g., face identity).


## 10 | **Conclusions**

11 | In summary, the present study analyzed publicly available MEG data from 83  
12 | [participants](#) while they performed an N-back task. We investigated the relationship between  
13 | neural activity (low- and high frequency oscillations) and [overt performance](#) (task condition  
14 | and [RT](#)). Increased working memory demands resulted in alpha-beta and gamma power  
15 | modulations in early visual cortex and brain regions in the dorsal and ventral visual [streams](#).  
16 | [Robust](#) cross-frequency power-to-power interactions between these brain areas and  
17 | between alpha-beta and gamma activity during WM scaled with WM demands. Granger  
18 | [causality](#) interactions were most prominent in the alpha band, from dorsal stream to ventral  
19 | stream areas. Overall, these data confirm earlier findings [and](#) provide additional support for  
20 | the notion that fluctuations in band-limited neural activity reflect behaviorally relevant local  
21 | and [inter-regional](#) neural processing. In addition, this work shows the utility of employing  
22 | publicly available, task-based MEG data for exploratory and confirmatory purposes.

## 23 | **Acknowledgments**

24 | [The authors thank Gregory A. Miller for providing valuable discussion and comments on](#)  
25 | [earlier versions of the manuscript](#). T.P. was supported by the German national funding  
26 | agency Deutsche Forschungsgemeinschaft (Ro805/17). O.J. was supported by the James S.  
27 | McDonnell Foundation Understanding Human Cognition Collaborative Award Grant No.  
28 | 220020448. J.-M.S. was supported by the Dutch Organization for Scientific Research (NWO  
29 | VIDI: 864.14.011).

|



|

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50

## References

- Adrian ED, Matthews BHC (1934) The Berger rhythm: potential changes from the occipital lobes in man. *Brain* 57:355-384.
- Allen EA, Liu J, Kiehl KA, Gelernter J, Pearlson GD, Perrone-Bizzozero NI, Calhoun VD (2011) Components of cross-frequency modulation in health and disease. *Frontiers in systems neuroscience* 5:59.
- Axmacher N, Henseler MM, Jensen O, Weinreich I, Elger CE, Fell J (2010) Cross-frequency coupling supports multi-item working memory in the human hippocampus. *Proceedings of the National Academy of Sciences of the United States of America* 107:3228-3233.
- Bastos AM, Vezoli J, Bosman CA, Schoffelen JM, Oostenveld R, Dowdall JR, De Weerd P, Kennedy H, Fries P (2015) Visual Areas Exert Feedforward and Feedback Influences through Distinct Frequency Channels. *Neuron* 85:390-401.
- Bauer M, Stenner MP, Friston KJ, Dolan RJ (2014) Attentional modulation of alpha/beta and gamma oscillations reflect functionally distinct processes. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 34:16117-16125.
- Berman JJ, Liu S, Bloy L, Blaskey L, Roberts TP, Edgar JC (2015) Alpha-to-gamma phase-amplitude coupling methods and application to autism spectrum disorder. *Brain Connect* 5:80-90.
- Buchholz VN, Jensen O, Medendorp WP (2014) Different roles of alpha and beta band oscillations in anticipatory sensorimotor gating. *Frontiers in human neuroscience* 8:446.
- Buzsaki G, Draguhn A (2004) Neuronal oscillations in cortical networks. *Science* 304:1926-1929.
- Canolty RT, Knight RT (2010) The functional role of cross-frequency coupling. *Trends in cognitive sciences* 14:506-515.
- Chen Y, Bressler SL, Ding M (2006) Frequency decomposition of conditional Granger causality and application to multivariate neural field potential data. *Journal of neuroscience methods* 150:228-237.
- Christophel TB, Klink PC, Spitzer B, Roelfsema PR, Haynes JD (2017) The Distributed Nature of Working Memory. *Trends in cognitive sciences* 21:111-124.
- Cooper NR, Croft RJ, Dominey SJ, Burgess AP, Gruzeliier JH (2003) Paradox lost? Exploring the role of alpha oscillations during externally vs. internally directed attention and the implications for idling and inhibition hypotheses. *International journal of psychophysiology : official journal of the International Organization of Psychophysiology* 47:65-74.
- de Lange FP, Jensen O, Bauer M, Toni I (2008) Interactions between posterior gamma and frontal alpha/beta oscillations during imagined actions. *Frontiers in human neuroscience* 2:7.
- Dhamala M, Rangarajan G, Ding M (2008) Estimating Granger causality from fourier and wavelet transforms of time series data. *Phys Rev Lett* 100:018701.
- Ding M, Chen Y, Bressler S (2006) Granger Causality: Basic Theory and Application to Neuroscience. In: *Handbook of Time Series Analysis*, (Schelter B, Winterhalder M, Timmer J, eds), pp 451-474: Wiley-VCH Verlage.
- Florin E, Baillet S (2015) The brain's resting-state activity is shaped by synchronized cross-frequency coupling of neural oscillations. *NeuroImage* 111:26-35.



- 1 Foxe JJ, Simpson GV, Ahlfors SP (1998) Parieto-occipital approximately 10 Hz activity  
2 reflects anticipatory state of visual attention mechanisms. *Neuroreport* 9:3929-3933.
- 3 Fries P (2005) A mechanism for cognitive dynamics: neuronal communication through  
4 neuronal coherence. *Trends in cognitive sciences* 9:474-480.
- 5 Gevins A, Smith ME, McEvoy L, Yu D (1997) High-resolution EEG mapping of cortical  
6 activation related to working memory: effects of task difficulty, type of processing,  
7 and practice. *Cerebral cortex* 7:374-385.
- 8 Geweke J (1982) Measurement of linear dependence and feedback between multiple time  
9 series. *J Am Stat Assoc* 77:304-313.
- 10 Glasser MF, Coalson TS, Robinson EC, Hacker CD, Harwell J, Yacoub E, Ugurbil K,  
11 Andersson J, Beckmann CF, Jenkinson M, Smith SM, Van Essen DC (2016) A multi-  
12 modal parcellation of human cerebral cortex. *Nature* 536:171-178.
- 13 Gonzalez Andino SL, Michel CM, Thut G, Landis T, Grave de Peralta R (2005) Prediction of  
14 response speed by anticipatory high-frequency (gamma band) oscillations in the  
15 human brain. *Human brain mapping* 24:50-58.
- 16 Granger CWJ (1969) Investigating Causal Relations by Econometric Models and Cross-  
17 spectral Methods. *Econometrica* 37:424-438.
- 18 Gross J, Kujala J, Hamalainen M, Timmermann L, Schnitzler A, Salmelin R (2001) Dynamic  
19 imaging of coherent sources: Studying neural interactions in the human brain.  
20 *Proceedings of the National Academy of Sciences of the United States of America*  
21 98:694-699.
- 22 Haegens S, Handel BF, Jensen O (2011) Top-down controlled alpha band activity in  
23 somatosensory areas determines behavioral performance in a discrimination task.  
24 *The Journal of neuroscience : the official journal of the Society for Neuroscience*  
25 31:5197-5204.
- 26 Haegens S, Osipova D, Oostenveld R, Jensen O (2010) Somatosensory working memory  
27 performance in humans depends on both engagement and disengagement of regions  
28 in a distributed network. *Human brain mapping* 31:26-35.
- 29 Haufe S, Nikulin VV, Muller KR, Nolte G (2013) A critical assessment of connectivity  
30 measures for EEG data: a simulation study. *NeuroImage* 64:120-133.
- 31 Hoogenboom N, Schoffelen JM, Oostenveld R, Fries P (2010) Visually induced gamma-band  
32 activity predicts speed of change detection in humans. *NeuroImage* 51:1162-1167.
- 33 Hoogenboom N, Schoffelen JM, Oostenveld R, Parkes LM, Fries P (2006) Localizing human  
34 visual gamma-band activity in frequency, time and space. *NeuroImage* 29:764-773.
- 35 Jensen O, Colgin LL (2007) Cross-frequency coupling between neuronal oscillations. *Trends*  
36 *in cognitive sciences* 11:267-269.
- 37 Jensen O, Mazaheri A (2010) Shaping functional architecture by oscillatory alpha activity:  
38 gating by inhibition. *Frontiers in human neuroscience* 4:186.
- 39 Jung TP, Makeig S, McKeown MJ, Bell AJ, Lee TW, Sejnowski TJ (2001) *Imaging Brain*  
40 *Dynamics Using Independent Component Analysis*. *Proceedings of the IEEE Institute*  
41 *of Electrical and Electronics Engineers* 89:1107-1122.
- 42 Kaminski M, Ding M, Truccolo WA, Bressler SL (2001) Evaluating causal relations in neural  
43 systems: granger causality, directed transfer function and statistical assessment of  
44 significance. *Biol Cybern* 85:145-157.
- 45 Kirihara K, Rissling AJ, Swerdlow NR, Braff DL, Light GA (2012) Hierarchical organization of  
46 gamma and theta oscillatory dynamics in schizophrenia. *Biological psychiatry* 71:873-  
47 880.
- 48 Klimesch W, Sauseng P, Hanslmayr S (2007) EEG alpha oscillations: the inhibition-timing  
49 hypothesis. *Brain research reviews* 53:63-88.
- 50 Kujala J, Jung J, Bouvard S, Lecaigard F, Lothe A, Bouet R, Ciumas C, Ryvlin P, Jerbi K  
51 (2015) Gamma oscillations in V1 are correlated with GABA(A) receptor density: A

1 multi-modal MEG and Flumazenil-PET study. *Sci Rep* 5:16347.  
2 Leszczynski M, Fell J, Axmacher N (2015) Rhythmic Working Memory Activation in the  
3 Human Hippocampus. *Cell Rep* 13:1272-1282.  
4 Lisman JE, Jensen O (2013) The theta-gamma neural code. *Neuron* 77:1002-1016.  
5 Maris E, Oostenveld R (2007) Nonparametric statistical testing of EEG- and MEG-data.  
6 *Journal of neuroscience methods* 164:177-190.  
7 Marshall TR, Bergmann TO, Jensen O (2015a) Frontoparietal Structural Connectivity  
8 Mediates the Top-Down Control of Neuronal Synchronization Associated with  
9 Selective Attention. *PLoS biology* 13:e1002272.  
10 Marshall TR, O'Shea J, Jensen O, Bergmann TO (2015b) Frontal eye fields control attentional  
11 modulation of alpha and gamma oscillations in contralateral occipitoparietal cortex.  
12 *The Journal of neuroscience : the official journal of the Society for Neuroscience*  
13 35:1638-1647.  
14 Mazaheri A, van Schouwenburg MR, Dimitrijevic A, Denys D, Cools R, Jensen O (2014)  
15 Region-specific modulations in oscillatory alpha activity serve to facilitate processing  
16 in the visual and auditory modalities. *NeuroImage* 87:356-362.  
17 Michalareas G, Vezoli J, van Pelt S, Schoffelen JM, Kennedy H, Fries P (2016) Alpha-Beta and  
18 Gamma Rhythms Subserve Feedback and Feedforward Influences among Human  
19 Visual Cortical Areas. *Neuron*.  
20 Miskovic V, Moscovitch DA, Santesso DL, McCabe RE, Antony MM, Schmidt LA (2011)  
21 Changes in EEG cross-frequency coupling during cognitive behavioral therapy for  
22 social anxiety disorder. *Psychol Sci* 22:507-516.  
23 Mitra PP, Pesaran B (1999) Analysis of dynamic brain imaging data. *Biophysical journal*  
24 76:691-708.  
25 Muthukumaraswamy SD, Singh KD (2013) Visual gamma oscillations: the effects of stimulus  
26 type, visual field coverage and stimulus motion on MEG and EEG recordings.  
27 *NeuroImage* 69:223-230.  
28 [Nolte G \(2003\) The magnetic lead field theorem in the quasi-static approximation and its](#)  
29 [use for magnetoencephalography forward calculation in realistic volume conductors.](#)  
30 [Physics in medicine and biology 48:3637-3652.](#)  
31 Oldfield RC (1971) The assessment and analysis of handedness: the Edinburgh inventory.  
32 *Neuropsychologia* 9:97-113.  
33 Oostenveld R, Fries P, Maris E, Schoffelen JM (2011) FieldTrip: Open source software for  
34 advanced analysis of MEG, EEG, and invasive electrophysiological data. *Computational*  
35 *intelligence and neuroscience* 2011:156869.  
36 Park H, Kang E, Kang H, Kim JS, Jensen O, Chung CK, Lee DS (2011) Cross-frequency  
37 power correlations reveal the right superior temporal gyrus as a hub region during  
38 working memory maintenance. *Brain Connect* 1:460-472.  
39 Park H, Lee DS, Kang E, Kang H, Hahn J, Kim JS, Chung CK, Jiang H, Gross J, Jensen O  
40 (2016) Formation of visual memories controlled by gamma power phase-locked to  
41 alpha oscillations. *Sci Rep* 6:28092.  
42 Perry G, Hamandi K, Brindley LM, Muthukumaraswamy SD, Singh KD (2013) The properties  
43 of induced gamma oscillations in human visual cortex show individual variability in  
44 their dependence on stimulus size. *NeuroImage* 68:83-92.  
45 Popov T, Popova P (2015) Same clock, different time read-out: Spontaneous brain  
46 oscillations and their relationship to deficient coding of cognitive content.  
47 *NeuroImage* 119:316-324.  
48 Popov T, Kastner S, Jensen O (2017) FEF-Controlled Alpha Delay Activity Precedes  
49 Stimulus-Induced Gamma-Band Activity in Visual Cortex. *The Journal of*  
50 *neuroscience : the official journal of the Society for Neuroscience* 37:4117-4127.  
51 Roux F, Uhlhaas PJ (2014) Working memory and neural oscillations: alpha-gamma versus

1 theta-gamma codes for distinct WM information? Trends in cognitive sciences 18:16-  
2 25.

3 Roux F, Wibral M, Mohr HM, Singer W, Uhlhaas PJ (2012) Gamma-band activity in human  
4 prefrontal cortex codes for the number of relevant items maintained in working  
5 memory. The Journal of neuroscience : the official journal of the Society for  
6 Neuroscience 32:12411-12420.

7 Saalman YB, Pinsk MA, Wang L, Li X, Kastner S (2012) The pulvinar regulates information  
8 transmission between cortical areas based on attention demands. Science 337:753-  
9 756.

10 Schoffelen JM, Oostenveld R, Fries P (2005) Neuronal coherence as a mechanism of effective  
11 corticospinal interaction. Science 308:111-113.

12 Schoffelen JM, Hulten A, Lam N, Marquand AF, Udden J, Hagoort P (2017) Frequency-  
13 specific directed interactions in the human brain network for language. Proceedings  
14 of the National Academy of Sciences of the United States of America 114:8083-8088.

15 Spaak E, Bonnefond M, Maier A, Leopold DA, Jensen O (2012) Layer-specific entrainment of  
16 gamma-band neural activity by the alpha rhythm in monkey visual cortex. Current  
17 biology : CB 22:2313-2318.

18 Staresina BP, Bergmann TO, Bonnefond M, van der Meij R, Jensen O, Deuker L, Elger CE,  
19 Axmacher N, Fell J (2015) Hierarchical nesting of slow oscillations, spindles and  
20 ripples in the human hippocampus during sleep. Nat Neurosci 18:1679-1686.

21 Staudigl T, Hanslmayr S (2013) Theta oscillations at encoding mediate the context-  
22 dependent nature of human episodic memory. Current biology : CB 23:1101-1106.

23 Swettenham JB, Muthukumaraswamy SD, Singh KD (2009) Spectral properties of induced  
24 and evoked gamma oscillations in human early visual cortex to moving and stationary  
25 stimuli. Journal of neurophysiology 102:1241-1253.

26 Tallon-Baudry C, Bertrand O (1999) Oscillatory gamma activity in humans and its role in  
27 object representation. Trends in cognitive sciences 3:151-162.

28 Tallon-Baudry C, Bertrand O, Peronnet F, Pernier J (1998) Induced gamma-band activity  
29 during the delay of a visual short-term memory task in humans. The Journal of  
30 neuroscience : the official journal of the Society for Neuroscience 18:4244-4254.

31 Tort AB, Komorowski R, Eichenbaum H, Kopell N (2010) Measuring phase-amplitude  
32 coupling between neuronal oscillations of different frequencies. Journal of  
33 neurophysiology 104:1195-1210.

34 Tort AB, Komorowski RW, Manns JR, Kopell NJ, Eichenbaum H (2009) Theta-gamma  
35 coupling increases during the learning of item-context associations. Proceedings of  
36 the National Academy of Sciences of the United States of America 106:20942-20947.

37 Tort AB, Kramer MA, Thorn C, Gibson DJ, Kubota Y, Graybiel AM, Kopell NJ (2008)  
38 Dynamic cross-frequency couplings of local field potential oscillations in rat striatum  
39 and hippocampus during performance of a T-maze task. Proceedings of the National  
40 Academy of Sciences of the United States of America 105:20517-20522.

41 van Ede F, de Lange FP, Maris E (2012) Attentional cues affect accuracy and reaction time via  
42 different cognitive and neural processes. The Journal of neuroscience : the official  
43 journal of the Society for Neuroscience 32:10408-10412.

44 van Ede F, Szebenyi S, Maris E (2014) Attentional modulations of somatosensory alpha, beta  
45 and gamma oscillations dissociate between anticipation and stimulus processing.  
46 NeuroImage 97:134-141.

47 van Ede F, de Lange F, Jensen O, Maris E (2011) Orienting attention to an upcoming tactile  
48 event involves a spatially and temporally specific modulation of sensorimotor alpha-  
49 and beta-band oscillations. The Journal of neuroscience : the official journal of the  
50 Society for Neuroscience 31:2016-2024.

51 van Kerkoerle T, Self MW, Dagnino B, Gariel-Mathis MA, Poort J, van der Togt C,

1 Roelfsema PR (2014) Alpha and gamma oscillations characterize feedback and  
2 feedforward processing in monkey visual cortex. *Proceedings of the National*  
3 *Academy of Sciences of the United States of America* 111:14332-14341.

4 Van Veen BD, van Drongelen W, Yuchtman M, Suzuki A (1997) Localization of brain  
5 electrical activity via linearly constrained minimum variance spatial filtering. *IEEE*  
6 *transactions on bio-medical engineering* 44:867-880.

7 van Wijk BC, Beudel M, Jha A, Oswal A, Foltynie T, Hariz MI, Limousin P, Zrinzo L, Aziz TZ,  
8 Green AL, Brown P, Litvak V (2016) Subthalamic nucleus phase-amplitude coupling  
9 correlates with motor impairment in Parkinson's disease. *Clin Neurophysiol*  
10 127:2010-2019.

11 Wang L, Hagoort P, Jensen O (2017) Language Prediction Is Reflected by Coupling between  
12 Frontal Gamma and Posterior Alpha Oscillations. *Journal of cognitive neuroscience*:1-  
13 16.

14 Weisz N, Hartmann T, Muller N, Lorenz I, Obleser J (2011) Alpha rhythms in audition:  
15 cognitive and clinical perspectives. *Frontiers in psychology* 2:73.

16 Wen X, Rangarajan G, Ding M (2013) Multivariate Granger causality: an estimation  
17 framework based on factorization of the spectral density matrix. *Philosophical*  
18 *transactions Series A, Mathematical, physical, and engineering sciences* 371:20110610.

19 Whittingstall K, Logothetis NK (2009) Frequency-band coupling in surface EEG reflects  
20 spiking activity in monkey visual cortex. *Neuron* 64:281-289.

21 Wilson GT (1972) The factorization of matrical spectral densities. *SIAM J Appl Math* 23:420-  
22 426.

23 Winkler I, Haufe S, Porbadnigk AK, Muller KR, Dahne S (2015) Identifying Granger causal  
24 relationships between neural power dynamics and variables of interest. *NeuroImage*  
25 111:489-504.

26