

Tilings in randomly perturbed dense graphs

Balogh, Jozsef; Treglown, Andrew; Wagner, Adam Zsolt

DOI:

[10.1017/S0963548318000366](https://doi.org/10.1017/S0963548318000366)

License:

Other (please specify with Rights Statement)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Balogh, J, Treglown, A & Wagner, AZ 2019, 'Tilings in randomly perturbed dense graphs', *Combinatorics, Probability and Computing*, vol. 28, no. 2, pp. 159-176. <https://doi.org/10.1017/S0963548318000366>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Checked for eligibility: 05/06/2018

© Cambridge University Press 2018

Published in *Combinatorics, Probability and Computing* on 16/07/2018

DOI: <https://doi.org/10.1017/S0963548318000366>

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Tilings in randomly perturbed dense graphs

József Balogh*, Andrew Treglown[†] and Adam Zsolt Wagner[‡]

June 1, 2018

Abstract

A perfect H -tiling in a graph G is a collection of vertex-disjoint copies of a graph H in G that together cover all the vertices in G . In this paper we investigate perfect H -tilings in a random graph model introduced by Bohman, Frieze and Martin [6] in which one starts with a dense graph and then adds m random edges to it. Specifically, for *any* fixed graph H , we determine the number of random edges required to add to an arbitrary graph of linear minimum degree in order to ensure the resulting graph contains a perfect H -tiling with high probability. Our proof utilises Szemerédi's Regularity lemma [29] as well as a special case of a result of Komlós [18] concerning almost perfect H -tilings in dense graphs.

MSC2000: 5C35, 5C70, 5C80.

1 Introduction

Embedding problems form a central part of both extremal and random graph theory. Indeed, many results in extremal graph theory concern minimum degree conditions that force some spanning substructure. For example, a foundation stone in the subject is Dirac's theorem [10] from 1952 which states that every graph G on $n \geq 3$ vertices and with minimum degree $\delta(G) \geq n/2$ is Hamiltonian. More recently, an important paper of Böttcher, Schacht and Taraz [9] resolved the Bollobás–Komlós conjecture; specifically, it provided a minimum degree condition which ensures a graph contains every r -partite spanning subgraph of bounded degree and small bandwidth.

Recall that the Erdős–Rényi random graph $G_{n,p}$ consists of vertex set $[n] := \{1, \dots, n\}$ where each edge is present with probability p , independently of all other choices. In this

*Department of Mathematical Sciences, University of Illinois at Urbana-Champaign, Urbana, Illinois 61801, USA jobal@math.uiuc.edu. Research is partially supported by NSF Grant DMS-1500121 and Arnold O. Beckman Research Award (UIUC Campus Research Board 15006).

[†]University of Birmingham, United Kingdom, a.c.treglown@bham.ac.uk. Research supported by EPSRC grant EP/M016641/1.

[‡]University of Illinois at Urbana-Champaign, Urbana, Illinois 61801, USA, zawagne2@illinois.edu.

setting, a key question is to establish the *threshold* at which $G_{n,p}$, with high probability, contains some spanning subgraph. In the case of Hamilton cycles, Pósa [27] showed that if $p \gg \log n/n$ then asymptotically almost surely (a.a.s.) $G_{n,p}$ is Hamiltonian whilst if $p \ll \log n/n$ a.a.s. $G_{n,p}$ is not Hamiltonian. In general though few thresholds for embedding a fixed spanning subgraph H in $G_{n,p}$ are known.

Bohman, Frieze and Martin [6] introduced a model which in some sense connects the two aforementioned questions together. Indeed, in their model one starts with a dense graph and then adds m random edges to it. A natural problem in this setting is to determine *how many* random edges are required to ensure that the resulting graph a.a.s. contains a given graph H as a spanning subgraph. For example, the main result in [6] states that for every $\alpha > 0$, there is a $c = c(\alpha)$ such that if we start with an arbitrary n -vertex graph G of minimum degree $\delta(G) \geq \alpha n$ and add cn random edges to it, then a.a.s. the resulting graph is Hamiltonian. This result is best possible in the sense that there are graphs G of linear minimum degree that require a linear number of edges to be added to become Hamiltonian (for example, consider any complete bipartite graph with vertex classes of size an, bn where $0 < a < b < 1$ and $a + b = 1$). Recently, Krivelevich, Kwan and Sudakov [21] proved an analogous result where now we wish to embed a fixed spanning tree of bounded degree. Other properties of this model (embedding a fixed subgraph, the diameter, connectivity, Ramsey properties) have been studied, for example, in [5, 22]. In [20] the framework was also generalised to hypergraphs and a number of exact results concerning perfect matchings and cycles in hypergraphs and digraphs were proven. Further, since the paper was submitted, a range of other results in the area have been obtained [3, 4, 7, 8, 14, 17, 26].

Krivelevich, Kwan and Sudakov [21] raised the question of determining an analogue of the Böttcher–Schacht–Taraz theorem [9] in the setting of randomly perturbed dense graphs. In this paper we consider an important subcase of this problem; *perfect H -tilings*.

1.1 Perfect tilings in graphs and random graphs

Given two graphs H and G , an H -tiling in G is a collection of vertex-disjoint copies of H in G . An H -tiling is called *perfect* if it covers all the vertices of G . Perfect H -tilings are also referred to as H -factors or *perfect H -packings*. A seminal result in extremal graph theory is the celebrated Hajnal–Szemerédi theorem [13] which determines the minimum degree threshold that ensures a graph contains a perfect K_r -tiling. Building on this result, Kühn and Osthus [23, 24] characterised, up to an additive constant, the minimum degree which ensures that a graph G contains a perfect H -tiling for an *arbitrary* graph H .

The perfect tiling problem for random graphs has also received significant attention. The most striking result in the area is a theorem of Johansson, Kahn and Vu [16] which determines the threshold for the property that the Erdős–Rényi random graph $G_{n,p}$ a.a.s. contains a perfect H -tiling where H is any fixed *strictly balanced* graph. Recently, Gerke and McDowell [12] determined the corresponding threshold in the case when H is a *nonvertex-balanced* graph. A number of subcases for both of these aforementioned results had been earlier proved e.g. in [1, 11, 15, 25]. Since it is instructive to compare both of these results to our main result, we will formally state them below.

Suppose that H is a graph on at least two vertices. We write $e(H)$ and $|H|$ for the number of edges and vertices in G respectively. Define

$$d(H) := \frac{e(H)}{|H| - 1} \quad \text{and} \quad d^*(H) := \max\{d(H') : H' \subseteq H, |H'| \geq 2\}.$$

We say that H is *strictly balanced* if for every proper subgraph H' of H with at least two vertices, $d(H') < d(H)$. H is *balanced* if $d^*(H) = d(H)$. For $v \in V(H)$, let

$$d^*(v, H) := \max\{d(H') : H' \subseteq H, |H'| \geq 2, v \in V(H')\}.$$

A graph H is *vertex-balanced* if, for all $v \in V(H)$, $d^*(v, H) = d^*(H)$.

Notice that if H is balanced then it is vertex-balanced. Further, if H is nonvertex-balanced then it is not balanced and not strictly balanced. The following seminal result of Johansson, Kahn and Vu [16] determines the threshold for the property that $G_{n,p}$ a.a.s. contains a perfect H -tiling where H is an arbitrary fixed strictly balanced graph.

Theorem 1.1 (Johansson, Kahn and Vu [16]). *Let H be a strictly balanced graph with m edges and let $n \in \mathbb{N}$ be divisible by $|H|$.*

- *If $p \gg n^{-1/d(H)}(\log n)^{1/m}$ then a.a.s. $G_{n,p}$ contains a perfect H -tiling.*
- *If $p \ll n^{-1/d(H)}(\log n)^{1/m}$ then a.a.s. $G_{n,p}$ does not contain a perfect H -tiling.*

Note here (and elsewhere where we consider bounds on p), we write e.g. $p \ll n^{-1/d(H)}(\log n)^{1/m}$ to mean $p = o(n^{-1/d(H)}(\log n)^{1/m})$. (Later we also use the \ll notation in hierarchies of constants; this is defined in Section 1.3.)

Johansson, Kahn and Vu [16] conjectured that Theorem 1.1 can be generalised to *all* vertex-balanced graphs H : Given any vertex $v \in V(H)$, define

$$s_v := \min\{e(H') : H' \subseteq H, |H'| \geq 2, v \in V(H') \text{ and } d(H') = d^*(v, H)\}$$

and let s be the maximum of the s_v 's amongst all $v \in V(H)$. They conjectured that $n^{-1/d^*(H)}(\log n)^{1/s}$ is the threshold for the property that $G_{n,p}$ a.a.s. contains a perfect H -tiling where H is an arbitrary fixed vertex-balanced graph. Note that Theorem 1.1 is a special case of this conjecture.

For nonvertex-balanced graphs, the following result determines the corresponding threshold.

Theorem 1.2 (Gerke and McDowell [12]). *Let H be a nonvertex-balanced graph and let $n \in \mathbb{N}$ be divisible by $|H|$.*

- *If $p \gg n^{-1/d^*(H)}$ then a.a.s. $G_{n,p}$ contains a perfect H -tiling.*
- *If $p \ll n^{-1/d^*(H)}$ then a.a.s. $G_{n,p}$ does not contain a perfect H -tiling.*

1.2 Tilings in randomly perturbed dense graphs

The main result of this paper is to determine how many random edges one needs to add to a graph of linear minimum degree to ensure it contains a perfect H -tiling for *any* fixed graph H . Throughout the paper we assume that an n -vertex graph has vertex set $[n]$ and if G and G' are n -vertex graphs then we define $G \cup G'$ to be the n -vertex (simple) graph with edge set $E(G) \cup E(G')$. We are now ready to state our main result.

Theorem 1.3. *Let H be a fixed graph with at least one edge and let $n \in \mathbb{N}$ be divisible by $|H|$. For every $\alpha > 0$, there is a $c = c(\alpha, H) > 0$ such that if $p \geq cn^{-1/d^*(H)}$ and G is an n -vertex graph with minimum degree $\delta(G) \geq \alpha n$ then a.a.s. $G \cup G_{n,p}$ contains a perfect H -tiling.*

Theorem 1.3 is best-possible in the sense that, for any fixed graph H , there are n -vertex graphs G of linear minimum degree such that if $p \ll n^{-1/d^*(H)}$ then a.a.s. $G \cup G_{n,p}$ does not contain a perfect H -tiling. We explain this in more detail in Section 2.1.

In the case when H is strictly balanced notice that unlike Theorem 1.1, Theorem 1.3 does not involve a logarithmic term. Thus comparing our model with the Erdős–Rényi model, we see that starting with a graph of linear minimum degree instead of the empty graph saves a logarithmic factor in terms of how many random edges one needs to ensure the resulting graph a.a.s. contains a perfect H -tiling. This same phenomenon is also exhibited in the analogous problems for Hamilton cycles [6] and spanning trees [21], as well as for matchings and loose cycles in the hypergraph setting [20]. Further, if the Johansson, Kahn and Vu conjecture is true then together with Theorem 1.3 this shows that the same phenomenon occurs for perfect H -tilings for any vertex-balanced H . Interestingly though the threshold in Theorem 1.2 is the *same* as that in Theorem 1.3. In other words, if H is nonvertex-balanced, starting with a graph of linear minimum degree instead of the empty graph essentially provides *no benefit* in terms of how many random edges one needs to ensure the resulting graph a.a.s. contains a perfect H -tiling!

It is also instructive to compare Theorem 1.3 to the problem of finding an almost perfect tiling in the random graph: We say that $G_{n,p}$ has an *almost perfect H -tiling* if for every $\varepsilon > 0$, the probability that the largest H -tiling in $G_{n,p}$ covers less than $(1 - \varepsilon)n$ vertices tends to zero as n tends to infinity. Ruciński [28] proved that $n^{-1/d^*(H)}$ is the threshold for $G_{n,p}$ having an almost perfect H -tiling (for any fixed graph H). Thus, in Theorem 1.3 one can already guarantee an almost perfect H -tiling without using any of the edges from G . Hence (in the case when H is strictly balanced), the edges in G are necessary to ‘transform’ an almost perfect H -tiling in $G_{n,p}$ into a perfect H -tiling in $G \cup G_{n,p}$.

The proof of Theorem 1.3 utilises Szemerédi’s Regularity lemma [29] as well as a special case of a result of Komlós [18] concerning almost perfect H -tilings in dense graphs. We also draw on ideas from [2, 21]. In Section 2.2 we give an overview of the proof.

1.3 Notation

Let G be a graph. We write $V(G)$ for the vertex set of G , $E(G)$ for the edge set of G and define $|G| := |V(G)|$ and $e(G) := |E(G)|$. Given a subset $X \subseteq V(G)$, we write $G[X]$ for

the subgraph of G induced by X . Given some $x \in V(G)$ we write $G - x$ for the subgraph of G induced by $V(G) \setminus \{x\}$. The degree of x is denoted by $d_G(x)$ and its neighbourhood by $N_G(x)$. Given a vertex $x \in V(G)$ and a set $Y \subseteq V(G)$ we write $d_G(x, Y)$ to denote the number of edges xy where $y \in Y$. Given disjoint vertex classes $X, Y \subseteq V(G)$, we write $G[X, Y]$ for the bipartite graph with vertex classes X and Y whose edge set consists of all those edges in G with one endpoint in X and the other in Y ; we write $e_G(X, Y)$ for the number of edges in $G[X, Y]$. Given a set X and $t \in \mathbb{N}$, let $\binom{X}{t}$ denote the set of all subsets of X of size t .

We write $0 < \alpha \ll \beta \ll \gamma$ to mean that we can choose the constants α, β, γ from right to left. More precisely, there are increasing functions f and g such that, given γ , whenever we choose some $\beta \leq f(\gamma)$ and $\alpha \leq g(\beta)$, all calculations needed in our proof are valid. Hierarchies of other lengths are defined in the obvious way. Throughout the paper we omit floors and ceilings whenever this does not affect the argument.

The paper is organised as follows. In Section 2 we give an overview of the proof of Theorem 1.3; we also give an example that shows the bound on p in Theorem 1.3 is best possible up to a multiplicative constant. Szemerédi's Regularity lemma is presented in Section 3 and then we introduce some useful tools in Section 4. The proof of Theorem 1.3 is given in Section 5.

2 An extremal example and overview of the proof

2.1 An extremal example

In this subsection we prove that Theorem 1.3 is best-possible in the sense that, given any graph H , there exist (sequences of) n -vertex graphs G_n with linear minimum degree so that, if $p \ll n^{-1/d^*(H)}$, then a.a.s. $G_n \cup G_{n,p}$ does not contain a perfect H -tiling. For this we require the following result. Recall the definition

$$d^*(H) := \max \left\{ \frac{e(H')}{v(H') - 1} : H' \subseteq H, |H'| \geq 2 \right\}.$$

Theorem 2.1 ([15], part of Theorem 4.9). *For every graph H with at least one edge and for every $0 < \varepsilon < 1$ there is a positive constant $c = c(H, \varepsilon)$ such that if $p \leq cn^{-1/d^*(H)}$,*

$$\lim_{n \rightarrow \infty} \mathbb{P}(G_{n,p} \text{ contains an } H\text{-tiling covering at least } \varepsilon n \text{ vertices}) = 0.$$

Consider any fixed graph H with at least one edge and let $b > a > 0$ such that $a + b = 1$ and $b > a(|H| - 1)$. Set $\varepsilon := b - a(|H| - 1) > 0$ and define $c = c(H, \varepsilon)$ as in Theorem 2.1. Let $n \in \mathbb{N}$ be divisible by $|H|$ and let G_n be the complete bipartite graph with vertex classes X and Y of sizes an and bn respectively. Consider $G'_n := G_n \cup G_{n,p}$ where $p \leq c(bn)^{-1/d^*(H)}$. Notice that if G'_n contains a perfect H -tiling, then $G_{n,p}[Y] \cong G_{bn,p}$ must contain an H -tiling covering at least $\varepsilon n > \varepsilon(bn)$ vertices. However, by the choice of c , Theorem 2.1 implies that a.a.s. such an H -tiling in $G_{n,p}[Y]$ does not exist. Thus, a.a.s. G'_n does not contain a perfect H -tiling. So we have indeed shown that the bound on p in Theorem 1.3 is best-possible.

The above example shows that if $0 < \alpha < 1/|H|$ then we need the random edges with $p \geq Cn^{-1/d^*(H)}$ to force a perfect H -tiling. We also note that if $\alpha > 1 - \frac{1}{\chi(H)}$ and n is large then by a theorem of Komlós, Sárközy and Szemerédi [19], every n -vertex graph G_n of minimum degree αn contains a perfect H -tiling and thus there is no need for the random edges at all. More generally, given any graph H , Kühn and Osthus [23, 24] determined the smallest $\alpha^* = \alpha^*(H) > 0$ such that, given any $\alpha > \alpha^*$, every sufficiently large n -vertex graph G_n of minimum degree at least αn contains a perfect H -tiling. It may be of interest to investigate the following question: given fixed α with $1/|H| < \alpha < \alpha^*$, how large must p be to ensure that whenever G_n has minimum degree at least αn then $G_n \cup G_{n,p}$ a.a.s. contains a perfect H -tiling?

Another natural question is whether Theorem 1.3 holds if we replace αn with a sublinear term. Note that the approach we use is not suitable for attacking this problem (since we apply Szemerédi's Regularity lemma).

2.2 Overview of the proof of Theorem 1.3

The first step of the proof is to obtain some special structure within G to help us embed the perfect H -tiling. In particular, by applying Szemerédi's Regularity lemma (Lemma 3.1) and a theorem of Komlós (Theorem 4.1) we obtain a spanning subgraph of G which 'looks' like the blow-up of a collection of stars. More precisely, there is a spanning subgraph G' of G ; constants $k, t \in \mathbb{N}$, and; a partition of $V(G)$ into classes V_0 and $W_{i,j}$ (for all $0 \leq i \leq t$ and $1 \leq j \leq k$), such that:

- V_0 is 'small';
- Each cluster $W_{i,j}$ has the same size;
- For all $1 \leq i \leq t$, $1 \leq j \leq k$, the pair $(W_{0,j}, W_{i,j})_{G'}$ is 'super-regular'.

We remark that a similar structure was used in [21] (though the role of the blown-up stars was different there).

If $t = 1$, then the aforementioned structure would consist of a collection of disjoint super-regular pairs $(W_{0,j}, W_{1,j})_{G'}$ and an 'exceptional set' V_0 . We could then obtain a perfect H -tiling using the strategy described below.

The first step is to find a small H -tiling \mathcal{H} that covers all of V_0 but so that \mathcal{H} only intersects the super-regular pairs $(W_{0,j}, W_{1,j})_{G'}$ in a very small number of vertices. To see that such an H -tiling \mathcal{H} in $G \cup G_{n,p}$ exists, note that given any $v \in V_0$, $N_G(v)$ has linear size so a.a.s, $G_{n,p}$ contains many copies of $H - x$ in $N_G(v)$ (for some $x \in V(H)$). In particular, this implies that v lies in many copies of H in $G \cup G_{n,p}$. Thus, this property allows us to greedily construct \mathcal{H} (though some care is needed to ensure we do not use too many vertices in any one cluster $W_{i,j}$).

Now if we remove all the vertices lying in \mathcal{H} we still have that each $(W_{0,j}, W_{1,j})_{G'}$ is super-regular. This structure can then be used to find an H -tiling \mathcal{H}_j in $G \cup G_{n,p}$ covering precisely the vertices of $W_{0,j} \cup W_{1,j}$. Indeed, in this case we employ an approach very similar

to that used in [2]. Then $\mathcal{H}, \mathcal{H}_1, \dots, \mathcal{H}_k$ together form a perfect H -tiling in $G \cup G_{n,p}$, as desired.

In particular, note that roughly speaking the authors of [2] prove that if (A, B) is a (very dense) super-regular pair in a graph G of small independence number, then there is a triangle-tiling in G covering precisely the vertices of $A \cup B$. Here the small independence number ensures we have large matchings in both $G[A]$ and $G[B]$; then the edges between A and B can be used to extend such edges to triangles with at least one vertex in each class, and thus ultimately (with significant care) one obtains the desired triangle-tiling. In our setting, the edges from $G_{n,p}$ ensure that, given any super-regular pair (A, B) in G , we have large $(H - x)$ -tilings in both $G_{n,p}[A]$ and $G_{n,p}[B]$ (for some $x \in V(H)$). We then (again with some care) extend such copies of $H - x$ to copies of H using the edges between A and B in G , to obtain the desired H -tiling.

To employ the approach used in [2], we really do require that $t = 1$. That is, the structure in G looks like a blow-up of disjoint edges. However, since the minimum degree of G is typically very small we can only ensure a structure in G that looks like a blow-up of stars, each of which contains a huge constant number of leaves (i.e. t is large).

Instead, we have to first carefully choose a large H -tiling \mathcal{H}' in $G \cup G_{n,p}$ so that what remains uncovered by \mathcal{H}' is precisely a collection of clusters that, in G' , form disjoint super-regular pairs. We then can proceed as described above. To construct a suitable H -tiling \mathcal{H}' , our proof heavily uses the blown-up star structure we found initially in G .

3 The Regularity lemma

In the proof of our main result we will make use of the Szemerédi's Regularity lemma [29], hence in this section we introduce the necessary notation and set-up for this lemma. The *density* of a bipartite graph G with vertex classes A and B is defined as

$$d_G(A, B) := \frac{e(A, B)}{|A||B|}.$$

Given any $\varepsilon > 0$ we say that G is ε -regular if for all sets $X \subseteq A$ and $Y \subseteq B$ with $|X| \geq \varepsilon|A|$ and $|Y| \geq \varepsilon|B|$ we have $|d_G(A, B) - d_G(X, Y)| < \varepsilon$. In this case we also say that $(A, B)_G$ is an ε -regular pair. Given $d \in [0, 1)$ we say that G is (ε, d) -super-regular if all sets $X \subseteq A$ and $Y \subseteq B$ with $|X| \geq \varepsilon|A|$ and $|Y| \geq \varepsilon|B|$ satisfy $d_G(X, Y) > d$ and moreover $d_G(a) > d|B|$ and $d_G(b) > d|A|$ for all $a \in A$ and $b \in B$.

The following degree form of the Regularity lemma can be easily derived from the classical version and will be particularly useful for us.

Lemma 3.1 (Regularity lemma). *For every $\varepsilon > 0$ and each integer ℓ_0 there is an $M = M(\varepsilon, \ell_0)$ such that if G is any graph on at least M vertices and $d \in [0, 1)$, then there exists a partition of $V(G)$ into $\ell + 1$ classes V_0, V_1, \dots, V_ℓ , and a spanning subgraph $G' \subseteq G$ with the following properties:*

- (i) $\ell_0 \leq \ell \leq M$, $|V_0| \leq \varepsilon|G|$, $|V_1| = \dots = |V_\ell| =: L$;

- (ii) $d_{G'}(v) > d_G(v) - (d + \varepsilon)|G|$ for all $v \in V(G)$;
- (iii) $e(G'[V_i]) = 0$ for all $i \geq 1$;
- (iv) for all $1 \leq i < j \leq \ell$ the graph $(V_i, V_j)_{G'}$ is ε -regular and has density either 0 or greater than d .

The sets V_1, \dots, V_ℓ are called *clusters*, V_0 is called the *exceptional set* and the vertices in V_0 *exceptional vertices*. We refer to G' as the *pure graph* of G . The *reduced graph* R of G is the graph whose vertices are V_1, \dots, V_ℓ and in which V_i is adjacent to V_j whenever $(V_i, V_j)_{G'}$ is ε -regular and has density greater than d .

Next we see that given a regular pair we can approximate it by a super-regular pair. The following lemma can be found in e.g. [30].

Lemma 3.2. *If (A, B) is an ε -regular pair with density d in a graph G (where $0 < \varepsilon < 1/3$), then there exists $A' \subseteq A$ and $B' \subseteq B$ with $|A'| \geq (1 - \varepsilon)|A|$ and $|B'| \geq (1 - \varepsilon)|B|$, such that (A', B') is a $(2\varepsilon, d - 3\varepsilon)$ -super-regular pair.*

4 Some useful results

4.1 Almost perfect star tilings

An important result of Komlós [18] determines the minimum degree threshold that forces an *almost* perfect H -tiling in a graph, for any fixed graph H . As hinted at in the proof overview, we require that the reduced graph R of G contains an almost perfect tiling of stars. The following special case of Komlós' theorem ensures such a tiling exists.

Theorem 4.1 (Komlós [18]). *Given any $t \in \mathbb{N}$ and $\varepsilon > 0$, there is an integer $n_0 = n_0(t, \varepsilon)$ so that, if $n \geq n_0$ and G is a graph on n vertices with*

$$\delta(G) \geq \frac{n}{(t+1)},$$

then G contains a $K_{1,t}$ -tiling that covers all but at most εn vertices.

4.2 Embedding H in random graphs

Let $\eta > 0$. Given an n -vertex graph G and a graph H we write $G \in F_H(\eta)$ if every induced subgraph of G on at least ηn vertices contains H as a (not necessarily induced) subgraph. The next theorem will allow us to find almost perfect H -tilings in large subgraphs of $G_{n,p}$.

Theorem 4.2 ([15], part of Theorem 4.9). *For every graph H with at least one edge and for every $\eta > 0$ there is a positive constant $C = C(H, \eta)$ such that if $p \geq Cn^{-1/d^*(H)}$,*

$$\lim_{n \rightarrow \infty} \mathbb{P}(G_{n,p} \in F_H(\eta)) = 1.$$

We remark that the proofs of both Theorem 4.2 and the next two results follow from straightforward applications of Janson's inequality.

Let $\gamma > 0$ and H be a graph. Consider a graph G with vertex set $[n]$. Let \mathcal{H}_n be a collection of copies of H in K_n (where we view K_n to have vertex set $[n]$). We write $G \in F_H(\gamma, \mathcal{H}_n)$ if every induced subgraph of G on at least γn vertices contains a copy of H that is not an element of \mathcal{H}_n . The next theorem allows us to find copies of H in $G_{n,p}$ that avoid certain edge sets.

Theorem 4.3. *Let H be a graph with at least one edge and suppose $0 < \gamma_1 \ll \gamma_2 \ll 1/|H|$. Then there is a positive constant $D = D(H, \gamma_1, \gamma_2)$ such that the following holds. Let \mathcal{H}_n be a collection of at most $\gamma_1 \binom{n}{|H|}$ copies of H in K_n . If $p \geq Dn^{-1/d^*(H)}$, then*

$$\lim_{n \rightarrow \infty} \mathbb{P}(G_{n,p} \in F_H(\gamma_2, \mathcal{H}_n)) = 1.$$

The proof of Theorem 4.2 is a simple application of Theorem 3.9 from [15] (which in turn follows from Janson's inequality). To prove Theorem 4.3 one can follow precisely the same proof, however, by instead applying a version of Theorem 3.9 from [15] in the setting where now some copies of H are excluded; again to prove such a result one follows the proof of Theorem 3.9 from [15] precisely.

Similarly to the above, given an n -vertex graph G and a graph H we write $G \in F'_H(\eta)$ if for all ordered pairs of disjoint sets $A, B \subset V(G)$ of size $|A|, |B| \geq \eta n$ there exists a copy of H in G with precisely one vertex in A and $|H| - 1$ vertices in B . Again, the same argument as in the proof of Theorem 4.2 shows the following.

Theorem 4.4. *For every graph H with at least one edge and for every $\eta > 0$ there is a positive constant $C = C(H, \eta)$ such that if $p \geq Cn^{-1/d^*(H)}$,*

$$\lim_{n \rightarrow \infty} \mathbb{P}(G_{n,p} \in F'_H(\eta)) = 1.$$

5 Proof of Theorem 1.3

Let H and $\alpha > 0$ be as in the statement of the theorem. Note that it suffices to prove the theorem in the case when $\alpha \ll 1/|H|$. Define additional constants $\phi, \ell_0, \varepsilon, \varepsilon_1, \varepsilon_2, \varepsilon_3, \varepsilon_4, \varepsilon_5, d_1, d$ and apply the Regularity lemma (Lemma 3.1) with input ε, ℓ_0 to obtain $M = M(\varepsilon, \ell_0)$ and define $\eta > 0$ such that $\eta \ll 1/M$, so that we have

$$0 < \eta \ll 1/M \leq 1/\ell_0 \ll \varepsilon \ll \varepsilon_1 \ll \varepsilon_2 \ll \varepsilon_3 \ll \varepsilon_4 \ll \phi \ll \varepsilon_5 \ll d_1 \ll d \ll \alpha \ll 1/|H|. \quad (5.1)$$

Set

$$t := \lceil 2\alpha^{-1} \rceil$$

and let H' be some fixed induced subgraph of H on $|H|-1$ vertices. Let $c = c(\eta, \varepsilon_2, \varepsilon_3, M, t, H) = c(\alpha, H)$ be a positive constant such that: (i) on input H , η , the conclusion of both Theorem 4.2 and Theorem 4.4 hold with $c/4$ playing the role of C and; (ii) $c \geq 4D(4Mt)^{1/d^*(H)}$ where D is the output of Theorem 4.3 on input H' , ε_2 and ε_3 .

Let G be a sufficiently large n -vertex graph with n divisible by $|H|$ and $\delta(G) \geq \alpha n$. Set

$$p := cn^{-1/d^*(H)}.$$

We wish to show that $G \cup G_{n,p}$ a.a.s. contains a perfect H -tiling. Our first step towards proving this will be to switch our attention to an appropriate subgraph of G .

Claim 5.1. *There is a $k \in \mathbb{N}$, a partition of $V(G)$ into classes V_0 and $W_{i,j}$ (for all $0 \leq i \leq t$ and $1 \leq j \leq k$), and a spanning subgraph G' of G such that:*

- (i) $\frac{\ell_0}{2t} \leq k \leq \frac{M}{t+1}$;
- (ii) $|V_0| \leq 2t\varepsilon n$;
- (iii) Each cluster $W_{i,j}$ has the same size $L \in \mathbb{N}$ where $n/2M \leq L \leq n/\ell_0$;
- (iv) For all $1 \leq i \leq t$, $1 \leq j \leq k$, the pair $(W_{0,j}, W_{i,j})_{G'}$ is $(4\varepsilon, d/2)$ -super-regular;
- (v) $\delta(G') \geq (\alpha - 2d)n$.

Proof. Apply the Regularity lemma (Lemma 3.1) to G with parameters ε, d and ℓ_0 to obtain a partition V_0, V_1, \dots, V_ℓ of $V(G)$, pure graph G' of G and the reduced graph R of G . So $|V_0| \leq \varepsilon n$ and $(1 - \varepsilon)n/\ell \leq L' := |V_i| = |V_j| \leq n/\ell$ for all $i, j \geq 1$. It is a well-known fact that the reduced graph R of G ‘almost’ inherits the minimum degree of G (see e.g. [30, Lemma 3.7]). In particular, since $\delta(G) \geq \alpha n$ and $\varepsilon \ll d \ll \alpha$ we have that

$$\delta(R) \geq \frac{\alpha\ell}{2} \geq \frac{\ell}{t+1}.$$

By (5.1), $\ell \geq \ell_0$ is sufficiently large compared to $1/\varepsilon$ and t . Thus, Theorem 4.1 implies that R contains a $K_{1,t}$ -tiling \mathcal{K} that covers all but at most $\varepsilon\ell$ vertices of R . Let k denote the size of \mathcal{K} . Hence,

$$\frac{\ell_0}{2t} \leq \frac{(1 - \varepsilon)\ell}{t+1} \leq k \leq \frac{\ell}{t+1} \leq \frac{M}{t+1}.$$

Move to V_0 all those vertices that lie in clusters uncovered by \mathcal{K} . Hence

$$|V_0| \leq \varepsilon n + \varepsilon\ell \times L' \leq \varepsilon n + \varepsilon\ell \times \frac{n}{\ell} = 2\varepsilon n.$$

Consider a copy $K'_{1,t}$ of $K_{1,t}$ in \mathcal{K} . In G' , this copy of $K_{1,t}$ corresponds to a collection of $t+1$ clusters $V_{i_0}, V_{i_1}, \dots, V_{i_t}$ such that for all $1 \leq j \leq t$ the pair $(V_{i_0}, V_{i_j})_{G'}$ is ε -regular and has density greater than d . By repeatedly applying Lemma 3.2 it is easy to check that we can remove precisely $\varepsilon t|V_{i_j}| = \varepsilon tL'$ vertices from V_{i_j} (for each $0 \leq j \leq t$) so that now

$(V_{i_0}, V_{i_j})_{G'}$ is $(4\varepsilon, d/2)$ -super-regular for each $1 \leq j \leq t$. Add the vertices removed from these clusters into V_0 . Repeat this process for all copies of $K_{1,t}$ in \mathcal{K} . Thus, now

$$|V_0| \leq 2\varepsilon n + \varepsilon t L' \ell \leq 2t\varepsilon n.$$

Given the j th copy of $K_{1,t}$ in \mathcal{K} we relabel the clusters so that the root of this $K_{1,t}$ is $W_{0,j}$ and the leaves are $W_{1,j}, \dots, W_{t,j}$. Hence (iv) holds. Further, each of the clusters $W_{i,j}$ has the same size $L \in \mathbb{N}$ where

$$\frac{n}{2M} \leq \frac{(1-\varepsilon)(1-\varepsilon t)n}{\ell} \leq L = (1-\varepsilon t)L' \leq \frac{n}{\ell} \leq \frac{n}{\ell_0},$$

so (iii) holds. Note that (v) holds by Lemma 3.1(ii) and as $\varepsilon \ll d$. \square

So far we have only used the deterministic edges, i.e. edges in G , but recall that we are aiming to find a perfect H -tiling in $G \cup G_{n,p}$. Next we will use these random edges to find copies of $H' := H - x$ (for some $x \in V(H)$) in the neighbourhood of vertices in V_0 . To simplify the later calculations we now use the standard trick of decomposing the random edges into a few ‘buckets’. That is, let G_1, G_2, G_3, G_4 be independently chosen elements of $G_{n,p/4}$ and observe that $G_1 \cup G_2 \cup G_3 \cup G_4$ has the same distribution as $G_{n,p'}$ for some $p' \leq p$. Hence it suffices to consider the graph $G \cup (G_1 \cup G_2 \cup G_3 \cup G_4)$ instead of the graph $G \cup G_{n,p}$. Define $\mathcal{W} := \{W_{i,j} : 0 \leq i \leq t \text{ and } 1 \leq j \leq k\}$ and $k' := |\mathcal{W}| = k(t+1)$.

Claim 5.2. *Asymptotically almost surely there is a set $Z \subseteq V(G)$ such that:*

(i) $(G' \cup G_1)[Z]$ contains a perfect H -tiling \mathcal{H}_1 ;

(ii) $V_0 \subseteq Z$, and for all $W_{i,j} \in \mathcal{W}$ we have that $|Z \cap W_{i,j}| \leq \varepsilon_1 L$.

Proof. Define an auxiliary bipartite graph Q with vertex classes V_0 and \mathcal{W} in which a vertex $v \in V_0$ is adjacent to a cluster $W_{i,j}$ precisely if $d_{G'}(v, W_{i,j}) \geq \alpha L/4$. For each $v \in V_0$, $d_{G'}(v) \geq (\alpha - 2d)n$ and is adjacent to at most $2t\varepsilon n$ vertices in V_0 . Therefore,

$$3\alpha n/4 \stackrel{(5.1)}{\leq} d_{G'}(v) - 2t\varepsilon n \leq Ld_Q(v) + k'\alpha L/4 \leq Ld_Q(v) + \alpha n/4,$$

and so $d_Q(v) \geq \alpha k'/2$.

Hence we can find an assignment $f : V_0 \rightarrow \mathcal{W}$ such that $vf(v)$ is an edge in Q for all $v \in V_0$ and for any $W_{i,j} \in \mathcal{W}$,

$$|f^{-1}(W_{i,j})| \leq \frac{4t\varepsilon n}{\alpha k'} \stackrel{(5.1)}{\leq} \frac{\varepsilon_1 L}{|H|}. \quad (5.2)$$

Here in the first inequality we use that $|V_0| \leq 2t\varepsilon n$ and $d_Q(v) \geq \alpha k'/2$ for all $v \in V_0$; in the last inequality we use that $k'L = n - |V_0| \geq (1 - 2t\varepsilon)n$.

Enumerate the vertices v_1, \dots, v_s of V_0 . For each such v_i , we will obtain a copy H_i of H in $G \cup G_1$ so that:

- H_i contains v_i for each $1 \leq i \leq s$, and all the other vertices in H_i lie in the cluster $f(v_i) \in \mathcal{W}$;
- H_i and H_j are vertex-disjoint for all $1 \leq i \neq j \leq s$.

Note that finding such copies of H would immediately prove the claim. Indeed, we then define Z to consist of all the vertices in the H_i . In particular, (5.2) then implies that $|Z \cap W_{i,j}| \leq \frac{\varepsilon_1 L}{|H|} \times (|H| - 1) \leq \varepsilon_1 L$ for each $W_{i,j} \in \mathcal{W}$.

Suppose for some $1 \leq j < s$ we have constructed H_1, \dots, H_{s-1} as desired. Consider $v_s \in V_0$ and let $W_{i_s, j_s} := f(v_s)$. Then by the definition of Q and f , there is a set $W \subseteq W_{i_s, j_s}$ so that $|W| \geq \alpha L/4 - \varepsilon_1 L \geq \alpha L/5$; W is disjoint from H_1, \dots, H_{s-1} and; $W \subseteq N_{G'}(v_s)$.

Note that $H' := H - x$ (for some $x \in V(H)$) either consists of isolated vertices or $0 < d^*(H') \leq d^*(H)$. Further $\eta n \leq \alpha n/(10M) \leq \alpha L/5$ by (5.1) and Claim 5.1(iii). So Theorem 4.2 implies that a.a.s, $G_1[W]$ contains a copy of H' . Since in G' , v_s is adjacent to every vertex in W , this yields the desired copy H_s of H in $G \cup G_1$. \square

For all i, j set

$$V_{i,j} := W_{i,j} \setminus Z.$$

Since $(G' \cup G_1)[Z]$ contains a perfect H -tiling \mathcal{H}_1 , it suffices to prove that, a.a.s, there is an H -tiling in $G' \cup G_2 \cup G_3 \cup G_4$ that covers precisely the vertices in $V(G) \setminus Z$. Note that by Claim 5.1(iv) and Claim 5.2(ii) we have that $(V_{0,j}, V_{i,j})_{G'}$ is $(5\varepsilon, d/3)$ -super-regular for all $1 \leq i \leq t$ and $1 \leq j \leq k$. Further,

$$(1 - \varepsilon_1)L \leq |V_{i,j}| \leq L \tag{5.3}$$

for all $0 \leq i \leq t$ and $1 \leq j \leq k$.

For each $1 \leq j \leq k$, randomly partition $V_{0,j}$ into t vertex classes $S_{1,j}, \dots, S_{t,j}$ such that

$$|S_{i,j}| = \frac{|V_{0,j}|}{t} \tag{5.4}$$

for all $1 \leq i \leq t$. For every $1 \leq i \leq t$ and $1 \leq j \leq k$, randomly partition $V_{i,j}$ into two vertex classes $V'_{i,j}$ and $T'_{i,j}$ so that

$$|T'_{i,j}| = \frac{|V_{i,j}|}{t}. \tag{5.5}$$

Roughly speaking, the rest of the proof now proceeds as follows: For each of the clusters $V'_{i,j}$ we will find an almost perfect H -tiling in $G_2[V'_{i,j}]$. This will ensure that almost all of the uncovered vertices in $V(G)$ lie in the clusters $S_{i,j}$ and $T'_{i,j}$ (for all i, j). However, as the next claim shows, each $(S_{i,j}, T'_{i,j})_{G'}$ is ‘super-regular’. Then by modifying each of the super-regular pairs slightly, this structure will allow us to cover all remaining vertices in $V(G)$ with an H -tiling, thereby completing our perfect H -tiling.

Claim 5.3. *The following conditions hold a.a.s:*

(i) Let $1 \leq i \leq t$ and $1 \leq j \leq k$. For all sets $X \subseteq S_{i,j}$ and $Y \subseteq T'_{i,j}$ with $|X| \geq \varepsilon_1 |S_{i,j}|$ and $|Y| \geq \varepsilon_1 |T'_{i,j}|$, we have that $d_{G'}(X, Y) \geq d/3$;

(ii) Let $1 \leq j \leq k$. For every vertex $v \in V_{0,j}$, $d_{G'}(v, T'_{i,j}) > d|T'_{i,j}|/4$ for all $1 \leq i \leq t$;

(iii) Let $1 \leq i \leq t$ and $1 \leq j \leq k$. For every $v \in V_{i,j}$, $d_{G'}(v, S_{i,j}) > d|S_{i,j}|/4$.

In particular, (i)–(iii) imply that $(S_{i,j}, T'_{i,j})_{G'}$ is $(\varepsilon_1, d/4)$ -super-regular for all $1 \leq i \leq t$ and $1 \leq j \leq k$.

Proof. For (i), note that $X \subseteq V_{0,j}$ and $Y \subseteq V_{i,j}$ where $|X| \geq \varepsilon_1 |S_{i,j}| \stackrel{(5.4)}{=} \varepsilon_1 |V_{0,j}|/t \stackrel{(5.1)}{\geq} 5\varepsilon |V_{0,j}|$ and $|Y| \geq 5\varepsilon |V_{i,j}|$. So as $(V_{0,j}, V_{i,j})_{G'}$ is $(5\varepsilon, d/3)$ -super-regular, (i) follows immediately.

For (ii) note that such a $v \in V_{0,j}$ satisfies $d_{G'}(v, V_{i,j}) \geq d|V_{i,j}|/3$ for all $1 \leq i \leq t$. Thus by applying a Chernoff bound for the hypergeometric distribution (e.g. [15, Theorem 2.10]), a.a.s. (ii) holds. (iii) holds similarly. \square

We will now obtain an H -tiling \mathcal{H}_2 in G_2 that covers almost all of the vertices in the classes $V'_{i,j}$. Fix $1 \leq i \leq t$ and $1 \leq j \leq k$. By repeatedly applying Theorem 4.2 to G_2 , we obtain an H -tiling $\mathcal{H}_{i,j}$ in $G_2[V'_{i,j}]$ that covers all but at most ηn of the vertices in $V'_{i,j}$. Let \mathcal{H}_2 denote the H -tiling in G_2 obtained by taking the disjoint union of all of the $\mathcal{H}_{i,j}$, and write $Z' := V(\mathcal{H}_2)$.

For each $1 \leq i \leq t$ and $1 \leq j \leq k$, let $T_{i,j}$ be obtained from $T'_{i,j}$ by adding to it all the vertices in $V'_{i,j}$ that are uncovered by $\mathcal{H}_{i,j}$. Thus, $|T_{i,j}| \leq |T'_{i,j}| + \eta n$. Note that $|T'_{i,j}| \stackrel{(5.5)}{=} |V_{i,j}|/t \stackrel{(5.3)}{\geq} (1 - \varepsilon_1)L/t$. On the other hand, $\eta n \leq \varepsilon(1 - \varepsilon_1)L/t$ by (5.1) and Claim 5.1(iii). Hence, $|T_{i,j}| \leq (1 + \varepsilon)|T'_{i,j}| \stackrel{(5.3), (5.5)}{\leq} (1 + \varepsilon)L/t$. Together with Claim 5.3(i)–(iii) this implies that $(S_{i,j}, T_{i,j})_{G'}$ is $(2\varepsilon_1, d/5)$ -super-regular. To summarise, so far we have proved the following.

Claim 5.4. *Asymptotically almost surely, there is a partition of $V(G)$ into classes Z , Z' and $S_{i,j}$, $T_{i,j}$ for each $1 \leq i \leq t$ and $1 \leq j \leq k$, such that:*

(i) *There is a perfect H -tiling \mathcal{H}_1 in $(G' \cup G_1)[Z]$ and a perfect H -tiling \mathcal{H}_2 in $G_2[Z']$;*

(ii) *Each $(S_{i,j}, T_{i,j})_{G'}$ is $(2\varepsilon_1, d/5)$ -super-regular;*

(iii) $(1 - \varepsilon_1)L/t \leq |S_{i,j}|, |T_{i,j}| \leq (1 + \varepsilon)L/t$. \square

To finish the proof of Theorem 1.3 we wish to prove that a.a.s. there is a perfect H -tiling in each of the graphs $(G' \cup G_4)[S_{i,j} \cup T_{i,j}]$. However, for this we need that $|H|$ divides $|S_{i,j} \cup T_{i,j}|$. So we first modify the clusters $S_{i,j}, T_{i,j}$ slightly to ensure this.

Suppose that $1 \leq i \leq t$ and $1 \leq j \leq k$ are such that $|S_{i,j} \cup T_{i,j}|$ is not divisible by $|H|$. Then, since $|H|$ divides n , there are $1 \leq i' \leq t$ and $1 \leq j' \leq k$ with $(i, j) \neq (i', j')$ so that $|S_{i',j'} \cup T_{i',j'}|$ is also not divisible by $|H|$. Recall $|S_{i,j}|, |S_{i',j'}| \gg \eta n$. Thus, by repeated applications of Theorem 4.4, a.a.s. we can find a collection $\mathcal{S}_{i,j}$ of at most $|H| - 1$ disjoint

copies of H in G_3 so that: each such H has one vertex in $S_{i,j}$ and $|H| - 1$ vertices in $S_{i',j'}$ and; after removal of the vertices in $S_{i,j}$ we now have that $|S_{i,j} \cup T_{i,j}|$ is divisible by $|H|$. Continuing in this way, we obtain an H -tiling \mathcal{H}_3 in G_3 so that: \mathcal{H}_3 only covers vertices in the clusters $S_{i,j}$; $|\mathcal{H}_3| \leq |H|kt$ and; after removal of all those vertices in \mathcal{H}_3 from each of the $S_{i,j}$ we have that $|S_{i,j} \cup T_{i,j}|$ is divisible by $|H|$ for all $1 \leq i \leq t$ and $1 \leq j \leq k$.

In particular, we now have the following updated version of Claim 5.4.

Claim 5.5. *Asymptotically almost surely, there is a partition of $V(G)$ into classes Z , Z' , Z'' and $S_{i,j}$, $T_{i,j}$ for each $1 \leq i \leq t$ and $1 \leq j \leq k$, such that:*

- (i) *There is a perfect H -tiling \mathcal{H}_1 in $(G' \cup G_1)[Z]$; a perfect H -tiling \mathcal{H}_2 in $G_2[Z']$ and; a perfect H -tiling \mathcal{H}_3 in $G_3[Z'']$;*
- (ii) *Each $(S_{i,j}, T_{i,j})_{G'}$ is $(3\varepsilon_1, d/6)$ -super-regular;*
- (iii) $(1 - 2\varepsilon_1)L/t \leq |S_{i,j}|, |T_{i,j}| \leq (1 + \varepsilon)L/t$;
- (iv) $|S_{i,j} \cup T_{i,j}|$ is divisible by $|H|$ for all $1 \leq i \leq t$ and $1 \leq j \leq k$. □

Now all that remains to prove is that there is a perfect H -tiling in each of the graphs $(G' \cup G_4)[S_{i,j} \cup T_{i,j}]$. Let $(S, T)_{G'}$ be one of our super-regular pairs and recall that H' is an induced subgraph of H on $h := |H| - 1$ vertices.

Call an h -set $\{v_1, \dots, v_h\} \in \binom{T}{h}$ *good* if $|S \cap N_{G'}(v_1) \cap \dots \cap N_{G'}(v_h)| \geq d_1|S|$, otherwise we say $\{v_1, \dots, v_h\} \in \binom{T}{h}$ is *bad*.

Claim 5.6. *At least $(1 - 4\varepsilon_1)^{h-1} \binom{|T|}{h}$ h -sets in $\binom{T}{h}$ are good.*

Proof. Pick any vertex $v_1 \in T$ and let $N_1 := N_{G'}(v_1) \cap S$. Since $(S, T)_{G'}$ is $(3\varepsilon_1, d/6)$ -super-regular, we have that $|N_1| \geq d|S|/6$ and every set $Y \subseteq T$ of size at least $3\varepsilon_1|T|$ contains at least one vertex v_2 with $|N_{G'}(v_2) \cap N_1| \geq d|N_1|/6$. So in particular at least $(1 - 4\varepsilon_1)|T|$ vertices v_2 in T are such that $|N_2| \geq d|N_1|/6 \geq d^2|S|/36$ where $N_2 := N_{G'}(v_2) \cap N_1$. Pick such a vertex v_2 . Similarly, there are at least $(1 - 4\varepsilon_1)|T|$ vertices v_3 in T such that $|N_{G'}(v_3) \cap N_2| \geq d|N_2|/6 \geq d^3|S|/216$. Continuing in this way we conclude that there are at least

$$|T| \times (1 - 4\varepsilon_1)^{h-1} |T|^{h-1} \times \frac{1}{h!} \geq (1 - 4\varepsilon_1)^{h-1} \binom{|T|}{h}$$

h -sets $\{v_1, \dots, v_h\}$ in $\binom{T}{h}$ so that $|S \cap N_{G'}(v_1) \cap \dots \cap N_{G'}(v_h)| \geq d^h|S|/6^h \stackrel{(5.1)}{\geq} d_1|S|$, as required. □

Call a good h -set $\{v_1, \dots, v_h\} \in \binom{T}{h}$ *excellent* if in G_4 they span a copy of H' . We now claim that a.a.s. every subset of T of size at least $\varepsilon_3|T|$ contains an excellent h -set. If H' consists of isolated vertices then every good set is excellent. So we may assume that H' contains an edge. Recall that $G_4 \cong G_{n,p/4}$ and note that $0 < d^*(H') \leq d^*(H)$. So

$$\frac{p}{4} \geq \frac{c}{4} n^{-1/d^*(H)} \geq D \left(\frac{n}{4Mt} \right)^{-1/d^*(H)} \geq D|T|^{-1/d^*(H')}.$$

The latter inequality holds since $|T| \geq \frac{n}{4Mt}$ by Claim 5.5(iii) and Claim 5.1(iii). Let $K'_{|T|}$ be a copy of $K_{|T|}$ on vertex set T . By Claim 5.6 there are at most $\varepsilon_2 \binom{|T|}{h}$ copies of H' in $K'_{|T|}$ whose vertex set is a bad h -set. Thus, applying Theorem 4.3 to $G_4[T] \cong G_{|T|,p/4}$, implies that a.a.s. every subset of T of size at least $\varepsilon_3|T|$ contains an excellent h -set. We may similarly define a good, bad and excellent h -set in $\binom{S}{h}$ and conclude a.a.s. every subset of S of size at least $\varepsilon_3|S|$ contains an excellent h -set.

Let $G^* := (G' \cup G_4)[S, T]$. We say that a copy H^* of H in G^* is an S -copy if H^* contains precisely one vertex x from T and $H^* - x$ is a copy of H' . Similarly define a T -copy of H to be a copy of H in G^* intersecting T in a copy of H' and S in one vertex. Set $N := L/t$; so $(1 - 2\varepsilon_1)N \leq |S|, |T| \leq (1 + \varepsilon)N$ by Claim 5.5(iii). The following two claims finish our proof of Theorem 1.3 - the first deals with the case where H is a single edge and the second deals with general H . The latter claim is an analogue of Lemma 3.1 in [2], and indeed the proof is very similar; we include it for completeness.

Claim 5.7. *If $H = K_2$ is a single edge then a.a.s. there exists a perfect H -tiling in G^* .*

Proof. Without loss of generality we may assume $|S| \leq |T|$. Let $z := |T| - |S|$ and note that by Claim 5.5(iv) we have that z is even. By Theorem 4.2 we know that in G_4 a.a.s. every set $Q \subset T$ of size $|Q| = \eta n < |T|/2$ contains an edge. Hence we may greedily form a matching M in $G_4[T]$ consisting of $z/2$ edges. Let $T' := T \setminus V(M)$ so that $|T'| = |S|$.

Since $|V(M)| \leq 3\varepsilon_1|T|$ and $G'[S, T]$ is $(3\varepsilon_1, d/6)$ -super-regular, we have that $G'[S, T']$ is $(\varepsilon_2, d/8)$ -super-regular. As $\varepsilon_2 \ll d$, it is easy to see that Hall's condition is satisfied and thus $G'[S, T']$ contains a perfect matching M' . Then $M \cup M'$ forms the desired perfect H -tiling in G^* . \square

Claim 5.8. *If $|H| \geq 3$ the following statements hold a.a.s:*

- (a) *Provided $|S \setminus Q| + |T| + \lfloor \phi\varepsilon_5 N \rfloor$ is divisible by $|H|$, for every $Q \subseteq S$ of size $|Q| = \phi N$ there is an H -tiling in G^* which covers every vertex of $(S \cup T) \setminus Q$ and which covers precisely $\lfloor \phi\varepsilon_5 N \rfloor$ vertices of Q . Moreover, the same assertion holds if one replaces T with any subset $T' \subseteq T$ where $|T \setminus T'| \leq \varepsilon_5 N$.*
- (b) *G^* contains a perfect H -tiling.*

Proof. For (a), let $z := \lfloor \phi\varepsilon_5 N \rfloor$, $t := \lfloor z/h \rfloor$ and $z' := z - ht \in \{0, 1, \dots, h-1\}$. We will construct an H -tiling in G^* that covers all of $(S \setminus Q) \cup T$ and precisely $z = ht + z'$ vertices of Q .

Let $T'_1 \subseteq T$ consist of all vertices in T with fewer than $d_1|Q|$ neighbours in Q . Note $G'[S, T]$ is (ε_2, d_1) -super-regular so we have that $|T'_1| \leq 2\varepsilon_2 N$. Form T_1 by adding at most h arbitrarily selected vertices from $T \setminus T'_1$ to T'_1 so that $|T \setminus T_1| - t$ is divisible by $|H|$. Since $G'[S, T]$ is (ε_2, d_1) -super-regular, every vertex of T_1 has at least $d_1|S| - |Q| \geq \frac{d_1 N}{2} > h|T_1| + 2\varepsilon_3 N$ neighbours in $S \setminus Q$. As every subset of S of size $2\varepsilon_3 N$ contains an excellent h -set, we may greedily form an H -tiling \mathcal{T}_1 of S -copies in G^* of size $|T_1|$ which covers every vertex of T_1 and does not use any vertex from Q .

We now select uniformly at random a subset $T_2 \subseteq T \setminus T_1$ of size $|T_2| = t$. Since every vertex in S has at least $d_1|T| - |T_1| \geq \frac{d_1 N}{2}$ neighbours in $T \setminus T_1$, Chernoff's inequality for the hypergeometric distribution implies that, with probability $1 - o(1)$, every vertex of S has at least $\frac{\phi \varepsilon_5 d_1}{5h} N$ neighbours in T_2 . Fix a choice of T_2 for which this event occurs. Let Q' be an arbitrarily selected subset of Q of size z' , so that $|Q'| \leq h-1$ and let $S' := (S \setminus (Q \cup V(\mathcal{T}_1))) \cup Q'$ and $T' := T \setminus (T_1 \cup T_2)$. Recall that, by assumption, $|S \setminus Q| + |T| + z$ is divisible by $|H|$, so

$$|S'| + |T'| = |S \setminus Q| + z' + |T| - |T_2| - |V(\mathcal{T}_1)| = (|S \setminus Q| + |T| + z) - (|H|t + |V(\mathcal{T}_1)|)$$

is divisible by $|H|$. Since $|T'|$ is divisible by $|H|$ by our selection of T_1 and T_2 , it follows that $|S'|$ is divisible by $|H|$ as well.

Let $t_2 := \lfloor \frac{\phi \varepsilon_5 d_1}{10h^2} N \rfloor$, $a := \frac{h}{(h-1)|H|} |S'| - \frac{1}{(h-1)|H|} |T'|$ and $b := \frac{h}{(h-1)|H|} |T'| - \frac{1}{(h-1)|H|} |S'| - t_2$. Note that since $|H| \geq 3$ both a and b are positive.

Subclaim 5.9. *There is an H -tiling \mathcal{T}_2 in $G^*[S' \cup T']$ which consists of a S -copies and b T -copies. In particular, $S'' := S' \setminus V(\mathcal{T}_2)$ and $T'' := T' \setminus V(\mathcal{T}_2)$ have sizes precisely $|S''| = |S'| - (ha + b) = t_2$ and $|T''| = |T'| - (a + hb) = ht_2$.*

Proof. Note $G'[S', T']$ is $(2\varepsilon_2, d_1/2)$ -super-regular. The copies of H in \mathcal{T}_2 may be chosen greedily. Indeed, suppose that we have already chosen an H -tiling \mathcal{T} in $G^*[S', T']$ consisting of at most a S -copies and b T -copies, then \mathcal{T} covers at most $ha + b$ vertices of S , and at most $a + hb$ vertices of T . Taking $S^* := S' \setminus V(\mathcal{T})$ and $T^* := T' \setminus V(\mathcal{T})$, we find that $|S^*| \geq t_2 \geq \varepsilon_4 N$ and $|T^*| \geq ht_2 \geq \varepsilon_4 N$. Since $G'[S', T']$ is $(2\varepsilon_2, d_1/2)$ -super-regular, it follows that there is some vertex $x \in S^*$ having at least $d_1 \varepsilon_4 N / 2 > 2\varepsilon_3 N$ neighbours in T^* . Since every set of size at least $2\varepsilon_3 N$ in T^* contains an excellent h -set this gives a T -copy which can be added to \mathcal{T} . The same argument with the roles of S^* and T^* reversed yields instead an S -copy which may be added to \mathcal{T} . This proves the claim. \square

Since by the choice of T_2 each vertex of S'' has at least $\frac{\phi \varepsilon_5 d_1}{5h} N > h|S''| + 2\varepsilon_3 N$ neighbours in T_2 , we may greedily form an H -tiling \mathcal{T}_3 in $G^*[S'' \cup T_2]$ consisting of t_2 T -copies which covers every vertex of S'' and which covers precisely ht_2 vertices of T_2 . At this point we have obtained an H -tiling $\mathcal{T}_1 \cup \mathcal{T}_2 \cup \mathcal{T}_3$ in G^* which covers every vertex of S except for those in $Q \setminus Q'$ and every vertex of T except for the precisely ht_2 vertices in T'' and the precisely $t - ht_2$ vertices in $T_2 \setminus V(\mathcal{T}_3)$. Therefore, in total, precisely t vertices of T remain uncovered, each of which has at least $d_1|Q| - |Q'| > ht + 2\varepsilon_3 N$ neighbours in $Q \setminus Q'$ by the choice of T_1 . We may therefore greedily form an H -tiling \mathcal{T}_4 of S -copies in G^* which covers all the remaining uncovered vertices in T and precisely ht vertices of $Q \setminus Q'$. Then $\mathcal{T}_1 \cup \mathcal{T}_2 \cup \mathcal{T}_3 \cup \mathcal{T}_4$ is the desired H -tiling.

The moreover part of (a) follows by observing that since $G'[S, T]$ is $(3\varepsilon_1, d/6)$ -super-regular, $G'[S, T']$ is (ε_2, d_1) -super-regular for any subset $T' \subseteq T$ where $|T \setminus T'| \leq \varepsilon_5 N$. Thus, the precise argument above, with T' playing the role of T , implies we have the desired H -tiling.

For (b), we may assume without loss of generality that $|T| \geq |S|$. Since every set of at least $2\varepsilon_3 N$ vertices of T contains an excellent h -set we may greedily form an H' -tiling M of

size at least $(|T| - 2\varepsilon_3 N)/h \geq N/(h+1)$ in $G_4[T]$ such that the vertex set of each of these copies of H' are excellent. Fix such an H' -tiling M , and form an auxiliary bipartite graph K with vertex classes S and M where $a \in S$ and $e = (x_1 x_2 \dots x_h) \in M$ are adjacent if and only if $a \in N_{G^*}(x_1) \cap \dots \cap N_{G^*}(x_h)$. Note that for every H' -copy $e = (x_1 x_2 \dots x_h) \in M$ we have that

$$d_K(e) = |N_{G^*}(x_1) \cap \dots \cap N_{G^*}(x_h) \cap S| \geq d_1 |S|$$

by the definition of an excellent h -set, so K has density at least d_1 . We now apply the following lemma from [2].

Lemma 5.10 ([2], Lemma 2.6). *Suppose that $1/n \ll \phi' \ll \varepsilon' \ll d' \ll 1/h'$. Let F be a bipartite graph with vertex classes A and B such that $n/h' \leq |A|, |B| \leq n$ and $d_F(A, B) \geq d'$. Then there exist subsets $X \subseteq A$ and $Y \subseteq B$ of sizes $|X| = \phi' n$ and $|Y| = (1 - \varepsilon')\phi' n$ such that $F[X', Y]$ contains a perfect matching for every subset $X' \subseteq X$ with $|X'| = |Y|$.*

We remark that this lemma is not precisely as stated in [2], but this version can easily be deduced from the original.

Hence we may choose subsets $X \subseteq S$ and $M' \subseteq M$ such that $|X| = \phi N$, $|M'| = (1 - \varepsilon_5)\phi N$ and such that $K[X', M']$ contains a perfect matching for every subset $X' \subseteq X$ with $|X'| = |M'|$. Let $T' := T \setminus V(M')$. Then, since $\phi \ll \varepsilon_5$ we may apply (a) to $G^*[S \cup T']$ with X playing the role of Q to obtain an H -tiling \mathcal{T}_1 in G^* which covers every vertex of G^* except for the vertices of $V(M')$ and precisely $(1 - \varepsilon_5)\phi N$ vertices of X . So, taking X' to be the vertices of X not covered by \mathcal{T}_1 , we have $|X'| = |M'|$. By the choice of X and M' it follows that $K[X', M']$ contains a perfect matching, which corresponds to a perfect H -tiling \mathcal{T}_2 in $G^*[X' \cup V(M')]$. This gives a perfect H -tiling $\mathcal{T}_1 \cup \mathcal{T}_2$ in G^* . □

Acknowledgment

The authors are grateful to the BRIDGE strategic alliance between the University of Birmingham and the University of Illinois at Urbana-Champaign. This research was conducted as part of the ‘Building Bridges in Mathematics’ BRIDGE Seed Fund project.

The authors are also grateful to the referee for their careful and helpful review.

References

- [1] N. Alon and R. Yuster, Threshold functions for H -factors, *Combin. Probab. Comput.* **2** (1993), 137–144.
- [2] J. Balogh, A. McDowell, T. Molla and R. Mycroft, Triangle-tilings in graphs without large independent sets, *Combin. Probab. Comput.*, to appear.
- [3] W. Bedenknecht, J. Han, Y. Kohayakawa and G. O. Mota, Powers of tight Hamilton cycles in randomly perturbed hypergraphs, submitted.

- [4] P. Bennett, A. Dudek and A. Frieze, Adding random edges to create the square of a Hamilton cycle, submitted.
- [5] T. Bohman, A. Frieze, M. Krivelevich and R. Martin, Adding random edges to dense graphs, *Random Structures & Algorithms* **24** (2004), 105–117.
- [6] T. Bohman, A. Frieze and R. Martin, How many edges make a dense graph Hamiltonian?, *Random Structures & Algorithms* **22** (2003), 33–42.
- [7] J. Böttcher, J. Han, Y. Kohayakawa, R. Montgomery, O. Parczyk and Y. Person, Universality for bounded degree spanning trees in randomly perturbed graphs, submitted.
- [8] J. Böttcher, R. Montgomery, O. Parczyk and Y. Person, Embedding spanning bounded degree graphs in randomly perturbed graphs, submitted.
- [9] J. Böttcher, M. Schacht and A. Taraz, Proof of the bandwidth conjecture of Bollobás and Komlós, *Math. Ann.* **343** (2009), 175–205.
- [10] G.A. Dirac, Some theorems on abstract graphs, *Proc. London Math. Soc.* **2** (1952), 69–81.
- [11] P. Erdős and A. Rényi, On the existence of a factor of degree one of a connected random graph, *Acta Math. Acad. Sci. Hungar.* **17** (1966), 359–368.
- [12] S. Gerke and A. McDowell, Nonvertex-balanced factors in random graphs, *J. Graph Theory* **78** (2015), 269–286.
- [13] A. Hajnal and E. Szemerédi, Proof of a conjecture of Erdős, *Combinatorial Theory and its Applications vol. II* **4** (1970), 601–623.
- [14] J. Han and Y. Zhao, Hamiltonicity in randomly perturbed hypergraphs, submitted.
- [15] S. Janson, T. Łuczak and A. Ruciński, *Random Graphs*, Wiley, 2000.
- [16] A. Johansson, J. Kahn and V. Vu, Factors in random graphs, *Random Structures & Algorithms* **33** (2008), 1–28.
- [17] F. Joos and J. Kim, Spanning trees in randomly perturbed graphs, submitted.
- [18] J. Komlós, Tiling Turán Theorems, *Combinatorica* **20** (2000), 203–218.
- [19] J. Komlós, G. Sárközy and E. Szemerédi, Proof of the Alon–Yuster conjecture, *Discrete Math.* **235** (2001), 255–269.
- [20] M. Krivelevich, M. Kwan and B. Sudakov, Cycles and matchings in randomly perturbed digraphs and hypergraphs, *Combin. Probab. Comput.* **25** (2016), 909–927.
- [21] M. Krivelevich, M. Kwan and B. Sudakov, Bounded-degree spanning trees in randomly perturbed graphs, *SIAM J. Discrete Math.* **31** (2017), 155–171.

- [22] M. Krivelevich, B. Sudakov and P. Tetali, On smoothed analysis in dense graphs and formulas, *Random Structures & Algorithms* **29** (2006), 180–193.
- [23] D. Kühn and D. Osthus, Critical chromatic number and the complexity of perfect packings in graphs, *17th ACM-SIAM Symposium on Discrete Algorithms* (SODA 2006), 851–859.
- [24] D. Kühn and D. Osthus, The minimum degree threshold for perfect graph packings, *Combinatorica* **29** (2009), 65–107.
- [25] T. Łuczak and A. Ruciński, Tree-matchings in graph processes, *SIAM J. Discrete Math.* **4** (1991), 107–120.
- [26] A. McDowell and R. Mycroft, Hamilton ℓ -cycles in randomly perturbed hypergraphs, submitted.
- [27] L. Pósa, Hamiltonian circuits in random graphs, *Discrete Math.* **14** (1976), 359–364.
- [28] A. Ruciński, Matching and covering the vertices of a random graph by copies of a given graph, *Discrete Math.* **105** (1992), 185–197.
- [29] E. Szemerédi, Regular partitions of graphs, *Problèmes Combinatoires et Théorie des Graphes Colloques Internationaux CNRS 260* (1978), 399–401.
- [30] A. Treglown, The regularity lemma and applications to packings in graphs, MSci thesis, University of Birmingham, 2007.