UNIVERSITY^{OF} BIRMINGHAM University of Birmingham Research at Birmingham

Neutral pion form factor measurement by the NA62 experiment

NA62 Collaboration

DOI: 10.1088/1742-6596/873/1/012016

License: Creative Commons: Attribution (CC BY)

Document Version Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

NA62 Collaboration 2017, 'Neutral pion form factor measurement by the NA62 experiment', *Journal of Physics: Conference Series*, vol. 873, 012016. https://doi.org/10.1088/1742-6596/873/1/012016

Link to publication on Research at Birmingham portal

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

•Users may freely distribute the URL that is used to identify this publication.

•Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.

•User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?) •Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

PAPER • OPEN ACCESS

Neutral pion form factor measurement by the NA62 experiment

To cite this article: Michal Zamkovsky et al 2017 J. Phys.: Conf. Ser. 873 012016

View the article online for updates and enhancements.

Related content

- <u>The CHarged particle ANTIcounter for the</u> <u>NA62 experiment at CERN</u> Marco Mirra
- <u>New limits on Heavy Neutrino from NA62</u> Sergey Kholodenko and NA62 Collaboration
- <u>The NA62 experiment at CERN</u> Francesco Gonnella and NA62 collaboration

Neutral pion form factor measurement by the NA62 experiment

Michal Zamkovsky

Speaker, Institute of Particle and Nuclear Physics, Charles University E-mail: michal.zamkovsky@cern.ch

F Ambrosino, A Antonelli, G Anzivino, R Arcidiacono, W Baldini, S Balev, J R Batley, M Behler, S Bifani, C Biino, A Bizzeti, T Blazek, B Bloch-Devaux, G Bocquet, V Bolotov, F Bucci, N Cabibbo, M Calvetti, N Cartiglia, A Ceccucci, P Cenci, C Cerri, C Cheshkov, J B Chze, M Clemencic, G Collazuol, F Costantini, A Cotta Ramusino, D Coward, D Cundy, A Dabrowski, G DAgostini, P Dalpiaz, C Damiani, H Danielsson, M De Beer, G Dellacasa, J Derr, H Dibon, D Di Filippo, L DiLella, N Doble, V Duk, J Engelfried, K Eppard, V Falaleev, R Fantechi, M Fidecaro, L Fiorini, M Fiorini, T Fonseca Martin, P L Frabetti, A Fucci, S Gallorini, L Gatignon, E Gersabeck, A Gianoli, S Giudici, A Gonidec, E Goudzovski, S Goy Lopez, E Gushchin, B Hallgren, M Hita-Hochgesand, M Holder, P Hristov, E Iacopini, E Imbergamo, M Jeitler, G Kalmus, V Kekelidze, K Kleinknecht, M Koval, V Kozhuharov, W Kubischta, V Kurshetsov, G Lamanna, C Lazzeroni, M Lenti, E Leonardi, L Litov, N Lurkin, D Madigozhin, A Maier, I Mannelli, F Marchetto, G Marel, M Markytan, P Marouelli, M Martini, L Masetti, P Massarotti, E Mazzucato, A Michetti, I Mikulec, M Misheva, N Molokanova, E Monnier, U Moosbrugger, C Morales Morales, M Moulson, S Movchan, D J Munday, M Napolitano, A Nappi, G Neuhofer, A Norton, T Numao, V Obraztsov, V Palladino, M Patel, M Pepe, A Peters, F Petrucci, M C Petrucci, B Peyaud, R Piandani, M Piccini, G Pierazzini, I Polenkevich, I Popov, Y Potrebenikov, M Raggi, B Renk, F Retire, P Riedler, A Romano, P Rubin, G Ruggiero, A Salamon, G Saracino, M Savri, M Scarpa, V Semenov, A Sergi, M Serra, M Shieh, S Shkarovskiy, M W Slater, M Sozzi, T Spadaro, S Stoynev, E Swallow, M Szleper, M Valdata-Nappi, P Valente, B Vallage, M Velasco, M Veltri, S Venditti, M Wache, H Wahl, A Walker, R Wanke, L Widhalm, A Winhart, R Winston, M D Wood, S A Wotton, O Yushchenko, A Zinchenko and M Ziolkowski

Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI. Published under licence by IOP Publishing Ltd 1

Abstract. The NA62 experiment at CERN collected a large sample of charged kaon decays with a highly efficient trigger for decays into electrons in 2007. The kaon beam represents a source of tagged neutral pion decays in vacuum. A measurement of the electromagnetic transition form factor slope of the neutral pion in the time-like region from ~1 million fully reconstructed π^0 Dalitz decay is presented. The limits on dark photon production in π^0 decays from the earlier kaon experiment at CERN, NA48/2, are also reported.

Introduction

The NA48/2 experiment at the CERN SPS recorded a sample of 2×10^{11} charged kaon decays in 2003-04. The following NA62 experiment collected a ten times smaller sample of K[±] in 2007 and was using the NA48 detector [1] and a modified beam line [2]. This paper reports on the measurements of neutral pion physics on NA62 and NA48/2, namely the measurement of the slope of the π^0 electromagnetic transition form factor from a sample of $\pi^0 \to e^+e^-\gamma$ decays and the search for the dark photon in the $\pi^0 \to \gamma A'$ decay.

1. Experimental setup

Narrow momentum band K[±] beams with central momentum 60 (74) GeV/c, with $\delta P_K/P_K \sim 1\%$ (rms), were extracted from 400 GeV/c proton beam for NA48/2 (NA62) configuration. The kaon beams were focused at the first drift chamber with ~ 10 mm transverse size. The beam flux with an intensity of $2.5 \cdot 10^7$ protons per pulse entered a decay region, where ~ 18% of kaons decayed. The decay region was 113 m long, with a diameter from 1.92 m in the first 66 m increasing to 2.4 m in the last 47 m. It was contained in an evacuated cylindrical steel tank with a volume ~ 120 m³. The vacuum at a level $\leq 10^{-4}$ mbar avoided interactions of kaon decay products before detection. The downstream part of the vacuum tank was closed by convex hemispherical (0.8 mm thick, that corresponds to $3 \cdot 10^{-3} X_0$) Kevlar window with a radius of 1.3 m. The difference between the NA48/2 and NA62 detector setups was the presence of the KABES detector, which has been removed after NA48/2. In NA48/2 two simultaneous beams of K^+ and K^- were used, while in NA62 one beam was alternatively stopped. The fiducial region was followed by a magnetic spectrometer, consisting of four drift chambers (DCH), housed in a vessel filled with helium gas at nearly atmospheric pressure. Through the middle of DCH and following detector a beam pipe allowed the undecayed particles to continue their path in vacuum. Each DCH consisted of 8 octagonal planes with 256 grounded wires and was operated with a gas mixture of 50% Argon and 50% Ethan. The maximum drift time in a plane was 100 ns. The resolution on the spatial position of the reconstructed hit was 90 μ m and the time resolution was about 700 ps. A spectrometer magnet was located between the chambers DCH2 and DCH3. It provided a momentum kick of 120 (257) MeV/c. The resolution of momentum measurement was: $\sigma_p/p = (1.02 \oplus 0.044 \cdot p)\%$ and $\sigma_p/p = (0.48 \oplus 0.009 \cdot p)$ in 2007, p in GeV/c. The spectrometer was followed by a scintillator hodoscope consisting of two planes segmented into horizontal and vertical strips achieving a very good ~ 150 ps time resolution. Further downstream was a liquid krypton calorimeter (LKr) that measured the energy of electrons and photons. The transverse segmentation into 13248 2 cm \times 2 cm projective cells and the 27 radiation length thickness resulted in an energy resolution $\sigma(E)/E = (3.2/\sqrt{E} \oplus 9.0/E \oplus 0.42)\%$ (E in GeV/c) and a transverse position resolution of 1.5 mm for 10 GeV showers. The calorimeter was followed by a muon system consisting of three scintillator planes, with an iron wall installed in front of each plane. A detailed description of the beam line, detectors and trigger is available in [1].

IOP Conf. Series: Journal of Physics: Conf. Series 873 (2017) 012016 doi:10.1088/1742-6596/873/1/012016

2. Theory

The Dalitz decay $\pi_D^0 \to e^+e^-\gamma$ proceeds through the $\pi^0\gamma\gamma$ vertex with an off-shell photon. The commonly used independent kinematic variables x and y are defined in terms of particle four-momenta:

$$x = \left(\frac{M_{ee}}{m_{\pi^0}}\right)^2 = \frac{(p_{e^+} + p_{e^-})^2}{m_{\pi^0}^2}, \qquad y = \frac{2p_{\pi^0} \cdot (p_{e^+} - p_{e^-})}{m_{\pi^0}^2(1-x)}.$$
 (1)

The x variable is the normalized invariant mass squared of the electron-positron pair, while y is related to the angles between the final state particle momenta. The kinematic limits on the variables are given by

$$r^2 \le x \le 1, \quad -\beta \le y \le \beta, \quad \text{where} \quad r = \frac{2m_e}{m_{\pi_0}} \quad \text{and} \quad \beta = \sqrt{1 - \frac{r^2}{x}}.$$
 (2)

The normalized π_D^0 leading order (LO) differential decay width reads:

$$\frac{1}{\Gamma(\pi_{2\gamma}^0)}\frac{d^2\Gamma(\pi_D^0)}{dxdy} = \frac{\alpha}{4\pi}\frac{(1-x)^3}{x}(1+y^2+\frac{r^2}{x})|F(x)|^2.$$
(3)

In the allowed kinematic region for the π_D^0 decay, F(x) is expected to vary slowly and it is sufficient to use a linear approximation F(x) = 1 + ax, where *a* is the so-called π^0 transition form factor (TFF) slope parameter. The expected π^0 TFF slope value from Vector Meson Dominance model, dominated by the ρ and ω mesons, is $a \approx 0.03$. The theoretical models of π^0 TFF enter hadronic light-by-light scattering contribution to $(g-2)_{\mu}$. For a recent overview and references see [3]. The comparison of TFF slope predictions with a model independent measurement is an important test of the theory models.

2.1. Radiative Corrections

The radiation corrections are a crucial aspect of measuring the π^0 TFF, as their effect on the differential decay rate exceed that of the TFF. The first study of radiative corrections was done in [4] and extended in [5], where the following diagrams were considered:



 $\delta_{virt} + \delta_{BS}$ corrections.

The revisited and improved radiative corrections in [6] were largely triggered by and used for this analysis. They include additional second-order contributions to the internal bremsstrahlung:



 $\delta_{1\gamma IR}$ correction.

The contribution of the radiative corrections in the (x, y) plane together with the separate contributions from different origins are shown in Figure 1. They enter the differential decay width through a function $\delta(x, y) = \delta_{virt} + \delta_{BS} + \delta_{1\gamma IR}$ in the next-to-leading-order (NLO):

$$\frac{d^2\Gamma}{dxdy} = \frac{d^2\Gamma^{LO}}{dxdy} \left(1 + \delta(x, y)\right), \qquad \delta(x, y) = \frac{d^2\Gamma^{NLO}}{dxdy} \Big/ \frac{d^2\Gamma^{LO}}{dxdy}.$$
(4)

IOP Conf. Series: Journal of Physics: Conf. Series 873 (2017) 012016

doi:10.1088/1742-6596/873/1/012016



Figure 1. Left: Size of the radiative corrections to the π_D^0 decay in the x, y plane. Right: Size of the radiative corrections versus x. Black line represents the total radiative correction size; blue corresponds to its component first computed in [5]; red shows the new contribution, introduced in [6].

As the π^0 TFF does not depend on the y Dalitz variable, the slope parameter is extracted from the reconstructed distribution of the x Dalitz variable. One can therefore integrate the π_D^0 differential decay width in eq. 3 over y and obtain:

$$\frac{1}{\Gamma(\pi_{2\gamma}^{0})}\frac{d\Gamma(\pi_{D}^{0})}{dx} = \frac{2\alpha}{3\pi}\frac{(1-x)^{3}}{x}(1+\frac{r^{2}}{2x})\sqrt{1-\frac{r^{2}}{x}}(1+\delta(x))(1+ax)^{2},$$
(5)

where the TFF is described by the linear parametrization and the radiative corrections to the x distribution are included.

3. Event selection

The tagged π^0 s are originating from $K^{\pm} \to \pi^{\pm} \pi^0$ decays, which branching fraction is BR $(K_{2\pi}) \approx$ 21%. In 2007, the data taking conditions were optimized for e^{\pm} from $K^{\pm} \rightarrow e^{\pm}\nu_e$, which is favorable also for the $K^{\pm} \rightarrow \pi^{\pm}\pi^0$ with subsequent decay $\pi^0 \rightarrow \gamma e^+ e^-$. During 2007 a large and clean sample of π_D^0 decays was recorded, counting 1.05×10^6 events. The selection of signal events was based on the three-track topology, a photon cluster in LKr and full kinematic closure. The selection was restricted to area with Dalitz variable x > 0.01. Figure 2 shows the reconstructed and simulated invariant mass of π_D^0 , and the distribution of the Dalitz x variable.

4. TFF measurement

The TFF was obtained by adjusting the simulation to the data x spectrum. A χ^2 fit to the reconstructed x distribution of the π_D^0 event candidates distributed over equipopulated bins has been performed to extract the TFF slope value. The NA62 preliminary result on the π^0 TFF slope parameter is obtained as:

$$a = (3.70 \pm 0.53_{stat} \pm 0.36_{syst}) \times 10^{-2}.$$



Figure 2. Left: Reconstructed and simulated invariant mass of π_D^0 . Right: Reconstructed x Dalitz variable for data and MC components. The TFF slope parameter value used to generate MC events was $a_{sim} = 0.032$.

The fit result is illustrated in Figure 3 where the effect of a positive TFF slope is clearly seen from the ratio of the data and MC distribution reweighted to a value a = 0. The horizontal positions of black markers correspond to the barycenters of the data distributed over 20 equipopulated bins. The red solid line represents the TFF function with the slope value equal to the fit central value. Red dashed lines correspond to the 1 σ band. The breakdown of the uncertainties is presented in Table 1. The value of the slope parameter can be compared to the various TFF theory expectations:

- $a = (2.90 \pm 0.50) \times 10^{-2}, \chi PT [7];$
- $a = (3.07 \pm 0.06) \times 10^{-2}$, dispersion theory [8];
- $a = (2.92 \pm 0.04) \times 10^{-2}$, two-hadron saturation [9].

The results from other experiments are shown in Figure 3.

5. Search for the dark photon

The idea of a new U(1) gauge symmetry with a very light gauge boson, called a Dark Photon (DP, usually denoted as A' boson) has been proposed in [10], [11]. The DP could mediate the annihilation of Dark Matter particles and explain the excess of positrons in cosmic rays [12] and also explain the muon gyromagnetic ratio (q-2) measurement [13]. Dark photon presence could manifest itself in the NA48/2 data set by the process $\pi^0 \to \gamma A'$ decay followed by the prompt $A' \to e^+e^-$ decay. The signal decay chain has the same particles in the final state as the π^0 Dalitz decay. The DP is characterized by two a priori unknown parameters, the mass $m_{A'}$ and the mixing parameter ε^2 . The expected branching fraction of π^0 into the DP is [14]

$$\mathcal{B}(\pi^0 \to \gamma A') = 2\varepsilon^2 \left(1 - \frac{m_{A'}^2}{m_{\pi^0}^2}\right)^3 \mathcal{B}(\pi^0 \to \gamma \gamma),\tag{6}$$



Figure 3. Left: Fit result illustration. Right: Results on the TFF slope from π_D^0 measurements.

Source	$\delta a(\times 10^2)$
Statistical-Data	0.49
Statistical-MC	0.20
Beam momentum spectrum simulation	0.30
Spectrometer momentum scale	0.15
Spectrometer resolution	0.05
LKr non-linearity and energy scale	0.04
Particle mis-ID	0.08
Accidental background	0.08
Neglected π_D^0 sources in MC	0.01

Table 1. Summary of uncertainty sources.

which is kinematically suppressed as $m_{A'}$ approaches m_{π^0} . In the DP mass range $2m_e < m_{A'} < m_{\pi^0}$ accessible in pion decays, the only allowed tree-level decay into SM fermions is $A' \to e^+e^-$, while the loop-induced SM decays $(A' \to 3\gamma, A' \to \nu\bar{\nu})$ are highly suppressed. Therefore, for a DP decaying only into SM particles, $\mathcal{B}(A' \to e^+e^-) \approx 1$, the expected total decay width is [14]

$$\Gamma_{A'} \approx \Gamma(A' \to e^+ e^-) = \frac{1}{3} \alpha \varepsilon^2 m_{A'} \sqrt{1 - \frac{4m_e^2}{m_{A'}^2}} \left(1 + \frac{2m_e^2}{m_{A'}^2}\right).$$
(7)

It follows that, for $2m_e \ll m_{A'} < m_{\pi^0}$, the DP mean proper lifetime $\tau_{A'}$ satisfies the relation

$$c\tau_{A'} = hc/\Gamma_{A'} \approx 0.8\mu m \times \left(\frac{10^{-6}}{\varepsilon^2}\right) \times \left(\frac{100 \text{MeV}/c^2}{m_{A'}}\right).$$
 (8)

This analysis is performed assuming that the DP decays at the production point (prompt decay), which is valid for sufficiently large values of $m_{A'}$ and ε^2 . Then the DP has the same production and signature as the Dalitz decay $\pi_D^0 \to e^+ e^- \gamma$, that represents an irreducible but well controlled background.

The result of the analysis of the full data set of NA48/2 including a total of $1.69 \times 10^7 \pi_D^0$ reconstructed events is reported. The π_D^0 events are coming from kaon decays $K^{\pm} \to \pi^{\pm}\pi^0$ (K₂ π) and $K^{\pm} \to \pi^0 \mu^{\pm} \nu$ (K_{µ3}), followed by the prompt $\pi^0 \to \gamma A'$, $A \to e^+e^-$ decay chain. These two exclusive selections apply the following criteria:

 $K_{2\pi}$ selection:

- $|m_{\pi\gamma ee} m_K| < 20 \text{ MeV}/c^2;$
- $|m_{\gamma ee} m_{\pi^0}| < 8 \text{ MeV}/c^2;$
- no missing momentum.

 $K_{\mu3}$ selection:

- $m_{miss}^2 = (P_K P_\mu P_{\pi^0})^2$ compatible with 0;
- $|m_{\gamma ee} m_{\pi^0}| < 8 \text{ MeV}/c^2;$
- missing total and transverse momentum.

The numbers of reconstructed events from $K_{2\pi}$ ($K_{\mu3}$) are 1.38×10^7 (0.31×10^7), respectively. A search for resonances in the di-electron invariant mass spectrum has been performed in the mass range 9 MeV/ $c^2 \leq m_{A'} < 120$ MeV/ c^2 . The lower boundary of the mass range is determined by the limited accuracy of the π_D^0 background simulation at low e^+e^- mass. The obtained upper limits on the numbers of DP candidates and $\mathcal{B}(\pi^0 \to \gamma A')$ in each mass hypothesis considered are presented in Figure 4.

No signal was observed and upper limits on the DP mixing parameter ε^2 were improved in the DP mass range 9 MeV/ $c^2 < m_{A'} < 70 \text{ MeV}/c^2$, see Figure 5. If dark photon couples to quarks and decays mainly to SM fermions, it is ruled out as explanation for the anomalous $(g - 2)_{\mu}$ measurement.

Conclusion

A total of $1.05 \times 10^6 \pi^0$ Dalitz decays, originating from $K^{\pm} \to \pi^{\pm}\pi^0$ decays, have been fully reconstructed. A preliminary measurement of the π^0 transition form factor slope based on NA62 2007 data statistics was performed:

$$a = (3.70 \pm 0.53_{stat} \pm 0.36_{sust}) \times 10^{-2}.$$

The precision of the TFF measurement has been improved in the time-like momentum region. Final result is in preparation.

Improved limits (down to $\epsilon^2 = 2 \times 10^{-7}$) in the 9-70 MeV/ c^2 mass range of the dark photon search in π^0 decays has been achieved by exploring NA48/2 data. The whole region favored by $(g-2)_{\mu}$ is excluded, assuming that DP decays into SM fermions only.

Acknowledgments

The speaker received support from project GA UK no. 404716.





Figure 4. Left: Numbers of observed data events (N_{obs}) and expected π_D^0 background events (N_{exp}) passing the joint DP selection (indistinguishable in a logarithmic scale), estimated uncertainties $\delta N_{obs} = \delta N_{exp}$ and δN_{exp} , and obtained upper limits at 90% CL on the numbers of DP candidates (N_{DP}) for each DP mass vale $m_{A'}$. The contribution to δN_{exp} from the MC statistical uncertainty is shown separately (δN_{exp}^{MC}) . The remaining and dominant component is due to the statistical errors on the trigger efficiencies measured in the DP signal region. Right: Obtained upper limits on $\mathcal{B}(\pi^0 \to \gamma A')$ at 90% CL for each DP mass value $m_{A'}$.



Figure 5. Upper limits at 90% CL on the mixing parameter ε^2 versus the DP mass $m_{A'}$ obtained from $\pi^0 \to \gamma A'$ search by NA48/2, compared to other published exclusion limits from meson decay, beam dump and e^+e^- collider experiments.Further details can be found in [15].

References

- [1] V Fanti et al, Nucl. Instrum. Meth. A 574, 433 (2007).
- [2] J R Batley et al, Eur. Phys. J. C 52, 875 (2007).
- [3] A Nyffeler, arXiv:1602.03398 [hep-ph].

Fifth Symposium on Prospects in the Physics of Discrete Symmetries

IOP Conf. Series: Journal of Physics: Conf. Series 873 (2017) 012016

IOP Publishing

doi:10.1088/1742-6596/873/1/012016

- [4] B E Lautrup and J Smith, Phys. Rev. D 3, 1122 (1971).
- [5] Mikaelian and Smith, Phys. Rev. D 5, (1972) 1763.
- [6] T Husek et al, Phys. Rev. D 92, 054027 (2015).
- [7] K Kampf et al, EPJ C 46 (2006), 191.
- [8] M Hoferichter et al, EPJ C 74, (2014), 3180.
- [9] T Husek et al, EPJ C **75**, (2015) 12, 586.
- [10] P Fayet, *Phys.Lett.* B **95**, 285 (1980).
- [11] C Boehm, P Fayet, and J Silk, *Phys.Rev.* D 69, 101302 (2004).
- [12] M Aguilar et al, Phys.Rev.Lett. 110, 141102 (2013).
- [13] M Pospelov, *Phys. Rev.* D 80, (2009) 095002.
- [14] B Batell, M Pospelov and A Ritz, Phys. Rev. D 80, (2009) 095024.
- [15] J R Batley et al, Phys. Lett. B 746, (2015) 178.