

Bark and leaf chlorophyll fluorescence are linked to wood structural changes in *Eucalyptus saligna*

Johnstone, Denise; Tausz, Michael; Moore, Gregory; Nicolas, Marc

DOI:

[10.1093/aobpla/plt057](https://doi.org/10.1093/aobpla/plt057)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Johnstone, D, Tausz, M, Moore, G & Nicolas, M 2014, 'Bark and leaf chlorophyll fluorescence are linked to wood structural changes in *Eucalyptus saligna*', *AoB PLANTS*, vol. 6, plt057. <https://doi.org/10.1093/aobpla/plt057>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.



Research Article

Bark and leaf chlorophyll fluorescence are linked to wood structural changes in *Eucalyptus saligna*

Denise Johnstone^{1*}, Michael Tausz², Gregory Moore¹ and Marc Nicolas³¹ Department of Resource Management and Geography, University of Melbourne, Burnley Campus, Richmond 3012, Australia² Department of Forest and Ecosystem Science, University of Melbourne, Creswick Campus, Creswick 3363, Australia³ Department of Agriculture and Food Systems, University of Melbourne, Parkville Campus, Parkville 3010, Australia**Received:** 14 April 2013; **Accepted:** 9 December 2013; **Published:** 3 February 2014**Citation:** Johnstone D, Tausz M, Moore G, Nicolas M. 2014. Bark and leaf chlorophyll fluorescence are linked to wood structural changes in *Eucalyptus saligna*. *AoB PLANTS* 6: plt057; doi:10.1093/aobpla/plt057

Abstract. Wood structure and wood anatomy are usually considered to be largely independent of the physiological processes that govern tree growth. This paper reports a statistical relationship between leaf and bark chlorophyll fluorescence and wood density. A relationship between leaf and bark chlorophyll fluorescence and the quantity of wood decay in a tree is also described. There was a statistically significant relationship between the leaf chlorophyll fluorescence parameter F_v/F_m and wood density and the quantity of wood decay in summer, but not in spring or autumn. Leaf chlorophyll fluorescence at 0.05 ms (the O step) could predict the quantity of wood decay in trees in spring. Bark chlorophyll fluorescence could predict wood density in spring using the F_v/F_m parameter, but not in summer or autumn. There was a consistent statistical relationship in spring, summer and autumn between the bark chlorophyll fluorescence parameter F_v/F_m and wood decay. This study indicates a relationship between chlorophyll fluorescence and wood structural changes, particularly with bark chlorenchyma.

Keywords: Bark; chlorophyll fluorescence; photosynthesis; stress physiology; wood decay; wood structure.

Introduction

Tree physiology and wood structure and anatomy are often considered to be independent, as wood occurs primarily in what is sometimes described as the non-functioning heartwood of the tree (Zweifel *et al.* 2006). On the other hand, wood as a tissue (i.e. the secondary xylem of trees) determines long-distance water transport in trees. During water transport, if xylem vessels are under water stress, air bubbles in the xylem can expand due to tension, a process known as cavitation (Hacke *et al.* 2001; Taiz and Zeiger 2010). Once a xylem vessel cavitates it fills with water vapour and then forms an embolism in quick succession, slowing xylem hydraulic conductivity (Tyree and Sperry 1989). Therefore, wood density is increasingly

being measured in conjunction with water-use properties, as low stem wood density can make angiosperms more vulnerable to cavitation, especially during drought (Hacke *et al.* 2001; Holste *et al.* 2006; Bobich *et al.* 2010). However, conifers do not necessarily follow this pattern as their xylem conduits are shorter and narrower. In a study of *Picea abies* (Norway spruce), wood density was unrelated to xylem cavitation (Rosner *et al.* 2007). The relationship between wood decay and physiological measurements not directly related to water use has rarely been assessed. Wood structural changes are frequently caused by wood decay organisms (Rayner and Boddy 1988). Decayed wood shows decreased density as a result of degradation by fungi or bacteria (Harris *et al.* 2004).

* Corresponding author's e-mail address: denisej@unimelb.edu.au

Published by Oxford University Press on behalf of the Annals of Botany Company.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

Weight loss or dry weight is a common means by which to evaluate wood decay, particularly in the early stages of decay (Wilcox 1978; Pandey and Pitman 2003; Wei et al. 2010). Despite ongoing methodological difficulties, wood decay can be quantified by a variety of methods, such as with devices using electrical conductivity, drilling resistance, core sampling or acoustic methods (Johnstone et al. 2010a). It appears logical that wood decay, leading to decreased wood density, can affect tree water transport and, consequently, canopy physiology, mainly under periods of increased demand on water transport. Because wood decay involves invading organisms such as fungi or bacteria, it may also be speculated that biochemical changes (e.g. defence reactions) can affect the physiological function of other tissues.

Trees have chlorenchyma, i.e. photosynthetically active tissue, in their bark below the rhytidomal or outer peridermal layers (Strain and Johnson 1963; Pfanz et al. 2002). Such cortical or peridermal chlorenchyma is able to utilize CO₂ from gaseous xylem efflux and from mitochondrial respiration to photosynthesize (Wittmann et al. 2006; Pfanz 2008). Bark photosynthesis can be strongly shade adapted, particularly in deciduous trees (Pfanz et al. 2002; Damesin 2003; Manetas 2004). *Eucalyptus globulus* bark behaved as a shade leaf in a study by Eyles et al. (2009); however, Tausz et al. (2005) found that parts of sun-exposed *Eucalyptus nitens* bark had photosynthetic pigments of similar quantity and composition to that of sun leaves. Bark photosynthetic activity in stems is generally lower than in the leaves of broadleaf trees such as *Betula pendula*, *Quercus robur* and *Fagus sylvatica*, but it could be a way of improving the carbon balance of stems, particularly where water is limiting (Wittmann and Pfanz 2008b).

Chlorophyll fluorescence (CF) is an excellent tool to assess the physiological state of photosynthetic tissues (Govindjee 2004). F_v/F_m is the most commonly cited CF parameter, where F_v is the difference between maximum (F_m) and minimum (F_o) fluorescence (Maxwell and Johnson 2000). F_v/F_m is the theoretical measure of the quantum efficiency of photosystem II (PSII) if all the PSII reaction centres are open (Maxwell and Johnson 2000). The average F_v/F_m value for healthy tissues is believed to be around 0.83 (Bjorkman and Demmig 1987; Johnson et al. 1993). Decreased values indicating reduced maximum quantum efficiency commonly occur upon impact of environmental stress. F_v/F_m is therefore commonly used to assess stress impacts on plants (Maxwell and Johnson 2000).

The analysis of the intermediate data points of the fast fluorescence rise (i.e. the determination of F_m in the calculation of F_v/F_m) is called the O–J–I–P polyphasic fast fluorescence rise analysis or the O–K–J–I–P polyphasic

fast fluorescence rise analysis (Susplugas et al. 2000; Strasser and Stirbet 2001; Govindjee 2004; Strasser et al. 2004; Percival 2005). The phases are O at the origin (0.05 ms), K at ~0.2 ms, J at ~2 ms, I at ~20 ms and P at ~200 ms, depending on the curve (Strasser and Stirbet 2001). O or F_o fluorescence is measured when all the plastoquinone Q_A electron carrier molecules are in their oxidized state (Krause and Weis 1984; Percival 2005). The K step, not apparent in all cases, may be the result of an imbalance in electron flow coming to the reaction centre from PSII in some species of plants (Strasser et al. 2004). The O–J phase is believed to represent the reduction of the Q_A molecule from Q_A to Q_A^- (Hsu and Leu 2003; Strasser et al. 2004; Percival 2005). J–I may be fluorescence from the abaxial layer of the sample in some plants (Hsu and Leu 2003), or both the J–I and I–P phases could reflect the existence of fast and slow reducing plastoquinone centres (Percival 2005). P or F_m occurs when all the plastoquinone Q_A electron carrier molecules are in their reduced state (Krause and Weis 1984; Percival 2005). The characteristics of the fast fluorescence rise also change upon stress impact, and are therefore used to assess stress impacts on plants.

There is evidence that leaf photosynthetic capacity and the hydraulic properties of tree stems are related (Brodribb and Feild 2000; Brodribb et al. 2007), yet any direct relationship between wood properties or wood decay and photosynthetic properties has rarely been examined. The symptoms of ‘esca’ disease in *Vitis vinifera* (grapevines) and CF parameters have been linked (Christen et al. 2007). Esca disease infects the xylem and causes the white rot decay and/or necrosis of woody tissues and, subsequently, wilting of the leaves. However, no investigations using tree species prior to the current study have attempted to relate photosynthetic properties to wood decay.

In a previous study, the authors investigated a relationship between crown condition and leaf and bark CF (Johnstone et al. 2012). There was little evidence to support a relationship between leaf CF and crown condition. On the other hand, there was a strong relationship between bark CF and crown condition. The current study uses the leaf and bark CF data from the above-mentioned study, but compares it with wood density and wood decay. In this study, the relationship between CF and wood structural properties is examined, rather than CF and growth parameters.

The current study investigated plantation-grown *Eucalyptus saligna* trees exhibiting a range of wood decay from virtually none to moderately decayed. We chose trees already decayed as inducing decay in trees can be a slow process, dependent on tree species and the causal agent of decay (Schwarze 2008). Trees were chosen to

represent the best possible range of decay under otherwise uniform conditions. We examined the relationships between wood decay and density and CF in leaf and bark tissues to test the following hypotheses: (i) increasing wood decay is related to stress symptoms in leaves, particularly in summer when demand on xylem water transport is greatest, and (ii) increasing wood decay is related to stress symptoms in bark chlorenchyma.

Methods

The trees used in this study were *E. saligna* (Bateman's Bay). They were ~20 years old in 2008, between 17 and 27 m high, and with diameters at 1.3 m of between 142 and 318 mm. The 36 selected trees were part of a larger species/provenance study covering a total area of ~10 ha in a eucalypt plantation at Tostaree in rural Victoria, Australia (latitude 37°47'; longitude 148°11'). Sample trees were chosen to represent a range of wood decay and excluded any break or edge trees. In this investigation, CF measurements in both leaves and bark were compared with wood density and the percentage of decay over three seasons (spring, summer and autumn).

Chlorophyll fluorescence measurements

Chlorophyll fluorescence data were collected and analysed according to the method described in Johnstone et al. (2012). Branches ~10 mm in diameter were harvested from the upper canopy with a 12-gauge shotgun in the morning, between 0600 and 0800 h depending on the season. Leaf fluorescence measurements were taken between 13 September and 21 September 2007 (spring), 28 January and 1 February 2008 (summer) and 5 April and 13 April 2008 (autumn). Most eucalypts can have two or three different leaf ages present in the crown at any one season, with leaves lasting up to 18 months. Eucalypts have opportunistic crown phenology dependent on their environmental conditions (Jacobs 1955).

Leaf CF measurements were taken on mature sun leaves from upper canopy branches using a Hansatech-handly plant efficiency analyser (Hansatech Instruments, King's Lynn, Norfolk, UK). Ten leaves from each tree were dark adapted for 30 min with leaf clips. A saturating flash of red light onto the leaf after the period of darkness induced a time-dependent fluorescence kinetic known as the Kautsky effect (Govindjee 2004; Percival 2005). All trees were tested within 2–3 h of being harvested as recommended by Epron and Dreyer (1992).

Bark CF testing was performed in a 350-mm strip in a cross-section of the trunk on the north half of the trees, 35 mm apart. The test area on the bark was circular and 4.5 mm in diameter. Eight to 10 tests were performed on each tree after material had been dark adapted for

30 min. The bark was not damaged or removed in any way. Test results were excluded if the bark was damaged, decorticated or had only recently been exposed to sunlight. The height at which trees were measured was variable as it was necessary to measure above the sock of rough bark at the base. Bark fluorescence measurements were taken between 24 September and 28 September 2007 (spring), 22 January and 26 January 2008 (summer) and 31 March and 4 April 2008 (autumn).

The CF data were averaged from 8–10 measurements from each tree in each tissue (bark and leaf) and in each season. The ratio F_v/F_m was calculated from the raw CF data. F_v/F_m is a derived measure $F_v = F_m - F_o$, where F_v is the difference between maximum (F_m) and minimum (F_o) CF (Maxwell and Johnson 2000). In addition to calculating the F_v/F_m ratio, time data taken over a 1-s period were logarithmically transformed and the O–J–I–P CF phases were allocated following the method devised by Strasser and Stirbet (2001).

Each polyphasic increase in fluorescence was characterized by examining logarithmic graphs for each season and in both leaf and bark tissues. After an exponential rise in graphed data, each phase was deemed complete, with the next phase being deemed to start at the critical point (O, J, I or P). Every step is followed by a characteristic temporary decrease or dip (Strasser et al. 2004). There was no 'K' step observed on the graphs. 'O' was at the origin, taken at 0.05 ms, as in many other studies (Krause and Weis 1984; Susplugas et al. 2000; Strasser and Stirbet 2001; Govindjee 2004; Strasser et al. 2004; Percival 2005). The O–J phase was characterized as ending at 4 ms (J step). The 'I' step in leaf fluorescence data was observed at 60 ms and in bark at 90 ms. The 'P' step was observed at ~700 ms on leaf fluorescence graphs, previously observed at 200–300 ms in other studies. The 'P' step was not observed in bark fluorescence as the last recording point taken by the instrument was at 1000 ms, and fluorescence was still increasing at this time. The JIP test was not applied to the data; comparisons were made using the raw fluorescence values for O (0.05 ms all data), J (4 ms all data), I (60 ms leaf data, 90 ms bark data), P (700 ms leaf data) and the 1000 ms data point on bark.

Wood density measurement and wood decay estimation

The 36 *E. saligna* were tested for basic wood density from a small sample collected from the trunk at 1.5 m in height from the trees when they were felled in 2008. Basic wood density was estimated as oven dry mass of wood/volume of wood when 'green' (Walker et al. 1993). Wood decay in the trees was quantified using the Resi system utilizing the IML-Resi constant feed drill described in Johnstone

et al. (2007, 2010b). The method begins with cross-sectional drilling measurements of the trunk at 0.3 m. The method combines the IML-Resi raw data and Shigo's (1979) compartmentalization of decay in trees (CODIT) model to predict the quantity of wood decay beyond the linear drill locations of the IML-Resi. The method relied on the experienced use of the IML-Resi, knowledge of models of decay in trees and image analysis software (Johnstone *et al.* 2007, 2010b).

Statistical analysis of data

A comparison was made between spring, summer and autumn CF data and wood density and wood decay data using simple linear regression analysis. Simple linear regression analyses were performed using the software package SAS (Statistical Analysis System) version 9.2 (SAS Institute Inc., Cary, NC, USA). Although multiple comparisons were made, Bonferroni corrections were not applied in order to maximize statistical power and minimize Type II errors in the analysis (Moran 2003).

One tree had no leaves and could not be included in leaf CF analysis, and the bark of this tree had died by the autumn sampling date. Data more than two standard deviations away from the next nearest result were considered outliers and eliminated from analysis, resulting in 34–35 individual replicate trees for regression analysis.

Results

Comparing leaf and bark fluorescence and basic wood density

There was a statistically significant and positive relationship between summer leaf F_v/F_m and basic wood density (Table 1 and Fig. 1A). There was a statistically significant

Table 1. Summarized results from simple linear regression analyses comparing spring, summer and autumn leaf or bark F_v/F_m with basic wood density data. *n*, the number of samples; *P*, the probability for the *t*-test that the coefficient of the independent variable is equal to zero; r^2 , the variation in the dependent variable that can be explained by the fluorescence data. ^aThe dependent variable is the spring basic wood density data in all cases. ^bThe statistical relationship is significant and positive. Bold values indicate statistical significance.

Independent variable ^a	<i>n</i>	<i>P</i>	r^2
Spring leaf fluorescence— F_v/F_m	34	0.531	0.012
Summer leaf fluorescence— F_v/F_m	34	0.001^b	0.291
Autumn leaf fluorescence— F_v/F_m	35	0.387	0.023
Spring bark fluorescence— F_v/F_m	35	0.035^b	0.128
Summer bark fluorescence— F_v/F_m	35	0.512	0.013
Autumn bark fluorescence— F_v/F_m	35	0.249	0.040

and positive relationship between spring bark F_v/F_m and basic wood density (Table 1 and Fig. 1B). There was no statistical relationship between spring and autumn leaf F_v/F_m or summer and autumn bark F_v/F_m and basic wood density (Table 1). There was no statistical relationship between spring, summer and autumn leaf CF at the O, J, I or P step and basic wood density (Table 2). There was also no statistically significant relationship between spring, summer and autumn bark CF at the O, J, I or 1000 ms step and basic wood density (Table 2).

Comparing leaf and bark fluorescence and wood decay

There was a statistically significant and positive relationship between spring leaf CF at the O step and wood decay (Table 3 and Fig. 2A). There was a statistically significant and negative relationship between the summer leaf F_v/F_m ratio and wood decay (Table 3). There was a statistically significant and negative relationship between spring, summer and autumn bark F_v/F_m and wood decay (Table 3 and Fig. 2B).

There was no statistically significant relationship between the spring leaf F_v/F_m ratio or CF at the J, I and P step and wood decay (Table 4). There was no statistically significant relationship between summer leaf CF at the O, J, I, and P step and wood decay (Table 4). There was no statistically significant relationship between autumn leaf CF and wood decay (Table 4). There was no statistically significant relationship between bark CF at the O, J, I or 1000 ms step and wood decay, in spring, summer or autumn (Table 4).

Discussion

Weight loss or its corollary wood density has been used to assess wood decay for many years (Kennedy 1958; Wilcox 1978; Wei *et al.* 2010). Wood decay organisms can be responsible for weight losses as small as 5 % or less (Noguchi *et al.* 1986). In instances of very early decay, even a light microscope may not be able to detect wood decay visually (Wilcox 1978). Hence there is clearly a strong relationship between measured wood density and wood decay, even in assumed sound or intact wood.

There was a statistically significant and positive relationship between summer leaf F_v/F_m and basic wood density, but not in spring or autumn. In this study, the summer period of investigation coincided with maximum seasonal tree stress in southern Australia, when the mean average maximum temperature at the test site in January 2008 was 27 °C (minimum average 16 °C, Bureau of Metrology Australia 2008). However, summer predawn leaf and stem water potentials were not significantly different from spring values, although values of around




Figure 1. (A) Basic wood density in kg m^{-3} versus summer leaf F_v/F_m . Trend line = linear regression, $P = 0.001$, $r^2 = 0.291$. F_v/F_m ratio data begin at 0.820, and basic density data begin at 400 kg m^{-3} . (B) Basic wood density in kg m^{-3} versus spring bark F_v/F_m . F_v/F_m ratio data begin at 0.7900, and basic density data begin at 400 kg m^{-3} . Trend line = linear regression, $P = 0.035$, $r^2 = 0.128$.

1.2 MPa for all trees in summer indicated mild to moderate drought stress at this site (White et al. 2000; Johnstone et al. 2012). Predawn water potentials did not show a relationship with wood density or decay, but water potentials were not measured during the day when water deficit becomes more noticeable. Summer leaf CF at the O step also correlated with a visual vitality measurement in summer (Johnstone et al. 2012), which suggests that the trees were suffering some type of stress during the seasonal summer drought and that the leaf O step was sensitive to the stress. Wood density is sometimes measured in conjunction with other parameters for assessing the water status of trees (O'Grady et al. 2009; Gotsch et al. 2010). Low stem wood density in angiosperms is sometimes thought to be indicative of increased vulnerability to xylem cavitation during drought stress (Holste et al. 2006; Bobich et al. 2010). If cavitation is occurring in the xylem of the *E. saligna* in the current study, it could establish a favourable environment for fungal pathogens (Rayner and Boddy 1988), or the pathogens may assist with the cavitation process (Tyree and Sperry 1989; Tyree and Zimmermann 2002). This may explain why there is a relationship between leaf CF and wood decay in two seasons (spring and summer) rather than just one as was the case with wood density.

Changes in the availability of water for the *E. saligna* may have contributed directly to the relationship between photosynthetic efficiency and wood decay discovered in this study. The water saturation of wood has long been known to prevent the development of wood decay, and air is necessary for the development of decay in wood (Rayner and Boddy 1988). The barrier zones in Shigo's CODIT model (Shigo 1979) are said to be a response to xylem embolism by Rayner and Boddy (1988), rather than the incursion of decay organisms *per se*. Cavitation

during moisture stress is one way a xylem vessel can develop an embolism (Tyree and Sperry 1989). The direct relationship between xylem cavitation and wood decay has not been evaluated, but the introduction of a gaseous phase during the compartmentalization process, according to Rayner and Boddy (1988), is a primary component in the development of wood decay in trees. It is when tree wood dries out that compartmentalization barriers are breached (Rayner and Boddy 1988).

Trees with lower wood density have also been associated with an increased risk of cavitation (Holste et al. 2006; Bobich et al. 2010). Therefore, it is not surprising that in this study, in the hot Australian summer, *E. saligna* showed an inverse relationship between leaf CF and wood density, and an even stronger relationship between leaf CF and wood decay. Unlike many other studies, the link between moisture stress, cavitation, embolism and wood density/decay described here is a within-species effect, rather than the ecological inter-species effect of low wood density and water relations/growth discussed in other studies (Bucci et al. 2004; O'Grady et al. 2009). The lower density wood is produced due to stressful environmental conditions or is a result of very early wood decay in the *E. saligna*; there are no genetic differences or predispositions at play. The link between wood density, leaf CF and wood decay within species established in this study has not been previously reported.

There was a statistically significant and positive relationship between leaf CF values at the 'O' step and wood decay in spring. The O–J step is believed to represent the reduction of the plastoquinone Q_A molecule from Q_A to Q_A^- between PSII and photosystem I (PSI) (Hsu and Leu 2003; Strasser et al. 2004; Percival 2005); therefore, it appears that the reduction of Q_A between PSII and PSI during leaf photosynthesis is associated with wood decay in *E. saligna*. F_v/F_m is the theoretical

Table 2. Summarized results from simple linear regression analyses comparing spring, summer and autumn leaf or bark net chlorophyll fluorescence with basic wood density data. Results from these analyses were not significant. *n*, the number of samples; *P*, the probability for the *t*-test that the coefficient of the independent variable is equal to zero; *r*², the variation in the dependent variable that can be explained by the fluorescence data. ^aThe dependent variable is the spring basic wood density data in all cases.

Independent variable ^a	<i>n</i>	<i>P</i>	<i>r</i> ²
Spring leaf fluorescence—'O' step	34	0.741	0.004
Spring leaf fluorescence—'J' step	34	0.620	0.008
Spring leaf fluorescence—'I' step	34	0.462	0.017
Spring leaf fluorescence—'P' step	34	0.891	0.001
Spring bark fluorescence—'O' step	35	0.702	0.005
Spring bark fluorescence—'J' step	35	0.691	0.005
Spring bark fluorescence—'I' step	35	0.298	0.033
Spring bark fluorescence—1000 ms	35	0.173	0.056
Summer leaf fluorescence—'O' step	34	0.072	0.097
Summer leaf fluorescence—'J' step	34	0.085	0.090
Summer leaf fluorescence—'I' step	34	0.134	0.069
Summer leaf fluorescence—'P' step	34	0.913	0.000
Summer bark fluorescence—'O' step	35	0.309	0.031
Summer bark fluorescence—'J' step	35	0.832	0.001
Summer bark fluorescence—'I' step	35	0.256	0.039
Summer bark fluorescence—1000 ms	35	0.191	0.051
Autumn leaf fluorescence—'O' step	34	0.810	0.002
Autumn leaf fluorescence—'J' step	34	0.558	0.011
Autumn leaf fluorescence—'I' step	34	0.905	0.001
Autumn leaf fluorescence—'P' step	34	0.747	0.003
Autumn bark fluorescence—'O' step	35	0.427	0.020
Autumn bark fluorescence—'J' step	35	0.461	0.017
Autumn bark fluorescence—'I' step	35	0.734	0.004
Autumn bark fluorescence—1000 ms	35	0.828	0.002

measure of the quantum efficiency of PSII if all the PSII reaction centres are open (Maxwell and Johnson 2000). There was a significant and negative relationship between the leaf CF F_v/F_m ratio and wood decay in summer, suggesting that wood decay may also be associated with the quantum efficiency of PSII in leaves.

This study further emphasizes the link between the operation of photosynthesis in leaves and environmental stress. The O step in the OJIP fluorescence transient in leaves, which relates to the part of the photosynthetic light reaction where plastoquinone Q_A electron carrier molecules are in their oxidized state between PSII and PSI, is particularly affected by moisture stress in other

Table 3. Summarized results from simple linear regression analyses comparing spring, summer and autumn leaf or bark F_v/F_m and 'O' step fluorescence values with wood decay data. *n*, the number of samples; *P*, the probability for the *t*-test that the coefficient of the independent variable is equal to zero; *r*², the variation in the dependent variable that can be explained by the fluorescence data. ^aThe dependent variable is wood decay in all cases. ^bThe statistical relationship is significant and positive. ^cThe statistical relationship is significant and negative. Bold values indicate statistical significance.

Independent variable ^a	<i>n</i>	<i>P</i>	<i>r</i> ²
Spring leaf fluorescence— F_v/F_m	34	0.505	0.014
Spring leaf fluorescence—'O' step	34	0.004^b	0.230
Spring bark fluorescence— F_v/F_m	35	0.036^b	0.127
Spring bark fluorescence—'O' step	35	0.363	0.025
Summer leaf fluorescence— F_v/F_m	34	0.025^c	0.148
Summer leaf fluorescence—'O' step	34	0.080	0.093
Summer bark fluorescence— F_v/F_m	35	0.037^b	0.125
Summer bark fluorescence—'O' step	35	0.101	0.079
Autumn leaf fluorescence— F_v/F_m	35	0.853	0.001
Autumn leaf fluorescence—'O' step	34	0.870	0.001
Autumn bark fluorescence— F_v/F_m	35	0.034^b	0.129
Autumn bark fluorescence—'O' step	35	0.363	0.025

studies of trees (Epron *et al.* 1992; Percival and Sheriffs 2002). This study establishes a new link between the quantum efficiency of PSII (F_v/F_m) in leaves, wood density and wood decay. The study also establishes a new and consistent pattern of correlation between the quantum efficiency of PSII (F_v/F_m) in bark and environmental stress, wood decay and to a lesser extent wood density. Further research could examine the link between the quantum efficiency of PSII in bark in relation to other tree species, and other environmental stressors.

Christen *et al.* (2007) investigated the esca disease in *V. vinifera* (grapevines) and the relationship between the white rot decay and/or necrosis of woody tissues, the wilting of leaves and CF parameters. They used four categories of white rot decay and eight categories of necrosis, rather than percentages of decay. Necrosis and white rot were more widespread in Cabernet Sauvignon than in Merlot plants. The more decayed Cabernet Sauvignon plants showed decreased efficiency in PSII and the PI_{ABS} (performance index) value according to the CF results, compared with the Merlot population. However, the statistical relationship between CF and wood decay was only significant at a cultivar level in *V. vinifera*, rather than at an individual plant level, as was the case in the *E. saligna* from the current study.




Figure 2. (A) Percentage of decay using the Resi system versus spring leaf chlorophyll fluorescence at the 'O' step in millivolts. Chlorophyll fluorescence data begin at 100 mV. Trend line = linear regression, $P = 0.004$, $r^2 = 0.230$. (B) Percentage of decay using the Resi system versus summer bark F_v/F_m . F_v/F_m ratio data begin at 0.800. Trend line = linear regression, $P = 0.021$, $r^2 = 0.148$.

Table 4. Summarized results from simple linear regression analyses comparing spring, summer and autumn leaf or bark 'J', 'I', 'P' or 1000 ms fluorescence values with wood decay data. Results from these analyses were not significant. n , the number of samples; P , the probability for the t -test that the coefficient of the independent variable is equal to zero; r^2 , the variation in the dependent variable that can be explained by the fluorescence data. ^aThe dependent variable is the wood decay data in all cases.

Independent variable ^a	n	P	r^2
Spring leaf fluorescence—'J' step	34	0.076	0.095
Spring leaf fluorescence—'I' step	34	0.456	0.018
Spring leaf fluorescence—'P' step	34	0.158	0.062
Spring bark fluorescence—'J' step	35	0.207	0.048
Spring bark fluorescence—'I' step	35	0.617	0.008
Spring bark fluorescence—1000 ms	35	0.901	0.001
Summer leaf fluorescence—'J' step	34	0.104	0.081
Summer leaf fluorescence—'I' step	34	0.452	0.018
Summer leaf fluorescence—'P' step	34	0.660	0.006
Summer bark fluorescence—'J' step	35	0.095	0.082
Summer bark fluorescence—'I' step	35	0.295	0.033
Summer bark fluorescence—1000 ms	35	0.430	0.019
Autumn leaf fluorescence—'J' step	34	0.969	0.000
Autumn leaf fluorescence—'I' step	34	0.350	0.027
Autumn leaf fluorescence—'P' step	34	0.319	0.031
Autumn bark fluorescence—'J' step	35	0.478	0.015
Autumn bark fluorescence—'I' step	35	0.691	0.005
Autumn bark fluorescence—1000 ms	35	0.987	0.000

The question as to why trees with decreased photosynthetic efficiency are more decayed is not easy to answer. Lorio (1986) suggested that the production of oleoresin, a

protective agent against *Dendroctonus frontalis* (southern pine beetle) in *Pinus taeda* (loblolly pine), is lower in suppressed trees when the production of wood is depressed. Hence, one reason why *E. saligna* trees with decreased photosynthetic efficiency may be more decayed may be because when the growth of wood is depressed, the synthesis of protective chemical compounds produced in the wood is also decreased. In addition, research on the progression of wood decay in trees suggests that the origin of wood decay can be in the sapwood, rather than from saprotrophic growth in non-functioning heartwood (Boddy and Rayner 1983; Parfitt et al. 2010). It is possible that during the seasonal challenge inherent in hot summers, there is pressure on photosynthesis as a result of stomatal closure and the resultant high light stress on photosystems (Faria et al. 1998). Therefore, under such conditions trees with partly decayed primary xylem suffer more because they may have to close their stomata more or more often because their water transport system is less efficient.

Induction curves did not reach their maximum in the bark fluorescence measurements; however, high F_v/F_m ratios indicate that values were close to maximum and were not significantly biased by low light intensities (Johnstone et al. 2012). Bark F_v/F_m ratios were negatively correlated with wood density in *E. saligna* in spring only, and in this instance the bark CF statistical relationships were weaker than those for leaf CF (Tables 1 and 2). The statistical relationships between bark F_v/F_m ratios and wood decay were weak but more consistent over three seasons than correlations with leaf parameters (Tables 3 and 4). Therefore, PSII in leaves may be more sensitive to the immediate effects of water flow disruption than bark photosynthesis, but the longer-term sustained effects of moisture stress, such as cavitation and the subsequent

entry of wood decay pathogens, affect PSII in bark in a more consistent pattern than in the leaves. Stem photosynthesis is believed to use gaseous xylem efflux as a source of CO₂ (Pfanzen 2008); therefore, if the xylem is not fully functioning it may affect the health of bark chlorenchyma, and thus PSII. Some tree species have been found to have elevated CO₂ in decayed wood tissues, while CO₂ was depressed in winter in other species (Rayner and Boddy 1988); thus the complex interactions of the metabolism of xylem, bark and wood decay organisms warrant further investigation.

The transpirational xylem stream supplies inorganic nutrients (and water) to bark chlorenchyma (Pfanzen 2008), so if the xylem stream is disrupted it may affect stem photosynthesis. *Eucalyptus* sp. may be sensitive to factors that affect stem photosynthesis as stem photosynthesis may be a more important source of photosynthates for them than for other broadleaved trees, because they have a low leaf area index and are prone to defoliation by insects, diseases or drought (Tausz et al. 2005; Eyles et al. 2009). Interestingly, unlike the leaf CF measurements, only the quantum efficiency (F_v/F_m) of PSII within bark chlorenchyma was associated with wood decay; the reduction of the plastoquinone Q_A molecule between PSII and PSI (O–J step) was not affected.

The PI_{ABS} CF value has been used to successfully quantify drought stress in trees (Percival and AlBalushi 2007; Swoczyna et al. 2010). The PI_{ABS} value was not calculated in the current study, as it is not as widely used as the F_v/F_m value. Future studies could examine the effect of wood decay and wood density in trees on the PI_{ABS} value in relation to bark photosynthesis and other derived measures that form part of the ‘JIP test’, such as the apparent rates of photosynthetic electron transport and non-photochemical quenching (Lüttge et al. 2003).

Conclusions

The CF measurements in this study clearly support the hypothesis that there is a relationship between CF and wood structural changes. The results suggest that when photosynthesis is impaired, trees are more prone to wood decay and low wood density. Although chlorenchymes are present in bark and indeed in many woody tree tissues (Pfanzen et al. 2002), it may not be possible to test trunk tissue in many tree species due to peridermal thickening (Aschan et al. 2001). However in some studies photosynthesis has been measured successfully in stems (Damesin 2003; Wittmann and Pfanzen 2008a). This raises the possibility of testing larger branches on many temperate species where the bark is not as thick as on the trunk of the tree. This study showed that the reduced functioning of

PSII in bark chlorenchyma in particular is an indication that a tree may have a larger quantity of decay in the xylem tissues.

Sources of Funding

Our work was funded by the University of Melbourne, Australia.

Contributions by the Authors

D.J. performed all of the collection and analysis of data, was responsible for the experimental design and wrote most of the introduction, methods, results, discussion and conclusion. M.T. assisted in the design of the experiments and wrote sections of the introduction and discussion. G.M. offered advice on experimental design and assistance with the introduction and discussion sections. M.N. contributed to the experimental design.

Conflicts of Interest Statement

None declared.

Acknowledgements

We thank the arboricultural consultancy company ‘Tree Logic’ for the loan of the Handy PEA chlorophyll fluorescence meter.

Literature Cited

- Aschan G, Wittmann C, Pfanzen H. 2001. Age-dependent bark photosynthesis of aspen twigs. *Trees—Structure and Function* **15**: 431–437.
- Bjorkman O, Demmig B. 1987. Photon yield of O₂ evolution and chlorophyll fluorescence characteristics at 77-K among vascular plants of diverse origins. *Planta* **170**:489–504.
- Bobich EG, Barron-Gafford GA, Rascher KG, Murthy R. 2010. Effects of drought and changes in vapour pressure deficit on water relations of *Populus deltoides* growing in ambient and elevated CO₂. *Tree Physiology* **30**:866.
- Boddy L, Rayner A. 1983. Origins of decay in living deciduous trees: the role of moisture content and a reappraisal of the expanded concept of tree decay. *New Phytologist* **94**:623–641.
- Brodrick T, Feild T. 2000. Stem hydraulic supply is linked to leaf photosynthetic capacity: evidence from New Caledonian and Tasmanian rainforests. *Plant, Cell and Environment* **23**:1381–1388.
- Brodrick TJ, Feild TS, Jordan GJ. 2007. Leaf maximum photosynthetic rate and venation are linked by hydraulics. *Plant Physiology* **144**: 1890.
- Bucci S, Goldstein G, Meinzer F, Scholz F, Franco A, Bustamante M. 2004. Functional convergence in hydraulic architecture and water relations of tropical savanna trees: from leaf to whole plant. *Tree Physiology* **24**:891–899.
- Christen D, Schonmann S, Jermini M, Strasser RJ, Defago G. 2007. Characterization and early detection of grapevine (*Vitis vinifera*)

- stress responses to esca disease by *in situ* chlorophyll fluorescence and comparison with drought stress. *Environmental and Experimental Botany* **60**:504–514.
- Damesin C. 2003. Respiration and photosynthesis characteristics of current-year stems of *Fagus sylvatica*: from the seasonal pattern to an annual balance. *New Phytologist* **158**:465–475.
- Epron D, Dreyer E. 1992. Effects of severe dehydration on leaf photosynthesis in *Quercus petraea* (Matt) Liebl.: photosystem II efficiency, photochemical and nonphotochemical fluorescence quenching and electrolyte leakage. *Tree Physiology* **10**:273–284.
- Epron D, Dreyer E, Breda N. 1992. Photosynthesis of oak trees [*Quercus petraea* (Matt) Liebl] during drought under field conditions—diurnal course of net CO₂ assimilation and photochemical efficiency of photosystem-II. *Plant, Cell and Environment* **15**:809–820.
- Eyles A, Pinkard EA, O'Grady AP, Worledge D, Warren CR. 2009. Role of cortical photosynthesis following defoliation in *Eucalyptus globulus*. *Plant, Cell and Environment* **32**:1004–1014.
- Faria T, Silvério D, Breia E, Cabral R, Abadia A, Abadia J, Pereira JS, Chaves MM. 1998. Differences in the response of carbon assimilation to summer stress (water deficits, high light and temperature) in four Mediterranean tree species. *Physiologia Plantarum* **102**:419–428.
- Gotsch SG, Geiger EL, Franco AC, Goldstein G, Meinzer FC, Hoffmann WA. 2010. Allocation to leaf area and sapwood area affects water relations of co-occurring savanna and forest trees. *Oecologia* **163**:291–301.
- Govindjee. 2004. Chlorophyll a fluorescence: a bit of basics and history. In: Papageorgiou GC, Govindjee, eds. *Chlorophyll a fluorescence a signature of photosynthesis*. Dordrecht: Springer, 1–41.
- Hacke UG, Sperry JS, Pockman WT, Davis SD, McCulloh KA. 2001. Trends in wood density and structure are linked to prevention of xylem implosion by negative pressure. *Oecologia* **126**:457–461.
- Harris R, Clark J, Matheny N. 2004. *Arboriculture—integrated management of landscape trees, shrubs and vines*. New Jersey: Prentice Hall.
- Holste EK, Jerke MJ, Matzner SL. 2006. Long term acclimatization of hydraulic properties, xylem conduit size, wall strength and cavitation resistance in *Phaseolus vulgaris* in response to different environmental effects. *Plant, Cell and Environment* **29**:836–843.
- Hsu BD, Leu KL. 2003. A possible origin of the middle phase of polyphasic chlorophyll fluorescence transient. *Functional Plant Biology* **30**:571–576.
- Jacobs MR. 1955. *Growth Habits of the Eucalypts*. Canberra: Government Press.
- Johnson G, Young A, Scholes J, Horton P. 1993. The dissipation of excess excitation energy in British plant species. *Plant, Cell and Environment* **16**:673–679.
- Johnstone D, Moore G, Tausz M, Nicolas M. 2010a. The measurement of wood decay in landscape trees. *Arboriculture and Urban Forestry* **36**:121–127.
- Johnstone D, Tausz M, Moore G, Nicolas M. 2010b. Quantifying wood decay in Sydney bluegum (*Eucalyptus saligna*) trees. *Arboriculture and Urban Forestry* **36**:243–252.
- Johnstone D, Tausz M, Moore G, Nicolas M. 2012. Chlorophyll fluorescence of the trunk rather than leaves indicates visual vitality in *Eucalyptus saligna*. *Trees—Structure and Function* **26**:1565–1576.
- Johnstone DM, Ades PK, Moore GM, Smith IW. 2007. Predicting wood decay in eucalypts using an expert system and the IML-Resistograph drill. *Arboriculture and Urban Forestry* **33**:76–82.
- Kennedy R. 1958. Strength retention in wood decayed to small weight losses. *Forest Products Journal* **8**:308–314.
- Krause GH, Weis E. 1984. Chlorophyll fluorescence as a tool in plant physiology. *Photosynthesis Research* **5**:139–157.
- Lorio PL Jr. 1986. Growth-differentiation balance: a basis for understanding southern pine beetle-tree interactions. *Forest Ecology and Management* **14**:259–273.
- Lüttge U, Berg A, Fetene M, Nauke P, Peter D, Beck E. 2003. Comparative characterization of photosynthetic performance and water relations of native trees and exotic plantation trees in an Ethiopian forest. *Trees—Structure and Function* **17**:40–50.
- Manetas Y. 2004. Probing cortical photosynthesis through *in vivo* chlorophyll fluorescence measurements: evidence that high internal CO₂ levels suppress electron flow and increase the risk of photoinhibition. *Physiologia Plantarum* **120**:509–517.
- Maxwell K, Johnson GN. 2000. Chlorophyll fluorescence—a practical guide. *Journal of Experimental Botany* **51**:659.
- Metrology. 2008. Bureau of Metrology Commonwealth of Australia. <http://www.bom.gov.au/> (accessed 13 February 2008).
- Moran MD. 2003. Arguments for rejecting the sequential Bonferroni in ecological studies. *Oikos* **100**:403–405.
- Noguchi M, Nishimoto K, Imamura Y, Fijii Y, Okumura S, Miyauchi T. 1986. Detection of very early stages of decay in western hemlock wood using acoustic emissions. *Forest Products Journal* **36**:35–36.
- O'Grady A, Cook P, Eamus D, Duguid A, Wischusen JDH, Fass T, Worledge D. 2009. Convergence of tree water use within an arid-zone woodland. *Oecologia* **160**:643–655.
- Pandey K, Pitman A. 2003. FTIR studies of the changes in wood chemistry following decay by brown-rot and white-rot fungi. *International Biodeterioration & Biodegradation* **52**:151–160.
- Parfitt D, Hunt J, Dockrell D, Rogers HJ, Boddy L. 2010. Do all trees carry the seeds of their own destruction? PCR reveals numerous wood decay fungi latently present in sapwood of a wide range of angiosperm trees. *Fungal Ecology* **3**:338–346.
- Percival GC. 2005. *Identifying tree decline using 'fluorescence fingerprinting'*. Hobart: International Society of Arboriculture Australia Chapter.
- Percival GC, AlBalushi AMS. 2007. Paclobutrazol-induced drought tolerance in containerized English and evergreen oak. *Arboriculture and Urban Forestry* **33**:397–409.
- Percival GC, Sheriffs CN. 2002. Identification of drought-tolerant woody perennials using chlorophyll fluorescence. *Journal of Arboriculture* **28**:215–223.
- Pfanz H. 2008. Bark photosynthesis. *Trees—Structure and Function* **22**:137–138.
- Pfanz H, Aschan G, Langenfeld-Heyser R, Wittmann C, Loose M. 2002. Ecology and ecophysiology of tree stems: cortical and wood photosynthesis. *Naturwissenschaften* **89**:147–162.
- Rayner ADM, Boddy L. 1988. *Fungal decomposition of wood. Its biology and ecology*. Chichester: John Wiley & Sons Ltd.
- Rosner S, Klein A, Müller U, Karlsson B. 2007. Hydraulic and mechanical properties of young Norway spruce clones related to growth and wood structure. *Tree Physiology* **27**:1165.
- Schwarze FW. 2008. *Diagnosis and prognosis of the development of wood decay in urban trees*. Rowville: Enspect.

- Shigo AL. 1979. Tree decay an expanded concept. *USDA Forest Service Information Bulletin* **419**:72.
- Strain BR, Johnson PL. 1963. Corticular photosynthesis and growth in *Populus tremuloides*. *Ecology* **44**:581–584.
- Strasser RJ, Stirbet AD. 2001. Estimation of the energetic connectivity of PS II centres in plants using the fluorescence rise OJIP: fitting of experimental data to three different PS II models. *Mathematics and Computers in Simulation* **56**:451–462.
- Strasser RJ, Tsimilli-Michael M, Srivastava A. 2004. Analysis of the chlorophyll *a* fluorescence transient. In: Papageorgiou GC, Govindjee, eds. *Chlorophyll *a* fluorescence a signature of photosynthesis*. Dordrecht: Springer, 321–362.
- Susplugas S, Srivastava A, Strasser RJ. 2000. Changes in the photosynthetic activities during several stages of vegetative growth of *Spirodela polyrhiza*: effect of chromate. *Journal of Plant Physiology* **157**:503–512.
- Swoczyzna T, Kalaji HM, Pietkiewicz S, Borowski J, Zaras-Januszkiewicz E. 2010. Photosynthetic apparatus efficiency of eight tree taxa as an indicator of their tolerance to urban environments. *Dendrobiology* **63**:65–75.
- Taiz L, Zeiger E. 2010. *Plant physiology*. Massachusetts: Sinauer Associates.
- Tausz M, Warren CR, Adams MA. 2005. Is the bark of shining gum (*Eucalyptus nitens*) a sun or a shade leaf? *Trees—Structure and Function* **19**:415–421.
- Tyree MT, Sperry JS. 1989. Vulnerability of xylem to cavitation and embolism. *Annual Review of Plant Biology* **40**:19–36.
- Tyree MT, Zimmermann MH. 2002. *Xylem structure and the ascent of sap*. Berlin: Springer.
- Walker JCF, Butterfield B, Langrish T, Harris J, Uprichard J. 1993. *Primary wood processing*. London: Chapman and Hall.
- Wei D, Houtman CJ, Kapich AN, Hunt CG, Cullen D, Hammel KE. 2010. Laccase and its role in production of extracellular reactive oxygen species during wood decay by the brown rot basidiomycete *Postia placenta*. *Applied and Environmental Microbiology* **76**: 2091–2097.
- White DA, Turner NC, Galbraith JH. 2000. Leaf water relations and stomatal behavior of four allopatric *Eucalyptus* species planted in Mediterranean southwestern Australia. *Tree Physiology* **20**: 1157–1165.
- Wilcox WW. 1978. Review of literature on the effects of early stages of decay on wood strength. *Wood and Fiber Science* **9**: 252–257.
- Wittmann C, Pfan H. 2008a. Antitranspirant functions of stem periderms and their influence on corticular photosynthesis under drought stress. *Trees—Structure and Function* **22**:187–196.
- Wittmann C, Pfan H. 2008b. General trait relationships in stems: a study on the performance and interrelationships of several functional and structural parameters involved in corticular photosynthesis. *Physiologia Plantarum* **134**:636–648.
- Wittmann C, Pfan H, Loreto F, Centritto M, Pietrini F, Alessio G. 2006. Stem CO₂ release under illumination: corticular photosynthesis, photorespiration or inhibition of mitochondrial respiration? *Plant, Cell and Environment* **29**:1149–1158.
- Zweifel R, Zimmermann L, Zeugin F, Newbery DM. 2006. Intra-annual radial growth and water relations of trees: implications towards a growth mechanism. *Journal of Experimental Botany* **57**: 1454–1459.