

Stress, the Immune System and Healthy Ageing.

Whittaker, Anna

DOI:

[10.1007/978-3-319-97034-9](https://doi.org/10.1007/978-3-319-97034-9)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Whittaker, A 2018, Stress, the Immune System and Healthy Ageing. in E Peel, C Holland & M Murray (eds), *Psychologies of Ageing. : Theory, Research and Practice*. 1 edn, Palgrave Macmillan, Basingstoke, pp. 107-130. <https://doi.org/10.1007/978-3-319-97034-9>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Whittaker, Peel, Holland, Murray eds. in *Psychologies of Ageing: Theories, Research & Practice*, 2018, Palgrave Macmillan reproduced with permission of Palgrave Macmillan.

This extract is taken from the author's original manuscript and has not been edited. The definitive, published, version of record is available here: <https://www.palgrave.com/gb/book/9783319970332> for product on www.palgrave.com and www.link.springer.com].

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Psychologies of Ageing: Theory, Research and Practice.

Chapter 5 – Stress, the Immune System and Healthy Ageing

Professor Anna C. Whittaker (previously Phillips)

School of Sport, Exercise & Rehabilitation Sciences, University of Birmingham, UK

Anna C. Whittaker is Professor of Behavioural Medicine in the School of Sport, Exercise & Rehabilitation Sciences, University of Birmingham. She is an internationally renowned researcher and Registered Health Psychologist working in Psychoneuroimmunology and Psychophysiology particularly in the context of ageing. She has conducted award winning work on stress and vaccination response across the life course and has a strong profile of public engagement with science. She is also now developing her research into the area of physical activity interventions for older adults to improve health, function and wellbeing through leading the European Commission Physical Activity and Nutrition INfluences In ageing (PANINI) training network and consortium across Europe.

Abstract

With ageing, the immune system does not function as well as it did earlier in life, leaving individuals open to a range of diseases including infections. Negative psychological factors such as stress and depression can worsen immunity, and in some cases, this is particularly the case among older adults. However, there are also positive psychological factors that have been shown to relate to positive immune and health outcomes. This chapter will outline the links between psychological and behavioural factors and the immune system, particularly in the context of ageing. It will focus on key positive and negative psychological factors that can boost or be detrimental to our immunity, and briefly touch on the mechanisms by which these factors influence immunity and health in later life.

Keywords: ageing; bereavement; caregiving; cortisol:DHEAS ratio; immunity; life events; neutrophils; social support; stress; vaccination;

Introduction

The **stress response** in humans, although an adaptive mechanism initially, has the potential to be chronic and detrimental to the organism if too large and/or prolonged (Sapolsky, 2007). This particularly seems to be the case later in the lifespan; in fact, although ageing is a physiological process that is part of normal development (Cutler, 1991a, 1991b), some of the changes in older age mirror the chronic effects of psychological stress on several of the body's biological systems. This chapter will mainly focus on the impact of stress on the immune system and the implications for health in older age, as stress effects on all bodily systems are beyond the scope of one chapter. Further, as the immune system undergoes several changes with ageing, this results in increased susceptibility to infectious diseases, all of which are also influenced by stress. The chapter will begin with an introduction to psychological stress and its relationship to illness susceptibility, and a summary of the physiological responses to stress. It will then provide an introduction to the immune system and the effects of ageing on it, including the impacts of ageing on stress hormones. Following this background, the chapter will examine research evidence in this area, focusing on the potentially negative impacts of caregiver stress on immunity in ageing, and the potentially positive impacts of social support.

Psychological stress

Psychological stress can be characterised in a number of ways but is usually considered as something that exceeds our coping abilities (Lazarus & Folkman, 1984). One way of characterising psychological stress in order to understand its sources is to categorise it as cataclysmic events, stressful life events, or daily hassles. Cataclysmic events include natural disasters and war which can have considerable effects on health. One example would be the

impact of being a soldier during the Vietnam war, which led to the development of post-traumatic stress disorder (PTSD) in many veterans, although not all. Those veterans who developed PTSD were then at increased risk of a range of non-communicable diseases over the next 20 years compared to those who did not respond to the stress or war with PTSD (Boscarino, 1997). Stressful life events are occurrences that happen to most individuals but not necessarily regularly, and their impact can be months to years such as unemployment, bereavement, marriage/divorce/separation, and caregiving for a spouse or family member. An example of the impact of this type of stressor is illustrated in a study in which individuals who had suffered more stressful life events and generally felt more stressed were more susceptible to developing an upper respiratory tract infection after exposure to a rhinovirus (such as a common cold) than those who reported fewer stressful events and lower stress levels (Cohen, Tyrell, & Smith, 1991). The final category of stressful daily hassles includes a range of generally short-lived minor inconveniences such as misplacing something, or being late. Such hassles are common and regular but can also relate to health outcomes, particularly in the case of a build-up of many hassles over a short time period. For example, individuals who reported higher numbers of minor daily hassles on a particular day showed increased symptoms of malaise and illness four days later (Sheffield, McVey, & Carroll, 1996). Psychological stress can also be characterised as of different durations; acute stress lasts for minutes or hours whereas chronic stress lasts for days, weeks, months or even years, and the resultant physiological impacts on health differ considerably, as will be discussed in the next section.

The stress response

The physiological response to psychological stress was characterized by Walter Cannon (1929) as the 'fight or flight' response. The main function of this response is to maintain bodily homeostasis. Biologically, the key site involved in this process is the hypothalamus (Barrett, 2005), a part of the brain that communicates by sending nerve impulses to other parts of the body. In this way, the hypothalamus acts within seconds and via the sympathetic nervous system stimulates the medulla of the adrenal gland to release catecholamines (adrenalin and noradrenalin) which act on receptors throughout the body to result in several effects such as increased heart rate, blood pressure and respiration, activation of smooth muscle, and increased core temperature and pain threshold (Charmandari, Tsigos, & Chrousos, 2005). In addition, the hypothalamus also produces chemical messengers that act more slowly and in the next minutes travel through the hypothalamic-pituitary-adrenal (HPA) axis (Sapolsky, Romero, & Munck, 2000). Chemical messengers in this pathway include corticotrophin releasing hormone (CRH) which stimulates the anterior pituitary gland to release another hormone, adrenocorticotrophic hormone (ACTH), into circulation. The target organ of ACTH is again the adrenal gland, but this time stimulates the cortical cells that synthesise and release species-specific glucocorticoids (GC) into the blood. The tight control of these GC (mainly cortisol in humans) is sustained via negative feedback that controls and terminates the release of CRH (Griffin & Ojeda, 2004). Cortisol potentiates the effects of catecholamines within the body (Charmandari et al., 2005) as well as initially activating the immune system, then later working as an anti-inflammatory agent generally suppressing immune function, to prevent harmful over-activation (Munck, Guyre, & Holbrook, 1984). However, when stress is prolonged, chronic, or there are repeated exposures to stressful events, this puts the body out of homeostasis for too long and dysregulation of the stress response axis can occur resulting in detrimental effects throughout the body, for example, on the cardiovascular system (McEwen, 1998; Phillips, 2011; Sedova et al., 2004) and also by

suppression of the immune system (Sorrells & Sapolsky, 2007). Acute stress can in fact be beneficial for the immune system, leading to temporary upregulation of immune cells responses, and has been associated in some cases with improved antibody responses to vaccination, e.g., (Edwards et al., 2006). In contrast, chronic stress, such as caregiving for a child with developmental delay, has been shown to down-regulate immunity, resulting in a decrease in the antibody response to vaccination, e.g., (Gallagher, Phillips, Drayson, & Carroll, 2009a).

The immune system

The immune system can be considered as two distinct, but interconnected elements; the innate and the adaptive immune systems. The innate response is often referred to as the 'first line of defence' against infection as it comprises mechanisms that are the first to react to an infection. The adaptive immune system is slower to respond but has the advantages that it includes memory of pathogens encountered and that its response is specific for each pathogen, thus conferring a tailored and long lasting protection against further infection by the same pathogen. The innate immune system consists of soluble components, namely the complement system, which consists of a series of proteins in the blood which, when activated, stimulate, or complement, the work of the cellular elements. The cellular elements, including neutrophils which deal with rapidly dividing bacteria and fungi, eosinophils which respond to parasitic infections, macrophages which secrete soluble factors (such as the cytokines $\text{TNF}\alpha$, IL-1, IL-6) to co-ordinate and amplify the immune response and also provide immunity against intracellular bacteria, and Natural Killer (NK) cells which detect and kill virally-infected cells and tumour cells. Adaptive immunity is provided by T and B lymphocytes, which develop and mature in the thymus and bone marrow respectively. T

cells can be further classified into CD4 expressing helper cells (which in turn can be split into Th1 and Th2 types), CD8 expressing cytotoxic cells and CD25 expressing T regulatory cells, which have immune suppressive function. B cells, when presented with an antigen (by dendritic cells or with T cell help) produce **antibodies** to provide extended protection against infections. When a naïve T or B cell encounters a pathogen it will proliferate and differentiate into an effector cell or a memory cell, so that if the pathogen is encountered a second time a more rapid response can be achieved. Ageing is known to have deleterious effects upon both the innate and adaptive immune responses, though the latter is much better characterized (Phillips, Burns, & Lord, 2007).

Ageing and immunity

With ageing, the innate immune response goes through some changes. For example, complement activation appears to be unaffected, but neutrophil bactericidal and phagocytic function *in vitro* is dramatically reduced (Butcher et al., 2001). Macrophage function is also modified, although the literature is rather contradictory, including reports of reduced phagocytosis capacity to engulf pathogens, release chemicals such as and superoxides to kill pathogens, function as seen in neutrophils, but enhanced secretion of the cytokines IL-6 and IL-8 in response to pathogens. NK cells are also affected by ageing; while their numbers do not change with age, their cytotoxic capacity is reduced (Hazeldine, Hampson, & Lord, 2012) which has also been shown to relate to reduced survival in people aged over 75 years (Ogata et al., 2001). In the adaptive immune system, the thymus gland atrophies and thus fewer naïve T cells are produced. As the size of the T cell pool is maintained at a constant level, the proportion of T cells that are memory cells increases. Consequently, as we age, we are less able to deal with new pathogens. In addition to changes in the ratio of naïve to memory T cells, there is a shift from T-helper 1 to T-helper 2 cells, and the end result is reduced cell-

mediated, Th1-type, immunity. Finally, with ageing, antibody production in response to antigen declines; for example, older people produce a lower antibody titre (a measure of how much antibody a person has produced) in response to vaccination than younger individuals and the antibodies produced are of lower affinity (binds less well to the invading antigens). This is thought to be largely the result of a decline in T cell help for B cells in older adults.

Another concept that frequently appears in the literature when discussing the ageing of the immune system is inflammaging (Franceschi, Capri, Monti, Giunta, Olivieri, Sevini, Panourgia, et al., 2007). Inflammatory factors are damaging in excessive amounts and inflammaging indicates an imbalance between inflammatory factors necessary to fight the infection, and anti-inflammatory components that act as a counter weight. It has been suggested that ageing and longevity could, therefore, potentially be dependent on this balance (Franceschi, Capri, Monti, Giunta, Olivieri, Sevini, Panouraia, et al., 2007). This would mean that immunosenescence, together with inflammatory markers such as different cytokines (IL-6, IL-8 and IL-15) could contribute to the prediction predictors of the longevity of organisms.

Stress hormones and ageing

As outlined above, stress, whether physical or psychological, is broadly sensed by two systems within the hypothalamus, the HPA axis and the sympathetic-adrenal-medullary system. Stress induces the release of catecholamines from the adrenal medulla, and both cortisol and dehydroepiandrosterone (DHEA) from the adrenal cortex. Catecholamines and cortisol can both be immunosuppressive if chronically elevated. In contrast, DHEA is a

precursor to sex hormones and is considered to be immune enhancing (Butcher et al., 2005). Due to the impact of these hormones on immunity, any change in their production could therefore have significant health implications. In humans, the production of DHEA and its sulphated form, DHEAS, declines with age, a process termed the adrenopause (Orentreich, Brind, Rizer, & Vogelmann, 1984). The synthesis of DHEA is maximal in humans at age 20-30 and declines gradually thereafter, so that by the seventh decade levels of DHEA can be as low as 10% of that seen in young adulthood (Orentreich, Brind, Vogelmann, Andres, & Baldwin, 1992); this adrenopause occurs at similar rates in both males and females. However, although DHEA/S levels fall with age, the production of glucocorticoids such as cortisol is remarkably unaltered (Orentreich et al., 1992), resulting in a relative excess of cortisol over DHEA/S and an imbalance of immune suppression over immune enhancement. The age-related immunological and endocrinological changes outlined above may have implications for the ability to cope (physiologically) with stress in older adults. It is likely that the combination of adrenopause, leading to a relative preponderance of cortisol, and an already reduced immune defence against infection through immune senescence, may leave this population particularly vulnerable to the negative effects of stress on immunity (Graham, Christian, & Kiecolt-Glaser, 2006; Phillips, Burns, & Lord, 2007). However, younger adults are also susceptible to the impact of stress on immunity. For example, various studies have now shown reduced antibody responses to vaccination in those with greater self-reported stress levels (Burns, Carroll, Drayson, Whitham, & Ring, 2003; Phillips, Burns, Carroll, Ring, & Drayson, 2005). Nonetheless, when ageing is combined with stress, the effects on immunity are often more detrimental. For example, in one large study, caregivers, but only those aged over 60 years, showed lower levels of a particular antibody, salivary immunoglobulin A, which targets pathogens in biological fluids, particularly saliva at mucosal surfaces (Gallagher, Phillips, Evans, et al., 2008).

Caregiver stress and immunity in ageing

As indicated above, one commonly studied model of the impact of stress on immunity is the role of caregiving for another person, be it a spouse or child with a physical or mental illness or disability. Older caregivers have most commonly been studied in this context, using the model of family dementia caregiving (Gouin, Hantsoo, & Kiecolt-Glaser, 2008). Caregiving is now well established as having a serious effect on psychological wellbeing and physical health among caregivers when compared to matched non-caregiving individuals (Pinquart & Sorensen, 2003). Both innate and adaptive immunity are affected by chronic stress experienced by older adults. For example, wound healing was slower in older dementia caregivers when compared to age, sex and income-matched controls (Kiecolt-Glaser, Marucha, Malarkey, Mercado, & Glaser, 1995). Further, lower production of proinflammatory cytokines involved in the wound healing process such as IL-1 α , IL-8 (Glaser et al., 1999), as well as IL-1 β (Kiecolt-Glaser et al., 1995) has been observed in caregivers compared to controls. Natural killer (NK) cells are essential in targeting tumour or virally-infected cells. (Esterling, Kiecolt-Glaser, Bodnar, & Glaser, 1994 showed that NK cells from caregivers respond more weakly compared to NK cells from controls).

A further association with the chronic stress of caregiving was found for adaptive cell-mediated immunity; elevated cortisol levels as well as poorer proliferation to antigen and lower IL-2 production was shown in a caregiving group (Bauer et al., 2000). Caregiving stress in older adults has also been shown to be associated with the T-helper 1 to T-helper 2 shift in the type of cytokine responses, with the difference that in older stressed individuals this was driven purely by an increase in IL-10 production, with no difference in IFN- γ

production by Th1 cells (Glaser et al., 2001). It is likely that stress-induced changes in catecholamine levels (Elenkov & Chrousos, 2002) during the psychological stress response drive this cytokine-related behaviour.

Inflammaging, as observed in older adults, might also be more severe among chronically stressed older adults, such as dementia caregivers. Indeed, when compared to non-caregiving older adults who also had immunosenescence, not only did older caregivers show higher levels of IL-6 (von Kanel et al., 2006), but its rate of increase was four times higher than in non-caregiving older controls, leaving them particularly vulnerable to IL-6 related diseases such as frailty, cardiovascular diseases, osteoporosis and others (Ershler & Keller, 2000).

A novel approach for assessing the severity by which caregiving stress affects the immune system of older caregivers is that of studies of latent-virus antibody titres. It is known, for example, that reactivation of latent viral infections, such as those initiated by Herpes group (HSV-1, EBV, and CMV) is typical for immunosuppressed patients such as HIV and transplant patients (Rasmussen, 1991). Interestingly, older spousal caregivers had higher IgG antibody titres against EBV VCA (virus capsid antigen) compared to the matched controls, indicating poorer control of the latent infection in this group (Kiecolt-Glaser, Speicher, Glaser, Dura, & Trask, 1991). Together with the higher antibody titre to total viral antigen of HSV-1, caregivers also had a decreased virus-specific T cell response; another component of immune system necessary for controlling the infection (Glaser & Kiecolt-Glaser, 1997). Older caregivers have also been characterised by higher antibody titres against CMV when compared to the controls (Pariante et al., 1997). In this instance, showing higher antibody

levels against CMV is an indicator that the virus is not under control, and therefore indicates poorer immunity.

Vaccination responses are affected by increasing age which makes older adults particularly vulnerable to frequent infections such as pneumonia and influenza, among the top five causes of high morbidity and mortality in this age group (Thompson et al., 2003). It would be expected that this aspect of immune incompetence would be further exacerbated in older adults affected by the chronic stress of caregiving. This is indeed the case; a significantly lower percentage of older caregivers of dementia patients showed a four-fold increase in antibody titre in response to vaccination against the influenza virus, a response that is clinically considered to be protective against infection (Vedhara et al., 1999a). This was accompanied by higher salivary cortisol concentration in the caregiver group when compared to the controls, pointing again to the role of HPA axis in immune regulation among chronically stressed individuals. Most antigens, however, trigger both humoral, i.e., the antibody response which is generated by B lymphocytes, as well as cellular responses, mainly mediated by cytotoxic CD8⁺ T-cells (Glaser, Sheridan, Malarkey, MacCallum, & Kiecolt-Glaser, 2000a; Kiecolt-Glaser, Glaser, Gravenstein, Malarkey, & Sheridan, 1996; Siergist, 2008). In addition, CD4⁺ helper T-cells are necessary as mediators between those two. It has been shown that both the antibody response to medical vaccination against the influenza virus, as well as IL-2 production in response to antigen stimulation, was lower in caregivers comparing to the controls (Kiecolt-Glaser et al., 1996). In the case of the pneumococcal pneumonia vaccine, even though caregivers managed to exert an adequate immune response initially, shown as a rise in IgG antibody titre, it declined over time more rapidly in this group than in the group of matched controls, likely either as a consequence of decrease in number of antibody-specific B-cells, or their ability to produce antibody (Glaser et al., 2000a;

Vedhara et al., 1999a). However, this impact of caregiving stress on vaccination responses is not unique to older caregivers as was previously thought (Vedhara et al., 2002). One elegant study showed that if the caregiving stress is severe enough, particularly if the behaviours exhibited by the care recipient are challenging, then even younger caregivers can display decrements in the antibody response to vaccination. This was shown in young parental caregivers of children with a learning delay such as Autism, Down's Syndrome, and other less well known syndromes, in comparison to parents of typically developing children; parental caregivers showed a reduction in the production of antibodies to the influenza and pneumococcal vaccinations, and this was particularly marked among those whose children displayed severe challenging behaviours (Gallagher et al., 2009a; Gallagher, Phillips, Drayson, & Carroll, 2009b). However, it should be noted that there was a trend for an effect for ageing such that the lowest antibody titres were among parental caregivers at the older end of the age group (Gallagher et al., 2009a). These studies have been followed up with similar case-control design studies but including both younger and older adults to examine the interaction between ageing and caregiving stress. These studies showed that for some aspects of immunity, ~~neutrophil function~~, ageing or the interaction between stress and ageing were not the key predictors of poorer immune function, but the stress itself was. In these studies caregivers reporting higher psychological distress (depression, anxiety, and caregiving burden) showed the poorest neutrophil function independent of age ([Vitlic et al., 2015](#)), and individuals with higher levels of psychological distress generally also displayed a somewhat higher cortisol:DHEAS ratio within each age group (Vitlic et al., unpublished data). (~~Vitlic et al., 2015~~). Regardless of age?

Chronic stress and immunity in ageing

Given the importance of the caregiving model of stress, little attention has been given to other sources of stress among older adults, and immunity. Although one study has reported that perceived stress, measured using the perceived stress questionnaire (Cohen, Kamarck, & Mermelstein, 1983), was associated with a poorer antibody response to the influenza vaccine in older adults (Kohut, Cooper, Nickolaus, Russell, & Cunnick, 2002), another small scale study found no association between perceived stress and antibody status following this vaccination in ~~elderly~~ very old (mean age 84 years) nursing home residents (Moynihan et al., 2004). Very few studies have focused on stressful life events, despite these being a common means of assessing the impact of stress on immunity in younger samples (e.g., Burns et al., 2003; Phillips et al., 2005). However, a study published in 2006 examined overall stressful life events using a life events rating scale and showed that middle-aged and older adults with higher ratings of stress and disruptiveness for the stressful events they had experienced in the past two years showed lower levels of IgA in saliva (Phillips, Carroll, Evans, et al., 2006). Further, another study took advantage of the UK National Health Service vaccination programme whereby adults aged 65+ years are invited for an annual influenza vaccination. Stressful life events and social support were measured using standardised psychometric questionnaires and antibody titres were assessed at baseline prior to vaccination, and one month later. Although there was no overall effect of stress and social support, one particular stressor of import to older adults related to their immune responses; those who had suffered bereavement in the past year showed lower antibody titres to two of the three 'flu vaccine components (Phillips, Carroll, Bums, et al., 2006). The absence of an association between overall life events and antibody response to influenza vaccination in the older adults and vaccination study (Phillips, Carroll, Bums, et al., 2006) contrasts with the results of previous research on young participants (Burns et al., 2003; Phillips et al., 2005). However, in these student studies, the modal number of life events experienced in the past year was six, with no

participants reporting one or less events (Phillips et al., 2005), whereas in the older sample, the modal number of major life events in the year prior to vaccination was zero, with 31% of the sample reporting no events, and a further 17% reporting only one. This might be due to differences between student and older adult life event stress scales. In the student studies, less serious events were included in the stress scale, for example, getting an unjustified low mark on a test or minor financial problems, along with more major events, whereas the older adults life events scale tended to focus on exposure to major life events. Accordingly, the absence of an association between antibody response and overall life events in older adults may reflect the use of a scale including only serious, rarer life events. However, the results for bereavement would argue against this explanation. In addition, it is also possible that older people simply experience fewer general life events than younger samples. There is certainly evidence to this effect: older individuals encountered fewer major life events than middle-aged participants in a large cohort study in the west of Scotland using the same life events measure as the ~~present~~ [Phillips et al., \(2006\)](#) study, but retrospectively over two years. Middle-aged (mean age 44 years) participants identified a mean of 2.0 events whereas the mean number of events for the older group (mean age 63 years) was 1.7 (Carroll, Phillips, Ring, Der, & Hunt, 2005). These data also suggest that ~~our~~ [the Phillips et al \(2006\)](#) participants were not unusual in experiencing few life events, given that the mean number of events reported over one year was 2.9. Accordingly, it may be that individual differences in general life events exposure are less important for immunity as people age, whereas bereavement, a specific life event that older people are more likely to encounter than the young, assumes greater prominence.

The negative association between bereavement and antibody status following vaccination is in line with previous studies of bereavement and immune function. Bereavement has been associated with in vitro functional immune measures such as decreased natural killer cell cytotoxicity and poorer lymphocyte proliferation to antigen (Bartrop, Luckhurst, Lazarus, Kiloh, & Penny, 1977; Goodkin et al., 1996; Irwin, Daniels, Smith, Bloom, & Weiner, 1987; Kemeny et al., 1995; Schleifer, Keller, Camerino, Thornton, & Stein, 1983; Zisook et al., 1994). In follow-up work, focussing on the two-month period post-bereavement, it was shown that neutrophils' killing ability was suppressed in bereaved older adults, an effect that was accompanied by the increase in cortisol:DHEAS ratio (Khanfer, Lord, & Phillips, 2011). However, when this study was replicated and also included younger adults, the impact of bereavement stress on neutrophil function was only evident for older caregivers (Vitlic, Khanfer, Lord, Carroll, & Phillips, 2014). Previous work has suggested that psychological morbidity may be a mediator between the bereavement and both immune system effects and their impact on risk of morbidities such as influenza and pneumonia. For example, Zisook, Shuchter, Irwin, Darko and Sledge (1994) reported an effect on immune indices where the bereaved older adults were reporting more depressive symptoms. Other work has indicated that the influence of conjugal bereavement on mortality is modified by gender, existing cardiovascular disease status, and depression (Stahl, Arnold, Chen, Anderson & Schulz (2016).

Stress need not be only psychological stress, but could also be physiological stress such as physical disease or physical trauma such as a severe fracture or burn. Older adults who experienced the physical trauma or stress of hip fracture had higher cortisol:DHEA ratios than healthy controls and lower neutrophil function (Butcher et al., 2005), and individuals

with lower neutrophil function were more likely to succumb to infection post-fracture (Butcher, Killampalli, Chahal, Kaya Alpar, & Lord, 2003). Further, older adults who developed depression post-hip fracture showed the highest cortisol:DHEA ratio and poorest neutrophil function (Duggal, Upton, Phillips, & Lord, 2013), as well as worse frailty and slower physical recovery (Phillips, Upton, Duggal, Carroll, & Lord, 2013).

Social support and immunity in ageing

Given the substantial impact of various types of stress on immunity in older adults, as outlined above, understanding psychological factors that can help to enhance or improve immunity in this group is particularly important. Social support, or comfort, caring, esteem, or help provided by other people or social groups can be a key resource that helps individuals cope with life. It has also been shown to have a substantial impact on health; for example, individuals with low numbers of supportive relationships had two to three times the mortality risk compared to those with large social networks (Berkman & Syme, 1979). Indeed, social network size and quality and frequency of social support have been shown to impact on morbidity and mortality from serious diseases in many epidemiological studies, e.g., (Barger, 2013; House, Robbins, & Metzner, 1982; Kaplan et al., 1988). Social support has also been shown to relate to immune function. For example, whereas students who had seroconverted (developed antibodies) after the first injection of the standard three-dose hepatitis B vaccination were less anxious and reported lower stress levels than those who had not, those who reported greater social support demonstrated a stronger combined immune response to the booster third inoculation (Glaser et al., 1992). In another study of college first years, loneliness and smaller social network size were associated with a poorer antibody response to the A/New Caledonian strain of the influenza vaccination (Pressman et al., 2005). Finally,

higher social support scores, particularly higher frequency of tangible support, was related to an increased antibody response to the A/Panama component of the influenza vaccination, again in university students (Phillips et al., 2005).

In a study of social support in older adults, social support was negatively correlated with A/Panama influenza strain antibody status following vaccination, a finding which contradicts the studies with students above that even the authors found difficult to explain (Moynihan et al., 2004). In contrast, a larger study of older adults considered the actual vaccination response, i.e., the change in antibody levels from pre- to post-vaccination (Phillips, Carroll, Bums, et al., 2006). In this study, although social network size and functional social support were not related to antibody response, married/cohabiting participants showed a better antibody response to the A/Panama strain at one month than those who were not married, particularly widowed, participants. Also, for those who were married or cohabiting, higher marital satisfaction was related to higher titres to A/Panama at one month. This is not entirely surprising given that poorer marital quality, in terms of adjustment and negative marital interactions, is associated with inferior functional immunity evidenced through reduced proliferation to some antigens, poorer latent virus control (Kiecolt-Glaser et al., 1987; Kiecolt-Glaser et al., 1997; Kiecolt-Glaser et al., 1988; Kiecolt-Glaser et al., 1993), and weaker natural killer cell cytotoxicity (Miller, Dopp, Myers, Stevens, & Fahey, 1999) in the general population. Further, it is possible that the variations between older care-givers and controls in terms of vaccination response (Glaser, Kiecolt-Glaser, Malarkey, & Sheridan, 1998; Glaser, Sheridan, Malarkey, MacCallum, & Kiecolt-Glaser, 2000b; Kiecolt-Glaser, Glaser, Gravenstein, Malarkey, & Sheridan, 1996; Vedhara et al., 1999b) may be driven, at least in part, by the effects of care-giving on marital quality and satisfaction, although more specific measurement of stressful life events and marital parameters would be necessary to

support this speculation. Whatever the case, these findings resonate with the broad consensus that both marriage (Gordon & Rosenthal, 1995; House, Landis, & Umberson, 1988; Johnson, Backlund, Sorlie, & Loveless, 2000; Verbrugge, 1979), and marital satisfaction (Coyne & DeLongis, 1986; Kiecolt-Glaser & Newton, 2001; Robles & Kiecolt-Glaser, 2003) are beneficial for health. Further, it is possible that in an older aged population, general social support is less critical, whereas the specific social support resource of a happy marriage becomes more important for health, including susceptibility to infection.

Interestingly, the studies of social support and immunity show direct associations rather than an effect of social support via buffering the negative impact of stress. It is possible that for psychological and other health outcomes, social support can buffer stress effects (Lazarus & Folkman, 1984; Rosengren, Orth-Gomer, Wedel, & Wilhelmsen, 1993), whereas for immune function, social support might impact immunity independently. Certainly, social support has been shown to affect different types of vaccination responses to those impacted by stressful life events in various studies among younger adults (Gallagher, Carroll, & Phillips, 2007; Gallagher, Phillips, Ferraro, Drayson, & Carroll, 2008; Phillips et al., 2005).

Conclusion

This chapter has focused on the impact of ageing on stress hormones and immune function as well as the effects of stress on immunity and the role of stress hormones. It also briefly touched on a variable that is known to relate to stress often, social support, and its links with immune function in ageing. Ageing is considered as a normal process, but in some cases the normal processes of ageing (adrenopause, inflammaging, immune senescence) can contribute

to a reduced ability to deal physiologically with stress in later life. However, the research has also shown that stress can impact on stress hormone levels and certain aspects of immunity even in young healthy adults, thus, chronic stress throughout the life span will likely further impact on health and wellbeing as these individuals age. Consequently, individuals with a life history of fewer severe stress exposures may be at lower risk of a heightened cortisol:DHEA ratio, immune decrements, and greater inflammation, even in the presence of the normal hormonal and immune changes associated with ageing. These individuals are thus likely to be more resilient to stress or trauma if and when it does occur later in the lifespan, although longitudinal research would be needed to confirm this possibility.

Although avoiding stressful events themselves may not be a realistic undertaking for most individuals, certainly where stress levels can be reduced by healthy behaviours or seeking social support, these methods are likely to have positive psychological, immune and thus health impact throughout life not just in older age. Healthy behaviours with direct effects on both perceived stress levels and immune function can also be pursued in order to increase resilience in later life. These would include exercise or physical activity, adequate sleep, a balanced diet, not smoking or taking drugs, and moderation of alcohol intake, some of which is outlined in Chapter 6, but their effects on immunity and within healthy ageing warrant a separate chapter each.

A certain level of stress can be beneficial for health, and indeed the immune system. As described in this chapter, this occurs in the case of acute stress, where the immune system can demonstrate enhanced function in response in much the same way a vaccine challenges the immune system (Lewitus & Schwartz, 2009). The challenge for healthy ageing is for the

occurrence of acute stressors not to escalate to the extent where they lead to overload and individuals beginning to show the detrimental physiological effects of chronic stress.

References

- Barger, S. D. (2013). Social integration, social support and mortality in the US National Health Interview Survey. *Psychosomatic Medicine*, 75(5), 510-517.
doi:PSY.0b013e318292ad99 [pii]10.1097/PSY.0b013e318292ad99
- Barrett, E. J. (2005). The adrenal gland. In B. W.F., & E.L., Boulpaep (Ed.), *Medical physiology: A cellular and molecular approach* (pp. 1049-1065): Elsevier Inc.
- Bartrop, R. W., Luckhurst, E., Lazarus, L., Kiloh, L. G., & Penny, R. (1977). Depressed lymphocyte function after bereavement. *The Lancet*, 1(8016), 834-836.
- Bauer, M. E., Vedhara, K., Perks, P., Wilcock, G. K., Lightman, S. L., & Shanks, N. (2000). Chronic stress in caregivers of dementia patients is associated with reduced lymphocyte sensitivity to glucocorticoids. *Journal of Neuroimmunology*, 103(1), 84-92. doi:10.1016/s0165-5728(99)00228-3
- Berkman, L. F., & Syme, S. L. (1979). Social networks, host resistance, and mortality: a nine-year follow-up study of Alameda County residents. *American Journal of Epidemiology*, 109(2), 186-204.
- Boscarino, J. A. (1997). Diseases among men 20 years after exposure to severe stress: implications for clinical research and medical care. *Psychosomatic Medicine*, 59(6), 605-614.
- Burns, V. E., Carroll, D., Drayson, M., Whitham, M., & Ring, C. (2003). Life events, perceived stress and antibody response to influenza vaccination in young healthy adults. *Journal of Psychosomatic Research*, 55, 569-572.

- Butcher, S. K., Chahal, H., Nayak, L., Sinclair, A., Henriquez, N. V., Sapey, E., . . . Lord, J. M. (2001). Senescence in innate immune responses: reduced neutrophil phagocytic capacity and CD16 expression in elderly humans. *Journal of Leukocyte Biology*, 70(6), 881-886.
- Butcher, S. K., Killampalli, V., Chahal, H., Kaya Alpar, E., & Lord, J. M. (2003). Effect of age on susceptibility to post-traumatic infection in the elderly. *Biochemical Society Transactions*, 31(2), 449-451.
- Butcher, S. K., Killampalli, V., Lascelles, D., Wang, K., Alpar, E. K., & Lord, J. M. (2005). Raised cortisol:DHEAS ratios in the elderly after injury: potential impact upon neutrophil function and immunity. *Aging Cell*, 4(6), 319-324.
- Cannon, W. B. (1929). *Bodily changes in pain, hunger, fear, and rage*. New York: D. Appleton & Co.
- Carroll, D., Phillips, A. C., Ring, C., Der, G., & Hunt, K. (2005). Life events and hemodynamic stress reactivity in the middle-aged and elderly. *Psychophysiology*, 42, 269-276.
- Charmandari, E., Tsigos, C., & Chrousos, G. (2005). Endocrinology of the stress response *Annual Review of Physiology* (Vol. 67, pp. 259-284).
- Cohen, S., Kamarck, T., & Mermelstein, R. (1983). A global measure of perceived stress. *Journal of Health and Social Behavior*, 24, 385-396.
doi:<http://dx.doi.org/10.2307/2136404>
- Cohen, S., Tyrell, D. A. J., & Smith, A. P. (1991). Psychological stress and susceptibility to the common cold. *The New England Journal of Medicine*, 325, 606-612.
- Coyne, J. C., & DeLongis, A. (1986). Going beyond social support: the role of social relationships in adaptation. *Journal of Consulting and Clinical Psychology*, 54(4), 454-460.

- Cutler, R. G. (1991a). Antioxidants and aging. *American Journal of Clinical Nutrition*, 53(1 Suppl), 373S-379S.
- Cutler, R. G. (1991b). Antioxidants and aging. *American Journal of Clinical Nutrition*, 53(1), S373-S379.
- Duggal, N. A., Upton, J., Phillips, A. C., & Lord, J. M. (2013). Depression is associated with reduced neutrophil function in hip fracture patients. . *Brain, Behavior & Immunity*, 33, 173-182. doi:10.1016/j.bbi.2013.07.004
- Edwards, K. M., Burns, V. E., Reynolds, T., Carroll, D., Drayson, M., & Ring, C. (2006). Acute stress exposure prior to influenza vaccination enhances antibody response in women. *Brain, Behavior & Immunity*, 20, 159-168.
- Elenkov, I. J., & Chrousos, G. P. (2002). Stress hormones, proinflammatory and antiinflammatory cytokines, and autoimmunity. In M. Cutolo, J. W. J. Bijlsma, R. G. Lahita, A. T. Masi, R. H. Straub, & H. L. Bradlow (Eds.), *Neuroendocrine Immune Basis of the Rheumatic Diseases II, Proceedings* (Vol. 966, pp. 290-303).
- Ershler, W. B., & Keller, E. T. (2000). Age-associated increased interleukin-6 gene expression, late-life diseases, and frailty. *Annual Review of Medicine*, 51, 245-270. doi:10.1146/annurev.med.51.1.245
- Esterling, B. A., Kiecolt-Glaser, J. K., Bodnar, J. C., & Glaser, R. (1994). CHRONIC Stress, social support and persistent alterations in the natural-killer cell response to cytokines in older adults. *Health Psychology*, 13(4), 291-298. doi:10.1037//0278-6133.13.4.291
- Franceschi, C., Capri, M., Monti, D., Giunta, S., Olivieri, F., Sevini, F., . . . Salvioli, S. (2007). Inflammaging and anti-inflammaging: A systemic perspective on aging and longevity emerged from studies in humans. *Mechanisms of Ageing and Development*, 128(1), 92-105. doi:10.1016/j.mad.2006.11.016

Franceschi, C., Capri, M., Monti, D., Giunta, S., Olivieri, F., Sevini, F., . . . Salvioli, S.

(2007). Inflammaging and anti-inflammaging: a systemic perspective on aging and longevity emerged from studies in humans. *Mech Ageing Dev*, 128(1), 92-105.

doi:S0047-6374(06)00249-1 [pii]10.1016/j.mad.2006.11.016

Gallagher, S., Carroll, D., & Phillips, A. C. (2007). Life events stress and antibody response to hepatitis A vaccination. *Brain, Behavior and Immunity*, 21, e13. Abstract.

Gallagher, S., Phillips, A. C., Drayson, M. T., & Carroll, D. (2009a). Caregiving for children with developmental disabilities is associated with a poor antibody response to influenza vaccination. *Psychosomatic Medicine*, 71(3), 341-344.

doi:10.1097/PSY.0b013e31819d1910

Gallagher, S., Phillips, A. C., Drayson, M. T., & Carroll, D. (2009b). Parental caregivers of children with developmental disabilities mount a poor antibody response to pneumococcal vaccination. *Brain, Behavior & Immunity*, 23, 338-346.

doi:10.1016/j.bbi.2008.05.006

Gallagher, S., Phillips, A. C., Evans, P., Der, G., Hunt, K., & Carroll, D. (2008). Caregiving is associated with low secretion rates of immunoglobulin A in saliva. *Brain Behav Immun*, 22(4), 565-572. doi:S0889-1591(07)00309-1 [pii]10.1016/j.bbi.2007.11.007

Gallagher, S., Phillips, A. C., Ferraro, A. J., Drayson, M. T., & Carroll, D. (2008).

Psychosocial factors are associated with the antibody response to both thymus-dependent and thymus-independent vaccines. . *Brain Behaviour & Immunity*, 22, 456-460.

Glaser, R., & Kiecolt-Glaser, J. K. (1997). Chronic stress modulates the virus-specific immune response to latent herpes simplex virus type 1. *Annals of Behavioral Medicine*, 19(2), 78-82.

- Glaser, R., Kiecolt-Glaser, J. K., Bonneau, R. H., Malarkey, W., Kennedy, S., & Hughes, J. (1992). Stress-induced modulation of the immune response to recombinant hepatitis B vaccine. *Psychosomatic Medicine*, 54, 22-29.
- Glaser, R., Kiecolt-Glaser, J. K., Malarkey, W., & Sheridan, J. F. (1998). The influence of psychological stress on the immune response to vaccines. *Annals New York Academy of Sciences*, 840, 649-655.
- Glaser, R., Kiecolt-Glaser, P. K., Marucha, P. T., MacCallum, R. C., Laskowski, B. F., & Malarkey, W. B. (1999). Stress-related changes in proinflammatory cytokine production in wounds. *Archives of General Psychiatry*, 56(5), 450-456.
doi:10.1001/archpsyc.56.5.450
- Glaser, R., MacCallum, R. C., Laskowski, B. F., Malarkey, W. B., Sheridan, J. F., & Kiecolt-Glaser, J. K. (2001). Evidence for a shift in the Th-1 to Th-2 cytokine response associated with chronic stress and aging (vol 56, pg 477, 2001). *Journals of Gerontology Series a-Biological Sciences and Medical Sciences*, 56(11), M673-M673.
- Glaser, R., Sheridan, J., Malarkey, W. B., MacCallum, R. C., & Kiecolt-Glaser, J. K. (2000a). Chronic stress modulates the immune response to a pneumococcal pneumonia vaccine. *Psychosomatic Medicine*, 62(6), 804-807.
- Glaser, R., Sheridan, J. F., Malarkey, W., MacCallum, R. C., & Kiecolt-Glaser, J. K. (2000b). Chronic stress modulates the immune response to a pneumococcal pneumonia vaccine. *Psychosomatic Medicine*, 62, 804-807.
- Goodkin, K., Feaster, D. J., Tuttle, R., Blaney, N. T., Kumar, M., Baum, M. K., . . . Fletcher, M. A. (1996). Bereavement is associated with time-dependent decrements in cellular immune function in asymptomatic human immunodeficiency virus type-1 seropositive homosexual men. *Clinical Diagnostics and Laboratory Immunology*, 3, 109-118.

- Gordon, H. S., & Rosenthal, G. E. (1995). Impact of marital status on outcomes in hospitalized patients. Evidence from an academic medical center. *Archives of Internal Medicine*, 155(22), 2465-2471.
- Gouin, J. P., Hantsoo, L., & Kiecolt-Glaser, J. K. (2008). Immune Dysregulation and Chronic Stress among Older Adults: A Review. *Neuroimmunomodulation*, 15(4-6), 251-259. doi:10.1159/000156468
- Graham, J. E., Christian, L. M., & Kiecolt-Glaser, J. K. (2006). Stress, age, and immune function: toward a lifespan approach. *Journal of Behavioral Medicine*, 29(4), 389-400.
- Griffin, J. E., & Ojeda, S. R. (2004). *Textbook of endocrine physiology*. New York: Oxford University Press.
- Hazeldine, J., Hampson, P., & Lord, J. M. (2012). Reduced release and binding of perforin at the immunological synapse underlies the age-related decline in natural killer cell cytotoxicity. *Aging Cell*, 1-9.
- House, J. S., Landis, K. R., & Umberson, D. (1988). Social relationships and health. *Science*, 241(4865), 540-545.
- House, J. S., Robbins, C., & Metzner, H. L. (1982). The association of social relationships and activities with mortality: prospective evidence from the Tecumseh Community Health Study. *American Journal of Epidemiology*, 116(1), 123-140.
- Irwin, M., Daniels, M., Smith, T. L., Bloom, E., & Weiner, H. (1987). Impaired natural killer cell activity during bereavement. *Brain Behav Immun*, 1(1), 98-104. doi:[http://dx.doi.org/10.1016/0889-1591\(87\)90011-0](http://dx.doi.org/10.1016/0889-1591(87)90011-0)
- Johnson, N. J., Backlund, E., Sorlie, P. D., & Loveless, C. A. (2000). Marital status and mortality: the national longitudinal mortality study. *Annals of Epidemiology*, 10(4), 224-238.

- Kaplan, G. A., Salonen, J. T., Cohen, R. D., Brand, R. J., Syme, S. L., & Puska, P. (1988). Social connections and mortality from all causes and from cardiovascular disease: prospective evidence from eastern Finland. *American Journal of Epidemiology*, 128(2), 370-380.
- Kemeny, M. E., Weiner, H., Duran, R., Taylor, S. E., Visscher, B., & Fahey, J. L. (1995). Immune system changes after the death of a partner in HIV-positive gay men. *Psychosom Med*, 57(6), 547-554.
- Khanfer, R., Lord, J. M., & Phillips, A. C. (2011). Neutrophil function and cortisol:DHEAS ratio in bereaved older adults. *Brain Behavior and Immunity*, 25(6), 1182-1186. doi:10.1016/j.bbi.2011.03.008
- Kiecolt-Glaser, J. K., Fisher, L. D., Ogrocki, P., Stout, J. C., Speicher, C. E., & Glaser, R. (1987). Marital quality, marital disruption, and immune function. *Psychosomatic Medicine*, 49(1), 13-34.
- Kiecolt-Glaser, J. K., Glaser, R., Cacioppo, J. T., MacCallum, R. C., Snyder-Smith, M., Kim, C., & Malarkey, W. B. (1997). Marital conflict in older adults: endocrinological and immunological correlates. *Psychosomatic Medicine*, 59(4), 339-349.
- Kiecolt-Glaser, J. K., Glaser, R., Gravenstein, S., Malarkey, W. B., & Sheridan, J. (1996). Chronic stress alters the immune response to influenza virus vaccine in older adults. *Proceedings of the National Academy of Sciences of the United States of America*, 93(7), 3043-3047. doi:10.1073/pnas.93.7.3043
- Kiecolt-Glaser, J. K., Glaser, R., Gravenstein, S., Malarkey, W. B., & Sheridan, J. (1996). Chronic stress alters the immune response to influenza virus vaccine in older adults. *Proceedings of the National Academy of Science. USA*, 93, 3043-3047.

- Kiecolt-Glaser, J. K., Kennedy, S., Malkoff, S., Fisher, L., Speicher, C. E., & Glaser, R. (1988). Marital discord and immunity in males. *Psychosomatic Medicine*, 50(3), 213-229.
- Kiecolt-Glaser, J. K., Malarkey, W., Chee, M., Newton, T., Cacioppo, J. T., Mao, H. Y., & Glaser, R. (1993). Negative behavior during marital conflict is associated with immunological down-regulation. *Psychosomatic Medicine*, 55, 395-409.
- Kiecolt-Glaser, J. K., Marucha, P. T., Malarkey, W. B., Mercado, A. M., & Glaser, R. (1995). Slowing of wound-healing by psychological stress. *The Lancet*, 346(8984), 1194-1196.
- Kiecolt-Glaser, J. K., & Newton, T. L. (2001). Marriage and health: his and hers. *Psychological Bulletin*, 127(4), 472-503.
- Kiecolt-Glaser, J. K., Speicher, C. E., Glaser, R., Dura, J. R., & Trask, O. J. (1991). Spousal caregivers of dementia victims - longitudinal changes in immunity and health. *Psychosomatic Medicine*, 53(4), 345-362.
- Kohut, M. L., Cooper, M. M., Nickolaus, M. S., Russell, D. R., & Cunnick, J. E. (2002). Exercise and psychosocial factors modulate immunity to influenza vaccine in elderly individuals. *Journal of Gerontology*, 57, 557-562.
- Lazarus, R. S., & Folkman, S. (1984). *Stress, coping and adaptation*. New York: Springer.
- Lewitus, G. M., & Schwartz, M. (2009). Behavioral immunization: immunity to self-antigens contributes to psychological stress resilience. *Molecular Psychiatry*, 14(5), 532-536.
doi:mp2008103 [pii]
10.1038/mp.2008.103
- McEwen, B. S. (1998). Protective and damaging effects of stress mediators. *New England Journal of Medicine*, 338(3), 171-179.

- Miller, G. E., Dopp, J. M., Myers, H. F., Stevens, S. Y., & Fahey, J. L. (1999). Psychosocial predictors of natural killer cell mobilization during marital conflict. *Health Psychology, 18*(3), 262-271.
- Moynihan, J. A., Larson, M. R., Treanor, J., Duberstein, P. R., Power, A., Shore, B., & Ader, R. (2004). Psychosocial factors and the response to influenza vaccination in older adults. *Psychosomatic Medicine, 66*(6), 950-953.
- Munck, A., Guyre, P. M., & Holbrook, N. J. (1984). Physiological functions of glucocorticoids in stress and their relation to pharmacological actions. *Endocrine Reviews, 5*(1), 25-44.
- Ogata, K., An, E., Shioi, Y., Nakamura, K., Luo, S., Yokose, N., . . . Dan, K. (2001). Association between natural killer cell activity and infection in immunologically normal elderly people. *Clinical and Experimental Immunology, 124*(3), 392-397.
- Orentreich, N., Brind, J. L., Rizer, R. L., & Vogelman, J. H. (1984). Age-changes and sex-differences in serum dehydroepiandrosterone sulphate concentrations throughout adulthood. *Journal of Clinical Endocrinology & Metabolism, 59*(3), 551-555.
- Orentreich, N., Brind, J. L., Vogelman, J. H., Andres, R., & Baldwin, H. (1992). Long-term longitudinal measurements of plasma dehydroepiandrosterone sulfate in normal men. *Journal of Clinical Endocrinology & Metabolism, 75*(4), 1002-1004.
- Pariente, C. M., Carpiello, B., Orru, M. G., Sitzia, R., Piras, A., Farci, A. M. G., . . . Miller, A. H. (1997). Chronic caregiving stress alters peripheral blood immune parameters: The role of age and severity of stress. *Psychotherapy and Psychosomatics, 66*(4), 199-207.
- Phillips, A. C. (2011). Blunted cardiovascular reactivity relates to depression, obesity, and self-reported health. *Biological Psychology, 86*(2), 106-113.
doi:10.1016/j.biopsycho.2010.03.016

- Phillips, A. C., Burns, V. E., Carroll, D., Ring, C., & Drayson, M. (2005). The association between life events, social support and antibody status following thymus-dependent and thymus-independent vaccinations in healthy young adults. *Brain Behavior and Immunity*, 19, 325-333.
- Phillips, A. C., Burns, V. E., & Lord, J. M. (2007). Stress and exercise: getting the balance right for an ageing immune system. *Exercise and Sports Sciences Reviews*, 35, 35-39.
- Phillips, A. C., Carroll, D., Burns, V. E., Ring, C., Macleod, J., & Drayson, M. (2006). Bereavement and marriage are associated with antibody response to influenza vaccination in the elderly. *Brain Behavior and Immunity*, 20(3), 279-289.
doi:10.1016/j.bbi.2005.08.003
- Phillips, A. C., Carroll, D., Evans, P., Bosch, J. A., Clow, A., Hucklebridge, F., & Der, G. (2006). Stressful life events are associated with low secretion rates of immunoglobulin A in saliva in the middle aged and elderly. *Brain Behavior and Immunity*, 20, 191-197.
- Phillips, A. C., Upton, J., Duggal, N. A., Carroll, D., & Lord, J. M. (2013). Depression following hip fracture is associated with increased physical frailty in older adults: the role of the cortisol: dehydroepiandrosterone sulphate ratio. *BMC Geriatr*, 13(1), 60.
doi:1471-2318-13-60 [pii]10.1186/1471-2318-13-60
- Pinquart, M., & Sorensen, S. (2003). Differences between caregivers and noncaregivers in psychological health and physical health: a meta-analysis. *Psychology and Aging*, 18(2), 250-267. doi:<http://dx.doi.org/10.1037/0882-7974.18.2.250>
- Pressman, S. D., Cohen, S., Miller, G. E., Barkin, A., Rabin, B. S., & Treanor, J. J. (2005). Loneliness, social network size, and immune response to influenza vaccination in college freshmen. *Health Psychology*, 24(3), 297-306.

- Rasmussen, L. E. (1991). Gene products of cytomegalovirus and their immunologic significance. In H. M (Ed.), *Cytomegalovirus: biology and infection* (2nd ed.): New York: Plenum Medical Book Company.
- Robles, T. F., & Kiecolt-Glaser, J. K. (2003). The physiology of marriage: pathways to health. *Physiology and Behavior*, 79(3), 409-416.
- Rosengren, A., Orth-Gomer, K., Wedel, H., & Wilhelmsen, L. (1993). Stressful life events, social support, and mortality in men born in 1933. *British Medical Journal*, 307(6912), 1102-1105.
- Sapolsky, R. M. (2007). Robert Sapolsky discusses physiological effects of stress. *Stanford News*. Retrieved from <http://news.stanford.edu/news/2007/march7/sapolskysr-030707.html>.
- Sapolsky, R. M., Romero, L. M., & Munck, A. U. (2000). How do glucocorticoids influence stress responses? Integrating permissive, suppressive, stimulatory, and preparative actions. *Endocrinological Review*, 21(1), 55-89.
- Schleifer, S. J., Keller, S. E., Camerino, M., Thornton, J. C., & Stein, M. (1983). Suppression of lymphocyte stimulation following bereavement. *Jama-Journal of the American Medical Association*, 250(3), 374-377.
- Sedova, L., Berube, J., Gaudet, D., Dumont, M., Tremblay, J., Hamet, P., & Pausova, Z. (2004). Diet-induced obesity delays cardiovascular recovery from stress in spontaneously hypertensive rats. *Obesity Research*, 12(12), 1951-1958.
doi:10.1038/oby.2004.245
- Sheffield, D., McVey, C., & Carroll, D. (1996). Daily events and somatic symptoms: Evidence of a lagged relationship. *British Journal of Medical Psychology*, 69, 267-269.

- Siergist, C. A. (2008). Vaccine immunology. In S. Plotkin, Orenstein, W., & Offit, P. (Ed.), *Vaccines* (pp. 17-36): Saunders, Elsevier.
- Sorrells, S. F., & Sapolsky, R. M. (2007). An inflammatory review of glucocorticoid actions in the CNS. *Brain Behavior and Immunity*, *21*(3), 259-272.
doi:10.1016/j.bbi.2006.11.006
- Stahl, S.T., Arnold, A.M., Chen, J-Y, Anderson, S., Schulz, R. (2016) Mortality After Bereavement: The Role of Cardiovascular Disease and Depression. *Psychosomatic Medicine*, *78*(6), 697–703. doi: 10.1097/PSY.0000000000000317
- Thompson, W. W., Shay, D. K., Weintraub, E., Brammer, L., Cox, N., Anderson, L. J., & Fukuda, K. (2003). Mortality associated with influenza and respiratory syncytial virus in the United States. *Jama-Journal of the American Medical Association*, *289*(2), 179-186. doi:10.1001/jama.289.2.179
- Vedhara, K., Cox, N. K. M., Wilcock, G. K., Perks, P., Hunt, M., Anderson, S., . . . Shanks, N. M. (1999a). Chronic stress in elderly carers of dementia patients and antibody response to influenza vaccination. *The Lancet*, *353*(9153), 627-631.
doi:10.1016/s0140-6736(98)06098-x
- Vedhara, K., Cox, N. K. M., Wilcock, G. K., Perks, P., Hunt, M., Anderson, S., . . . Shanks, N. M. (1999b). Chronic stress in elderly carers of dementia patients and antibody response to influenza vaccination. *The Lancet*, *20*, 627-631.
- Vedhara, K., McDermott, M. P., Evans, T. G., Treanor, J. J., Plummer, S., Tallon, D., . . . Schifitto, G. (2002). Chronic stress in non-elderly caregivers: psychological, endocrine and immune implications. *Journal of Psychosomatic Research*, *53*(6), 1153-1161.

- Verbrugge, L. M. (1979). Marital status and health. *Journal of Marriage and Family*, 41, 267-285.
- Vitlic, A., Khanfer, R., Lord, J. M., Carroll, D., & Phillips, A. C. (2014). Bereavement reduces neutrophil oxidative burst only in older adults: Role of the HPA axis and immunesenescence. *Immunity and Ageing*, 11(1).
- Vitlic, A., Lord, J. M., Taylor, A. E., Arlt, W., Bartlett, D. B., Rossi, A., . . . Phillips, A. C. (2015). Neutrophil function in young and old caregivers. *British Journal of Health Psychology*. doi:10.1111/bjhp.12156
- von Kanel, R., Dimsdale, J. E., Mills, P. J., Ancoli-Israel, S., Patterson, T. L., Mausbach, B. T., & Grant, I. (2006). Effect of Alzheimer caregiving stress and age on frailty markers interleukin-6, C-reactive protein, and D-dimer. *Journals of Gerontology Series a-Biological Sciences and Medical Sciences*, 61(9), 963-969.
- Zisook, S., Shuchter, S. R., Irwin, M., Darko, D. F., Sledge, P., & Resovsky, K. (1994). Bereavement, depression, and immune function. *Psychiatry Research*, 52(1), 1-10.