

## Cortisol excess in patients with primary aldosteronism impacts on left ventricular hypertrophy

Adolf, Christian; Köhler, Anton; Franke, Anna; Lang, Katharina; Riester, Anna; Löw, Anja ; Heinrich, Daniel A.; Bidlingmaier, Martin; Treitl, Marcus; Ladurner, Roland; Beuschlein, Felix; Arlt, Wiebke; Reincke, Martin

DOI:

[doi.org/10.1210/jc.2018-00617](https://doi.org/10.1210/jc.2018-00617)

License:

Other (please specify with Rights Statement)

*Document Version*

Peer reviewed version

*Citation for published version (Harvard):*

Adolf, C, Köhler, A, Franke, A, Lang, K, Riester, A, Löw, A, Heinrich, DA, Bidlingmaier, M, Treitl, M, Ladurner, R, Beuschlein, F, Arlt, W & Reincke, M 2018, 'Cortisol excess in patients with primary aldosteronism impacts on left ventricular hypertrophy', *Journal of Clinical Endocrinology and Metabolism*, vol. 103, no. 12, pp. 4543-4552. <https://doi.org/doi.org/10.1210/jc.2018-00617>

[Link to publication on Research at Birmingham portal](#)

### **Publisher Rights Statement:**

This is a pre-copyedited, author-produced PDF of an article accepted for publication in *Journal of Clinical Endocrinology and Metabolism* following peer review. The version of record [insert complete citation information here] is available online at: xxxxxxx [insert URL that the author will receive upon publication here].

### **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

### **Take down policy**

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

1 **Cortisol excess in patients with primary aldosteronism impacts on left ventricular hypertrophy**

2 Christian Adolf<sup>1\*</sup>, Anton Köhler<sup>2\*</sup>, Anna Franke<sup>1</sup>, Katharina Lang<sup>3,4</sup>, Anna Riestler<sup>1</sup>, Anja Löw<sup>2\*</sup>, Daniel A  
3 Heinrich<sup>1</sup>, Martin Bidlingmaier<sup>1</sup>, Marcus Treitl<sup>5</sup>, Roland Ladurner<sup>6</sup>, Felix Beuschlein<sup>1,7</sup>, Wiebke Arlt<sup>3,4\*</sup>,  
4 Martin Reincke<sup>1\*</sup>.

5 \*CA, AK, WA and MR contributed equally to the study

6 <sup>1</sup>Medizinische Klinik und Poliklinik IV, Klinikum der Universität München, LMU München,  
7 Ziemssenstraße 1, 80336 Munich, Germany.

8 <sup>2</sup>Medizinische Klinik und Poliklinik I, Klinikum der Universität München, LMU München, Germany.

9 <sup>3</sup>Institute of Metabolism and Systems Research, University of Birmingham, Birmingham, B15 2TT United  
10 Kingdom.

11 <sup>4</sup>Centre for Endocrinology, Diabetes and Metabolism, Birmingham Health Partners, Birmingham, B15  
12 2TH, UK.

13 <sup>5</sup>Klinik und Poliklinik für Radiologie, Klinikum der Universität München, LMU München, Munich,  
14 Germany.

15 <sup>6</sup>Klinik für Viszeral- und Endokrine Chirurgie, Klinikum der Universität München, LMU München,  
16 Munich, Germany.

17 <sup>7</sup>Klinik für Endokrinologie, Diabetologie und Klinische Ernährung, Universitätsspital Zürich, Zürich,  
18 Switzerland.

19 **Running title:** Cortisol excess impacts on cardiac structure in PA

20 **Précis:** Left ventricular mass in patients with primary aldosteronism improves after medical and surgical  
21 treatment and is positively correlated with total glucocorticoid excretion.

22 **Word count:** Manuscript: 2959 (without abstract, acknowledgment, references, tables and figures)

23 **Figures:** 2      **Tables:** 4      **Supplementary Table:** 1      **Supplementary Figures:** 4

24 **Corresponding author:**

25 Prof. Dr. Martin. Reincke

26 Medizinische Klinik und Poliklinik IV

27 Klinikum der Universität München

28 Ziemssenstr. 1, 80336 München, Germany

29 Tel.: +49-89-4400-52100 | Fax: +49-89-4400-54428

30 E-Mail: martin.reincke@med.uni-muenchen.de

31 **Disclosure statement:** The authors have nothing to disclose

32 **Keywords:** primary aldosteronism, aldosterone, cortisol, cardiac remodeling, left ventricular hypertrophy

33

34 **Abstract**

35 **Context:** Primary aldosteronism (PA) represents the most frequent form of endocrine hypertension.  
36 Hyperaldosteronism and hypercortisolism both induce excessive left ventricular hypertrophy (LVH)  
37 compared to matched essential hypertensives. In recent studies frequent co-secretion of cortisol and  
38 aldosterone has been reported in PA patients.

39 **Objective:** Our aim was to investigate the impact of cortisol co-secretion on left ventricular hypertrophy  
40 in PA patients. We determined 24-h excretion of mineralocorticoids and glucocorticoids by gas  
41 chromatography-mass spectrometry and assessed cardiac remodeling using echocardiography initially and  
42 one year after initiation of treatment for PA.

43 **Patients:** We included 73 patients from the Munich center of the German Conn's registry; 45 with  
44 unilateral aldosterone-producing adenoma and 28 with bilateral adrenal hyperplasia.

45 **Results:** At the time of diagnosis, 85% of PA patients showed left ventricular hypertrophy according to  
46 left ventricular mass index (LVMI, median 62.4 g/m<sup>2</sup>). LVMI correlated positively with total  
47 glucocorticoid excretion ( $r^2=0.076$ ,  $p=0.018$ ) as well as with tetrahydroaldosterone excretion ( $r^2=0.070$ ,  
48  $p=0.024$ ). Adrenalectomy led to significantly reduced LVMI in aldosterone-producing adenoma ( $p<0.001$ )  
49 while mineralocorticoid receptor antagonist therapy in bilateral adrenal hyperplasia patients reduced  
50 LVMI to a lesser degree ( $p=0.024$ ). In multivariate analysis, the decrease in LVMI was positively  
51 correlated with total glucocorticoid excretion and systolic 24-hour blood pressure, but not with  
52 tetrahydroaldosterone excretion.

53 **Conclusion:** Cortisol excess appears to have an additional impact on cardiac remodeling in patients with  
54 PA. Treatment of PA by either adrenalectomy or mineralocorticoid receptor antagonist improves LVMI.  
55 This effect was most pronounced in patients with high total glucocorticoid excretion.

## 56 **Introduction**

57 Primary aldosteronism (PA) is the most frequent cause of endocrine hypertension affecting about  
58 5-10 % of patients with elevated blood pressure (1). PA was first described in 1955 by Jerome W. Conn  
59 (2) and characterized in its classical form as hypertension, hypokalemia and alkalosis. Patients with  
60 bilateral adrenal hyperplasia (BAH), the most common subtype, require lifelong treatment with a  
61 mineralocorticoid receptor antagonist (MRA), whereas patients with unilateral aldosterone-producing  
62 adenoma (APA) can be cured by adrenalectomy (ADX). Prolonged exposure to elevated aldosterone  
63 concentrations causes cardiac and renal damage independently of blood pressure (3). These changes may  
64 have adverse impact on clinical outcome.

65 One of the targets of aldosterone action are cardiomyocytes which express mineralocorticoid  
66 receptors (MR) (4). Indeed, MR activation has been shown to induce myocardial damage (3), including  
67 diffuse myocardial fibrosis, left ventricular hypertrophy (LVH) and left ventricular dilatation (4). In  
68 echocardiographic studies, LVH was more frequent and more progressive in PA patients in comparison to  
69 matched essential hypertensives (5). Likewise, a number of studies have highlighted an increased risk of  
70 stroke, myocardial infarction (MI) and atrial fibrillation in PA patients (6,7). LVH itself is one of the most  
71 important predictors for major cardiac events and mortality and is associated with an increased risk of  
72 arrhythmia, MI and stroke (8).

73 In a recent study, we identified cortisol co-secretion as a key feature of PA (9) associated with an  
74 adverse metabolic risk phenotype, providing a logical explanation for important comorbidities observed in  
75 patients with primary aldosteronism more aligned with the effects of glucocorticoid rather than  
76 mineralocorticoid excess, such as insulin resistance and type 2 diabetes (10-13), and osteoporosis (14).  
77 Interestingly, pronounced LVH has also been described in patients with Cushing's syndrome (CS), who  
78 are also afflicted by arterial hypertension, impaired glucose tolerance and serum electrolyte imbalance  
79 (15,16). High cortisol levels in patients with CS are associated with an increased mortality rate due to  
80 cardiovascular complications (17).

81           Therefore, we hypothesized that cortisol co-secretion observed in PA might have additional  
82 adverse effects on cardiac function and cardiovascular outcome in PA patients. Thus, our aim was to  
83 investigate the impact of increased cortisol secretion on echocardiographic findings in patients with APA  
84 or BAH.

## 85 **Methods**

### 86 *Patients*

87           During 2008 and 2013 we consecutively enrolled 210 patients with primary aldosteronism at the  
88 Munich center of the German Conn's Registry. In 89 patients a urinary steroid metabolite excretion in  
89 24-h urine was performed and of those 89, 73 patients with either APA (n=45) or BAH (n=28) had a  
90 technically accurate echocardiography examination, which represents the cohort included in this study.  
91 Analysis of urinary steroid metabolite excretion and echocardiography data represented post-hoc analyses.  
92 All patients gave written informed consent, and the protocol of the German Conn's registry was approved  
93 by the ethics committee of the University of Munich. At each visit, patients underwent standardized  
94 clinical phenotyping including collection of anthropometric data and clinical characteristics such as  
95 duration of hypertension and current medication.

96           The diagnostic procedures were performed according to the Endocrine Society Practice Guidelines  
97 (18,19). In short, PA was diagnosed by an elevated plasma aldosterone to renin ratio (ARR; cut-off 10.0  
98 ng/mU, sitting position) and an abnormal confirmatory test (e.g. salt loading test, captopril challenge test).  
99 Antihypertensive medication was stopped (n=5) whenever possible prior to testing. Otherwise it was  
100 replaced by alpha 1-adrenergic receptor (doxazosin) or calcium-channel blockers (verapamil) (n=68).  
101 Subtype differentiation between unilateral and bilateral disease was based on adrenal vein sampling  
102 (AVS). In case of APA only patients who underwent ADX were included in the analysis. BAH patients  
103 were treated with MRA. In most patients, spironolactone was used at a dose of 25-50 mg per day. All  
104 patients were re-evaluated one year after treatment in a standardized fashion.

105

### 106 *Laboratory analysis*

107           Blood samples were drawn in a fasting state in sitting position at 8.00 a.m. Plasma aldosterone  
108 concentration was measured using the radioimmunoassay "aldosterone Coat-a-Count" (Biermann DPC).  
109 Active renin concentration was measured by the Liaison chemiluminescence assay (Diasorin). All other  
110 analyses were performed in our central laboratory using standard methods. To determine urinary steroid

111 excretion, the patients conducted a 24-hour urine collection. Subsequently, gas chromatography-mass  
112 spectrometry (GC-MS) in selected-ion-monitoring (SIM) analysis mode was performed to determine the  
113 urinary steroid metabolite excretion as described previously (9), allowing the quantification of 40 different  
114 steroid metabolites, including 3 $\alpha$ ,5 $\beta$ -tetrahydroaldosterone (THAldo), the major mineralocorticoid  
115 metabolite. Total glucocorticoid excretion was calculated as the sum of quantified metabolites of cortisol  
116 and cortisone, comprising tetrahydrocortisol, 5 $\alpha$ -tetrahydrocortisol, tetrahydrocortisone,  $\alpha$ - and  $\beta$ -cortol,  
117  $\alpha$ - and  $\beta$ -cortolone, 6 $\beta$ -hydroxycortisol, and urinary cortisol and cortisone (9).

118

### 119 *Cardiac ultrasound examination*

120         Comprehensive echocardiographic examination was conducted by experienced sonographers from  
121 the department of internal medicine I (cardiology) from the Ludwig-Maximilians-Universität München.  
122 The sonographers were blinded with regard to diagnosis and clinical details. Commercially available high-  
123 quality ultrasound systems were used (GE Healthcare Vivid 7, Philips iE 33). The patients were lying  
124 down in the left lateral decubitus position. Echocardiography included two-dimensional, M-Mode and  
125 Doppler ultrasound recordings. Images were obtained in the parasternal (long and short axis) and apical  
126 views. The left ventricular internal dimension (LVID), interventricular septum (IVS), posterior wall  
127 thickness (PWT) and left atrial dimension in diastole (LAd) were measured via parasternal long axis view.  
128 Echocardiographic parameters were measured according to the recommendation of the American Society  
129 of Echocardiography (20,21). Echocardiography-based left ventricular mass (LVM) estimation is  
130 generally calculated as the difference between epicardium delimited volume and left ventricular chamber  
131 volume multiplied by an estimate of myocardial density (22). The LVM was calculated by the Penn  
132 Convention formula:  $LVM=1.04 \cdot [(LVIDd+PWTd+IVSd]^3-[LVIDd]^3)-13.6g$  (23,24). Obesity is  
133 independently associated to LVH (25). Because of higher average values of BMI in our study cohort,  
134 LVH was determined as the left ventricular mass index (LVMI). LVM was indexed by height<sup>2.7</sup> to obtain  
135 LVMI. By using the LVMI we minimized the interference of obesity in LVM estimation (26). LVH was  
136 prospectively defined as a value of LVMI  $\geq 50 g/m^{2.7}$  in males and  $\geq 47g/m^{2.7}$  in females (27). Relative wall



137 thickness (RWT) was calculated according to the following equation:  $RWT=(IVSd + PWT)/LVIDd$ . LVH  
138 was separated in concentric hypertrophy with  $RWT \geq 0.42$  and eccentric hypertrophy with  $RWT < 0.42$   
139 (21). Normal LVMI values and RWT values  $\geq 0.42$  were defined as concentric remodeling (28).

140

#### 141 *Statistical analysis*

142 All values are expressed as median and 25th and 75th percentile if not mentioned otherwise.  
143 Within-group changes from baseline to follow-up were calculated by Wilcoxon signed-rank test.  
144 Spearman's Rank Order was used to perform bivariate correlation analysis. Stepwise multiple regression  
145 analysis was used for multivariate analysis. Two-tailed probability values of  $< 5\%$  were considered to be  
146 statistically significant. Statistical analysis was performed using standard statistical software (SPSS 23,  
147 IBM, Chicago, Illinois).

148 **Results**

149 *Patient characteristics*

150 Clinical characteristics of the total cohort of 73 patients with PA are shown in **Table 1**, the  
151 comparison of APA and BAH subgroups in **Table 2**. APA and BAH were diagnosed in 45 and 28  
152 patients, respectively. As expected, at diagnosis, patients with APA had higher plasma aldosterone  
153 concentration ( $p=0.001$ ), higher urinary THAldo concentrations ( $p=0.001$ ) and lower potassium levels  
154 ( $p<0.001$ ), and more pronounced renal impairment according to globular filtration rate (GFR;  $p=0.043$ ).  
155 Total glucocorticoid excretion did not differ between groups ( $p=0.184$ ).

156 Patient characteristics one year after initiation of treatment (MRA treatment in BAH, ADX in  
157 APA) are listed in **Tables 1+2**. Systolic and diastolic blood pressure and serum potassium levels  
158 normalized in both subgroups. As expected, defined daily doses (DDD) of antihypertensive medication  
159 decreased significantly in the APA group, whereas in the BAH group there was only a trend towards  
160 lower DDDs. BMI remained stable in both groups and both groups had a decline in renal function.  
161 Triglyceride levels increased significantly in both groups, whereas HDL cholesterol levels decreased only  
162 in the BAH group ( $p=0.006$ ). Pro-BNP, a potential indicator of cardiac preload, improved in both groups.  
163 The decline was not due to optimized treatment for heart failure. In fact, DDDs of ACE inhibitors/  
164 angiotensin II receptor blockers (1.4 vs. 1.1;  $p=0.002$ ), beta blockers (0.4 vs. 0.2;  $p=0.001$ ) and diuretics  
165 (0.4 vs. 0.2;  $p<0.001$ ) have decreased significantly at follow-up.

166

167 *Echocardiographic findings in patients with unilateral aldosteronism and bilateral hyperplasia*

168 **Table 3** summarizes echocardiographic geometric characteristics in APA and BAH patients.  
169 LVMI was elevated, the IVS was thickened and the LA was enlarged in both groups at baseline. The  
170 overall prevalence of LVH was 85% before initiation of treatment, compared to 66% at follow-up. There  
171 was a shift from eccentric and concentric hypertrophy towards normal left ventricular geometry (**Suppl.**  
172 **Table 1**). As expected, LVMI improved in both APA ( $p<0.001$ ) and BAH ( $p=0.024$ ) patients with  
173 treatment (**Fig. 1**). The reduction of LVMI ( $\Delta$ LVMI) was numerically greater in APA patients with a

174 significant decrease of LVIDd ( $p < 0.001$ ), PWTd ( $p = 0.020$ ) and IVSd ( $p = 0.001$ ). In BAH patients only  
175 LVIDd ( $p = 0.001$ ) improved significantly.

176

177 ***THAldo, total glucocorticoid excretion, urinary sodium excretion and left ventricular structure***

178 As reported previously (9), both THAldo excretion and total glucocorticoid excretion were  
179 increased in PA patients (**Table 1**). In univariate analysis THAldo ( $p = 0.024$ ), urinary sodium excretion  
180 ( $p = 0.044$ ) and total glucocorticoid excretion ( $p = 0.018$ ) correlated with LVMI at time of diagnosis (**Suppl.**  
181 **Fig. 1+2**). In contrast, the relative changes in LVMI in response to treatment ( $\Delta$ LVMI) correlated with  
182 total glucocorticoid excretion ( $p = 0.042$ ), but neither with THAldo ( $p = 0.776$ ) nor with urinary sodium  
183 excretion ( $p = 0.214$ ) (**Suppl. Fig. 1+2**). Moreover, when arbitrarily dividing PA patients into low and high  
184 steroid secretors, according to whether their THAldo and total glucocorticoid excretion was below or  
185 above the median, high total glucocorticoid excretion, but not THAldo, predicted a reduction in LVMI  
186 (**Fig. 2**). Similarly, in multivariate analyses, total glucocorticoid excretion and 24-h systolic blood pressure  
187 were strong predictors of left ventricular geometry changes, whereas THAldo did not have a significant  
188 effect in this model (**Table 4**). One year after treatment we could detect a significant decrease of total  
189 glucocorticoid excretion in APA patients. In accordance with our findings in APA patients with complete  
190 biochemical remission a higher decrease in total glucocorticoid excretion was followed by a more distinct  
191 reduction of LVMI at follow-up ( $r^2 = 0.138$ ,  $p = 0.023$ ).

192

193

194 **Discussion**

195 This is the first study to evaluate the impact of glucocorticoid co-secretion on LVH in patients  
196 with PA. PA is characterized by increased aldosterone secretion, but in recent years a relevant cortisol co-  
197 secretion has been recognized in several case reports and small case series (29-32). We have recently  
198 reported that increased secretion of glucocorticoids is a major biochemical feature in a large proportion of  
199 patients with PA, whereas clinically overt signs of Cushing's syndrome are rare (9). We identified cortisol  
200 co-secretion in a substantial percentage of patients with APA and BAH and reported its association with  
201 parameters of the metabolic syndrome, such as BMI, HOMA-IR, and plasma lipids. Moreover, we found  
202 that after unilateral ADX glucocorticoid secretion normalized, followed by postoperative tertiary adrenal  
203 insufficiency in one third of patients. Based on this observation, we wondered whether total glucocorticoid  
204 excretion might also impact on the cardiac phenotype in PA.

205 Previous echocardiographic evaluations have already demonstrated excess LVM and more  
206 frequent LVH in patients with PA (5,33,34). Our patients with APA and BAH frequently had LVH  
207 according to LVMI; the most common left ventricular adaptation was eccentric hypertrophy. It has been  
208 described before, that high plasma aldosterone concentration results in eccentric changes in LV geometry  
209 (33,35) and that LVM is reduced by either MRA or ADX in patients with PA (36-39). A meta-analysis of  
210 four studies with an average follow-up of four years reported comparable effects for both treatment  
211 strategies (40). In line with this literature, our patients responded to treatment with a significant reduction  
212 in LVMI and an increase in the percentage of normal LV geometry.

213 At diagnosis, LVMI in the APA group was higher than in the BAH group. At follow-up, our APA  
214 patients showed a trend towards a more distinct reduction in LVMI compared to BAH patients without  
215 reaching statistical significance. Inhibition of MR-mediated aldosterone effects by specific medical  
216 treatment is an explanation for the comparable effects of MRA and ADX in LVM reduction. Our findings  
217 mirror previous studies reporting that the response of LVM reduction in adrenalectomized PA patients  
218 could occur earlier than in PA patients treated with MRA (36). Persistent hyperaldosteronemia with  
219 possible persistence of non-genomic effects of aldosterone has been proposed to potentially explain why

220 MRA treatment takes longer to show comparable effects than surgery (41). The results of our study  
221 suppose an impact of cortisol co-secretion on LVH and LV geometry. Total glucocorticoid excretion was  
222 positively correlated with baseline LVMI and  $\Delta$ LVMI at follow-up. Therefore, patients with higher total  
223 glucocorticoid excretion showed higher decreases in LVMI after one year of treatment. In line with these  
224 findings, the decrease of glucocorticoid excretion correlated with the improvement of LVMI in our APA  
225 patients with biochemical remission.

226         Chronic cortisol hypersecretion, e. g. in patients with CS, is known to cause amongst others,  
227 truncal obesity, arterial hypertension, impaired glucose tolerance and dyslipidemia (42). However, also a  
228 variety of alterations in cardiac structure and function have been reported, including increased LVM,  
229 increased interventricular septum thickness and concentric hypertrophy or remodeling (15,16,43,44). CS is  
230 known to be associated with elevated cardiovascular morbidity and mortality (45). Cardiac dysfunction  
231 itself represents one of the most important cardiovascular complications affecting mortality. The two  
232 major forms of cardiac dysfunction in CS are cardiac hypertrophy and congestive heart failure (46,47). It  
233 is thought that the main cortisol effects leading to LVH are hypertension, potentiation of noradrenalin and  
234 angiotensin II responsiveness of the cardiomyocytes and cardiomyocyte proliferation and hypertrophy  
235 (48). MRs have similar affinity for aldosterone and cortisol. Glucocorticoid excess impairs conversion of  
236 cortisol to its MR-inactive cortisone by  $11\beta$ -hydroxysteroid dehydrogenase type 2 ( $11\beta$ -HSDS2) in  
237 classical aldosterone target tissues as the distal nephron, leading to glucocorticoid-mediated  
238 mineralocorticoid effects (49,50). In the cardiomyocytes,  $11\beta$ -HSDS2 is not expressed at relevant levels.  
239 Therefore, in physiologic circumstances the MR in cardiomyocytes is mostly occupied by cortisol, which  
240 circulates in much higher concentrations than aldosterone. In the event of cardiac tissue damage, cortisol  
241 acts as an MR agonist as shown by Mihailidou et al. in ischemia-reperfusion studies in rat heart  
242 Langendorff preparations (49). This effect can be blocked by spironolactone but not by the  
243 glucocorticoid/progesterone antagonist RU486. However, the abnormalities in LV structure and function  
244 have been reported to ameliorate (44) or even to be reversible upon normalisation of glucocorticoid excess  
245 (51).

246 Therefore, treatment with ADX or MRA should both be effective against the MR mediated effects  
247 of glucocorticoid excess on myocardial tissue. To our knowledge no glucocorticoid dependent MR-  
248 mediated effects on lipid and glucose metabolism have been reported so far. Therefore, MRA treatment  
249 could be inferior in this regard when increased glucocorticoid secretion in BAH remains unblocked during  
250 MRA treatment in glucocorticoid target organs, such as the endocrine pancreas, abdominal fat tissue and  
251 the liver. In the current study, however, the improvement of glucose metabolism following adrenalectomy  
252 was minimal, but a clinical relevant improvement in glucose homeostasis has been shown previously in  
253 several cohorts (52).

254 We acknowledge the limitation that steroid metabolite analysis and echocardiography were post  
255 hoc investigations and that our study was neither powered nor planned to examine differences between the  
256 two treatment strategies of PA, ADX and MRA treatment. However, following one year of MRA  
257 treatment, 54% of our BAH patients still had concentric remodeling or concentric hypertrophy as  
258 compared to 42% in APA patients. Therefore, this finding might be explained by ongoing glucocorticoid  
259 effects in BAH patients or inadequate MRA dosage. Unfortunately, we have only limited data for follow-  
260 up of urinary glucocorticoid excretion in BAH patients. In addition, one year of follow-up might be too  
261 short to detect full treatment effects of MRA treatment. Therefore, additional studies are required to  
262 address this issue.

263 It is known that patients with essential hypertension obtain LVM reduction by treatment with  
264 either ACE inhibitor or MRA and that a combination has an additive effect (53). Blood pressure levels and  
265 antihypertensive medication according to DDD's were reduced in both groups one year after start of  
266 treatment. Although we detected comparable effects on 24-h blood pressure levels, the DDD's showed  
267 numerically a greater reduction in adrenalectomized APA patients. A contributing factor can be the rather  
268 low dose of spironolactone of an average of 42 mg/d, in accordance with Endocrine Society Practice  
269 Guidelines which can be explained by clinical side effects including gynecomastia preventing further dose  
270 escalation. Similar to previous studies, most of the patients after ADX (58%) still needed  
271 antihypertensives because of residual hypertension (36,54).

272 The strengths of our study include the prospective standardized collection of all data and  
273 biomaterial within the context of the German Conn's registry, the homogeneously characterized study  
274 population, and the subtyping of all patients by adrenal vein sampling. A major limitation of the study is  
275 the very limited follow-up data for urinary steroid excretion (n=5) for BAH patients. Another limitation is  
276 the examination of left ventricular geometry by echocardiography. The main limitation of this technique is  
277 an inappropriate acoustic window, limiting patient inclusion to suitable candidates. Secondly, left  
278 ventricular parameters for estimation of LVM were generated by M-mode measurement. This is less  
279 accurate compared to real time three-dimensional echocardiography or 3D imaging by MRI (55). Thirdly,  
280 echocardiographic examinations in this study were performed by different experienced investigators,  
281 which could have had an influence on wall thicknesses estimation and therefore LVM and LVMI  
282 estimation. However, the investigators were blinded with regard to the underlying cause of disease, which  
283 we consider a strength of the study approach.

284

## 285 **Conclusions**

286 In this study, we investigated the effects of increased glucocorticoid secretion in 73 PA patients  
287 (45 APA, 28 BAH) on cardiac geometry. Our data show that total glucocorticoid excretion is associated  
288 with LVH independent of mineralocorticoid excess. Moreover, high total glucocorticoid excretion, but not  
289 THAldo predicted the  $\Delta$ LVMI after adrenalectomy in APA patients. In summary, our data suggest a  
290 relevant role of glucocorticoid secretion in PA on LV geometry, pointing out the relevance of cortisol co-  
291 secretion in the context of PA.

292

293 **Acknowledgements**

294 This work has been supported by the Else Kröner-Fresenius Stiftung (German Conn's Registry-Else-  
295 Kröner Hyperaldosteronism Registry; 2013\_A182 and 2015\_A171, to M.R.), the Medical Research  
296 Council UK (Strategic Biomarker Grant G0801473, to W.A.; Clinical Research Training Fellowship  
297 MR/R002339/1, to K.L.), the Deutsche Forschungsgemeinschaft (RE 752/20–1, to M.R., and BE2177/13-  
298 1, to F.B), the European Research Council (grant number 694913 [PAPA] to M.R.), and the European  
299 Commission Horizon 2020 Program under grant agreement 633983 (ENSAT-HT, to W.A., M.R., and  
300 F.B.).

301 The study was only feasible due to the support of our clinical PA team (Nina Nirschl, Lisa Sturm,  
302 Friederike Konrad), the Endocrine laboratory teams in Munich and Birmingham and our collaborators  
303 from the European Network for the Study of Adrenal Tumors.

304

305

306

307

308

309

310

311

312

313

314

315

316

317

318



319 **Table 1. Baseline and 1 year follow-up characteristics of all patients with primary aldosteronism.**

320 Data are given as median and 25<sup>th</sup> and 75<sup>th</sup> percentile in square brackets. Asterisk indicates significance.

321 Comparisons were performed by Wilcoxon signed rank test.

Patient characteristics (n=73)	n	baseline	after treatment	p
<b>Gender</b> [f/m]	73	28/45	--	n.c.
<b>Age</b> [years]	73	53 [43; 58]	--	n.c.
<b>Duration of hypertension</b> [months]	73	129 [52; 267]	--	n.c.
<b>Antihypertensive agents</b> [DDD]	73	3.0 [2.0; 4.3]	2.0 [0.1; 3.7]	0.002*
<b>BMI</b> [kg/m <sup>2</sup> ]	73	27.7 [24.9; 30.9]	27.5 [24.9; 30.3]	0.705
<b>Aldosterone</b> [ng/l]	73	219.0 [148.0; 346.5]	124.5 [39.8; 198.8]	0.003*
<b>Plasma renin</b> [mU/l]	73	4.9 [2.2; 13.4]	18.9 [8.9; 44.6]	<0.001*
<b>Serum potassium</b> [mmol/l]	73	3.4 [3.0; 3.7]	4.2 [3.9; 4.5]0	<0.001*
<b>Serum creatinine</b> [mg/dl]	73	0.8 [0.7; 1.0]	1.0 [0.8; 1.2]	<0.001*
<b>Serum urea</b> [mg/dl]	73	13 [11; 16]	19 [15; 27]	<0.001*
<b>GFR</b> [ml/min/1.73 m <sup>2</sup> ]	73	88.4 [72.3; 100.8]	73.9 [56.3; 85.9]	<0.001*
<b>SBP</b> [mmHg]	73	151 [138; 174]	133 [124; 142]	<0.001*
<b>DBP</b> [mmHg]	73	93 [86; 103]	87 [80; 93]	<0.001*
<b>24h-SBP</b> [mmHg]	57	144 [134; 157]	132 [125; 140]	<0.001*
<b>24h-DBP</b> [mmHg]	57	91 [84; 98]	84 [77; 88]	<0.001*
<b>FPG</b> [mg/dl]	73	102 [93; 113]	99 [91; 107]	0.033*
<b>HDL-C</b> [mg/dl]	70	56 [47; 70]	49 [40; 62]	<0.001*
<b>LDL-C</b> [mg/dl]	70	124 [92; 151]	126 [100; 142]	0.740
<b>Triglycerides</b> [mg/dl]	70	105 [72; 134]	114 [86; 178]	<0.001*
<b>Total cholesterol</b> [mg/dl]	70	197 [174; 227]	200 [174; 219]	0.995
<b>Diabetes mellitus</b> [n]	73	8 (11%)	8 (11%)	n.c.
<b>Total glucocorticoid excretion</b> [μg/24h]; median of healthy controls: 8262 [6380; 11044]	73	11807 [8270; 15266]	--	n.c.
	44	12772 [9000; 16131]§	9458 [5578; 13460]§	<0.001*
<b>Tetrahydroaldosterone excretion</b> [μg/24h]; median of healthy controls: 30 [22; 44]	73	82 [52; 128]	--	n.c.
	44	88 [57; 131]§	21 [14; 39]§	<0.001*

322 Abbreviations: FPG: fasting plasma glucose; DDD: defined daily dose; SBP: systolic blood pressure;

323 DBP: diastolic blood pressure; §: dataset of 44 patients with complete follow-up data, n.c.: not calculated.

324 **Table 2. Baseline and 1 year follow-up characteristics of patients with primary aldosteronism according to subtype.**

325 Data are given as median and 25<sup>th</sup> and 75<sup>th</sup> percentile in square brackets. Asterisk indicates significance. Comparisons were performed by  
 326 Wilcoxon signed-rank-test and by McNemar-test. Differences between baseline values of both group were marked with # for p<0.05 and ## for  
 327 p<0.0001.

Patient characteristics	Aldosterone producing adenoma (n=45)		p	Bilateral adrenal hyperplasia (n=28)		p
	baseline	after ADX		baseline	after MRA	
Time of assessment						
Age [years]	54 [46; 60]	--	n.c.	49 [41; 58]	--	n.c.
Sex [f/m]	17/28	--	n.c.	11/17	--	n.c.
Duration of hypertension [months]	130 [55; 250]	--	n.c.	130 [35; 283]	--	n.c.
BMI [kg/m <sup>2</sup> ]	27.7 [24.9; 32.1]	28.1 [25.2; 30.3]	0.803	27.6 [25.0; 30.7]	27.1 [24.3; 30.4]	0.323
Aldosterone [ng/l]	242.0 [181.6; 427.9]	53.0 [35.0; 129.5]	<0.001*	159.5 [110.5; 247.5]#	216.0 [160.0; 388.6]	<0.001*
Plasma renin [mU/l]	5.5 [2.1; 12.1]	17.0 [9.1 45.1]	<0.001*	4.7 [2.5; 15.8]	19.9 [7.7; 45.0]	<0.001*
Serum potassium [mmol/l]	3.2 [2.8; 3.5]	4.2 [3.9; 4.5]	<0.001*	3.6 [3.2; 3.8]##	4.2 [3.9; 4.5]	<0.001*
Serum creatinine [mg/dl]	0.8 [0.8; 1.0]	1.1 [0.9; 1.3]	<0.001*	0.8 [0.7; 0.9]	0.8 [0.7; 1.1]	0.016*
Serum urea [mg/dl]	13 [11; 16]	22 [16; 29]	<0.001*	13 [10; 17]	16 [13; 24]	0.003*
GFR [ml/min/1.73 m <sup>2</sup> ]	79.4 [69.9; 99.6]	68.8 [54.5; 77.8]	<0.001*	93.9 [84.1; 104.3]#	81.3 [71.5; 105.9]	0.031*
SBP [mmHg]	152 [139; 172]	133 [126; 144]	<0.001*	146 [134; 175]	132 [123; 137]	0.005*
DBP [mmHg]	93 [87; 105]	87 [80; 94]	0.001*	92 [86; 101]	88 [79; 93]	0.106
24-SBP [mmHg] †	149 [139; 157]	132 [125; 140]	<0.001*	144 [131; 155]	132 [123; 139]	0.032*
24-DBP [mmHg] †	91 [86; 99]	83 [77; 87]	<0.001*	91 [83; 98]	84 [76; 88]	0.007*
FPG [mg/dl]	102 [94; 112]	99 [91; 106]	0.043*	101 [91; 119]	100 [90; 112]	0.692
Diabetes mellitus [n]	5 (11%)	5 (11%)	n.c.	3 (11%)	3 (11%)	n.c.
HDL-C [mg/dl] †	54 [47; 69]	49 [42; 64]	0.076	56 [47; 72]	51 [39; 62]	0.006*
LDL-C [mg/dl] †	133 [92; 151]	126 [97; 145]	0.861	123 [92; 152]	126 [101; 142]	0.409
Triglycerides [mg/dl] †	103 [66; 134]	113 [79; 169]	0.001*	107 [80; 135]	117 [95; 190]	0.005*

<b>Total cholesterol</b> [mg/dl] †	203 [174; 231]	200 [170; 221]	0.480	192 [173; 221]	197 [175; 219]	0.402
<b>Antihypertensive agents</b> [DDD]	3.5 [2.0; 4.7]	1.5 [.0; 3.2]	<0.001*	3.0 [1.6; 3.5]	2.7 [1.0; 4.6]	0.848
<b>Tetrahydroaldosterone excretion</b> [µg/24h]	95 [63; 140]		n.c.	55 [40; 97]#		n.c.
	95 [62; 147]§	18 [13; 31]§	<0.001*	54 [39; 77]§§	60 [38; 70]§§	n.c.
<b>Total glucocorticoid excretion</b> [µg/24h]	12980 [9200; 15266]		n.c.	9908 [6328; 16109]		n.c.
	12563 [9015; 15516]§	9304 [5522; 11300]§	<0.001*	19943 [8573; 24897]§§	17073 [6603; 17581]§§	n.c.
<b>Total sodium excretion</b> [mmol/24h]	204 [136; 238]	164 [131; 257]	0.200	186 [150; 224]	196 [115; 240]	0.713
<b>Pro-BNP</b> [pg/ml]	117 [52; 297]	68 [32; 124]	<0.001*	96 [61; 167]	65 [41; 117]	0.023*

328 Abbreviations: ADX: adrenalectomy; MRA: mineralocorticoid receptor antagonist treatment; FPG: fasting plasma glucose; DDD: defined daily  
329 dose; SBP: systolic blood pressure; DBP: diastolic blood pressure; §: dataset of 44 patients with complete follow-up data, §§: dataset of 5 patients  
330 with complete follow-up data, n.c.: not calculated, †: Due to incomplete data the calculations for 24-SBP and 24-DBP (APA n=35, BAH n=22),  
331 and HDL-C, LDL-C, Triglycerides, Total cholesterol (APA n=43, BAH n=27) were performed with a reduced number of patients as listed in  
332 brackets.

333 **Table 3. Echocardiographic characteristics of patients with primary aldosteronism according to subtype.**

334 Data are given as median and 25th and 75th percentile in square brackets. Asterisk indicates significance. Comparisons were performed by  
 335 Wilcoxon signed-rank-test.

Left ventricular parameters	Aldosterone producing adenoma (n=45)		p	Bilateral adrenal hyperplasia (n=28)		p
	baseline	after ADX		baseline	after MRA	
<b>LVMI</b> [g/m <sup>2.7</sup> ]	64.6 [54.7; 71.4]	56.5 [42.7; 63.6]	<0.001*	57.8 [50.3; 70.1]	53.5 [42.1; 65.2]	0.024*
<b>ΔLVMI</b> [g/m <sup>2.7</sup> ]		8.0 [1.5;18.3]	n.c.		6.4 [-2.0; 14.4]	n.c.
<b>LVIDd</b> [mm]	52 [49; 56]	50 [47; 53]	<0.001*	52 [47; 55]	49 [44; 53]	0.001*
<b>LVIDs</b> [mm]	31 [29; 36]	30 [27; 33]	0.128	31 [28; 34]	29 [26; 34]	0.083
<b>PWTd</b> [mm]	10 [9; 12]	10 [9; 11]	0.020*	10 [8; 11]	10 [9; 12]	0.319
<b>IVSd</b> [mm]	12 [11; 13]	11 [10; 13]	0.001*	12 [10; 13]	12 [10; 13]	0.072
<b>RWTd</b> [cm]	0.40 [0.35; 0.47]	0.40 [0.36; 0.46]	0.775	0.39 [0.33; 0.44]	0.43 [0.38; 0.50]	0.014*
<b>LAd</b> [mm]	42 [38; 46]	39 [36; 43]	<0.001*	42 [36; 47]	41 [33; 46]	0.090

336 Abbreviations: ADX: adrenalectomy; MRA: mineralocorticoid receptor antagonist treatment; LVMI: left ventricular mass indexed for height to the  
 337 2.7 power; ΔLVMI: reduction of left ventricular mass indexed for height to the 2.7 power after treatment; LVM: left ventricular mass, LVIDd: left  
 338 ventricular internal dimension in diastole; LVIDs: left ventricular internal dimension in systole; PWTd: posterior wall thickness in diastole; IVSd:  
 339 interventricular septum thickness in diastole; RWTd: relative wall thickness in diastole; LAd: left atrial internal dimension in diastole, n.c.: not  
 340 calculated.

341 **Table 4. Uni- and multivariate analyses of the associations with echocardiographic parameters in all**  
 342 **patients with primary aldosteronism.**

343 Data are given as p values. Asterisk indicates significance. Correlation analysis was performed using  
 344 Spearman's Rank-Order test and stepwise multiple regression analysis.

	<b>THAldo</b>	<b>TGE</b>	<b>24h-SBP</b>	<b>THAldo, TGE, 24h-SBP</b>
<b>Left ventricular parameters</b>	<b>Univariate, p</b>	<b>Univariate, p</b>	<b>Univariate, p</b>	<b>Multivariate</b>
<b>LVMI [g/m<sup>2.7</sup>]</b>	0.024*	0.018*	<0.001*	24h-SBP*
<b>ΔLVMI [g/m<sup>2.7</sup>]</b>	0.776	0.042*	0.008*	24h-SBP*
<b>LVIDd [mm]</b>	0.531	0.003*	0.052	TGE*
<b>IVSd [mm]</b>	0.119	0.105	0.003*	24h-SBP*
<b>PWTd [mm]</b>	0.277	0.008*	0.003*	TGE*, 24h-SBP*
<b>LAd [mm]</b>	0.523	0.026*	0.008*	n.s.

345 Abbreviations: LVMI: left ventricular mass indexed for height to the 2.7 power; ΔLVMI: reduction of left  
 346 ventricular mass indexed for height to the 2.7 power after treatment; LVM: left ventricular mass,  
 347 LVIDd: left ventricular internal dimension in diastole; PWTd: posterior wall thickness in diastole;  
 348 IVSd: interventricular septum thickness in diastole; LAd: left atrial internal dimension in diastole;  
 349 24h-SBP: 24-hour systolic blood pressure, THAldo: tetrahydroaldosterone; TGE: total glucocorticoid  
 350 excretion; n.s.: no significant results.

351

352 **Supplementary Table 1. Changes in left ventricular geometry of all patients with primary**  
 353 **aldosteronism after one year of treatment.**

<b>Left ventricular geometry</b>	<b>Aldosterone producing adenoma (n=45)</b>		<b>Bilateral adrenal hyperplasia (n=28)</b>	
	<b>baseline</b>	<b>after ADX</b>	<b>baseline</b>	<b>after MRA</b>
<b>Normal (%)</b>	13	22	18	25
<b>Concentric Remodeling (%)</b>	0	11	0	11
<b>Eccentric Hypertrophy (%)</b>	51	36	43	21
<b>Concentric Hypertrophy (%)</b>	36	31	39	43

354 Abbreviations: APA: unilateral disease; BAH: bilateral disease; ADX: adrenalectomy;

355 MRA: mineralocorticoid receptor antagonist treatment.

356

357 **Figure Legends**

358

359 **Figure 1: LVMI at baseline and after treatment in BAH and APA patients treated with either MRA**  
360 **or ADX.**

361 Median and 95 per cent confidence interval of LVMI are shown before (white bar) and after treatment  
362 (checkered bar). Asterisk indicates significance.

363 Abbreviations: APA: unilateral disease; BAH: bilateral disease; LVMI: left ventricular mass indexed for  
364 height to the 2.7 power.

365 **Figure 2: Reduction of LVMI following specific treatment according to baseline total glucocorticoid**  
366 **excretion and THAldo levels.**

367 Median and 95 per cent confidence interval of baseline glucocorticoid and THAldo excretion of PA  
368 patients with low and high total glucocorticoid (8390 µg/24h; 15266 µg/24h) or THAldo (52 µg/24h; 121  
369 µg/24h) excretion are shown. Asterisk indicates significance.

370 Abbreviations: APA: unilateral disease; BAH: bilateral disease, LVMI: left ventricular mass indexed for  
371 height to the 2.7 power; TGE: total glucocorticoid excretion; THAldo: tetrahydroaldosterone.

372

373

374

375

376

377

378


379

380

381

382


383 **Figure 1:**  
384



385  
386



387 **Figure 2:**  
388



389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419

420 **Supplementary Figure Legends**

421

422 **Supplementary Figure 1:** Correlation of total glucocorticoid excretion with left ventricular mass indexed  
423 for height to the 2.7 power at baseline in patients with primary aldosteronism.

424 **Supplementary Figure 2:** Correlation of tetrahydroaldosterone excretion with left ventricular mass  
425 indexed for height to the 2.7 power at baseline in patients with primary aldosteronism.

426 **Supplementary Figure 3:** Correlation of baseline glucocorticoid excretion with reduction of left  
427 ventricular mass indexed for height to the 2.7 power in patients with primary aldosteronism.

428 **Supplementary Figure 4:** Correlation of baseline tetrahydroaldosterone excretion with reduction of left  
429 ventricular mass indexed for height to the 2.7 power in patients with primary aldosteronism.

430

431

432

433

434

435

436

437

438

439

440


441

442

443

444

445 **Supplementary Figure 1:**



446

447

448

449

450

451

452

453


454

455

456


457

458 **Supplementary Figure 2:**




459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472

473 **Supplementary Figure 3:**



474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486

487 **Supplementary Figure 4:**



488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499

500 **References:**

- 501 **1.** Monticone S, Burrello J, Tizzani D, Bertello C, Viola A, Buffolo F, Gabetti L, Mengozzi G, Williams  
502 TA, Rabbia F, Veglio F, Mulatero P. Prevalence and Clinical Manifestations of Primary  
503 Aldosteronism Encountered in Primary Care Practice. *J Am Coll Cardiol* 2017; 69:1811-1820
- 504 **2.** Conn JW. Primary aldosteronism. *J Lab Clin Med* 1955; 45:661-664
- 505 **3.** Marney AM, Brown NJ. Aldosterone and end-organ damage. *Clin Sci (Lond)* 2007; 113:267-278
- 506 **4.** Catena C, Colussi G, Marzano L, Sechi LA. Aldosterone and the heart: from basic research to  
507 clinical evidence. *Horm Metab Res* 2012; 44:181-187
- 508 **5.** Rossi GP, Sacchetto A, Visentin P, Canali C, Graniero GR, Palatini P, Pessina AC. Changes in left  
509 ventricular anatomy and function in hypertension and primary aldosteronism. *Hypertension*  
510 1996; 27:1039-1045
- 511 **6.** Reincke M, Fischer E, Gerum S, Merkle K, Schulz S, Pallauf A, Quinkler M, Hanslik G, Lang K,  
512 Hahner S, Allolio B, Meisinger C, Holle R, Beuschlein F, Bidlingmaier M, Endres S, German Conn's  
513 Registry-Else Kroner-Fresenius-Hyperaldosteronism R. Observational study mortality in treated  
514 primary aldosteronism: the German Conn's registry. *Hypertension* 2012; 60:618-624
- 515 **7.** Mulatero P, Monticone S, Bertello C, Viola A, Tizzani D, Iannaccone A, Crudo V, Burrello J, Milan  
516 A, Rabbia F, Veglio F. Long-term cardio- and cerebrovascular events in patients with primary  
517 aldosteronism. *J Clin Endocrinol Metab* 2013; 98:4826-4833
- 518 **8.** Shenasa M, Shenasa H. Hypertension, left ventricular hypertrophy, and sudden cardiac death. *Int*  
519 *J Cardiol* 2017; 237:60-63
- 520 **9.** Arlt W, Lang K, Sitch AJ, Dietz AS, Rhayem Y, Bancos I, Feuchtinger A, Chortis V, Gilligan LC,  
521 Ludwig P, Riester A, Asbach E, Hughes BA, O'Neil DM, Bidlingmaier M, Tomlinson JW, Hassan-  
522 Smith ZK, Rees DA, Adolf C, Hahner S, Quinkler M, Dekkers T, Deinum J, Biehl M, Keevil BG,  
523 Shackleton CHL, Deeks JJ, Walch AK, Beuschlein F, Reincke M. Steroid metabolome analysis  
524 reveals prevalent glucocorticoid excess in primary aldosteronism. *JCI Insight* 2017; 2
- 525 **10.** Hanslik G, Wallaschofski H, Dietz A, Riester A, Reincke M, Allolio B, Lang K, Quack I, Rump LC,  
526 Willenberg HS, Beuschlein F, Quinkler M, Hannemann A, participants of the German Conn's R.  
527 Increased prevalence of diabetes mellitus and the metabolic syndrome in patients with primary  
528 aldosteronism of the German Conn's Registry. *Eur J Endocrinol* 2015; 173:665-675
- 529 **11.** Wu VC, Chueh SJ, Chen L, Chang CH, Hu YH, Lin YH, Wu KD, Yang WS, Group TS. Risk of new-  
530 onset diabetes mellitus in primary aldosteronism: a population study over 5 years. *J Hypertens*  
531 2017; 35:1698-1708
- 532 **12.** Beuschlein F, Reincke M, Arlt W. The impact of Connshing's syndrome - mild cortisol excess in  
533 primary aldosteronism drives diabetes risk. *J Hypertens* 2017; 35:2548
- 534 **13.** Hundemer GL, Curhan GC, Yozamp N, Wang M, Vaidya A. Cardiometabolic outcomes and  
535 mortality in medically treated primary aldosteronism: a retrospective cohort study. *Lancet*  
536 *Diabetes Endocrinol* 2018; 6:51-59
- 537 **14.** Wu VC, Chang CH, Wang CY, Lin YH, Kao TW, Lin PC, Chu TS, Chang YS, Chen L, Wu KD, Chueh SJ.  
538 Risk of Fracture in Primary Aldosteronism: A Population-Based Cohort Study. *J Bone Miner Res*  
539 2017; 32:743-752
- 540 **15.** Muiesan ML, Lupia M, Salvetti M, Grigoletto C, Sonino N, Boscaro M, Rosei EA, Mantero F, Fallo  
541 F. Left ventricular structural and functional characteristics in Cushing's syndrome. *J Am Coll*  
542 *Cardiol* 2003; 41:2275-2279
- 543 **16.** Takagi S, Tanabe A, Tsuiki M, Naruse M, Takano K. Hypokalemia, diabetes mellitus, and  
544 hypercortisolemia are the major contributing factors to cardiac dysfunction in adrenal Cushing's  
545 syndrome. *Endocr J* 2009; 56:1009-1018
- 546 **17.** Etxabe J, Vazquez JA. Morbidity and mortality in Cushing's disease: an epidemiological approach.  
547 *Clin Endocrinol (Oxf)* 1994; 40:479-484

- 548 **18.** Funder JW, Carey RM, Fardella C, Gomez-Sanchez CE, Mantero F, Stowasser M, Young WF, Jr.,  
549 Montori VM, Endocrine S. Case detection, diagnosis, and treatment of patients with primary  
550 aldosteronism: an endocrine society clinical practice guideline. *J Clin Endocrinol Metab* 2008;  
551 93:3266-3281
- 552 **19.** Funder JW, Carey RM, Mantero F, Murad MH, Reincke M, Shibata H, Stowasser M, Young WF, Jr.  
553 The Management of Primary Aldosteronism: Case Detection, Diagnosis, and Treatment: An  
554 Endocrine Society Clinical Practice Guideline. *J Clin Endocrinol Metab* 2016; 101:1889-1916
- 555 **20.** Lang RM, Bierig M, Devereux RB, Flachskampf FA, Foster E, Pellikka PA, Picard MH, Roman MJ,  
556 Seward J, Shanewise JS, Solomon SD, Spencer KT, Sutton MS, Stewart WJ, Chamber  
557 Quantification Writing G, American Society of Echocardiography's G, Standards C, European  
558 Association of E. Recommendations for chamber quantification: a report from the American  
559 Society of Echocardiography's Guidelines and Standards Committee and the Chamber  
560 Quantification Writing Group, developed in conjunction with the European Association of  
561 Echocardiography, a branch of the European Society of Cardiology. *J Am Soc Echocardiogr* 2005;  
562 18:1440-1463
- 563 **21.** Lang RM, Bierig M, Devereux RB, Flachskampf FA, Foster E, Pellikka PA, Picard MH, Roman MJ,  
564 Seward J, Shanewise J, Solomon S, Spencer KT, St John Sutton M, Stewart W, American Society of  
565 Echocardiography's N, Standards C, Task Force on Chamber Q, American College of Cardiology  
566 Echocardiography C, American Heart A, European Association of Echocardiography ESoC.  
567 Recommendations for chamber quantification. *Eur J Echocardiogr* 2006; 7:79-108
- 568 **22.** Foppa M, Duncan BB, Rohde LE. Echocardiography-based left ventricular mass estimation. How  
569 should we define hypertrophy? *Cardiovasc Ultrasound* 2005; 3:17
- 570 **23.** Devereux RB, Alonso DR, Lutas EM, Gottlieb GJ, Campo E, Sachs I, Reichek N. Echocardiographic  
571 assessment of left ventricular hypertrophy: comparison to necropsy findings. *Am J Cardiol* 1986;  
572 57:450-458
- 573 **24.** Devereux RB. Detection of left ventricular hypertrophy by M-mode echocardiography. Anatomic  
574 validation, standardization, and comparison to other methods. *Hypertension* 1987; 9:119-26
- 575 **25.** Kuch B, Hense HW, Gneiting B, Doring A, Muscholl M, Brockel U, Schunkert H. Body composition  
576 and prevalence of left ventricular hypertrophy. *Circulation* 2000; 102:405-410
- 577 **26.** Cuspidi C, Meani S, Negri F, Giudici V, Valerio C, Sala C, Zanchetti A, Mancina G. Indexation of left  
578 ventricular mass to body surface area and height to allometric power of 2.7: is the difference  
579 limited to obese hypertensives? *J Hum Hypertens* 2009; 23:728-734
- 580 **27.** Webster A, Brady W, Morris F. Recognising signs of danger: ECG changes resulting from an  
581 abnormal serum potassium concentration. *Emerg Med J* 2002; 19:74-77
- 582 **28.** Drazner MH. The progression of hypertensive heart disease. *Circulation* 2011; 123:327-334
- 583 **29.** Spath M, Korovkin S, Antke C, Anlauf M, Willenberg HS. Aldosterone- and cortisol-co-secreting  
584 adrenal tumors: the lost subtype of primary aldosteronism. *Eur J Endocrinol* 2011; 164:447-455
- 585 **30.** Yoon V, Heyliger A, Maekawa T, Sasano H, Carrick K, Woodruff S, Rabaglia J, Auchus RJ, Ghayee  
586 HK. Benign adrenal adenomas secreting excess mineralocorticoids and glucocorticoids.  
587 *Endocrinol Diabetes Metab Case Rep* 2013; 2013:130042
- 588 **31.** Hiraishi K, Yoshimoto T, Tsuchiya K, Minami I, Doi M, Izumiyama H, Sasano H, Hirata Y.  
589 Clinicopathological features of primary aldosteronism associated with subclinical Cushing's  
590 syndrome. *Endocr J* 2011; 58:543-551
- 591 **32.** Tong A, Liu G, Wang F, Jiang J, Yan Z, Zhang D, Zhang Y, Cai J. A Novel Phenotype of Familial  
592 Hyperaldosteronism Type III: Concurrence of Aldosteronism and Cushing's Syndrome. *J Clin*  
593 *Endocrinol Metab* 2016; 101:4290-4297
- 594 **33.** Muiesan ML, Salvetti M, Paini A, Agabiti-Rosei C, Monteduro C, Galbassini G, Belotti E, Aggiusti C,  
595 Rizzoni D, Castellano M, Agabiti-Rosei E. Inappropriate left ventricular mass in patients with  
596 primary aldosteronism. *Hypertension* 2008; 52:529-534



- 597 **34.** Galetta F, Bernini G, Franzoni F, Bacca A, Fivizzani I, Tocchini L, Bernini M, Fallahi P, Antonelli A,  
598 Santoro G. Cardiac remodeling in patients with primary aldosteronism. *J Endocrinol Invest* 2009;  
599 32:739-745
- 600 **35.** Iwashima Y, Horio T, Kuroda S, Takishita S, Kawano Y. Influence of plasma aldosterone on left  
601 ventricular geometry and diastolic function in treated essential hypertension. *Hypertens Res*  
602 2002; 25:49-56
- 603 **36.** Catena C, Colussi G, Lapenna R, Nadalini E, Chiuch A, Gianfagna P, Sechi LA. Long-term cardiac  
604 effects of adrenalectomy or mineralocorticoid antagonists in patients with primary  
605 aldosteronism. *Hypertension* 2007; 50:911-918
- 606 **37.** Pringle SD, Macfarlane PW, Isles CG, Cameron HL, Brown IA, Lorimer AR, Dunn FG. Regression of  
607 electrocardiographic left ventricular hypertrophy following treatment of primary  
608 hyperaldosteronism. *J Hum Hypertens* 1988; 2:157-159
- 609 **38.** Degre S, Detry JM, Unger P, Cosyns J, Brohet C, Kormoss N. Effects of spironolactone-altizide on  
610 left ventricular hypertrophy. *Acta Cardiol* 1998; 53:261-267
- 611 **39.** Lin YH, Huang KH, Lee JK, Wang SM, Yen RF, Wu VC, Chung SD, Liu KL, Chueh SC, Lin LY, Ho YL,  
612 Chen MF, Wu KD, group Ts. Factors influencing left ventricular mass regression in patients with  
613 primary aldosteronism post adrenalectomy. *J Renin Angiotensin Aldosterone Syst* 2011; 12:48-53
- 614 **40.** Marzano L, Colussi G, Sechi LA, Catena C. Adrenalectomy is comparable with medical treatment  
615 for reduction of left ventricular mass in primary aldosteronism: meta-analysis of long-term  
616 studies. *Am J Hypertens* 2015; 28:312-318
- 617 **41.** Rossi G, Boscaro M, Ronconi V, Funder JW. Aldosterone as a cardiovascular risk factor. *Trends*  
618 *Endocrinol Metab* 2005; 16:104-107
- 619 **42.** Arnaldi G, Angeli A, Atkinson AB, Bertagna X, Cavagnini F, Chrousos GP, Fava GA, Findling JW,  
620 Gaillard RC, Grossman AB, Kola B, Lacroix A, Mancini T, Mantero F, Newell-Price J, Nieman LK,  
621 Sonino N, Vance ML, Giustina A, Boscaro M. Diagnosis and complications of Cushing's syndrome:  
622 a consensus statement. *J Clin Endocrinol Metab* 2003; 88:5593-5602
- 623 **43.** Kamenicky P, Redheuil A, Roux C, Salenave S, Kachenoura N, Raissouni Z, Macron L, Guignat L,  
624 Jublanc C, Azarine A, Brailly S, Young J, Mousseaux E, Chanson P. Cardiac structure and function  
625 in Cushing's syndrome: a cardiac magnetic resonance imaging study. *J Clin Endocrinol Metab*  
626 2014; 99:E2144-2153
- 627 **44.** Toja PM, Branzi G, Ciambellotti F, Radaelli P, De Martin M, Lonati LM, Scacchi M, Parati G,  
628 Cavagnini F, Pecori Giraldi F. Clinical relevance of cardiac structure and function abnormalities in  
629 patients with Cushing's syndrome before and after cure. *Clin Endocrinol (Oxf)* 2012; 76:332-338
- 630 **45.** Lindholm J, Juul S, Jorgensen JO, Astrup J, Bjerre P, Feldt-Rasmussen U, Hagen C, Jorgensen J,  
631 Kosteljanetz M, Kristensen L, Laurberg P, Schmidt K, Weeke J. Incidence and late prognosis of  
632 cushing's syndrome: a population-based study. *J Clin Endocrinol Metab* 2001; 86:117-123
- 633 **46.** Mancini T, Kola B, Mantero F, Boscaro M, Arnaldi G. High cardiovascular risk in patients with  
634 Cushing's syndrome according to 1999 WHO/ISH guidelines. *Clin Endocrinol (Oxf)* 2004; 61:768-  
635 777
- 636 **47.** Petramala L, Battisti P, Lauri G, Palleschi L, Cotesta D, Iorio M, De Toma G, Sciomer S, Letizia C.  
637 Cushing's syndrome patient who exhibited congestive heart failure. *J Endocrinol Invest* 2007;  
638 30:525-528
- 639 **48.** Pivonello R, De Martino MC, Iacuanello D, Simeoli C, Muscogiuri G, Carlomagno F, De Leo M,  
640 Cozzolino A, Colao A. Metabolic Alterations and Cardiovascular Outcomes of Cortisol Excess.  
641 *Front Horm Res* 2016; 46:54-65
- 642 **49.** Mihailidou AS, Loan Le TY, Mardini M, Funder JW. Glucocorticoids activate cardiac  
643 mineralocorticoid receptors during experimental myocardial infarction. *Hypertension* 2009;  
644 54:1306-1312

- 645 **50.** Young MJ, Moussa L, Dilley R, Funder JW. Early inflammatory responses in experimental cardiac  
646 hypertrophy and fibrosis: effects of 11 beta-hydroxysteroid dehydrogenase inactivation.  
647 *Endocrinology* 2003; 144:1121-1125
- 648 **51.** Pereira AM, Delgado V, Romijn JA, Smit JWA, Bax JJ, Feelders RA. Cardiac dysfunction is reversed  
649 upon successful treatment of Cushing's syndrome. *European Journal of Endocrinology* 2010;  
650 162:331-340
- 651 **52.** Remde H, Hanslik G, Rayes N, Quinkler M. Glucose Metabolism in Primary Aldosteronism. *Horm*  
652 *Metab Res* 2015; 47:987-993
- 653 **53.** Pitt B, Reichek N, Willenbrock R, Zannad F, Phillips RA, Roniker B, Kleiman J, Krause S, Burns D,  
654 Williams GH. Effects of eplerenone, enalapril, and eplerenone/enalapril in patients with essential  
655 hypertension and left ventricular hypertrophy: the 4E-left ventricular hypertrophy study.  
656 *Circulation* 2003; 108:1831-1838
- 657 **54.** Blumenfeld JD, Sealey JE, Schlussek Y, Vaughan ED, Jr., Sos TA, Atlas SA, Muller FB, Acevedo R,  
658 Ulick S, Laragh JH. Diagnosis and treatment of primary hyperaldosteronism. *Ann Intern Med*  
659 1994; 121:877-885
- 660 **55.** Pouleur AC, le Polain de Waroux JB, Pasquet A, Gerber BL, Gerard O, Allain P, Vanoverschelde JL.  
661 Assessment of left ventricular mass and volumes by three-dimensional echocardiography in  
662 patients with or without wall motion abnormalities: comparison against cine magnetic resonance  
663 imaging. *Heart* 2008; 94:1050-1057

664