

# High proportion of drug hypersensitivity reactions to sulfasalazine following its use in anti-PD-1 associated inflammatory arthritis

Ford, Mark; Sahbudin, Ilfita; Filer, Andrew; Steven, Neil; Fisher, Benjamin

DOI:

[10.1093/rheumatology/key234](https://doi.org/10.1093/rheumatology/key234)

License:

Other (please specify with Rights Statement)

*Document Version*

Peer reviewed version

*Citation for published version (Harvard):*

Ford, M, Sahbudin, I, Filer, A, Steven, N & Fisher, B 2018, 'High proportion of drug hypersensitivity reactions to sulfasalazine following its use in anti-PD-1 associated inflammatory arthritis', *Rheumatology*.  
<https://doi.org/10.1093/rheumatology/key234>

[Link to publication on Research at Birmingham portal](#)

## **Publisher Rights Statement:**

Checked for eligibility 18/07/2018

This is a pre-copyedited, author-produced version of an article accepted for publication in *Rheumatology* following peer review. The version of record [insert complete citation information here] is available online at: xxxxxx [insert URL and DOI of the article on the OUP website].

## **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## **Take down policy**

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

# High proportion of drug hypersensitivity reactions to sulfasalazine following its use in anti-PD-1 associated inflammatory arthritis

*Mark Ford<sup>1</sup>, Ifita Sahbudin<sup>1,2,3</sup>, Andrew Filer<sup>1,2,3</sup>, Neil Steven<sup>4,5</sup>, and Benjamin A. Fisher<sup>1,2,3</sup>.*

1. Rheumatology Department, University Hospitals Birmingham NHS Foundation Trust, Birmingham, UK.
2. Rheumatology Research Group and Arthritis Research UK Rheumatoid Arthritis Pathogenesis Centre of Excellence (RACE), Institute of Inflammation and Ageing, University of Birmingham, Birmingham, UK
3. National Institute for Health Research (NIHR) Birmingham Biomedical Research Centre
4. CR-UK Clinical Trials Unit, Institute of Immunology and Immunotherapy, University of Birmingham, Birmingham, UK
5. Department of Oncology, University Hospitals Birmingham NHS Foundation Trust, Birmingham, UK.

Correspondence:

Dr Benjamin Fisher ([b.fisher@bham.ac.uk](mailto:b.fisher@bham.ac.uk)),

Rheumatology Research Group,

Institute of Inflammation and Ageing,

University of Birmingham,

Birmingham

B15 2TT, UK

**Key words:** Anti-PD-1, anti-PD-L1, pembrolizumab, melanoma, immune-related adverse event, arthritis.

**Key messages:**

Sulfasalazine use in anti-PD-1 associated inflammatory arthritis may be associated with more frequent hypersensitivity reactions

SIR, Checkpoint blockade has revolutionised the management of several cancers by targeting the coinhibitory pathways that down-modulate T cell receptor-mediated T cell activation. However the resulting enhanced T cell activation also gives rise to immune-related adverse events (IrAEs). Arthritis is an increasingly recognised IrAE following treatment with the licensed antibodies to programmed cell death protein-1 (PD-1) and may persist following discontinuation of anti-PD-1.[1]. [1] Treatment often consists of corticosteroids, with escalation if required to disease modifying anti-rheumatic drugs (DMARDs). However uncertainty exists about the possible impact of potent DMARD therapy on durability of cancer response with some oncologists reluctant to consider use of methotrexate.[2]

Sulfasalazine combines salicylic acid with the anti-microbial sulfapyridine and has been suggested as one possible therapy for IrAE.[1, 2] Sulfasalazine can cause a drug hypersensitivity reaction that rarely may be systemic and potentially life-threatening.[3] Importantly such drug hypersensitivity reactions are T cell mediated.[4, 5]

We reviewed all cases of anti-PD-1 associated inflammatory arthritis treated with sulfasalazine at our centre. Patient characteristics for the four patients are presented in Table 1. All had metastatic

melanoma treated with the anti-PD-1 antibody pembrolizumab and three had had prior checkpoint blockade with the anti-CTLA4 antibody ipilimumab.

In Case 1, the initial management of knee and then ankle monoarthritis was with intraarticular steroid but sulfasalazine was introduced after progression to involve both knees and ankles. Ten days later, he was admitted with pyrexia (37.7°C) and a widespread erythematous maculopapular rash. Blood tests revealed an acute rise in alanine transferase (66 U/L) and C-Reactive Protein (CRP) (189 mg/L) and new lymphopenia ( $0.8 \times 10^9/L$ ). He improved with 3 days of intravenous methylprednisolone followed by a reducing course of oral prednisolone. Sulfasalazine and pembrolizumab were both discontinued.

In case 2, sulfasalazine was commenced for initial monoarthritis of the knee following only transient improvement with intra-articular steroid. Five days after starting sulfasalazine, he was admitted with fever (38.5°C) and cough with unremarkable chest radiograph and a raised CRP (78 mg/L). He was treated with antibiotics for a presumed infection of unknown origin (PUO). Sulfasalazine was suspended for one week and future pembrolizumab infusions were deferred. Eleven days after restarting sulfasalazine, he was readmitted with fevers and again treated for presumed PUO and sulfasalazine was stopped. A month later, sulfasalazine was restarted but nine days later he was admitted with low grade fever, non-productive cough, and raised CRP (170 mg/L). No source of infection was identified and he was treated with oral prednisolone and sulfasalazine was stopped. Five weeks later, an escalating dose of sulfasalazine was reintroduced, but on reduction in prednisolone he developed nausea, diarrhoea, deranged liver function tests (ALT 288 U/L) and raised CRP. Sulfasalazine was stopped and his symptoms resolved.

Case 3 had rash, facial swelling and fever but did not seek medical attention as her symptoms settled on stopping sulfasalazine.

Pembrolizumab was suspended in case 4 due to the development of a symmetrical polyarthritis involving the shoulders, knees and hands. Four months after the final pembrolizumab infusion, an escalating dose of sulfasalazine was introduced and five weeks later, he was admitted with a five day history of profound loose stool successfully treated with high dose steroids. Sulfasalazine was discontinued.

We observed a high proportion of adverse reactions with sulfasalazine in the context of anti-PD-1 associated inflammatory arthritis and sulfasalazine was discontinued in all cases. Drug hypersensitivity reactions are mediated by drug-reactive T lymphocytes,[5] and the risk may increase with higher drug levels, or with alteration in T cell function, as exemplified by the association of amoxicillin-clavulanate hypersensitivity with the PTPN22 locus,[6] which encodes a regulator of T cell function. It is therefore plausible that inhibition of PD-1 might also increase risk and this might also apply to other drugs strongly associated with T cell mediated drug hypersensitivity syndromes such as carbamazepine and allopurinol.

In RA clinical trials the withdrawal rate for sulfasalazine was as high as 25%, but this was mostly due to the common side effects of nausea, headache and dizziness, with a drug rash occurring in only 4-5% and other hypersensitivity manifestations being much rarer.[3] Cutaneous and other IrAEs are common with checkpoint blockade, and could give rise to diagnostic uncertainty as illustrated in the second of our cases. However the temporal association, not observed in our experience with methotrexate, the high prevalence and similarity to the recognised sulfasalazine hypersensitivity syndrome, and the reoccurrence upon rechallenge, all point towards a genuine drug association. Our fourth case was different and characterised by colitis, but in this case pembrolizumab was discontinued four months before sulfasalazine was started. Colitis is a common IrAE following checkpoint blockade, and rare following sulfasalazine, suggesting this to be unrelated. However recent data indicate that the gut microbiome can influence emergence of IrAEs following checkpoint

blockade,[7] and since sulfasalazine has an antibiotic component, we cannot exclude an indirect association through alteration of faecal flora.[8]

In conclusion we observed a high proportion of hypersensitivity reactions following use of sulfasalazine, and suggest caution in its use to manage anti-PD-1 related inflammatory arthritis.

### **Acknowledgements**

BAF has received support from the NIHR Birmingham Biomedical Research Centre and the NIHR/Wellcome Trust Birmingham Clinical Research Facility. The views expressed in this publication are those of the authors and not necessarily those of the NHS, the National Institute for Health Research or the Department of Health. AF is supported by the MRC-ARUK Centre for Musculoskeletal Ageing Research and the Kennedy Trust funded Arthritis Therapy Acceleration Programme.

### **Conflicts of Interest**

BAF has received consultancy fees from Novartis, Roche and BMS. AF has received consultancy fees from Novartis and Chugai.

## References

1. Naidoo J, Cappelli LC, Forde PM, Marrone KA, Lipson EJ, Hammers HJ, Sharfman WH, Le DT, Baer AN, Shah AA *et al*: **Inflammatory Arthritis: A Newly Recognized Adverse Event of Immune Checkpoint Blockade**. *The oncologist* 2017, **22**(6):627-630.
2. Chan MM, Kefford RF, Carlino M, Clements A, Manolios N: **Arthritis and tenosynovitis associated with the anti-PD1 antibody pembrolizumab in metastatic melanoma**. *Journal of immunotherapy (Hagerstown, Md : 1997)* 2015, **38**(1):37-39.
3. Box SA, Pullar T: **Sulphasalazine in the treatment of rheumatoid arthritis**. *British journal of rheumatology* 1997, **36**(3):382-386.
4. Yang F, Gu B, Zhang L, Xuan J, Luo H, Zhou P, Zhu Q, Yan S, Chen SA, Cao Z *et al*: **HLA-B\*13:01 is associated with salazosulfapyridine-induced drug rash with eosinophilia and systemic symptoms in Chinese Han population**. *Pharmacogenomics* 2014, **15**(11):1461-1469.
5. Pavlos R, Mallal S, Ostrov D, Buus S, Metushi I, Peters B, Phillips E: **T cell-mediated hypersensitivity reactions to drugs**. *Annual review of medicine* 2015, **66**:439-454.
6. Lucena MI, Molokhia M, Shen Y, Urban TJ, Aithal GP, Andrade RJ, Day CP, Ruiz-Cabello F, Donaldson PT, Stephens C *et al*: **Susceptibility to amoxicillin-clavulanate-induced liver injury is influenced by multiple HLA class I and II alleles**. *Gastroenterology* 2011, **141**(1):338-347.
7. Dubin K, Callahan MK, Ren B, Khanin R, Viale A, Ling L, No D, Gobourne A, Littmann E, Huttenhower C *et al*: **Intestinal microbiome analyses identify melanoma patients at risk for checkpoint-blockade-induced colitis**. *Nature communications* 2016, **7**:10391.
8. Zheng H, Chen M, Li Y, Wang Y, Wei L, Liao Z, Wang M, Ma F, Liao Q, Xie Z: **Modulation of Gut Microbiome Composition and Function in Experimental Colitis Treated with Sulfasalazine**. *Frontiers in microbiology* 2017, **8**:1703.

Table 1. Patient characteristics.

Case	Gender	Age	Previous immunotherapy	Cancer Response to anti-PD-1	Initial Joint presentation	Subsequent joint presentation	ANA	Months from Anti-PD1 to start of arthritis	Duration on sulfasalazine (days)	Reason for stopping SSZ	Management of Adverse event
1	Male	57	Ipilimumab. (Stopped due to hypophysitis and disease progression)	Stable	Monoarthritis	Symmetrical Oligoarthritis (knees and ankles)	Negative	4	10	fever, erythematous maculopapular rash across the face, neck, chest and back, acute rise in ALT and CRP and lymphopenia	Discontinuation IV and oral corticosteroid
2	Male	51	Ipilimumab. (Stopped due to hypophysitis and hepatitis)	Remission	Monoarthritis	Symmetrical Medium/large joint polyarthritis	Negative	7	a) 5 b) 11 c) 9 d) 21 (concomitant steroid)	a) fever, cough, raised CRP. b) fever, raised CRP. c) fever, non-productive cough, raised CRP. d) nausea, diarrhoea, raised ALT and CRP	a) Discontinuation b) Discontinuation c) Discontinuation and oral corticosteroid d) Discontinuation and oral corticosteroid
3	Female	50	Nil	Slight progression	Monoarthritis	Symmetrical Polyarthritis	1:400 <sup>1</sup>	5	10	rash over chest, swelling around eyes and face, profound lethargy and fever.	Discontinuation
4	Male	77	Ipilimumab. (Stopped due to progression)	Remission	Symmetrical Polyarthritis	Symmetrical Polyarthritis	Negative	2	35	acute late onset colitis	Discontinuation IV and oral corticosteroid

Footnote: All cases were anti-CCP antibody and rheumatoid factor negative. ALT, alanine aminotransferase. CRP, C-reactive protein. <sup>1</sup>Predated introduction of sulfasalazine.