

# Glucocorticoids and bone: consequences of endogenous and exogenous excess and replacement therapy

Hardy, Rowan; Zhou, Hong; Seibel, Markus J; Cooper, Mark S.

DOI:

[10.1210/er.2018-00097](https://doi.org/10.1210/er.2018-00097)

License:

None: All rights reserved

*Document Version*

Peer reviewed version

*Citation for published version (Harvard):*

Hardy, R, Zhou, H, Seibel, MJ & Cooper, MS 2018, 'Glucocorticoids and bone: consequences of endogenous and exogenous excess and replacement therapy', *Endocrine Reviews*. <https://doi.org/10.1210/er.2018-00097>

[Link to publication on Research at Birmingham portal](#)

## **Publisher Rights Statement:**

Checked for eligibility 14/09/2018

This is a pre-copyedited, author-produced version of an article accepted for publication in *Endocrine Reviews* following peer review. The version of record Hardy et al Glucocorticoids and bone: consequences of endogenous and exogenous excess and replacement therapy is available online at: <https://doi.org/10.1210/er.2018-00097>.

## **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## **Take down policy**

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

1 Title: **Glucocorticoids and bone: consequences of endogenous and**  
2 **exogenous excess**

3

4 Authors:

5 Rowan Hardy<sup>1</sup>, Hong Zhou<sup>2</sup>, Markus J. Seibel<sup>2,3,5</sup>, Mark S Cooper<sup>3,4,5</sup>

6

7 Affiliations:

8 1 University of Birmingham, Birmingham, UK

9 2 Bone Research Program, ANZAC Research Institute, Sydney, Australia

10 3 Department of Endocrinology and Metabolism, Concord Repatriation General Hospital, Sydney,  
11 Australia

12 4 Adrenal Steroid Laboratory, ANZAC Research Institute, Sydney, Australia

13 5 Concord Clinical School, The University of Sydney, Sydney, Australia

14

15 Short title: **Glucocorticoids and bone**

16

17 Keywords: Glucocorticoids, cortisol, osteoblasts, bone, osteoporosis

18

19 Corresponding Author:

20 Mark S Cooper, Adrenal Steroid Laboratory, ANZAC Research Institute, University of Sydney, Concord

21 Repatriation General Hospital, Concord, Australia

22 Email: [mark.cooper@sydney.edu.au](mailto:mark.cooper@sydney.edu.au)

23 TEL: +61 (2) 97676775

24 FAX: +61 (2) 97677603

25

26 Contact for reprints:

27 Mark S Cooper

28

29 Funding:

30 This work was not directly supported by any grants or fellowships

31

32 Disclosure summary:

33 None of the authors have anything to disclose

34

35 Abstract:

36 Osteoporosis associated with long-term glucocorticoid therapy remains a common and serious bone  
37 disease. In addition, in recent years it has become clear that more subtle states of *endogenous*  
38 glucocorticoid excess may have a major impact on bone health. Adverse effects can be seen with  
39 mild systemic glucocorticoid excess but there is also evidence of tissue-specific regulation of  
40 glucocorticoid action within bone as a mechanism of disease. This review article will examine a) the  
41 role of endogenous glucocorticoids in normal bone physiology, b) the skeletal effects of endogenous  
42 glucocorticoid excess in the context of endocrine conditions such as Cushing's disease and  
43 autonomous cortisol secretion (subclinical Cushing's syndrome), and c) the actions of therapeutic  
44 (exogenous) glucocorticoids on bone. We will review the extent to which the effect of  
45 glucocorticoids on bone is influenced by variations in tissue metabolising enzymes and  
46 glucocorticoid receptor expression and sensitivity. We will consider how the effects of therapeutic  
47 glucocorticoids on bone are complicated by the effects of the underlying inflammatory disease being  
48 treated. We will also examine the impact that glucocorticoid replacement regimens have on bone in  
49 the context of primary and secondary adrenal insufficiency.

50

51 Precis:

52 We reviewed literature relating to the effects of glucocorticoids on bone. This included the impact of  
53 endogenously synthesised and therapeutically administered glucocorticoids on bone and bone cells.

54

55 **I. Introduction**

56 Glucocorticoid induced osteoporosis (GIOP) remains an important and common clinical problem.

57 GIOP was first recognised in patients with Cushing’s disease or other states of endogenous  
58 glucocorticoid excess.<sup>1</sup> However, since the introduction of therapeutic glucocorticoids over 60 years  
59 ago, GIOP is now much more commonly seen in people treated with therapeutic glucocorticoids.<sup>2</sup> It  
60 is well established that therapeutic glucocorticoid treatment is associated with significant loss of  
61 bone density, deterioration of bone structure and substantial increases in fracture risk.<sup>3,4</sup> The  
62 condition appears to behave in many ways distinct to that of age-related or postmenopausal  
63 osteoporosis and, as such, is regarded as a distinct metabolic bone disease.<sup>5</sup>

64 The study of GIOP is complicated by the almost universal involvement of an underlying, usually  
65 inflammatory disease, as the reason for glucocorticoid treatment in the first place.<sup>5,6</sup> These  
66 underlying illnesses are rarely incorporated into animal models examining the pathogenesis of GIOP.  
67 Various treatments have been evaluated for GIOP in the clinical setting but usually only after these  
68 have proven effective in the context of postmenopausal osteoporosis. Trials in GIOP are generally  
69 powered based on BMD changes rather than fracture risk reduction. All current treatments for GIOP  
70 have significant limitations in terms of effectiveness and risk of adverse effects.

71 In addition to the clear evidence that high levels of therapeutic glucocorticoids do harm to bone  
72 there is increasing evidence that more subtle states of endogenous of glucocorticoid excess  
73 detrimentally impact on bone.<sup>7</sup> The main focus of this research has been the impact of subclinical  
74 endogenous hypercortisolism (also known as sub-clinical Cushing’s syndrome or autonomous  
75 cortisol secretion), a condition characterised by autonomous cortisol secretion usually by one or  
76 more adrenal cortex nodules. There is current debate regarding how prevalent this condition is and  
77 how significant its impact is on bone but many studies indicate that the effects on bone can be  
78 substantial.<sup>7</sup> It is less clear how to investigate and manage bone loss and extra-skeletal  
79 manifestations in subclinical endogenous hypercortisolism.

Comment [M1]: ? add graphical abstract here?

80 There is now considerable evidence that glucocorticoid action can be modulated by various  
81 mechanisms at a tissue level. These mechanisms include variations in the expression and sensitivity  
82 of the glucocorticoid receptors<sup>8</sup>, export of steroids out of the cell by transmembrane transporters<sup>9</sup>  
83 and enzymatic metabolism of glucocorticoids to more or less active forms.<sup>10</sup> In particular, there has  
84 been interest in the role of the 11 $\beta$ -hydroxysteroid dehydrogenases (11 $\beta$ -HSDs) which interconvert  
85 the active glucocorticoids cortisol and corticosterone with their inactive counterparts cortisone and  
86 dehydrocorticosterone.<sup>10</sup> These enzymes appear to influence bone cell differentiation and function  
87 and changes in enzyme expression have been implicated in the development of some aspects of  
88 glucocorticoid induced bone loss. Excessive tissue glucocorticoid action in the presence of normal  
89 circulating levels of glucocorticoids might thus play a more generalised role in other forms of  
90 osteoporosis not traditionally associated with glucocorticoid excess.

Comment [RSH2]: Could use a figure here

91 The issue of whether glucocorticoid levels are sufficient, inadequate or excessive for bone health is  
92 relevant to the treatment of states of adrenal insufficiency such as Addison's disease or  
93 hypopituitarism. Evidence suggests that historically, glucocorticoid replacement regimens were  
94 excessive in many people and this is likely to have detrimentally impacted on bone health. More  
95 contemporary (and lower) replacement glucocorticoid doses appear to have less of an adverse  
96 impact on bone in terms of bone density and biochemical markers. However, whether this translates  
97 into reduced fracture risk is unclear.

98 This review will therefore examine the role endogenous glucocorticoids play in normal bone  
99 physiology, examine the skeletal effects of endogenous glucocorticoid excess in the context of  
100 endocrine conditions such as Cushing's disease and autonomous cortisol secretion, and explore the  
101 actions of therapeutic glucocorticoids on bone. Based on a Medline<sup>TM</sup> publication search within the  
102 last five years (to February 2018) supplemented by earlier studies of continuing significance we  
103 review how the effect of glucocorticoids on bone is influenced by tissue metabolising enzymes and  
104 glucocorticoid receptor expression. We will consider how these effects are complicated by

105 inflammation. We will additionally examine the impact that glucocorticoid replacement has on bone  
106 in the context of adrenal insufficiency.

107

108 **II Mechanisms of action of glucocorticoids on bone**

109 This section outlines how glucocorticoids have their effect on bone. Evidence primarily based on  
110 mouse models suggests that the main adverse effects of high levels of glucocorticoids on bone are  
111 through direct effects on cells involved in bone remodelling; osteoblasts, osteocytes and osteoclasts.  
112 Mechanisms such as impaired cellular proliferation, increased apoptosis, altered autophagy and  
113 changes in RANKL/OPG, wnts/sclerostin expression have all been proposed to be important  
114 mediators of these effects. These mechanisms have been examined in animal models of disease and  
115 to some extent in human samples. We will conclude this section by discussing evidence that some of  
116 the adverse effects of glucocorticoids on systemic fuel metabolism are mediated through the  
117 skeleton.

118 An important consideration when interpreting the literature relating to glucocorticoid effects on  
119 bone is to appreciate the significance and relationships of the various forms of glucocorticoids that  
120 have been studied or implicated in disease. The main glucocorticoid secreted from the adrenal  
121 cortex in humans is cortisol. When administered therapeutically, cortisol is referred to as  
122 hydrocortisone. A smaller amount of corticosterone (about 5-10% that of cortisol) is also secreted  
123 from the human adrenal cortex.<sup>11</sup> Although traditionally considered to have a minor role in human  
124 physiology recent work examining the selective export of cortisol and corticosteroids from the cell  
125 suggests that corticosterone secretion could be important over and above the secretion of cortisol.<sup>9</sup>  
126 In the mouse and rat corticosterone is the main glucocorticoid secreted from the adrenal due to the  
127 absence of the 17 $\alpha$ -hydroxylase enzyme in the adult adrenal gland in rodents.<sup>12</sup> Cortisol and  
128 corticosterone have direct and similar actions at the glucocorticoid and mineralocorticoid receptors  
129 but in classical mineralocorticoid target tissues (kidney, colon, salivary and sweat glands) these  
130 glucocorticoids are inactivated by the enzyme 11 $\beta$ -hydroxysteroid dehydrogenase type 2 (11 $\beta$ -HSD2)  
131 to cortisone and dehydrocorticosterone respectively. These steroids lack activity at the level of the  
132 GR or MR but can be reactivated to cortisol and corticosterone by 11 $\beta$ -hydroxysteroid

**Comment [RSH3]:** Again a figure here could be useful. Could we tie it into HPA axis, with relative circulating levels and the 1<sup>st</sup> pass metabolism of pred in liver

Agree we need a figure



133 dehydrogenase type 1 (11 $\beta$ -HSD1) enzyme which is expressed in a range of tissues, in particular in  
134 the liver but also in bone. Prednisolone and prednisone are the most widely used oral  
135 glucocorticoids. As with cortisol and cortisone these compounds differ by just a hydroxylation at  
136 position 11 of the steroid ring with prednisolone being the active form and prednisone the inactive  
137 form. In practice orally administered prednisone and prednisolone have similar properties in vivo  
138 since prednisone is efficiently converted to prednisolone by hepatic 11 $\beta$ -HSD1 activity on first pass  
139 metabolism in the liver.

140 Endogenous glucocorticoids have the potential to bind to either the classical glucocorticoid receptor  
141 (GR) or the mineralocorticoid receptor (MR). The MR is also referred to as the type 1 or high affinity  
142 GR since the affinity of the MR for cortisol and corticosterone is 10 times higher than that of the  
143 GR.<sup>13</sup> As discussed above, the main factor influencing endogenous glucocorticoid binding to the MR  
144 is the presence of 11 $\beta$ -HSD2. As a further complication, the synthetic glucocorticoid dexamethasone  
145 only binds to the GR and not the MR. These differences have important implications when  
146 interpreting differences between studies examining the mechanisms underlying GIOP.

147 The downstream cellular consequences of glucocorticoid receptor binding have been reviewed in  
148 detail elsewhere.<sup>14,15</sup> These mechanisms will be discussed in each section where specifically relevant  
149 to glucocorticoid actions on bone.

150

## 151 II.I Effects on osteoblasts, osteocytes and osteoclasts

152 Glucocorticoids have direct effects on specific tissues but also exert their effects through indirect  
153 mechanisms, e.g. through the regulation of endocrine signalling pathways. The extent to which  
154 glucocorticoid induced bone loss is mediated through direct effects on the cells which coordinate  
155 bone metabolism (osteoblasts, osteocytes, osteoclasts and their respective precursors) has been  
156 debated. Bone cells are clearly very sensitive to glucocorticoids in vitro and in vivo. In vivo mouse

157 models that have attempted to examine this question show that the effects of glucocorticoids on  
158 bone are primarily through direct actions on bone cells and bone remodelling. The situation  
159 regarding the important clinical manifestations of GIOP in humans, fractures, might be different  
160 since glucocorticoids can influence falls related factors such as muscle strength that are difficult to  
161 replicate in mice.

162 Whether or not there is a single primary or dominant target of glucocorticoids accounting for the  
163 effects on bone remains unclear. This is not helped by the wide variety of mouse models which vary  
164 according to strain; age; sex; skeletal site examined; and type, dose, duration and route of  
165 glucocorticoid administration.<sup>16,17</sup> Furthermore, the effects of glucocorticoids are not consistent  
166 across skeletal sites and surfaces. Although glucocorticoid treatment of mouse models mimics some  
167 of the findings seen with clinical use, the extent to which non-human models mirror the  
168 pathophysiology in humans remains unclear.

169 In the following section the individual effects of glucocorticoids on 1, osteoblasts, 2, osteocytes, and  
170 3, osteoclasts will be discussed in terms of in vitro and in vivo actions. These sections reflect the  
171 majority of recent studies investigating how glucocorticoids affect bone. There is, however, a small  
172 number of studies that report consequences possibly mediated by other cells. For example, a  
173 preliminary report indicated that mice that lack lymphocytes are protected against glucocorticoid  
174 induced changes in bone density, suggesting a possible role for these cells in GIOP.<sup>18</sup>

175

## 176 II.I.I Effects on osteoblasts

177 In vitro effects:

178 In contrast to the clearly detrimental impact that therapeutic glucocorticoids have on bone in the  
179 clinical setting in many in vitro situations, glucocorticoids have an important and positive role in the  
180 commitment and differentiation of cells of the osteoblast lineage. Glucocorticoids have a stimulatory

181 role in the differentiation of uncommitted mesenchymal precursor cells to the osteoblastic lineage  
182 and high doses of glucocorticoids are generally part of the differentiation medium in protocols for  
183 the differentiation of these cells.<sup>19,20</sup> Glucocorticoids demonstrate clear stimulatory activity on the  
184 expression of a range of cellular markers related to osteoblast function, including osteocalcin and  
185 alkaline phosphatase.<sup>21-24</sup> Glucocorticoids show inconsistent effects on cellular proliferation but in  
186 general high doses of glucocorticoids slows the proliferation rate of mature osteoblast like cells in  
187 culture.<sup>25</sup> The observation that glucocorticoids in vivo usually result in a dramatic decrease in bone  
188 formation but in vitro actions are largely stimulatory has been difficult to explain. It is possible that  
189 there is dose dependency with low levels of glucocorticoids being stimulatory and high doses being  
190 inhibitory for osteoblasts.<sup>26</sup> Other lines of evidence suggest that in vitro effects of glucocorticoids in  
191 culture are more complex than previously considered. For example, within a single primary culture  
192 of osteoblastic cells there are various different populations present. It has been suggested that more  
193 mature osteoblasts have stimulatory paracrine functions on less mature osteoblast precursors and  
194 that these are glucocorticoid dependent. For example, disruption of glucocorticoid signalling by the  
195 artificial introduction of 11 $\beta$ -HSD2 into mature osteoblasts results in reduced differentiation of less  
196 mature osteoblasts within the same culture, an effect likely due to alterations of expression of wnt  
197 or wnt-related genes.<sup>27</sup> These results indicate that the communication between various types of  
198 bone cells at different stages of differentiation is likely to be complex and glucocorticoids appear  
199 important in these communication pathways. A recent review focussing on the various mechanisms  
200 by which glucocorticoids affect osteoblast function has been published by Frenkel et al.<sup>28</sup>

201 Glucocorticoids influence the proliferation, differentiation or function of osteoblasts but most  
202 dramatically they influence their survival and death. It is now clear that osteoblast apoptosis has an  
203 important role in bone physiology.<sup>29</sup> Glucocorticoids stimulate osteoblast apoptosis in vitro,  
204 triggering the rapid activation of the kinases Pyk2 and JNK and increasing reactive oxygen species  
205 (ROS) in primary cultures.<sup>30,31</sup> Glucocorticoids can increase apoptosis via increased endoplasmic  
206 reticulum stress and glucocorticoid actions through this pathway synergise with TNFa.<sup>32</sup>

207 Glucocorticoids also regulate the expression and activity of pro-apoptotic factors of the Bcl2 family  
208 such as Bim.<sup>33</sup> Knockdown of Bim in osteoblasts protects against glucocorticoid induced apoptosis  
209 and silencing of E4BP4 attenuates Bim expression and also blocks glucocorticoid induced apoptosis  
210 in osteoblasts.<sup>34</sup> Glucocorticoids also increase expression of Bak (another Bcl2 family member) and  
211 decrease expression of Bcl-XL, a pro-survival Bcl2 protein.<sup>35</sup> Dexamethasone can induce Bcl2  
212 mediated cell death via induction of p53.<sup>36</sup> As such there appears to be multiple pathways by which  
213 glucocorticoids induce apoptosis of osteoblasts.

214 A vast range of studies have attempted to identify specific pathways by which glucocorticoids act on  
215 osteoblasts in culture. The most prominent targets proposed include: effectors of apoptosis,  
216 RANKL/OPG signalling, wnts and their inhibitors, microRNAs, IL-11 and BMP/notch signalling.

217 Glucocorticoids stimulate expression of RANKL and suppress expression of OPG in primary cultures  
218 of osteoblasts and osteoblast like cell lines.<sup>37-39</sup> These changes would be expected to generate a pro-  
219 osteoclastogenic signal. The significance of osteoblast expressed RANKL has recently been  
220 questioned with the osteocyte now considered to be the most important source of RANKL in normal  
221 physiology.<sup>40,41</sup>

222 Wingless (wnt) signalling is firmly established as a critical mediator of many of the anabolic and  
223 catabolic signalling pathways in bone.<sup>42</sup> Glucocorticoids have dramatic impacts on a range of wnt  
224 related genes. At low doses glucocorticoids promote the secretion of wnt9a and wnt10b. At higher  
225 doses glucocorticoids suppress intracellular wnt signalling in osteoblasts resulting in a suppression of  
226 osteoblast differentiation.<sup>43,44</sup> Significant interest has focussed on the synthesis of wnt inhibitors  
227 such as DKK-1 and sclerostin. Whereas sclerostin will be discussed in the next section since it is  
228 expressed exclusively in osteocytes, DKK1 is an important wnt inhibitor which is expressed in  
229 osteoblasts and reported to be positively regulated by glucocorticoids.<sup>45</sup> DKK1 appears to have a  
230 negative impact on bone formation but also causes a reduction in the expression of OPG by

231 osteoblasts,<sup>46</sup> which would favour an increase in osteoclastogenesis and bone resorption, in addition  
232 to the reported suppression of anabolic osteoblast behaviour.

233 Other osteoblastic signalling pathways targeted by glucocorticoids include insulin-like growth  
234 factors, transforming growth factors, basic fibroblast growth factor, and platelet-derived growth  
235 factors. In vitro, glucocorticoids suppress the expression of IGF I and PDGF, which possess anabolic  
236 mitogenic actions in osteoblasts, whilst reducing the anabolic actions of TGF $\beta$ .<sup>47-49</sup> Novel cellular  
237 targets in osteoblasts which appear to be influenced by glucocorticoids in vitro include IL-11<sup>50</sup>, E3  
238 ubiquitin ligases<sup>51</sup> and microRNA-199a.<sup>25</sup>

239

240 In vivo effects:

241 Much of what we know about the impact of glucocorticoid on osteoblast function comes from  
242 animal models of GIOP. Although these models appear to recapitulate some of the features seen in  
243 human GIOP there is considerable diversity and variability in the phenotypes seen with the models  
244 used. This raises questions as to the applicability of these models to the human situation. Mouse  
245 models vary in terms of the strain used; the animal age and gender; the glucocorticoid type, route  
246 and dose employed; and the skeletal site examined. The C57/B6 mouse strain is increasingly  
247 employed as this background is generally most efficient when using tissue targeted transgenic  
248 models via the Cre/lox approach. Recent evidence suggests that this mouse strain is relatively  
249 resistant to the effects of glucocorticoids on bone compared to other mouse strains.<sup>52</sup> It is not clear  
250 whether this reflects a bone specific difference in glucocorticoid sensitivity or a more generalised  
251 disparity. Although this variability between animal strains in terms of glucocorticoid sensitivity and  
252 mechanisms of glucocorticoid adverse consequences makes conclusions more difficult to draw, it is  
253 possible that what is seen in mice is some of the clinical variability seen in humans. The findings from  
254 mice models might need to be considered in their aggregate form rather than depending too much

255 on individual models. A comprehensive review of the various non-human models used to study GIOP  
256 has recently been published.<sup>16</sup> In addition to being useful for the study of GIOP these models have  
257 also indicated that glucocorticoid signalling is important for normal mineralisation of vertebral bones  
258 and bone growth at some surfaces.

259 The acceptance of mice as models for human GIOP started with a highly influential study which  
260 examined the Swiss Webster mouse strain treated with subcutaneous pellets of prednisolone.<sup>53</sup>  
261 These studies demonstrated that glucocorticoids could induce osteoblast and osteocyte apoptosis in  
262 vivo. The effects of glucocorticoids on osteoblasts appeared to dominate those on osteoclasts and  
263 bone resorption. The direct cellular targets of glucocorticoids within bone have been examined in  
264 subsequent studies utilising C57/B6 mice. These studies include tissue selective blockade of  
265 glucocorticoid signalling in specific cell lineages using the 11 $\beta$ -HSD2 enzyme or selective deletion of  
266 glucocorticoid receptors. A potentially important distinction between these approaches is that  
267 models using 11 $\beta$ -HSD2 will have reduced glucocorticoid signalling through both GR and MR if these  
268 receptors are present within target cells whereas selective GR or MR deletion will only target these  
269 aspects of glucocorticoid signalling. Another important caveat using these approaches is that it is  
270 now clear that the promoters used to drive selective expression within bone have some limitations  
271 in that expression in tissues of interest is not normally complete and expression of transgenes can  
272 occur to a limited extent in off target tissues, for instance in selective regions of the brain during  
273 development.<sup>54-56</sup> This off target expression may differ between strains.

274 11 $\beta$ -HSD2 is a glucocorticoid inactivating enzyme which has expression primarily in MR expressing  
275 tissues such as kidney, colon and salivary gland. There is additionally some expression within the  
276 brain and in fetal tissues.<sup>57,58</sup> Osteoblastic cells (osteoblasts and osteocytes) in the adult mouse do  
277 not express 11 $\beta$ -HSD2 although the enzyme has been detected in fetal bone.<sup>59</sup> 11 $\beta$ -HSD2 is highly  
278 effective at reducing glucocorticoid signalling and when expressed in osteoblasts appears to entirely  
279 block the effect of physiological concentrations of endogenous glucocorticoids.<sup>60</sup> The enzyme is also

280 effective at blocking the action of prednisolone.<sup>61</sup> As such, transgenic expression of 11 $\beta$ -HSD2 within  
281 osteoblasts has been utilised to examine the impact of glucocorticoids on these cells. Expression of  
282 11 $\beta$ -HSD2 within osteoblasts has been reported in two different strains of mice with different  
283 promoters. In the C57/B6 strain, expression of 11 $\beta$ -HSD2 gene was under the control of the  
284 osteocalcin promoter and as such would be expected to be expressed in mature osteoblasts and  
285 osteocytes.<sup>61</sup> These mice did not have an obvious basal phenotype but were protected against the  
286 actions of glucocorticoids on osteoblast apoptosis and loss of bone density. In the CD1 strain, 11 $\beta$ -  
287 HSD2 has been driven under the control of the 2.3Kb Col1A1 promoter.<sup>62,63</sup> This truncated form of the  
288 full type I collagen promoter is expressed in mature osteoblasts and osteocytes but not in other cell  
289 types that normally produce type I collagen.<sup>64</sup> These mice had a subtle basal phenotype with  
290 reduced bone density of the vertebrae implying an impairment of bone mineralisation.<sup>63</sup> These mice  
291 also had delayed ossification of the cranial bones and reduced periosteal circumference of long  
292 bones indicating reduced periosteal apposition of bone.<sup>65,66</sup> These mice were also protected against  
293 the adverse effects of glucocorticoids on bone, specifically the reduction in bone formation rate and  
294 increase in endosteal bone resorption seen in controls.<sup>67</sup>

295 Mice with targeted deletion of the GR in osteoblastic cells have also been generated. These mice had  
296 Cre driven by the Runx2 promoter which is expressed in cells throughout the osteoblast lineage.<sup>50</sup>  
297 These mice had a basal phenotype characterised by mildly reduced bone size and reduced bone  
298 density. As with the mice above this strongly indicated that endogenous glucocorticoids are not  
299 essential for bone formation but do have a mild anabolic effect on bone. When treated with  
300 prednisolone these mice did not demonstrate the bone loss seen in their wild type equivalents.  
301 These negative effects of glucocorticoids still occurred in mice where the GR was modified such that  
302 it was not able to form dimeric complexes (the dim-dim mice). This implies that the actions of  
303 glucocorticoids on osteoblasts are mediated primarily by the monomeric form of the receptor (a  
304 mechanism associated with transrepression and typically associated with anti-inflammatory actions)  
305 rather than the dimeric form traditionally thought of as mediating the beneficial metabolic actions of

306 glucocorticoids. This study identified suppression of IL-11 as an important mediator of the adverse  
307 effects of glucocorticoids on bone.

308 Although it has been generally assumed that the GR is the most important target of glucocorticoids  
309 in the osteoblast it has been reported that the MR is also expressed in these cells.<sup>68</sup> In mouse models  
310 blockade of MR signalling either through the use of the MR antagonist spironolactone or through  
311 transgenic deletion of the MR results in some protection against the effects of therapeutic  
312 glucocorticoids.<sup>69</sup> The relative importance of GR and MR signalling in bone and whether there are  
313 interactions between the GR and MR signalling pathways has not been determined.

314

315 II.I.II Effects on osteocytes:

316 Our understanding of the role that osteocytes play in the coordination of bone remodelling has  
317 developed rapidly over the last two decades. A role for osteocytes in the development of GIOP was  
318 suggested at an early stage and in particular a role for osteocyte apoptosis was demonstrated in  
319 animal models.<sup>53</sup> Thus, osteocyte appears to be extremely sensitive to glucocorticoids.

320

321 Evidence for osteocyte apoptosis.

322 Evidence for a role of glucocorticoids in osteocyte function initially came from animal models of  
323 glucocorticoid treatment in which apoptosis of osteocytes could be demonstrated.<sup>53,70</sup> These  
324 observations were supported by studies examining human bone from individuals that had been  
325 exposed to high levels of glucocorticoids where signs of osteocyte apoptosis were also seen.<sup>53,71</sup>  
326 Osteocytes are thought to be long-lived cells and it is uncertain whether osteocytes that have  
327 apoptosed can be replaced by new osteocytes. As such death of osteocytes would be likely to have  
328 prolonged consequences for the organism. Apoptosis of osteocytes was also demonstrated in mice

**Comment [RSH4]:** Is it worth having a table summarising the GC targeted inhibition in OBs and reporting the phenotype?

Yes definitely



329 treated with a high dose of prednisolone (2.4mg/kg/d over 28 days) while osteocytes remained  
330 unaffected at a lower dose of 1.4mg/kg/d, suggesting that there may be a threshold for the  
331 development of osteocyte apoptosis.<sup>72</sup>

332 Various factors have been found to protect against glucocorticoid induced apoptosis of osteocytes in  
333 animal models. These include PTH, bisphosphonates, calcitonin and OPG.<sup>73-76</sup> Whether this  
334 mechanism contributes to the therapeutic efficacy of some of these agents in human GIOP is unclear  
335 and difficult to test clinically.

336 It should be noted that osteocyte apoptosis in GIOP has not been a universal finding. Indeed, there  
337 was no evidence of osteoblast or osteocyte apoptosis in control mice treated with glucocorticoids in  
338 the study examining the effects of osteoblast/osteocyte specific deletion of GR on the sensitivity of  
339 bone to glucocorticoids.<sup>50</sup> This lack of osteocyte apoptosis was manifest despite glucocorticoids  
340 having a clearly detrimental effect on bone formation and bone strength. It is possible that this  
341 observation reflects differences between strains and glucocorticoid dosing but it also implies that  
342 glucocorticoid induced apoptosis of osteocytes is not an essential step for glucocorticoids to have  
343 their negative effect on bone. In this context there may be a parallel with the clinical situations of  
344 osteoporosis and osteonecrosis. Therapeutic glucocorticoid administration can cause both but the  
345 development of clinically significant osteonecrosis is generally much rarer than that of osteoporosis.

346

347 Glucocorticoid treatment has been demonstrated to induce significant structural changes in the  
348 environment of the osteocyte. Glucocorticoids adversely affect fluid flow in the canalicular network.  
349 <sup>77</sup> This effect would be expected to have detrimental effects on osteocyte health but could also  
350 directly influence bone mechanical strength through effects on bone tissue hydration. Glucocorticoid  
351 treatment is also associated with an increase in mean osteocyte lacunar size.<sup>78</sup> There is, in addition,  
352 a reduction in mineralisation in the bone adjacent to the osteocytes, a phenomenon referred to as

353 'osteocytic osteolysis'.<sup>78</sup> This suggests that part of the anatomical pathology involved GIOP could be  
354 microscopic changes to bone mineral properties through periacinar osteolysis or  
355 hypomineralisation. The mechanism by which glucocorticoids cause these changes is not  
356 established. These changes could explain the relatively rapid change in fracture risk during  
357 glucocorticoid treatment and its reversibility. In addition, this type of microarchitectural change  
358 would reduce bone strength disproportionately to the change in BMD measured by DXA, thus  
359 accounting for the increased fracture risk observed for the same level of BMD in GIOP.

360 The group that described the microscopic changes in osteocyte lacunae discussed above have  
361 attempted to define the cellular processes that are responsible for these changes. They found that  
362 glucocorticoid treatment was associated with expression of a range of genes in osteocytes  
363 associated with autophagy.<sup>72,79</sup> Autophagy is a cellular pathway designed to maintain cellular  
364 homeostasis by degrading damaged organelles through formation of autophagosomes. Osteocytes  
365 are reported to respond to glucocorticoid treatment with an increase in autophagy markers and the  
366 accumulation of autophagosome vacuoles. It was hypothesised that autophagy maintained cell  
367 viability in the presence of glucocorticoids.<sup>79</sup> The balance between protective autophagy clearance of  
368 damaged organelles and destruction of key cellular components may shift across tissue sites and  
369 with different glucocorticoid doses making its contribution to glucocorticoid mediated suppression  
370 of osteoblasts in vivo difficult to truly appreciate. Piemontese et al. tested whether genetic  
371 suppression of autophagy was associated with increased sensitivity of osteocytes to  
372 glucocorticoids.<sup>80</sup> They deleted autophagy related gene 7 (Atg7), a gene central to the autophagy  
373 process, from osteocytes using the Dmp1Cre promoter. In control mice glucocorticoids stimulated  
374 autophagy in osteocytes and this was blocked in transgenic mice. However, there was no impact of  
375 autophagy suppression on the effects of glucocorticoids on bone. Interestingly, chemical inhibitors  
376 of autophagy have demonstrated protection against glucocorticoid induced bone loss and  
377 maintained bone formation.<sup>81</sup> However, autophagy had now been reported to occur in osteoclasts  
378 exposed to glucocorticoids.<sup>82,83</sup> Selective deletion of Atg7 in osteoclast precursors suppressed

379 glucocorticoid induced increases in bone resorption and bone loss in mice without any impact on  
380 osteoblast differentiation.<sup>82</sup> Currently it appears that glucocorticoids induce autophagy in both  
381 osteocytes and osteoclasts but that the process in osteoclasts but not osteocytes impacts on bone  
382 strength.

383 The osteocyte is now clearly established as being central to the process of bone remodelling through  
384 secretion of several key regulators of bone physiology.<sup>84</sup> Osteocytes can generate OPG and RANKL.  
385 Glucocorticoids down regulate the production of OPG in osteocytes whereas the expression of  
386 RANKL appears unchanged.<sup>85,86</sup> Osteocyte secretion of RANKL appears to be a requirement for loss  
387 of cortical bone in mice treated with glucocorticoids.<sup>86</sup> In this study glucocorticoids did not directly  
388 regulate RANKL in osteocytes but rather reduced the expression of OPG which allowed greater  
389 activity of the RANKL present. The osteocyte is also an important producer of wnt signalling  
390 antagonists such as sclerostin and DKK1. Glucocorticoids appear to increase the production of both  
391 sclerostin and DKK1 by osteocytes.<sup>51</sup> The role of sclerostin in glucocorticoid induced bone loss has  
392 been examined in studies using anti-sclerostin antibodies and animals with genetic knockout of  
393 sclerostin. Treatment of mice with anti-sclerostin antibodies prevented the glucocorticoid induced  
394 reduction in bone formation seen with placebo treated mice.<sup>87</sup> One study reported that these  
395 antibodies protected mice from glucocorticoid induced osteocyte apoptosis.<sup>88</sup> A study using  
396 sclerostin/Sost knockout mice found that sclerostin deficiency protected against glucocorticoid  
397 induced bone loss but did not protect against a decrease in bone formation or an increase in  
398 osteoblast/osteocyte apoptosis.<sup>85</sup> The protection appeared due to preservation of OPG levels and a  
399 protection against increased bone resorption that was seen in wild type mice.<sup>85</sup>

400 These studies suggest that all of these molecules may have a role in different aspects of the effects  
401 of glucocorticoids depending on the model used. With regards to sclerostin and DKK1, it is currently  
402 unclear which particular pathway is most relevant in animals and humans. In particular it is not clear  
403 whether they both have essential roles or if there is compensation or redundancy between them.

404

405 Controversies in the field.

406 A major issue is that osteocyte apoptosis should leave long standing consequences on bone since  
407 these cells are thought to be long lived. However, epidemiological studies indicate that the increased  
408 risk of fracture during treatment with glucocorticoids declines rapidly when the treatment is  
409 discontinued. It is possible that in humans there is a spectrum of osteocytic damage that can  
410 manifest as osteoporosis if the degree of osteocyte damage is modest, but as frank osteonecrosis if  
411 the degree of osteocyte damage is more extensive. It is also not clear why some studies in mice do  
412 not show any evidence of osteocyte apoptosis even when there are clearly negative effects of  
413 glucocorticoid treatment on other aspects of bone health.

414 A further limitation is that the transgenic models discussed above which target glucocorticoid  
415 receptor signalling in osteoblasts also disrupt glucocorticoid signalling in osteocytes. To date, no  
416 osteocyte or osteoblast specific GR deletion model has been produced and evaluated in the context  
417 of GIOP. As such the relative contributions of osteoblasts and osteocytes to the observed  
418 phenotypes are not clear. It is possible that there may be independent contributions from both of  
419 these cell types which require a fuller exploration.

420

421 II.I.III Effects on osteoclasts:

422 The effects of glucocorticoids on osteoclasts have been examined in vitro and in vivo with indirect  
423 inferences being made through clinical studies. The examination of the role of osteoclasts in GIOP  
424 has been complicated since glucocorticoids appear to have direct effects on osteoclasts or their  
425 precursors but also have powerful indirect influences on osteoclastogenesis and osteoclast function  
426 via effects on osteoblasts and osteocytes.

427 In vitro studies have shed light on the direct actions of glucocorticoids in human osteoclasts where  
428 they increase resorption activity and pit formation.<sup>89,90</sup> High doses of glucocorticoids are used in  
429 culture media to promote the growth and differentiation of osteoclasts.<sup>91</sup> Greater mechanistic  
430 insights into these observations have come from murine osteoclast culture studies where addition of  
431 glucocorticoids prolongs longevity through their activation of the GR receptor.<sup>92,93</sup> However, these  
432 same studies have identified that therapeutic glucocorticoids are also able to suppress osteoclast  
433 differentiation and activation in vitro by increasing apoptosis and interfering with cytoskeletal  
434 reorganisation and rendering them less responsive to the pro-osteoclastogenic actions of M-CSF.<sup>92,93</sup>  
435 Similarly, osteoclast activity is suppressed within cultures of osteoclast in rats as a result of increased  
436 apoptosis in response to glucocorticoids.<sup>94</sup> Overall, similar to what has been found in osteoblasts,  
437 the effects of glucocorticoids on osteoclast formation and bone resorbing capacity appear to be dose  
438 dependent with mostly stimulatory actions at low concentrations and inhibitory effects at very high  
439 concentrations.

440 The effects of glucocorticoids on osteoclasts in vivo have been examined in murine models of  
441 glucocorticoid excess.<sup>93,95,96</sup> Bone resorption/osteoclast activity is increased during early treatment  
442 with glucocorticoids supporting in vitro observations that glucocorticoids increase the survival of  
443 osteoclasts. Targeted abrogation of glucocorticoid signalling in osteoclasts using 11 $\beta$ -HSD2  
444 expression resulted in protection against this initial increase in osteoclast activity in mice treated  
445 with prednisolone.<sup>93</sup> However, with prolonged exposure to high levels of glucocorticoids the number  
446 of osteoclasts is reduced due to a delay in the differentiation of new osteoclasts.<sup>92</sup>

447 The most provocative studies in this area examined the deletion of the GR in osteoclasts using the  
448 LysM<sup>CRE</sup> transgene.<sup>92</sup> LysM<sup>CRE</sup> is expressed in cells of the monocyte/macrophage lineage including  
449 osteoclasts. The mouse strain had a mixed 129/C57 genetic background. Rather than generating an  
450 osteoclastic phenotype the main consequence of osteoclast GR deletion was unexpectedly protected  
451 against the fall in bone formation during treatment with dexamethasone (10mg/kg daily injections)

452 as assessed by mineral apposition rate and serum osteocalcin levels. The osteoblasts did not appear  
453 to be protected against glucocorticoid-induced apoptosis. This study implies that at least some of  
454 the effects of glucocorticoids on the osteoblast might be mediated via the osteoclast. No mechanism  
455 for such communication was identified. Although not examined in this context several established  
456 signalling pathways by which osteoclasts can potentially suppress bone formation have been  
457 reported<sup>97,98</sup> giving these findings plausibility despite their sharp contrast with most of the existing  
458 literature.

459 Similar studies using osteoclast GR deletion have failed to show the same effect. Prednisolone  
460 treatment of another mouse (Balb/c background) with GR knockout using the LysM<sup>CRE</sup> transgene did  
461 not demonstrate any protection against the reduction in bone formation as assessed by bone  
462 formation rate.<sup>50</sup> Likewise, the expression of 11 $\beta$ -HSD2 within osteoclasts using the tartrate resistant  
463 acid phosphatase (TRAP) promoter in the FVB/N mouse strain failed to protect mice against a  
464 decrease in bone formation as assessed by serum osteocalcin levels in response to treatment with  
465 slow release prednisolone pellets.<sup>93</sup> It is possible that these differences relate to subtle differences in  
466 strain, glucocorticoid dose or experimental set up. For instance the data regarding the effect of  
467 osteoclasts on bone formation examined the growth of the calvarial bone surface.<sup>92</sup> As discussed  
468 earlier the outer cortex of bone seems to respond differently to glucocorticoids<sup>67</sup> and thus the  
469 choice of surface may be an important factor in these results. Overall, given that there is only one  
470 study in support, on the current balance of evidence a significant role for osteoclasts in  
471 glucocorticoid induced suppression of bone formation throughout the skeleton appears unlikely.

472

473 Summary:

474 Glucocorticoids have multiple effects on osteoblasts, osteocytes and osteoclasts (summarised in  
475 figure...). Many of these effects appear to be individually very powerful in determining specific

Comment [M5]: Need to add figure.

476 phenotypes when examined in mouse models and this implies that no single mechanism is likely to  
477 mediate all of the effects seen in the clinical setting. Reduced bone formation at trabecular bone  
478 sites and increased endocortical resorption appear to be the most consistent pathological findings.  
479 The results appear to indicate that the osteocyte is the most important target of glucocorticoids but  
480 several major signalling pathways and cellular processes are all affected simultaneously. It should  
481 also be noted that none of the studies described above examined the effects of glucocorticoids in  
482 the context of inflammation. Clinical studies of patients treated with therapeutic glucocorticoids for  
483 various conditions consistently demonstrate that inflammation and the activity of the underlying  
484 disease being treated can have a substantial effect on bone independent of glucocorticoid use or  
485 more likely through complex interactions between glucocorticoids, the underlying illness and bone  
486 metabolism.

487

## 488 II.II Other endocrine and non-endocrine effects on bone

489 Glucocorticoids have effects on bone independent of their direct actions on bone cells. These effects  
490 are however difficult to study in animal models, particularly those that do not simulate an underlying  
491 disease being treated. Therapeutic glucocorticoids are well known to reduce sex steroid levels and  
492 this could have an adverse impact on bone.<sup>99</sup> The reduction in sex steroid levels is likely to be greater  
493 in people with serious inflammatory illness which in itself is likely to impact on the hypothalamo-  
494 pituitary-gonadal axis.<sup>100</sup> Evidence in support of this notion comes from clinical trial data which  
495 indicate that premenopausal women are relatively protected against the effects of glucocorticoids  
496 on fracture risk.<sup>101</sup> Clinical studies have indicated that estrogen treatment of post-menopausal  
497 women<sup>102,103</sup> or testosterone (but not nandrolone) treatment of men<sup>104</sup> taking glucocorticoids results  
498 in an increase in spine but not hip bone density (as measured by DXA). No fracture data are  
499 available. In women taking glucocorticoids who are already taking HRT there does not appear to be

500 any increase in bone density with continued use whereas the addition of intermittent PTH injection  
501 substantially improves BMD at the spine.<sup>105</sup>

502 Glucocorticoids also have complex effects on calcium, vitamin D metabolism and parathyroid  
503 hormone. The literature relating to these actions is relatively old but indicates that glucocorticoids  
504 interfere with intestinal calcium absorption and increases renal calcium excretion.<sup>106</sup> Early research  
505 also suggested a role for altered parathyroid hormone levels in the pathogenesis of GIOP.<sup>107</sup>  
506 However, a comprehensive review failed to find strong evidence for a role of parathyroid hormone  
507 in the detrimental effects of glucocorticoids on bone.<sup>108</sup>

508 The effects of glucocorticoids are also attributable to changes in other circulating or locally produced  
509 hormones. The GH/IGF1 axis is known to have anabolic effects on bone growth and bone density.<sup>109</sup>  
510 These hormones are suppressed by high levels of glucocorticoids. Many in vitro studies have  
511 indicated that the suppressive effects of glucocorticoids on osteoblast function can be partially  
512 reversed by GH and/or IGF1 treatment.<sup>109</sup> However, there are very few clinical studies examining this  
513 issue. Small studies in which children taking glucocorticoids for inflammatory bowel disease or  
514 arthritis were treated with GH indicated that GH therapy could improve some measures of bone  
515 formation and reverse effects of glucocorticoids on growth but these studies lacked control  
516 groups.<sup>110,111</sup>

517 Glucocorticoids also have adverse effects on muscle strength, which known to influence bone  
518 strength through mechanical loading<sup>112</sup>. This association of glucocorticoids with muscle strength is  
519 well characterised in Cushing's disease where proximal myopathy is a characteristic and relatively  
520 specific feature of glucocorticoid excess. These actions are mediated through inhibition of  
521 myogenesis and increased proteolysis and atrophy of muscle fibres.<sup>113-115</sup> As a consequence,  
522 glucocorticoid treatment appears to be a risk factor for falls. However, as discussed elsewhere, in a  
523 disease situation it is very difficult to disentangle the effect of glucocorticoid treatment from that of  
524 the underlying disease. Indeed, in certain inflammatory myopathies including polymyositis and



525 dermatomyositis the application of therapeutic glucocorticoids protects against muscle wasting  
526 through the suppression of disease activity.<sup>116</sup> Some evidence supporting a positive role of  
527 endogenous glucocorticoids in maintaining muscle mass during inflammatory disease comes from  
528 mice in which 11 $\beta$ -HSD1 expression in muscle has been deleted.<sup>117</sup> These mice have normal muscle  
529 size and characteristics in the basal state but in response to inflammation, muscle loss is much  
530 greater in 11 $\beta$ -HSD1 deficient mice. These data highlight the clinical dilemmas in treating  
531 inflammatory muscle and joint diseases with glucocorticoids, where such treatment may result in  
532 detrimental, neutral or strongly positive effects on muscle strength and falls risk, depending on the  
533 impact of glucocorticoids on the underlying illness.

534

### 535 II.III Metabolic consequences mediated through bone cells

536 In addition to the deleterious effects of therapeutic glucocorticoids on bone, these medications are  
537 also associated with an increased risk of impaired glucose tolerance, diabetes or, in people with pre-  
538 existing diabetes, worsening of diabetic control.<sup>118,119</sup> These effects have previously been assumed to  
539 be due to actions of glucocorticoids on tissues classically associated with insulin secretion or  
540 sensitivity such as the liver, muscle and pancreas.<sup>118</sup> However, recent studies suggest that there may  
541 in addition be a role for bone in the development of dysmetabolism associated with glucocorticoid  
542 treatment<sup>120</sup>.

543 Multiple studies in mice indicate that osteocalcin has metabolic effects. In particular osteocalcin, and  
544 in particular the uncarboxylated form of osteocalcin, appear to improve glycemic control and insulin  
545 sensitivity through effects on insulin secretion and insulin sensitivity.<sup>121</sup> Cross-sectional studies in  
546 humans also demonstrate correlations between serum undercarboxylated osteocalcin and diabetes  
547 risk.<sup>122</sup> The exact molecular pathways affected are unclear particularly since the identity of the

548 osteocalcin receptor(s) is still uncertain, although some candidates have been proposed such as the  
549 GPRC6A receptor.<sup>123</sup>

550 Given that the level of osteocalcin in the circulation is dramatically reduced by therapeutic  
551 glucocorticoids it was hypothesised that some of the effects of glucocorticoids on systemic  
552 metabolism might be mediated by the glucocorticoid induced reduction of circulating osteocalcin  
553 concentrations. Studies using 11 $\beta$ -HSD2 transgenic mice which, as discussed above, selectively  
554 express 11 $\beta$ -HSD2 in osteoblasts and osteocytes, demonstrated that the glucocorticoid induced  
555 reduction in serum osteocalcin levels was substantially reduced when glucocorticoid-signalling in  
556 osteoblasts was disrupted. Furthermore, these mice had preserved glucose tolerance compared to  
557 littermate control mice that did not have 11 $\beta$ -HSD2 expression in bone.<sup>121</sup> This protection against  
558 the effects of glucocorticoids on glucose tolerance was also seen in wild type mice in which  
559 osteocalcin was heterotopically and constitutively expressed in the liver. This rescuing of the  
560 dysglycemic phenotype was seen with heterotopic expression of either wild type osteocalcin (which  
561 would be expected to be carboxylated in vivo) or a mutant form of osteocalcin which lacked the  
562 ability to be carboxylated. It is unclear whether similar protection might exist in humans.

563 An alternative approach has been to examine any possible impact of bone active treatments in  
564 patients treated with glucocorticoids. It is known that bisphosphonates reduce, and teriparatide  
565 stimulates osteocalcin synthesis and as such these treatments might result in differences in glycemic  
566 control when used in the treatment of GIOP. One small prospective study involving 111 people  
567 taking glucocorticoids that were treated with either bisphosphonates or teriparatide reported a  
568 small but significant decrease in HbA1c in people who took teriparatide whereas there was no  
569 change in HbA1c in those that took bisphosphonates or just calcium and vitamin D.<sup>124</sup>

570 Although osteocalcin has been the most studied mediator of effects of bone on systemic metabolism  
571 it is likely that other pathways exist. These pathways have been reviewed elsewhere but have not  
572 yet been examined in the context of glucocorticoids and bone.<sup>125,126</sup>

573 Given that there is a role for excess glucocorticoids in the development of systemic dysmetabolism  
574 there is a possibility that endogenous glucocorticoids have a similar influence on systemic  
575 metabolism via an action on bone. Circumstantial evidence for this exists in that transgenic deletion  
576 of 11 $\beta$ -HSD1 globally protects mice against the adverse effects of glucocorticoids on energy  
577 metabolism.<sup>127</sup> However, deletion of 11 $\beta$ -HSD1 in classical target tissues of glucocorticoids such as  
578 liver, fat or muscle failed to prevent these effects suggesting that other tissues also contribute to the  
579 adverse metabolic phenotype seen with glucocorticoid exposure.

580

581 **III Endogenous glucocorticoids and bone**

582 In this section we will review data relating to the effects of endogenous glucocorticoid excess on  
583 bone. We will review the bone phenotype in Cushing's disease but also more subtle states of  
584 autonomous states of circulating glucocorticoid excess. We will then examine the role of tissue  
585 specific changes in glucocorticoid action focussing primarily on the role of 11 $\beta$ -hydroxysteroid  
586 dehydrogenase enzymes.

587

588 III.I States of circulating glucocorticoid excess, Cushing's disease and autonomous cortisol  
589 production

590 III.I.I Bone disease in Cushing's disease/syndrome.

591 As discussed in section II.1, there is strong evidence that endogenous glucocorticoids are required  
592 for normal bone metabolism and osteoblastogenesis.<sup>63,128</sup> In contrast, in Cushing's disease and other  
593 clear cut forms of endogenous circulating glucocorticoid excess, there is normally a substantial  
594 negative impact on bone (reviewed in Toth and Grossman<sup>129</sup>). This was recognised early on by  
595 Cushing, and bone related complications of Cushing's disease are clearly evident in clinical practice.<sup>1</sup>  
596 More recent studies that have attempted to quantify the bone effects of endogenous Cushing's  
597 syndrome have generally been small (up to around 180 patients) and varied according to the  
598 number of patients with each underlying cause of Cushing's (pituitary, adrenal, ectopic,  
599 adrenocortical cancer etc.), and length of time before diagnosis was made. Despite this diversity, the  
600 studies have been very consistent in indicating a substantially increased risk of fracture (typically a  
601 fracture prevalence of 50% is reported) and a greater chance of having very low bone density (below  
602 a T-score of -2.5) when assessed by DXA. The incidence and prevalence figures depend on the extent  
603 to which fractures are searched for. In a self-report survey of 125 patients with endogenous  
604 Cushing's syndrome and age and sex matched controls, fracture risk appeared to be elevated

605 substantially in the 2 years prior to diagnosis with an incidence rate ratio of 6 in patients with  
606 Cushing's syndrome.<sup>130</sup> Interestingly there was no evidence of an increased risk of fracture prior to 2  
607 years before the diagnosis. Additionally, after successful treatment the reported fracture rate was  
608 also no different from that of controls. Although an important study with a high (83%) response rate,  
609 the data was limited by the study's focus on clinical fractures.

610 This type of analysis cannot accurately determine the risk of vertebral fractures. Vertebral fractures  
611 are frequently misdiagnosed or missed. As a consequence, the overall rate of vertebral fractures of  
612 any origin is usually grossly underestimated unless examined for specifically.<sup>131</sup> Vertebral fractures,  
613 even if asymptomatic, are amongst the strongest risk factors for further fracture and premature  
614 mortality.<sup>131,132</sup> The standard approach to the diagnosis of vertebral fractures is to examine for loss  
615 of vertebral height on spine radiographs using the Genant classification (with a fracture defined as a  
616 loss of anterior vertebral height of 20% or more)<sup>133</sup>. A study by Tauchmanova et al. focussed  
617 particularly on the risk of spine fractures in patients with endogenous Cushing's syndrome of various  
618 etiologies and examined spine radiographs in cases and controls.<sup>134</sup> In an analysis of 80 patients and  
619 80 controls, vertebral fractures were present in a remarkable 76% of patients with Cushing's. In an  
620 equally remarkable 85% of patients with a vertebral fracture, multiple fractures were present. Only  
621 24% of spine fractures were known to the patient.

622 A comprehensive and contemporary analysis of bone disease in a cohort of patients with  
623 endogenous Cushing's syndrome was reported by Belaya et al.<sup>135</sup> All patients had chest radiographs  
624 and AP and lateral spine radiographs. In 182 patients studied, 81 patients had fractures. 70 of these  
625 patients had fractures of the spine. 53 out of these 70 patients had multiple vertebral fractures. Out  
626 of over 150 fractures just 7 were non rib, non-vertebral fractures. These figures indicate that  
627 prevention of spine fractures should be the major skeletal priority in patients with endogenous  
628 Cushing's syndrome.

629 Although Cushing's syndrome is associated with bone loss (as assessed by DXA) and osteoporotic  
630 fractures, the utility of bone mineral density scans in predicting fracture is limited. In the studies  
631 described above, fractures (and in particular spine fractures) occurred in some patients with  
632 relatively well preserved BMD.<sup>134</sup> In the largest study, bone density measured by DXA was not  
633 predictive of fracture in a multivariable model that took into account the severity of  
634 hypercortisolaemia.<sup>135</sup> The only predictor of fracture in this study was the severity of Cushing's.  
635 However in an earlier study spine BMD was a predictor of vertebral fracture in Cushing's.<sup>134</sup> Since the  
636 severity of Cushing's is associated with reductions in bone mass it is likely that the severity of  
637 Cushing's and the decrease in measured BMD both provide clinically useful information in the  
638 assessment of fracture risk in these patients. Trabecular bone score (TBS; a non-invasive measure of  
639 trabecular bone architecture derived from spine DXA scans) has also been evaluated in patients with  
640 Cushing's.<sup>135</sup> Values were found to be significantly reduced (indicating impaired trabecular bone  
641 structure) but the scores did not have predictive value in estimating the risk of vertebral bone  
642 fracture. Advanced imaging techniques such as high resolution peripheral quantitative CT (HR-pQCT)  
643 and hrQCT of vertebral bone, and techniques for in vivo examination of material properties such as  
644 microindentation, which have all been used in patients treated with glucocorticoids, have not yet  
645 been reported in patients with Cushing's.

646 Other potential predictors of fracture have also been examined in Cushing's. In one study fracture  
647 risk at the spine appeared to be independent of the presence of menstrual irregularities with  
648 amenorrheic women having a similar risk of fracture and BMD to those with eumenorrhea.<sup>134</sup>  
649 Another study however suggested that reduction in BMD was more likely in women with estrogen  
650 deficiency.<sup>136</sup> Fracture risk was higher in patients with ectopic ACTH syndrome, presumably as a  
651 result of the higher cortisol levels usually found in this condition.<sup>134</sup> Serum osteocalcin levels have  
652 also been associated with fracture risk but again this relationship appears to be mediated by the  
653 levels of cortisol present.<sup>135</sup> In terms of prediction of changes in BMD, the correlation between the  
654 extent of reduction in BMD and degree of cortisol excess has been reported in eumenorrheic women

655 with Cushing's.<sup>137</sup> The extent of reduction of BMD in patients with Cushing's has also been  
656 associated with the duration of disease.<sup>136</sup>

657 Whereas fracture risk in Cushing's has been quantified in only a small number of studies, changes in  
658 biochemical markers of bone turnover have been assessed in at least 16 reports (reviewed in<sup>129</sup>). A  
659 finding in all but one of these studies is that serum osteocalcin levels are considerably decreased in  
660 Cushing's. The results for other formation markers (PINP, PICP, alkaline phosphatase) show less, if  
661 any, change. Bone resorption markers do not appear to change in a consistent fashion in Cushing's.  
662 The sensitivity of osteocalcin expression to glucocorticoids is well known and in this situation serum  
663 osteocalcin levels might be viewed as a marker of bone tissue glucocorticoid exposure rather than a  
664 true bone formation marker. The relationship of low serum osteocalcin with excessive cortisol levels  
665 is so strong that serum osteocalcin has been proposed as a diagnostic marker of Cushing's  
666 syndrome.<sup>138</sup> In a group of patients with Cushing's syndrome serum osteocalcin levels were found to  
667 be highly correlated with serum cortisol measured at 0800 hrs, 2400 hrs and after a low dose  
668 dexamethasone suppression test. In a follow up study, the diagnostic utility of serum osteocalcin in  
669 patients presenting with obesity and risk factors for Cushing's syndrome was evaluated.<sup>139</sup> It was  
670 found that osteocalcin had a sensitivity of 74% and a specificity of 97% for the identification of  
671 Cushing's syndrome. Additional prospective studies will be required to fully evaluate the clinical  
672 utility of osteocalcin as a diagnostic tool in Cushing's syndrome.

673 The changes in bone status in response to successful therapy have also been evaluated. In the self-  
674 report survey of patients with Cushing's syndrome described above the risk of fracture did not  
675 appear to be elevated after treatment.<sup>130</sup> This study was likely to have modest sensitivity in terms of  
676 fracture detection given the relatively small number of patients available and the lack of detailed  
677 analysis of spine fractures. Studies consistently report a rise in BMD after successful treatment.<sup>140-143</sup>  
678 Although pre-disease BMD is clearly not available in the majority of people it is reported that the  
679 deficit in bone mass is largely reversible, at least in younger patients.<sup>142</sup> These changes appear

680 relatively complex and vary between skeletal sites. Successful treatment is associated with an  
681 improvement in spine areal BMD but also an increase in bone area.<sup>140</sup> This suggests the possibility of  
682 new bone being laid down on the outside of the vertebral bones (periosteal apposition) when  
683 glucocorticoid levels are restored to normal. Intriguingly, after successful treatment bone density  
684 and bone area at the wrist were actually reported to decrease.<sup>140</sup> Although the authors proposed  
685 that this reflects a redistribution of bone from the appendicular to the axial skeleton it is unclear  
686 how such redistribution might occur, particular in relation to bone area, as this would require  
687 removal of bone from the outer cortex of the bone. The results are in keeping with the findings in  
688 mice that formation of bone at the outer cortex of some bones of the peripheral skeleton is actually  
689 stimulated by glucocorticoids rather than being suppressed.<sup>66</sup> Serum osteocalcin also increases  
690 rapidly after treatment.<sup>138</sup> Whereas there is no correlation between serum osteocalcin and other  
691 bone markers prior to treatment, shortly after successful treatment a strong correlation between  
692 osteocalcin and bone resorption markers develops (as is normally seen in populations of healthy  
693 people).<sup>138</sup> As such, serum osteocalcin levels appear to primarily reflect cortisol levels in patients  
694 with Cushing's prior to treatment, in treated patients they behave more like a traditional marker of  
695 bone formation. Although the data in general suggest a reversal of bone disease in patients  
696 successfully cured caution should be taken if patients need long term glucocorticoid replacement  
697 after cure. In a group of patients successfully cured the continuing use of glucocorticoid replacement  
698 was associated with reductions in BMD, BMC and osteocalcin compared to matched controls.<sup>136</sup>  
699 These effects were most evident in women with coexisting estrogen deficiency. This exaggerated  
700 sensitivity of estrogen deficient women is in keeping with the greater risk of fracture of post-  
701 menopausal women treated with therapeutic glucocorticoids (discussed in section IV.1). As such,  
702 glucocorticoid replacement must be particularly carefully monitored in this group.

703

704 III.I.II Bone disease is autonomous cortisol secretion.



705 More subtle states of glucocorticoid excess also appear to detrimentally impact on bone. Most  
706 attention has focussed on the concept of subclinical endogenous hypercortisolism, also referred to  
707 as subclinical Cushing's and more recently autonomous cortisol secretion.<sup>144,145</sup> This condition is in  
708 principle defined by abnormal cortisol secretion in the absence of clinical features of glucocorticoid  
709 excess. It is usually associated with nodules of the adrenal cortex (adrenal incidentaloma, AI). The  
710 condition is controversial and the best diagnostic criteria have yet to be established. In various  
711 studies the criteria differ but the most common component of the diagnosis is failure to suppress  
712 serum cortisol after a 1mg dexamethasone suppression test (DST).<sup>144</sup> Depending on the definition  
713 the condition appears to be relatively common and is driven by the background prevalence of AIs.  
714 The prevalence of AIs based on radiographic series depends heavily on age but it is estimated that  
715 3% of people aged 50 have an adrenal nodule whereas up to 10% of elderly individuals may have  
716 AIs.<sup>144</sup> It has been estimated that up to 30% of patients with AIs have some degree of autonomous  
717 cortisol secretion<sup>7</sup> and as such up to 1-3% of the population aged 50 and above might have  
718 autonomous cortisol secretion.

719 The research examining the relationship between bone health and the presence of autonomous  
720 cortisol secretion (usually in the context of patients known to have AI) is dominated by the studies of  
721 Chiodini and colleagues. These include cross-sectional, longitudinal, retrospective and prospective  
722 studies examining bone density, bone markers and fracture prevalence and incidence in these  
723 individuals. Most studies reported a reduction in BMD at the spine as assessed by either DXA<sup>146,147</sup> or  
724 qCT.<sup>148,149</sup> Trabecular bone score has also been reported to be lower and to predict the development  
725 of fracture in this group of patients.<sup>147</sup> As with bone changes in Cushing's disease the data relating to  
726 the change in BMD at the hip is less clear with some studies indicating a reduction in BMD and some  
727 no change. Differences between studies are likely due to the relatively small number of patients  
728 examined in most studies and heterogeneity in the proportions of men, pre-menopausal and post-  
729 menopausal women. Again, in a similar fashion to that seen with endogenous Cushing's syndrome,  
730 autonomous cortisol secretion is reported to be associated with a decrease in blood osteocalcin

731 levels but no consistent changes in other markers or bone formation or markers of bone resorption.

732 These studies were however relatively small with typically less than 50 patients.

733 The most dramatic findings in these studies are the presence of vertebral fractures. A recent meta-

734 analysis of these studies found that the prevalence of radiographically identified vertebral fractures

735 was 63.6% (CI 56-71%) in patients with autonomous cortisol secretion compared to a prevalence of

736 16% (CI 5-28) in controls.<sup>7</sup> Interestingly, patients known to have AI that do not meet the criteria for

737 autonomous cortisol secretion were reported to have a higher prevalence of spine fractures (28%)

738 than controls without AI (20-35). This suggested to the authors that some patients with AI with

739 excessive production of cortisol might not be detected by current tests and by implication that all AIs

740 are a risk factor for fracture. The authors of the meta-analysis could not identify any patient related

741 factors that predicted the development of fractures beyond the presence of autonomous cortisol

742 secretion. A study of 570 patients with AIs attempted to determine the threshold cortisol level post

743 1mg DST (using an Abbott TDxFLs cortisol assay) which is best able to predict the presence and the

744 future development of vertebral fractures.<sup>150</sup> It was found that a post DST cortisol level of greater

745 than 2.0 microgram per decilitre (55 nmol/L) was the best criteria in both situations with sensitivities

746 and specificities between 68 and 80%. The presence of cortisol levels above this threshold was

747 associated with an odds ratio of fracture of over 10.

748 These prevalence and incidence rates of vertebral fracture in people with AI are extremely high and

749 could represent a large burden of disease that is currently not being addressed. However, the

750 proportion of these fractures that actually cause symptoms or otherwise impact on patient well-

751 being is unknown. Although difficult to perform, future trials would ideally aim to determine

752 whether AIs (with or without autonomous cortisol secretion) are associated with a greater risk of

753 clinical vertebral fractures, height loss, kyphosis development or reduced quality of life relating to

754 musculoskeletal health. An alternative way of assessing whether autonomous cortisol secretion

755 relating to AIs is associated with clinically significant vertebral fracture is to examine the prevalence

756 of these abnormalities in patients presenting with clinical vertebral fracture. In one study 7 out of 65  
757 patients presenting with osteoporosis and spine fracture were found to have subclinical  
758 hypercortisolism.<sup>151</sup> In a subsequent study of over 600 patients with osteoporosis and no apparent  
759 cause the rate of subclinical hypercortisolemia was significantly lower at 1.3%.<sup>152</sup> These patients  
760 however had a relatively low rate of reported fracture and in particular of clinical vertebral fracture.  
761 On the basis of what is known about endogenous Cushing's and subclinical Cushing's it is reasonable  
762 to assume that the development of a vertebral fracture rather than just a low BMD by DXA would be  
763 a more sensitive indicator of the presence of abnormal cortisol secretion.

764 Remaining questions in this area include the most appropriate treatment approach to a patient with  
765 bone disease related to autonomous cortisol secretion by AI and when and how to investigate for  
766 the presence of autonomous cortisol secretion in patients presenting with bone disease. A recent  
767 study suggested that adrenalectomy was effective at reducing the risk of new vertebral fracture over  
768 a follow up period of 28-40 months.<sup>153</sup> This study was limited by a lack of randomisation. In the  
769 absence of randomised clinical trials it would be reasonable to consider the option of adrenalectomy  
770 in patients with AI, autonomous cortisol secretion and bone disease, particularly in the presence of  
771 other conditions that might be exacerbated by cortisol excess such as hypertension and diabetes. An  
772 additional option is the use of medications that are proven to be effective in the treatment of  
773 idiopathic osteoporosis or GIOP associated with therapeutic glucocorticoid use. There is additionally  
774 data based on a small number of people that indicates that the bisphosphonate clodronate is  
775 effective in increasing BMD at the lumbar spine in subclinical Cushing's.<sup>154</sup> No guidelines are  
776 available for use of possible medical therapies in this particular situation. The use of treatment  
777 should also consider that the most common site of fracture in this condition is the spine, BMD can  
778 be selectively reduced at the spine and bone density may not fully predict fracture risk associated  
779 with glucocorticoid excess. Fracture risk calculators such as FRAX and the Garvan Fracture Risk  
780 calculator are based on hip density and might underestimate the risk of vertebral fractures in this  
781 condition.

782 It remains unclear whether patients with post-menopausal and age related osteoporosis and no  
783 symptoms of hypercortisolemia should be tested routinely for the condition. A pragmatic approach  
784 at the current time would be to test those individuals who have a higher likelihood of cortisol excess  
785 e.g. people presenting with vertebral fractures, people with BMD values that are highly discordant  
786 between spine and hip with spine being low, and people with non-traumatic fractures that occurred  
787 in the context of relatively normal bone density. The most appropriate test to identify people with  
788 excess cortisol secretion that is likely to impact on bone would appear to be the 1mg DST with a cut-  
789 off of 2 micrograms per decilitre (55 nmol/L) (although these values should be adjusted based on the  
790 performance of the local cortisol assay).<sup>7,150,151</sup> Clearly the distinction between autonomous cortisol  
791 secretion and overt Cushing's syndrome might be difficult in these situations where there is clear cut  
792 bone disease in association with abnormal cortisol secretion. In these situations additional  
793 investigations are required to determine the basis for the abnormal cortisol levels.

794

795 III.I.III Bone impact of physiological variation in the HPA axis.

796 It is possible that individual variations in the circulating level of endogenous glucocorticoids might  
797 also have an impact on bone even in the absence of any disease or condition affecting the HPA axis.  
798 By examining healthy post-menopausal women before and after treatment with the adrenal  
799 corticosteroid synthesis inhibitor metyrapone it has been established that the circadian variation in  
800 serum osteocalcin is influenced by adrenal cortisol secretion.<sup>155</sup> In the same study, other bone  
801 formation or resorption markers did not appear to be influenced by adrenal function suggesting a  
802 specific sensitivity of osteocalcin to glucocorticoids independent of its role as a marker of bone  
803 formation. Whether variations in adrenal cortisol secretion impacts on bone health has been  
804 primarily examined in studies looking at serum or salivary cortisol levels and differences in bone  
805 health (mostly assessed as BMD by DXA) during ageing. These studies have generally found weak  
806 associations between levels of circulating glucocorticoids at various time of day and either current

807 bone density or change in bone density over time. The results also appear to differ depending on  
808 whether women or men are studied. In the a study of 228 elderly community dwelling people  
809 salivary cortisol levels at 2300 were negatively associated with lumbar spine BMD in women  
810 whereas in men 0700 salivary cortisol levels negatively correlated with spine BMD.<sup>156</sup> In 34 healthy  
811 elderly men that had frequent serum cortisol measurements over a 24 hour period the integrated  
812 serum cortisol level over the 24 hour period was negatively associated with lumbar spine BMD.  
813 Additionally, trough cortisol predicted the rate of bone loss at the spine and femoral neck over the  
814 subsequent 4 years.<sup>157</sup> In a study of over 500 men and women from the Longitudinal Ageing Study  
815 Amsterdam serum fasting cortisol was associated with lower BMD at the femoral neck after  
816 adjustment for age and BMI.<sup>158</sup> A study of 135 elderly women and 171 men examined the  
817 relationship between serum cortisol, serum cortisone, bone markers and BMD.<sup>159</sup> It was found that  
818 serum cortisol had no relationship with any bone measurements but serum cortisone was negatively  
819 associated with serum osteocalcin levels and spine BMD. These relationships were independent of  
820 the levels of cortisol. This study suggests that the relationship between adrenal corticosteroid  
821 production and bone health may, at least in part, be mediated via cortisone. The only other study  
822 that explored the role of cortisone in bone health performed a comprehensive analysis of adrenal  
823 corticosteroid output and metabolism in young males in relation to bone development at the  
824 proximal radius. In this study the level of urinary cortisone metabolites was independently and  
825 negatively associated with reduced bone density.<sup>160</sup>

826 A significant limitation of these studies is their lack of information relating to fractures. Two studies  
827 have however provided information in relation to fracture risk and adrenal corticosteroid  
828 production. A sub-study of the MacArthur Study of Successful Ageing measured overnight (between  
829 2000 to 0800 hrs) urinary free cortisol excretion in 684 men and women aged 70-79 at baseline.<sup>161</sup>  
830 Higher baseline UFC was significantly associated with the incidence of self-reported fractures over  
831 the next 4 years. These relationships appeared to be relatively strong e.g. in the highest quartile of  
832 UFC the adjusted odds of a fracture was over 5. A more recent cross-sectional study examined the

833 relationship between salivary cortisol measurements taken at various times of the day and the TBS  
834 and presence of vertebral fractures.<sup>162</sup> The study involved over 600 women and vertebral fractures  
835 were defined on the basis of Genant grade 2 or greater fractures on spine radiographs. This criteria  
836 (a loss of height of greater than 25%) is more stringent than that typically used in the studies  
837 examining spine fractures in people with subclinical Cushing's described above and would be  
838 expected to increase the clinical significance of these fractures. It was found that salivary cortisol  
839 levels at 2000 hrs were associated with the presence of vertebral fractures and that this relationship  
840 was independent of age and BMD. A negative linear association between 2000 hrs salivary cortisol  
841 and TBS values was also observed. Morning salivary cortisol levels were not found to be associated  
842 with fracture prevalence. In multivariable models both evening salivary cortisol levels and TBS scores  
843 independently predicted the presence of a spine fracture. Although methodologically very different  
844 these two studies strongly support the idea that high exposure to endogenous cortisol levels in the  
845 evening and overnight, even within the normal range, is associated with an increased risk of  
846 fracture.

847

#### 848 III.I.IV Bone impact of variation in glucocorticoid receptor expression.

849 A further possible way in which endogenous glucocorticoid action within bone could be amplified is  
850 an alteration in the sensitivity or number of the glucocorticoid (or mineralocorticoid) receptors  
851 within the cell or an alteration in post-receptor signalling. Several studies have examined the  
852 influence of GR gene (NC3R1) polymorphisms on the sensitivity of bone to glucocorticoids. These  
853 have generally been small and either negative or reported weak and inconsistent associations.  
854 Huizenga et al examined the influence of the N363S polymorphism of the GR gene on various  
855 aspects of glucocorticoid sensitivity and bone composition.<sup>163</sup> Heterozygous carriers of this  
856 polymorphism had greater suppression of serum cortisol levels during a 0.25mg overnight DST  
857 implying greater sensitivity at the level of the GR. In terms of bone density there was a non-

858 significant difference of approximately 0.5 of a Z-score at the spine ( $p=0.08$ ) but no suggestion of a  
859 difference at the hip. This study was additionally limited by the low number of people with the  
860 N363S polymorphism at just 10 compared to over 100 controls without. The gene for the GR has not  
861 been linked to osteoporosis or fracture risk in genome wide association studies suggesting that  
862 variation in the GR is unlikely to be a major factor in the development of these conditions. A possible  
863 reason for this lack of association is that relatively modest changes in GR sensitivity are unlikely to  
864 have consequences as long as normal HPA negative feedback is intact. Any difference in sensitivity  
865 would be expected to be compensated for by small changes in circulating levels of cortisol.

866 GR gene variants that influence glucocorticoid sensitivity could influence the degree of bone damage  
867 that occurs in people with excessive adrenal cortisol production due to disease states or exogenous  
868 glucocorticoid usage. In these situations the HPA negative feedback would be unable to adjust for  
869 difference in glucocorticoid sensitivity. Studies in these situations have suggested a possible impact  
870 of GR variants. Szappanos et al. examined several GR gene variants (N363S, BclI, ER22/23EK and  
871 A3669G) in 60 people with endogenous Cushing's syndrome and 129 healthy controls.<sup>164</sup> They found  
872 that individuals with Cushing's syndrome that were homozygous for the BclI polymorphism had  
873 reduced BMD at the hip by DXA and an increased level of serum betaCTx (a bone resorption marker).  
874 The other polymorphisms did not appear to influence bone. Koetz et al. examined the influence of  
875 the BclI polymorphism in 112 patients with adrenal insufficiency.<sup>165</sup> Patients homozygous for the G  
876 variant (which would be expected to increase cellular glucocorticoid sensitivity) were found to have  
877 greater serum betaCTx and greater urinary NTx. However there was no difference in BMD at hip or  
878 spine. Interestingly these patients were treated with significantly lower doses of replacement  
879 glucocorticoids. This lower dose may have offset the increased tissue glucocorticoid sensitivity.

880 It might also be hypothesised that variants in GR sensitivity would predict the effects of therapeutic  
881 glucocorticoids on bone. However, these studies are likely to be complicated by any impact that  
882 variation in GR sensitivity might have on the activity of the underlying disease being treated. For

883 example, carriers of the N363S or Bcll minor variants (which predict increased glucocorticoid  
884 sensitivity) are reported to have a lower risk of developing rheumatoid arthritis.<sup>166</sup> Likewise, patients  
885 with rheumatoid arthritis that are carriers of the Bcll or N363S variants have lower levels of baseline  
886 disease activity even in the absence of glucocorticoid treatment.<sup>167</sup> It seems likely that any  
887 differences in GR sensitivity through genetic polymorphism will alter bone sensitivity in parallel to  
888 that of the underlying disease requiring treatment.

889

890 Another mechanism by which glucocorticoid action could be altered at a tissue level is through the  
891 active transport of glucocorticoids across cell membranes. Several membrane transporters can  
892 remove certain types of glucocorticoids from the cytoplasm. This is best exemplified by the active  
893 transport of the synthetic glucocorticoid dexamethasone by cells of the blood brain barrier.<sup>168</sup> More  
894 recently the selective transport out of certain tissues of cortisol and corticosterone by ABC  
895 transporters has been highlighted.<sup>9</sup> The clinical relevance of these effects is yet to be fully  
896 established and they have not yet been examined in the context of bone cells or GIOP.

897

### 898 III.II Tissue specific amplification of glucocorticoid action

899 Traditionally glucocorticoid action at a tissue level has been assumed to be closely linked with the  
900 levels of glucocorticoids in the circulation. More recently it has become apparent that there are  
901 additional potential levels of regulation between the circulation and action at a tissue level. The  
902 most extensively examined of these levels is that of tissue 'pre-receptor' glucocorticoid metabolism.  
903 Various enzymes capable of glucocorticoid metabolism are present within bone. The enzymes that  
904 have previously been examined include the 11 $\beta$ -HSDs and 5 $\alpha$ -reductases. Although expression of  
905 the 5 $\alpha$ -reductase type 1 enzyme has been reported<sup>169</sup> the activity of this enzyme in human bone  
906 appears modest and more attention has focussed on the 11 $\beta$ -HSDs.<sup>10,170</sup> There are two 11 $\beta$ -HSD



907 enzymes. 11 $\beta$ -HSD1 is an intrinsically bidirectional enzyme which interconverts hormonally inactive  
908 cortisone (human) and dehydrocorticosterone (DHC) (rodent) with their active counterparts cortisol  
909 and corticosterone respectively.<sup>10</sup> Although bidirectional in most situations in vivo the enzyme acts  
910 principally as an activating enzyme due to the presence of a cofactor generating enzyme hexose-6-  
911 phosphate dehydrogenase.<sup>171</sup> This enzyme provides a supply of NADPH within the endoplasmic  
912 reticulum where 11 $\beta$ -HSD1 is located. 11 $\beta$ -HSD2 by contrast is a powerful glucocorticoid inactivating  
913 enzyme converting active cortisol and corticosterone to inactive cortisone and DHC. 11 $\beta$ -HSD2 is  
914 normally expressed in classical mineralocorticoid sensitive tissues such as kidney, colon and  
915 pancreas, where it protects the MR from binding by glucocorticoids, whilst 11 $\beta$ -HSD1 is more widely  
916 expressed in tissues such as liver, adipose and skin. In terms of expression within bone, 11 $\beta$ -HSD  
917 activity was first recognised in cultured osteosarcoma cells and primary cultures of osteoblasts.<sup>172,173</sup>  
918 In osteosarcoma cells 11 $\beta$ -HSD2 mRNA and activity were detected whereas primary cultures of bone  
919 demonstrated exclusive expression of 11 $\beta$ -HSD1<sup>173</sup>. It is now known that 11 $\beta$ -HSD2 is expressed in a  
920 range of malignant tissues and its presence in osteosarcoma cells is thought to reflect their  
921 malignant status rather than being a feature of bone cells.<sup>60,174</sup> Studies in adult mouse and human  
922 bone demonstrate expression of 11 $\beta$ -HSD1 but not 11 $\beta$ -HSD2.<sup>59,175</sup> Immunohistochemistry and in  
923 situ hybridisation studies demonstrated that the main cell type expressing 11 $\beta$ -HSD1 in bone were  
924 osteoblasts and osteocytes<sup>175</sup>. 11 $\beta$ -HSD1 expression was seen to a lesser degree in osteoclasts.<sup>175</sup> In  
925 vitro expression appeared to vary across osteoblast differentiation with levels being low in immature  
926 cells, rising and reaching a peak in mature osteoblasts.<sup>176</sup> The functional significance of 11 $\beta$ -HSD  
927 expression in bone cells was examined by transfection and stable expression of these enzymes in  
928 osteosarcoma cell lines which do not normally have 11 $\beta$ -HSD activity.<sup>177</sup> Whereas empty vector cells  
929 were unresponsive to cortisone, expression of 11 $\beta$ -HSD1 rendered cells sensitive to cortisone in  
930 terms of reduced proliferation and expression of glucocorticoid responsive bone cell markers.

Comment [M6]: Should have a figure here.

931 The expression and activity of 11 $\beta$ -HSD1 have been shown to be regulated by age, cell  
932 differentiation status, inflammation and by glucocorticoids themselves. Primary cultures of human  
933 osteoblasts demonstrated greater ability to generate cortisol from cortisone when cells were grown  
934 from older compared to younger donors.<sup>178</sup> This relationship was also observed in mice where mRNA  
935 for 11 $\beta$ -HSD1 was increased in bones obtained from old compared to young mice.<sup>77</sup> The  
936 inflammatory cytokines TNF $\alpha$  and IL-1 $\beta$  are powerful stimulators of 11 $\beta$ -HSD1 activity in  
937 mesenchymal derived cell populations such as osteoblasts, and have been proposed as potential  
938 mediators of increased 11 $\beta$ -HSD1 activity in aging.<sup>179,180</sup> This upregulation appears to be via an NF- $\kappa$ B  
939 dependent mechanism, although CCAAT/enhancer-binding protein (C/EBP)  $\beta$  has also been shown to  
940 play a role in this inflammatory induction of 11 $\beta$ -HSD1.<sup>180-182</sup> Glucocorticoids themselves also cause a  
941 modest increase in 11 $\beta$ -HSD1 activity and expression in osteoblasts and they can synergise with pro-  
942 inflammatory cytokines to cause a more dramatic increase in 11 $\beta$ -HSD1 expression.<sup>178,183</sup>

943 Clinical studies also indicate the presence of 11 $\beta$ -HSD1 within bone. In a cohort of elderly subjects  
944 the level of cortisone in the circulation was a significant negative predictor of the blood level of  
945 osteocalcin whereas cortisol was not.<sup>159</sup> This suggested that 11 $\beta$ -HSD1 within osteoblasts is a  
946 regulator of osteocalcin synthesis. A number of relatively small genetic association studies have  
947 suggested that polymorphisms in the 11 $\beta$ -HSD1 gene (HSD11B1) might contribute to the  
948 development of osteoporosis, regulate the level of serum osteocalcin, or increase the risk of  
949 fracture.<sup>184-187</sup> However, the gene has not been identified as a candidate in large GWAS studies. It is  
950 possible that these polymorphisms might be important in some ethnic groups but not others. It also  
951 needs to be considered that 11 $\beta$ -HSD1 is also expressed in other tissues and any associations could  
952 be mediated indirectly e.g. through an effect on the regulation of the degree of inflammation, rather  
953 through an effect on bone cells themselves.

954 The functional role of 11 $\beta$ -HSD1 in bone has been examined in some animal models. In mice with  
955 global deletion of 11 $\beta$ -HSD1 there is no alteration in bone density or structure.<sup>188</sup> However, on the

956 background examined the 11 $\beta$ -HSD1 global knock out mice have an alteration in feedback regulation  
957 of the HPA axis leading to a high level of corticosterone in the circulation.<sup>189</sup> It is possible that this  
958 high circulating level might offset any tissue level reduction in glucocorticoid levels. The global  
959 knockout mouse has not been evaluated in the context of old age or in models of glucocorticoid  
960 excess and inflammation associated osteoporosis. Certainly, in the context of glucocorticoid induced  
961 muscle wasting, skin thinning and hepatic steatosis, global deletion of 11 $\beta$ -HSD1 results in almost  
962 complete protection raising the possibility that these mice will also be protected from glucocorticoid  
963 induced osteoporosis.

964 The 11 $\beta$ -HSD enzymes also regulate the activity of the most widely used oral glucocorticoids  
965 prednisone and prednisolone.<sup>190</sup> 11 $\beta$ -HSD1 converts inactive prednisone to active prednisolone with  
966 similar enzyme kinetics to that of the conversion of cortisone to cortisol. In healthy males the  
967 baseline level of 11 $\beta$ -HSD1 (measured as the ratio of corticosteroid metabolites on a 24 hour urine  
968 collection) predicted the response of bone formation markers to a short course (7 days) of oral  
969 prednisolone. High baseline 11 $\beta$ -HSD1 activity was associated with the greatest falls in serum  
970 osteocalcin and PINP levels. This relationship was independent of the circulating levels of prednisone  
971 or prednisolone. The conclusions from this study are limited due to the activity being measured in  
972 the total body rather than in the bone itself. Additionally, even though the predictive ability of total  
973 measures of 11 $\beta$ -HSD1 activity are predictive of the response of bone to glucocorticoids these  
974 relationships may not persist in patients treated with glucocorticoids for inflammatory disease since  
975 inflammation itself is associated with a tissue specific increase in 11 $\beta$ -HSD1 activity.<sup>191</sup> In patients  
976 with inflammatory bowel disease baseline measures of 11 $\beta$ -HSD1 activity on a urine sample were  
977 not predictive of the change in bone density in response to oral glucocorticoid treatment.<sup>192</sup>

978 Given that inflammation increases 11 $\beta$ -HSD1 activity and activation of therapeutic glucocorticoids  
979 within bone cells, the potential exists for locally activated steroids to both abrogate inflammatory  
980 bone loss whilst directly contributing to glucocorticoid mediated bone loss. Clinical data suggest that

981 the reality may lie somewhere in-between the two, with therapeutic glucocorticoids partially  
982 suppressing disease activity in patients with chronic inflammatory disease and reducing immediate  
983 bone loss whilst ultimately contributing to glucocorticoid-induced osteoporosis with prolonged  
984 use.<sup>193,194</sup> These data support the idea that a rapid and marked increase in 11 $\beta$ -HSD1 in response to  
985 inflammation is an important part of the host response to inflammation, with elevated  
986 glucocorticoid activation preventing inflammatory bone loss in an acute setting. This situation is  
987 complicated in chronic inflammation, where prolonged increases in 11 $\beta$ -HSD1 may begin to promote  
988 bone loss in a similar fashion as seen with long term therapeutic glucocorticoid application.

989 If correct, the targeted inhibition of 11 $\beta$ -HSD1 in an inflammatory context may be highly  
990 disadvantageous, in a similar manner as reported in the muscles of mice with systemic inflammation  
991 on a 11 $\beta$ -HSD1 KO background.<sup>195</sup> Here, the reduction in local steroid activation within muscle  
992 greatly increases systemic inflammation and local muscle inflammatory cytokine production,  
993 increasing inflammatory catabolic and anti-anabolic muscle wasting. In a similar manner, systemic  
994 inhibition of 11 $\beta$ -HSD1 may exacerbate inflammatory bone loss. Instead, alternative approaches, for  
995 example, targeted inhibition of pro-inflammatory NF- $\kappa$ B (or any other tissue specific regulator of  
996 11 $\beta$ -HSD1 activity) or bone selective inhibition of 11 $\beta$ -HSD1 may be a more effective approach.

997

#### 998 **IV Therapeutic glucocorticoid excess and bone**

999 This section reviews data relating to the epidemiology and treatment of iatrogenic GIOP.  
1000 Importantly, as discussed throughout this article, the term GIOP in this context could be misleading.  
1001 This is because therapeutic glucocorticoids might indeed lead to bone loss through their direct  
1002 actions but are also likely to have complex interactions with the underlying disease being treated.<sup>5,6</sup>  
1003 Glucocorticoids in some situations might magnify the amount of damage being done to bone  
1004 through worsening of imbalances between bone formation and resorption but in conditions

1005 characterised by systemic inflammation, glucocorticoids, particularly when used at modest doses  
1006 might be 'bone sparing' through their anti-inflammatory actions.<sup>196,197</sup> In the later situation bone  
1007 disease might be present in the context of prolonged glucocorticoid use but the damage would not  
1008 be truly 'glucocorticoid-induced'. It is generally thought that most systemic inflammatory illnesses  
1009 cause bone loss primarily through increased bone resorption with a relative suppression or restraint  
1010 on bone formation.<sup>6,198</sup> The effectiveness of different treatment approaches (anti-resorptive agents  
1011 targeting osteoclast activity vs. anabolic drugs targeting bone formation) might depend on the  
1012 extent to which bone disease is secondary to inflammation or to the glucocorticoids needed to  
1013 control the underlying disease.

#### 1014 IV.I Epidemiology of glucocorticoid use and impact on bone

1015 The use of therapeutic glucocorticoids in the community is still high and may even be increasing.  
1016 Studies around the turn of the century from the UK reported that up to 1% of the population were  
1017 taking oral glucocorticoids on a long term basis.<sup>199,200</sup> This figure rose to almost 3% in the elderly.  
1018 Data from the US based on the NHANES database between 1999 and 2008 estimated that the  
1019 prevalence of long term use was 1.2%.<sup>201</sup> In the Global Longitudinal Study of Osteoporosis in Women  
1020 (GLOW) the rate of glucocorticoid usage at baseline study visit in this post-menopausal population  
1021 was 4.6%.<sup>202</sup> Studies based on UK databases indicate that the rate of long term glucocorticoid use is  
1022 gradually increasing.<sup>203</sup> A recent study based on the population of Denmark reported that 3% of the  
1023 Danish population filled at least one prescription for a systemically administered therapeutic  
1024 glucocorticoid. In the Danish elderly population this figure rose to around 8-10%.<sup>204</sup> As such a  
1025 significant proportion of the global population is exposed to therapeutic glucocorticoids.  
1026 Several studies have attempted to estimate the fracture risk associated with long term  
1027 glucocorticoid use. In some groups of patients treated long-term with oral glucocorticoids, the risk of  
1028 developing osteoporosis and vertebral fractures was estimated at 50% or more.<sup>4</sup> These rates will  
1029 depend on the specific disease being treated and the age and gender profile of the populations

1030 studied. Population based studies have similarly indicated that glucocorticoid usage is associated  
1031 with an increased risk of fracture.<sup>3</sup> Importantly, risks of fracture were increased at the hip (relative  
1032 risk increase 1.6) and spine (relative risk increase 2.6) as well as an increased risk of non-vertebral  
1033 fractures (relative risk 1.3). Even relatively modest doses of glucocorticoids were associated with a  
1034 significantly increased fracture risk, with doses as low as 2.5mg/day being linked to spine fractures.  
1035 Risk of fractures was also associated with daily dose with a 20% increased risk of fracture seen at 5  
1036 mg/day of prednisolone, increasing to 60% at 20 mg/day. The time of onset and offset of fracture  
1037 risk was particularly instructive. In the study by van Staa et al. the risk of fracture increased rapidly  
1038 within a short time of commencing glucocorticoid therapy.<sup>3</sup> The risk of fracture remained elevated  
1039 while glucocorticoids were continued but fell rapidly after glucocorticoids were ceased. The  
1040 mechanisms for these rapid changes in fracture risk are unclear but changes in bone density alone  
1041 are an unlikely explanation. An increased risk of falls due to myopathy associated with glucocorticoid  
1042 use might be part of the explanation. The risk of fracture also appeared to increase in glucocorticoid  
1043 users even before therapy was initiated, indicating that the indication for treatment is an important  
1044 component of the increase in risk of fracture during therapy. As such glucocorticoid treatment is  
1045 likely to be a marker of the presence of a disease associated with increased fracture risk as well as an  
1046 independent factor itself. Support for this involvement of underlying disease in fracture risk comes  
1047 from studies of patients taking inhaled glucocorticoids for respiratory disease, who had a higher  
1048 fracture risk than healthy matched controls.<sup>205</sup> However, patients taking inhaled bronchodilators but  
1049 no inhaled glucocorticoids had a similar increase in fracture risk compared to controls, a finding that  
1050 implies that the underlying respiratory condition was the most significant contributor to fracture  
1051 risk.

1052 Subsequent studies have attempted to determine whether fracture risk was associated with  
1053 cumulative or daily dose. In general both dose and duration influence fracture risk and these two  
1054 factors are difficult to separate out in clinical practice.<sup>206,207</sup> The question was recently addressed in a  
1055 population based cohort of over 50,000 patients from Canada which examined the relative

**Comment [M7]:** Table for epidemiology studies?

1056 importance of recent or remote and short or prolonged use of glucocorticoids on bone density and  
1057 fracture incidence.<sup>208</sup> In this cohort only recent prolonged glucocorticoid use was associated with  
1058 reduced femoral neck T-scores and a BMD-independent increase in the risk of major and hip  
1059 fractures. However, most other studies provide reassurance that glucocorticoid use is only harmful  
1060 to bone when used for relatively long durations. Occasional intermittent use of high dose  
1061 glucocorticoids has been reported to be relatively safe in terms of bone health.<sup>209</sup> However, a recent  
1062 retrospective cohort study based on private insurance claims from the US demonstrated an  
1063 association between short-term glucocorticoid use (less than 30 days) with various types of harm  
1064 including a 1.8 fold increased risk of fracture.<sup>210</sup> Such short-term use of glucocorticoids was common  
1065 at 20% of the population over the 3-year period examined, suggesting that intermittent  
1066 glucocorticoid use could contribute more to population fracture risk than previously thought.

1067 Another important source of information relating to the epidemiology of GIOP are the placebo arms  
1068 of the initial RCTs that examined the impact of various treatments on the development of GIOP.<sup>211-</sup>  
1069 <sup>213</sup> This approach has the advantage of having greater sensitivity for determining the impact of  
1070 glucocorticoids on vertebral fracture risk as spine radiographs were typically taken in these trials.

1071 Overall, the placebo arms of these studies indicate that glucocorticoid treatment is associated with a  
1072 high risk of fracture, particularly vertebral fracture in post-menopausal women and older men.<sup>101,214</sup>  
1073 However, the risk of fracture in pre-menopausal women and younger men appeared to be very low.  
1074 Where fractures did occur the BMD T-score tended to be below a level of -1.5. Whether these  
1075 younger patient groups have reduced absolute risks of fracture by virtue just of their age or whether  
1076 there are independent age related protective factors such as sex steroid levels remains unclear. A  
1077 recent formal meta-regression of data from the placebo arms of these studies has reported annual  
1078 incidence rates of vertebral fracture of 5.1% and 3.2% for patients initiating or continuing  
1079 glucocorticoid treatment respectively.<sup>214</sup> The corresponding rates of non-vertebral fractures were  
1080 2.5% and 3.0%.

1081 Other studies have attempted to define the effect of glucocorticoid treatment on bone density and  
1082 architecture over time. Zhu et al. performed a carefully controlled prospective study of women with  
1083 SLE on long term glucocorticoids followed up for over 2 years.<sup>215</sup> Areal BMD by DXA at multiple sites  
1084 and microstructural analysis by high resolution peripheral QCT (HR-pQCT) at the distal radius were  
1085 examined at baseline, 12 and 24 months. In premenopausal women the changes in aBMD by DXA  
1086 over the two years were very similar between cases and controls. There was however a significant  
1087 decrease in cortical area and thickness and an increase in cortical porosity in cases compared to  
1088 controls. In post-menopausal women, again there was no significant difference in changes in BMD  
1089 between cases and controls but by HR-pQCT there was a significant decrease in volumetric BMD at  
1090 the cortex and more substantial decreases in cortical thickness and increases in cortical porosity. The  
1091 increases in porosity seen were double in post-menopausal women compared to premenopausal  
1092 women and the decrease in cortical thickness 10 fold greater. As such, at least in SLE, cortisol bone  
1093 loss is significant during glucocorticoid treatment with the magnitude of changes being substantially  
1094 greater in post-menopausal women. This data indicate that prolonged use of glucocorticoids leads to  
1095 deterioration in bone, and in particular cortical architecture, even in the absence of any changes in  
1096 BMD measured by DXA. A similar DXA independent deterioration in bone quality may also occur at  
1097 the spine. Paggiosi et al. compared spine aBMD by DXA and TBS values in 484 women treated with  
1098 or without glucocorticoids.<sup>216</sup> There was no difference in aBMD between groups but the TBS Z-score  
1099 was 0.8 lower in the group treated with glucocorticoids. Whether the change in TBS is predictive of  
1100 fractures in this situation is not yet clear.

1101 An important consideration in these studies is whether the changes in trabecular or cortical bone  
1102 seen in people treated with glucocorticoids is due to the glucocorticoids or to the underlying disease  
1103 being treated. Some studies suggest that the underlying disease itself could be contributing more to  
1104 the adverse effects on bone than glucocorticoids. For example Olsson et al. examined the effect of  
1105 short term, high dose glucocorticoids on bone in patients with multiple sclerosis.<sup>217</sup> No independent  
1106 association was found between glucocorticoid usage and BMD. However, disease activity was



1107 strongly associated with decreases in spine and hip BMD. In a population based cohort study  
1108 examining 1 million patients with or without COPD, COPD severity was strongly related to an  
1109 increased risk of osteoporosis and fracture.<sup>218</sup> However, prednisolone use and inhaled corticosteroid  
1110 use were associated with a reduced rather than increased risk of osteoporosis. Although this type of  
1111 study could be influenced by confounding this is further evidence of the importance of the  
1112 underlying disease on fracture risk in glucocorticoid treated patients. Even though reductions in  
1113 BMD in patients treated with glucocorticoids may be more strongly related to the underlying disease  
1114 in some circumstances than the glucocorticoids themselves, from a practical point of view BMD  
1115 measurements are still clinically useful as these patients are likely to be at an increased risk of  
1116 fracture and likely to benefit from treatment.

1117

#### 1118 IV.II Risk stratification in the clinical setting

1119 As with guidelines relating to other forms of osteoporosis there has been a shift away from fixed cut  
1120 off values of BMD by DXA as the basis for estimation of risk to the use of calculators such as FRAX<sup>TM</sup>  
1121 <sup>219</sup>and the Garvan Fracture Risk calculator<sup>220</sup> that include multiple aspects of fracture risk. Currently  
1122 guidelines for GIOP risk stratification attempt to focus on estimated absolute risk but usually include  
1123 additional 'red flags' that would prompt treatment even if the estimated risk is below the  
1124 intervention threshold. A question relating to exposure to glucocorticoids is incorporated into  
1125 FRAX<sup>TM</sup>, the most widely used fracture risk calculator, but not other fracture risk calculators. The  
1126 glucocorticoid question requires a yes/no answer with the yes answer indicated if the patient is  
1127 currently exposed to oral glucocorticoids or has been exposed to oral glucocorticoids for more than  
1128 3 months at a dose of prednisolone of 5mg daily or more. As discussed earlier, the risks associated  
1129 with glucocorticoids in epidemiological surveys indicate that current treatment with glucocorticoids  
1130 is a more powerful risk factor for fracture than remote use and the relationship is also, to some  
1131 extent, dose dependent. To address this additional guidance is now available on how to adjust the

1132 FRAX output manually based on glucocorticoid dose and recency of exposure.<sup>219</sup> A more  
1133 fundamental limitation of FRAX and other calculators in the context of GIOP is the focus on hip and  
1134 non-vertebral fractures with little emphasis on spine fractures. This extends to the requirement to  
1135 enter femoral neck BMD with no possibility of entering spine BMD scores. As such in younger  
1136 patients with relatively preserved BMD it is unlikely that high fracture risk values will be generated  
1137 even if the risk of spine fractures was clinically significant. To address this, many guidelines also  
1138 suggest treatment to be considered if the patient has a low traumatic fracture, or has a low BMD e.g.  
1139 less than -1.5 T score by DXA, particularly at the spine. Although not evaluated in the context of  
1140 clinical studies it would seem reasonable clinical practice to have a lower threshold for obtaining  
1141 spine radiographs or VFA by DXA (if available) in patients at risk of GIOP.

1142

#### 1143 IV.III Treatment strategies

1144 There are a range of guidelines and recommendations published for the pharmacological treatment  
1145 of GIOP.<sup>221-223</sup> Intervention thresholds and rules for treatment use and patient reimbursement vary  
1146 considerably between countries. These guidelines will not be reviewed extensively here but rather  
1147 the evidence related to the effectiveness of various treatments specifically on fracture risk are  
1148 described below.

1149 A number of RCTs have been performed in the context of GIOP. These are generally much smaller in  
1150 size than the RCTs that demonstrated the effectiveness of these medications in post-menopausal  
1151 osteoporosis and generally were not powered for fracture. The main RCTs that compared  
1152 treatments to placebo include those that evaluated etidronate,<sup>212</sup> alendronate<sup>213</sup> and risedronate,<sup>211</sup>  
1153 whereas zoledronic acid<sup>224</sup> and teriparatide<sup>225</sup> evaluated medications in non-inferiority studies.  
1154 These trials consistently demonstrated improvements in BMD by DXA relative to placebo in patients  
1155 treated shortly after initiation of glucocorticoids and patients treated with glucocorticoids prior to

1156 initiation of therapy. Although not powered for fracture risk reduction, post-hoc analyses have  
1157 indicated the likely impact of these treatments on fracture risk, particularly at the spine.<sup>226,227</sup> Other  
1158 studies have examined the effect of vitamin D or its metabolites but these have generally been  
1159 smaller and found less substantial changes in BMD.<sup>228-230</sup> Due to heterogeneity of inclusion criteria,  
1160 baseline fracture risk and methods of ascertainment of BMD and fracture incidence it is difficult to  
1161 compare between treatments. However, the effectiveness of alendronate and teriparatide has  
1162 been compared in the context of non-inferiority. Saag et al. compared teriparatide with alendronate  
1163 treatment over 18 months in a randomised double blind study of 428 women and men receiving  
1164 therapeutic glucocorticoids.<sup>225</sup> Spine BMD by DXA increased substantially more in the teriparatide  
1165 group than in the alendronate group. Most importantly, in a pre-specified secondary analysis, the  
1166 risk of developing new morphometric vertebral fractures were considerably lower in patients taking  
1167 teriparatide (0.6%) than alendronate (6.1%). These results are dramatic especially taking into account  
1168 that alendronate substantially reduces the risk of new morphometric vertebral fractures when  
1169 compared to placebo.<sup>226</sup> The superiority of teriparatide compared to bisphosphonates for increasing  
1170 bone strength at the spine is also supported by an open label randomised trial of teriparatide versus  
1171 and risedronate in men treated with glucocorticoids.<sup>231</sup> This smaller RCT (92 men in total) focussed  
1172 primarily on spine BMD and structural parameters using DXA, QCT of L1-3, and high resolution QCT  
1173 or the T12 vertebra. Both treatments led to improvements in various parameters of bone density  
1174 and strength but these were in general much greater with teriparatide. Estimates of vertebral  
1175 strength using finite element analysis (FEA) demonstrated clear superiority of teriparatide over  
1176 risedronate. Even though small, there was also a trend towards fewer spine fractures in men treated  
1177 with teriparatide compared to those treated with risedronate. The effects of alendronate versus  
1178 teriparatide on spine TBS have also been compared in a secondary analysis of a RCT of these  
1179 treatments in GIOP.<sup>232</sup> It was found that alendronate had no impact on TBS despite increasing aBMD  
1180 whereas teriparatide caused a significant increase in TBS in addition to positive impact on aBMD.  
1181 These data collectively indicate that the bisphosphonates and teriparatide are effective in reducing

1182 fracture risk at the spine in GIOP but teriparatide appears superior over 18 months. An important  
1183 caveat regarding teriparatide is that it is generally only licenced for a duration of 18 months  
1184 (depending on country) and subsequent anti-resorptive medications are typically needed after this  
1185 treatment ends.

1186 The small size of these RCTs and relative rarity of non-vertebral and hip fractures compared to  
1187 morphometric spine fractures means that the effect of these drugs on non-vertebral and hip fracture  
1188 risk is not possible to determine. Three recent publications have used retrospective database  
1189 analysis to try to determine the real world impact of these medications on overall fracture risk and  
1190 hip fracture risk specifically. Overman et al. examined the effect of anti-osteoporotic medications  
1191 (AOMs, including bisphosphonates, teriparatide and denosumab) collectively on the risk of clinical  
1192 fractures in women aged 50 plus taking oral glucocorticoids included in the MarketScan databases.  
1193 The analysis included 7885 women with 12.1% of them treated with AOMs. It was found that AOM  
1194 use was associated with significantly reduced hazard ratios (HRs) for fracture of 0.52 at 1 year and  
1195 0.68 at 3 years.<sup>233</sup> Axelsson et al. examined the association between alendronate use and hip  
1196 fracture risk in women and men using glucocorticoids in a national Swedish database including  
1197 433,195 patients 65 years and older.<sup>234</sup> The use of alendronate was associated with a significantly  
1198 reduced risk of hip fracture (HR 0.35). A third study by Bergman et al. also used Swedish national  
1199 databases which included over 3 million people to compare the impact of alendronate on risk of  
1200 fracture in patients treated with glucocorticoids.<sup>235</sup> They found that alendronate use was associated  
1201 with a 16% reduction in non-vertebral fracture and a 34% reduction in hip fractures compared to  
1202 non-users. These studies clearly have intrinsic limitations due to their non-randomised nature and  
1203 could be influenced by confounding by indication. However, sensitivity analyses within these studies  
1204 consistently supported a strong and clinically important reduction in non-vertebral and hip fracture  
1205 risk in patients taking glucocorticoids who are treated with anti-osteoporosis medications.  
1206 Importantly, these studies also failed to detect any evidence of harm relating to gastrointestinal  
1207 adverse reactions with these medications. These reductions in non-vertebral fracture risk imply that

1208 treatments for GIOP might also improve the strength of cortical bone. A recent study directly  
1209 examined in vivo changes in bone tissue properties in glucocorticoid treated patients commencing  
1210 various osteoporosis treatments.<sup>236</sup> Reference point indentation (a form of microindentation) was  
1211 performed on the tibia using a hand held device under local anaesthetic. This technique measures  
1212 the resistance of cortical bone to indentation and thus provides a measure of tissue properties. Over  
1213 periods of 7 and 20 weeks treatment with calcium and vitamin had no impact on material properties  
1214 (the Bone Material Strength Index; BMSi). However, treatment with risedronate over 20 weeks and  
1215 treatment with denosumab or teriparatide over 7 and 20 weeks resulted in a significant increase in  
1216 BMSi. There was however no change in BMD by DXA in any group. Although the clinical utility of  
1217 microindentation techniques remains uncertain these preliminary studies strongly support a rapid  
1218 and beneficial effect of osteoporosis medications on cortical bone properties in patients at risk of  
1219 GIOP.

1220 A concern with anti-resorptive drugs is that they predispose to the development of atypical femoral  
1221 fractures (AFFs). Although the pathogenesis of AFFs is still unclear decreased bone turnover is  
1222 implicated. Since both anti-resorptive medications and glucocorticoids decrease bone turnover  
1223 prolonged use of both might theoretically increase the risk of AFFs. Although early reports<sup>237</sup>  
1224 suggested a possible link between glucocorticoids and AFFs in patients taking bisphosphonates more  
1225 recent studies do not support such an association.<sup>238</sup>

1226 A 2 year double blind placebo controlled non-inferiority RCT examining the effectiveness of  
1227 denosumab compared to risedronate in over 700 women and men with GIOP has been completed  
1228 but not published. Data from the 1<sup>st</sup> year of treatment has been reported in abstract form and the  
1229 drug appears to have a positive effect on bone mineral density with increases at spine and hip in  
1230 excess of those seen with risedronate. Small observational studies of the use of denosumab in GIOP  
1231 suggest that the drug is effective in maintaining or increasing BMD.<sup>239-241</sup> Formal licencing of  
1232 denosumab for this indication is expected in the future.

1233

1234 **V Skeletal impact of glucocorticoid replacement**

1235 This section will review the data relating to the skeletal impact of glucocorticoid replacement for  
1236 primary or secondary adrenal insufficiency. The amount of literature in this area is modest but there  
1237 have been some recent contributions and the subject is clearly of relevance to an endocrine  
1238 audience. When interpreting the literature relating to the impact of glucocorticoid replacement on  
1239 bone it is important to consider that patients are treated for long periods of time with some patients  
1240 treated for several decades, that treatment regimens have varied over time with a trend to using  
1241 lower doses of glucocorticoids in more recent years, and that glucocorticoid replacement may be  
1242 only one aspect of their underlying condition that might impact on bone. For example, patients  
1243 treated for Addison's disease will additionally have adrenal androgen deficiency and a greater  
1244 chance of previous thyrotoxicosis; patients with hypopituitarism will commonly have coexisting  
1245 growth hormone deficiency and patients with CAH will commonly have differences in height and  
1246 bone structure that make comparisons to controls difficult. Much of the literature focusses on BMD  
1247 by DXA but there is some fracture risk data.

1248 The epidemiology of hip fractures in patients with Addison's disease has been examined in a  
1249 population based analysis from Sweden.<sup>242</sup> Using hospital database, information relating to the  
1250 diagnosis of autoimmune adrenal insufficiency and hip fracture of 3,219 patients were identified and  
1251 compared to over 31,000 age and sex matched controls from the background population. The risk of  
1252 hip fracture was found to be substantially increased with a hazard ratio of 1.8 (CI 1.6-2.1). The  
1253 relative risk increase was independent of age or sex. The risk of any fracture was also increased to a  
1254 similar degree. The relationship between risk of hip fracture and the time of diagnosis of Addison's  
1255 disease was also explored. The relative risk of fracture was most increased in the first year after the  
1256 diagnosis but was elevated at all time points. Interestingly, the risk of fracture was also increased by  
1257 almost 3-fold in the year prior to the diagnosis, indicating that the cause of the increased fracture

1258 incidence, at least at this time point is not glucocorticoid replacement. It is possible that  
1259 glucocorticoid deficiency has a major negative effect on bone in keeping with some of the  
1260 observations of an anabolic effect of physiological levels of glucocorticoid action within bone.<sup>50,63,66</sup>  
1261 However, there are additionally many other reasons for this increased fracture risk not least an  
1262 increased risk of falls arising from the often severe myopathy seen in untreated adrenal failure.<sup>243</sup>  
1263 Nevertheless the data support the notion that fracture risk might be increased in patients that are  
1264 undertreated as well as those that are over-treated with replacement glucocorticoids.

1265 Other studies have focussed on bone density in Addison's disease. These were mostly cross-sectional  
1266 and are difficult to interpret since some of the patients included had been exposed to higher doses  
1267 of glucocorticoids than typically used now for prolonged periods of time. In general these studies  
1268 reported that adrenal insufficiency was associated with a reduction in BMD at the spine and the hip  
1269 and that this reduction was greater with more prolonged use. These studies are summarised in a  
1270 recent review by Lee and Greenfield.<sup>244</sup> One relatively recent cross-sectional study of patients with  
1271 Addison's disease or CAH treated with lower glucocorticoid replacement doses failed to find a  
1272 reduction in BMD as assessed by DXA suggesting that replacement regimens adopted more recently  
1273 have less negative impact on bone.<sup>245</sup> However, another study of 87 patients with Addison's disease  
1274 and 81 age and sex matched controls found a higher than expected prevalence of spine fractures  
1275 (using DXA based vertebral fracture assessment (VFA)) in patients with adrenal insufficiency despite  
1276 there being no difference in BMD.<sup>246</sup> Using the Genant criteria 31% of Addison's patients had at least  
1277 one vertebral abnormality compared to 12.8% of controls. Suggesting that these fractures might be  
1278 related to treatment the risk of fracture appeared greater in those with a longer duration of disease.  
1279 Interestingly mineralocorticoid replacement was associated with the presence of a higher BMD.

1280 A recent prospective study examined the impact of targeted reduction in glucocorticoid replacement  
1281 dose in patients with Addison's disease and CAH.<sup>247</sup> In patients where a reduction in glucocorticoid  
1282 replacement appeared justified there was a significant increase in spine and hip Z scores over a 2

1283 year follow up period. This suggests that bone health can be improved by careful attention to  
1284 glucocorticoid replacement doses but the actual impact of these changes on risk of fractures has not  
1285 been examined.

1286 Several studies have examined the bone health of individuals with CAH and the likely impact of  
1287 glucocorticoid replacement.<sup>248-252</sup> These studies are complicated by the differences in height and  
1288 bone size, which tend to overstate reductions in BMD. Additionally, excessive androgen exposure  
1289 resulting from inadequate glucocorticoid dosing might have an anabolic effect on bone. However  
1290 despite this it appears that bone density is reduced at all skeletal sites in CAH using both DXA and  
1291 spinal QCT. The reduction has been correlated with cumulative glucocorticoid exposure in some  
1292 studies<sup>249</sup> but not others.<sup>248</sup> Two small studies by the same research group found that adult women  
1293 with CAH had an increased risk of fracture<sup>251</sup> but men with CAH did not.<sup>252</sup>

1294



1295 **VI Future directions**

1296 Currently it is unclear how to separate the anabolic versus catabolic actions of glucocorticoids on  
1297 bone. Endogenous glucocorticoids have powerful effects on bone health which appear to some  
1298 extent to be regional and surface dependent. In scientific studies there is no consistency in the  
1299 regions and surfaces examined which makes comparisons between studies difficult.

1300 There are also limitations with current animal models as it is unclear to what extent they adequately  
1301 model the human situation. Few examine the sensitivity of bone in the context of an underlying  
1302 illness which is being treated with glucocorticoids. Animal studies examining interventions to protect  
1303 against GIOP need to be assessed in inflammatory disease models to determine whether these  
1304 interventions have independent actions on the underlying illness being treated in the first place.

1305 It remains unclear if there is a single target for glucocorticoid action within bone that can be  
1306 targeted therapeutically. Although there have been attractive targets such as the IGF1 system,  
1307 osteoblast apoptosis, IL-11, autophagy, OPG/RANKL and wnts/DKK1/sclerostin no single system  
1308 appears to account for all of the actions of glucocorticoids on bone. As discussed above, even if a  
1309 single target was identified it would need to be clear that this target was not involved in the  
1310 beneficial effects of glucocorticoids on the immune system. Currently the most likely candidates in  
1311 this respect are approaches that target sclerostin and/or DKK1 (the other major antagonist of wnt  
1312 signalling). The feasibility of such an approach has been demonstrated in principle outside the  
1313 context of glucocorticoid therapy in rodents and non-human primates.<sup>253</sup>

1314 It should also be remembered that glucocorticoid excess is associated with many adverse effects  
1315 outside of bone and these are currently not addressed well. An ideal treatment approach would  
1316 target multiple components of risk. Being able to do this without blocking the anti-inflammatory  
1317 actions of glucocorticoids has proven difficult. One approach to achieve this goal that has been an  
1318 active topic of research for several decades is the development of selective glucocorticoid receptor

1319 modulators. There are various theoretical underpinnings of these molecules largely based on the  
1320 concept of dissociating effects of glucocorticoids which are mediated by transactivation and  
1321 transrepression.<sup>254,255</sup> These concepts now appear overly simplistic and furthermore the actions of  
1322 glucocorticoids on bone appear to be through transrepression rather than transactivation.<sup>50</sup> There  
1323 have been some interesting compounds examined, in particular 'compound A', which exhibits some  
1324 useful features although it's relatively narrow therapeutic range is likely to limit human use.<sup>256</sup> Also,  
1325 these agents are complicated by the likelihood that they could interfere with HPA axis regulation  
1326 leading to low levels of endogenous cortisol. This could create a mixture of excessive glucocorticoid  
1327 action in some tissues but an absence of glucocorticoid action in others. Moreover, these  
1328 compounds would not exhibit the same properties of selective activation by tissue specific enzymes  
1329 and would be unlikely to have effects on the MR (which could play a role in some inflammatory  
1330 situations or even modulate the impact of more conventional glucocorticoids on bone). Given these  
1331 complexities selective glucocorticoid receptor modulators are unlikely to be developed but if they  
1332 are they would probably need to be evaluated in specific inflammatory conditions rather than for  
1333 inflammatory disease in general.

1334 There also remains uncertainty regarding what is actually being treated in patients with GIOP. In  
1335 some contexts glucocorticoids are likely to be detrimental but in other situations altered bone  
1336 remodelling due to the underlying inflammatory disease might be more important. Glucocorticoids  
1337 might therefore have an important role in controlling inflammation related bone loss and thus be  
1338 bone 'sparing' rather than negative to bone. If this distinction can be made it might be possible to  
1339 independently target glucocorticoid induced and inflammation induced bone disease separately or  
1340 synergistically.

1341

1342

1343 References:

- 1344 1. Cushing HW. The basophil adenomas of the pituitary body and their clinical manifestations  
1345 (pituitary basophilism). Bull Johns Hopkins Hosp 1932;50:137-95.
- 1346 2. Strehl C, Bijlsma JW, de Wit M, et al. Defining conditions where long-term glucocorticoid  
1347 treatment has an acceptably low level of harm to facilitate implementation of existing  
1348 recommendations: viewpoints from an EULAR task force. Ann Rheum Dis 2016;75:952-7.
- 1349 3. van Staa TP, Leufkens HG, Abenham L, Zhang B, Cooper C. Use of oral corticosteroids and  
1350 risk of fractures. J Bone Miner Res 2000;15:993-1000.
- 1351 4. Walsh LJ, Lewis SA, Wong CA, et al. The impact of oral corticosteroid use on bone mineral  
1352 density and vertebral fracture. American journal of respiratory and critical care medicine  
1353 2002;166:691-5.
- 1354 5. Hardy R, Cooper MS. Bone loss in inflammatory disorders. J Endocrinol 2009;201:309-20.
- 1355 6. Briot K, Geusens P, Em Bultink I, Lems WF, Roux C. Inflammatory diseases and bone fragility.  
1356 Osteoporosis international : a journal established as result of cooperation between the European  
1357 Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA 2017;28:3301-  
1358 14.
- 1359 7. Chiodini I, Vainicher CE, Morelli V, et al. MECHANISMS IN ENDOCRINOLOGY: Endogenous  
1360 subclinical hypercortisolism and bone: a clinical review. European journal of endocrinology /  
1361 European Federation of Endocrine Societies 2016;175:R265-R82.
- 1362 8. Manenschijn L, van den Akker EL, Lamberts SW, van Rossum EF. Clinical features associated  
1363 with glucocorticoid receptor polymorphisms. An overview. Ann NY Acad Sci 2009;1179:179-98.
- 1364 9. Nixon M, Mackenzie SD, Taylor AI, et al. ABCC1 confers tissue-specific sensitivity to cortisol  
1365 versus corticosterone: A rationale for safer glucocorticoid replacement therapy. Science translational  
1366 medicine 2016;8:352ra109.

- 1367 10. Gathercole LL, Lavery GG, Morgan SA, et al. 11beta-hydroxysteroid dehydrogenase 1:  
1368 translational and therapeutic aspects. *EndocrRev* 2013;34:525-55.
- 1369 11. Raubenheimer PJ, Young EA, Andrew R, Seckl JR. The role of corticosterone in human  
1370 hypothalamic-pituitary-adrenal axis feedback. *Clinical endocrinology* 2006;65:22-6.
- 1371 12. Keeney DS, Jenkins CM, Waterman MR. Developmentally regulated expression of adrenal  
1372 17 $\alpha$ -hydroxylase cytochrome P450 in the mouse embryo. *Endocrinology* 1995;136:4872-9.
- 1373 13. Gomez-Sanchez E, Gomez-Sanchez CE. The multifaceted mineralocorticoid receptor.  
1374 *Comprehensive Physiology* 2014;4:965-94.
- 1375 14. Vandewalle J, Luybaert A, De Bosscher K, Libert C. Therapeutic Mechanisms of  
1376 Glucocorticoids. *Trends in endocrinology and metabolism: TEM* 2018;29:42-54.
- 1377 15. Cohen DM, Steger DJ. Nuclear Receptor Function through Genomics: Lessons from the  
1378 Glucocorticoid Receptor. *Trends in endocrinology and metabolism: TEM* 2017;28:531-40.
- 1379 16. Wood CL, Soucek O, Wong SC, et al. Animal models to explore the effects of glucocorticoids  
1380 on skeletal growth and structure. *The Journal of endocrinology* 2018;236:R69-R91.
- 1381 17. Gasparini SJ, Weber MC, Henneicke H, Kim S, Zhou H, Seibel MJ. Continuous corticosterone  
1382 delivery via the drinking water or pellet implantation: A comparative study in mice. *Steroids*  
1383 2016;116:76-82.
- 1384 18. Grahne L, Jochems C, Andersson A, et al. Possible role of lymphocytes in glucocorticoid-  
1385 induced increase in trabecular bone mineral density. *The Journal of endocrinology* 2015;224:97-108.
- 1386 19. Jaiswal N, Haynesworth SE, Caplan AI, Bruder SP. Osteogenic differentiation of purified,  
1387 culture-expanded human mesenchymal stem cells in vitro. *Journal of cellular biochemistry*  
1388 1997;64:295-312.
- 1389 20. Pittenger MF, Mackay AM, Beck SC, et al. Multilineage potential of adult human  
1390 mesenchymal stem cells. *Science* 1999;284:143-7.
- 1391 21. Bellows CG, Aubin JE, Heersche JN. Physiological concentrations of glucocorticoids stimulate  
1392 formation of bone nodules from isolated rat calvaria cells in vitro. *Endocrinology* 1987;121:1985-92.

- 1393 22. Cheng SL, Yang JW, Rifas L, Zhang SF, Avioli LV. Differentiation of human bone marrow  
1394 osteogenic stromal cells in vitro: induction of the osteoblast phenotype by dexamethasone.  
1395 *Endocrinology* 1994;134:277-86.
- 1396 23. Leboy PS, Beresford JN, Devlin C, Owen ME. Dexamethasone induction of osteoblast mRNAs  
1397 in rat marrow stromal cell cultures. *J Cell Physiol* 1991;146:370-8.
- 1398 24. Buttery LD, Bourne S, Xynos JD, et al. Differentiation of osteoblasts and in vitro bone  
1399 formation from murine embryonic stem cells. *Tissue Eng* 2001;7:89-99.
- 1400 25. Shi C, Huang P, Kang H, et al. Glucocorticoid inhibits cell proliferation in differentiating  
1401 osteoblasts by microRNA-199a targeting of WNT signaling. *J Mol Endocrinol* 2015;54:325-37.
- 1402 26. Mak W, Shao X, Dunstan CR, Seibel MJ, Zhou H. Biphasic glucocorticoid-dependent  
1403 regulation of Wnt expression and its inhibitors in mature osteoblastic cells. *Calcified tissue*  
1404 *international* 2009;85:538-45.
- 1405 27. Zhou H, Mak W, Zheng Y, Dunstan CR, Seibel MJ. Osteoblasts directly control lineage  
1406 commitment of mesenchymal progenitor cells through Wnt signaling. *Journal of Biological Chemistry*  
1407 2008;283:1936-45.
- 1408 28. Frenkel B, White W, Tuckermann J. Glucocorticoid-Induced Osteoporosis. *Advances in*  
1409 *experimental medicine and biology* 2015;872:179-215.
- 1410 29. Manolagas SC. Birth and death of bone cells: basic regulatory mechanisms and implications  
1411 for the pathogenesis and treatment of osteoporosis. *EndocrRev* 2000;21:115-37.
- 1412 30. Plotkin LI, Manolagas SC, Bellido T. Glucocorticoids induce osteocyte apoptosis by blocking  
1413 focal adhesion kinase-mediated survival. Evidence for inside-out signaling leading to anoikis. *J Biol*  
1414 *Chem* 2007;282:24120-30.
- 1415 31. Plotkin LI, Weinstein RS, Parfitt AM, Roberson PK, Manolagas SC, Bellido T. Prevention of  
1416 osteocyte and osteoblast apoptosis by bisphosphonates and calcitonin. *J Clin Invest* 1999;104:1363-  
1417 74.

- 1418 32. Almeida M, Han L, Ambrogini E, Weinstein RS, Manolagas SC. Glucocorticoids and tumor  
1419 necrosis factor alpha increase oxidative stress and suppress Wnt protein signaling in osteoblasts. The  
1420 Journal of biological chemistry 2011;286:44326-35.
- 1421 33. Espina B, Liang M, Russell RG, Hulley PA. Regulation of bim in glucocorticoid-mediated  
1422 osteoblast apoptosis. Journal of cellular physiology 2008;215:488-96.
- 1423 34. Chen F, Zhang L, OuYang Y, Guan H, Liu Q, Ni B. Glucocorticoid induced osteoblast apoptosis  
1424 by increasing E4BP4 expression via up-regulation of Bim. Calcified tissue international 2014;94:640-  
1425 7.
- 1426 35. Chang JK, Li CJ, Liao HJ, Wang CK, Wang GJ, Ho ML. Anti-inflammatory drugs suppress  
1427 proliferation and induce apoptosis through altering expressions of cell cycle regulators and pro-  
1428 apoptotic factors in cultured human osteoblasts. Toxicology 2009;258:148-56.
- 1429 36. Li H, Qian W, Weng X, et al. Glucocorticoid receptor and sequential P53 activation by  
1430 dexamethasone mediates apoptosis and cell cycle arrest of osteoblastic MC3T3-E1 cells. PloS one  
1431 2012;7:e37030.
- 1432 37. Brandstrom H, Bjorkman T, Ljunggren O. Regulation of osteoprotegerin secretion from  
1433 primary cultures of human bone marrow stromal cells. Biochemical and biophysical research  
1434 communications 2001;280:831-5.
- 1435 38. Hofbauer LC, Gori F, Riggs BL, et al. Stimulation of osteoprotegerin ligand and inhibition of  
1436 osteoprotegerin production by glucocorticoids in human osteoblastic lineage cells: potential  
1437 paracrine mechanisms of glucocorticoid-induced osteoporosis. Endocrinology 1999;140:4382-9.
- 1438 39. Vidal NO, Brandstrom H, Jonsson KB, Ohlsson C. Osteoprotegerin mRNA is expressed in  
1439 primary human osteoblast-like cells: down-regulation by glucocorticoids. J Endocrinol 1998;159:191-  
1440 5.
- 1441 40. Nakashima T, Hayashi M, Fukunaga T, et al. Evidence for osteocyte regulation of bone  
1442 homeostasis through RANKL expression. Nat Med 2011;17:1231-4.

- 1443 41. Xiong J, Piemontese M, Onal M, et al. Osteocytes, not Osteoblasts or Lining Cells, are the  
1444 Main Source of the RANKL Required for Osteoclast Formation in Remodeling Bone. PLoS one  
1445 2015;10:e0138189.
- 1446 42. Baron R, Kneissel M. WNT signaling in bone homeostasis and disease: from human  
1447 mutations to treatments. Nat Med 2013;19:179-92.
- 1448 43. Ohnaka K, Tanabe M, Kawate H, Nawata H, Takayanagi R. Glucocorticoid suppresses the  
1449 canonical Wnt signal in cultured human osteoblasts. Biochemical and biophysical research  
1450 communications 2005;329:177-81.
- 1451 44. Morimoto E, Li M, Khalid AB, Krum SA, Chimge NO, Frenkel B. Glucocorticoids Hijack Runx2  
1452 to Stimulate Wif1 for Suppression of Osteoblast Growth and Differentiation. Journal of cellular  
1453 physiology 2017;232:145-53.
- 1454 45. Ohnaka K, Taniguchi H, Kawate H, Nawata H, Takayanagi R. Glucocorticoid enhances the  
1455 expression of dickkopf-1 in human osteoblasts: novel mechanism of glucocorticoid-induced  
1456 osteoporosis. Biochemical and biophysical research communications 2004;318:259-64.
- 1457 46. Diarra D, Stolina M, Polzer K, et al. Dickkopf-1 is a master regulator of joint remodeling.  
1458 Nature Medicine 2007;13:156-63.
- 1459 47. Centrella M, McCarthy TL, Canalis E. Glucocorticoid regulation of transforming growth factor  
1460 beta 1 activity and binding in osteoblast-enriched cultures from fetal rat bone. Mol Cell Biol  
1461 1991;11:4490-6.
- 1462 48. Bennett A, Chen T, Feldman D, Hintz RL, Rosenfeld RG. Characterization of insulin-like  
1463 growth factor I receptors on cultured rat bone cells: regulation of receptor concentration by  
1464 glucocorticoids. Endocrinology 1984;115:1577-83.
- 1465 49. McCarthy TL, Centrella M, Canalis E. Cortisol inhibits the synthesis of insulin-like growth  
1466 factor-I in skeletal cells. Endocrinology 1990;126:1569-75.
- 1467 50. Rauch A, Seitz S, Baschant U, et al. Glucocorticoids suppress bone formation by attenuating  
1468 osteoblast differentiation via the monomeric glucocorticoid receptor. Cell Metab 2010;11:517-31.

- 1469 51. Sato AY, Richardson D, Cregor M, et al. Glucocorticoids Induce Bone and Muscle Atrophy by  
1470 Tissue-Specific Mechanisms Upstream of E3 Ubiquitin Ligases. *Endocrinology* 2017;158:664-77.
- 1471 52. Ersek A, Santo AI, Vattakuzhi Y, George S, Clark AR, Horwood NJ. Strain dependent  
1472 differences in glucocorticoid-induced bone loss between C57BL/6J and CD-1 mice. *Sci Rep*  
1473 2016;6:36513.
- 1474 53. Weinstein RS, Jilka RL, Parfitt AM, Manolagas SC. Inhibition of osteoblastogenesis and  
1475 promotion of apoptosis of osteoblasts and osteocytes by glucocorticoids. Potential mechanisms of  
1476 their deleterious effects on bone. *JClinInvest* 1998;102:274-82.
- 1477 54. Scheller EL, Leininger GM, Hankenson KD, Myers MG, Jr., Krebsbach PH. Ectopic expression  
1478 of Col2.3 and Col3.6 promoters in the brain and association with leptin signaling. *Cells, tissues,*  
1479 *organs* 2011;194:268-73.
- 1480 55. Terasawa M, Shimokawa R, Terashima T, Ohya K, Takagi Y, Shimokawa H. Expression of  
1481 dentin matrix protein 1 (DMP1) in nonmineralized tissues. *Journal of bone and mineral metabolism*  
1482 2004;22:430-8.
- 1483 56. Camerino C, Conte E, Cannone M, Caloiero R, Fonzino A, Tricarico D. Nerve Growth Factor,  
1484 Brain-Derived Neurotrophic Factor and Osteocalcin Gene Relationship in Energy Regulation, Bone  
1485 Homeostasis and Reproductive Organs Analyzed by mRNA Quantitative Evaluation and Linear  
1486 Correlation Analysis. *Frontiers in physiology* 2016;7:456.
- 1487 57. Wyrwoll CS, Holmes MC, Seckl JR. 11beta-hydroxysteroid dehydrogenases and the brain:  
1488 from zero to hero, a decade of progress. *Frontiers in neuroendocrinology* 2011;32:265-86.
- 1489 58. Condon J, Gosden C, Gardener D, et al. Expression of type 2 11 $\beta$ -hydroxysteroid  
1490 dehydrogenase and corticosteroid hormone receptors in early human fetal life.  
1491 *JClinEndocrinolMetab* 1998;83:4490-7.
- 1492 59. Woitge H, Harrison J, Ivkovic A, Krozowski Z, Kream B. Cloning and in vitro characterization of  
1493 alpha 1(I)-collagen 11 beta-hydroxysteroid dehydrogenase type 2 transgenes as models for  
1494 osteoblast-selective inactivation of natural glucocorticoids. *Endocrinology* 2001;142:1341-8.



- 1495 60. Rabbitt E, Lavery GG, Walker EA, Cooper MS, Stewart PM, Hewison M. Pre-receptor  
1496 regulation of glucocorticoid action by 11 $\beta$ -hydroxysteroid dehydrogenase: a novel determinant of  
1497 cell proliferation. *FASEB J* 2002;16:36-44.
- 1498 61. O'Brien CA, Jia D, Plotkin LI, et al. Glucocorticoids Act Directly on Osteoblasts and Osteocytes  
1499 to Induce Their Apoptosis and Reduce Bone Formation and Strength. *Endocrinology* 2004;145:1835-  
1500 41.
- 1501 62. Dunstan CR, Zhou H, Brennan K, Zheng Y, Seibel MJ. Osteoblast Targeted Overexpression of  
1502 Hydroxysteroid Dehydrogenase Type 2 Induces Delayed Calvarial Development in Transgenic Mice.  
1503 *ASBMR Meeting Nashville 2005*:SU520.
- 1504 63. Sher LB, Woitge HW, Adams DJ, et al. Transgenic expression of 11 $\beta$ -hydroxysteroid  
1505 dehydrogenase type 2 in osteoblasts reveals an anabolic role for endogenous glucocorticoids in  
1506 bone. *Endocrinology* 2004;145:922-9.
- 1507 64. Kalajzic I, Kalajzic Z, Kaliterna M, et al. Use of type I collagen green fluorescent protein  
1508 transgenes to identify subpopulations of cells at different stages of the osteoblast lineage. *JBone*  
1509 *MinerRes* 2002;17:15-25.
- 1510 65. Zhou H, Mak W, Kalak R, et al. Glucocorticoid-dependent Wnt signaling by mature  
1511 osteoblasts is a key regulator of cranial skeletal development in mice. *Development* 2009;136:427-  
1512 36.
- 1513 66. Kalak R, Zhou H, Street J, et al. Endogenous glucocorticoid signalling in osteoblasts is  
1514 necessary to maintain normal bone structure in mice. *Bone* 2009;45:61-7.
- 1515 67. Henneicke H, Herrmann M, Kalak R, et al. Corticosterone selectively targets endo-cortical  
1516 surfaces by an osteoblast-dependent mechanism. *Bone* 2011;49:733-42.
- 1517 68. Beavan S, Horner A, Bord S, Ireland D, Compston J. Colocalization of glucocorticoid and  
1518 mineralocorticoid receptors in human bone. *JBone MinerRes* 2001;16:1496-504.

1519 69. Fumoto T, Ishii KA, Ito M, Berger S, Schutz G, Ikeda K. Mineralocorticoid receptor function in  
1520 bone metabolism and its role in glucocorticoid-induced osteopenia. *Biochem Biophys Res Commun*  
1521 2014;447:407-12.

1522 70. Eberhardt AW, Yeager-Jones A, Blair HC. Regional trabecular bone matrix degeneration and  
1523 osteocyte death in femora of glucocorticoid- treated rabbits. *Endocrinology* 2001;142:1333-40.

1524 71. Weinstein RS, Nicholas RW, Manolagas SC. Apoptosis of osteocytes in glucocorticoid-induced  
1525 osteonecrosis of the hip. *JClinEndocrinolMetab* 2000;85:2907-12.

1526 72. Jia J, Yao W, Guan M, et al. Glucocorticoid dose determines osteocyte cell fate. *FASEB*  
1527 *journal* : official publication of the Federation of American Societies for Experimental Biology  
1528 2011;25:3366-76.

1529 73. Plotkin LI, Weinstein RS, Parfitt AM, Roberson PK, Manolagas SC, Bellido T. Prevention of  
1530 osteocyte and osteoblast apoptosis by bisphosphonates and calcitonin. *JClinInvest* 1999;104:1363-  
1531 74.

1532 74. Jilka RL, Weinstein RS, Bellido T, Roberson P, Parfitt AM, Manolagas SC. Increased bone  
1533 formation by prevention of osteoblast apoptosis with parathyroid hormone. *JClinInvest*  
1534 1999;104:439-46.

1535 75. Weinstein RS, O'Brien CA, Almeida M, et al. Osteoprotegerin prevents glucocorticoid-  
1536 induced osteocyte apoptosis in mice. *Endocrinology* 2011;152:3323-31.

1537 76. Weinstein RS, Jilka RL, Almeida M, Roberson PK, Manolagas SC. Intermittent parathyroid  
1538 hormone administration counteracts the adverse effects of glucocorticoids on osteoblast and  
1539 osteocyte viability, bone formation, and strength in mice. *Endocrinology* 2010;151:2641-9.

1540 77. Weinstein RS, Wan C, Liu Q, et al. Endogenous glucocorticoids decrease skeletal  
1541 angiogenesis, vascularity, hydration, and strength in aged mice. *Aging Cell* 2010;9:147-61.

1542 78. Lane NE, Yao W, Balooch M, et al. Glucocorticoid-treated mice have localized changes in  
1543 trabecular bone material properties and osteocyte lacunar size that are not observed in placebo-  
1544 treated or estrogen-deficient mice. *JBone MinerRes* 2006;21:466-76.

- 1545 79. Xia X, Kar R, Gluhak-Heinrich J, et al. Glucocorticoid-induced autophagy in osteocytes. *J Bone*  
1546 *Miner Res* 2010;25:2479-88.
- 1547 80. Piemontese M, Onal M, Xiong J, et al. Suppression of autophagy in osteocytes does not  
1548 modify the adverse effects of glucocorticoids on cortical bone. *Bone* 2015;75:18-26.
- 1549 81. Dai W, Jiang L, Lay YA, et al. Prevention of glucocorticoid induced bone changes with beta-  
1550 ecdysone. *Bone* 2015;74:48-57.
- 1551 82. Lin NY, Chen CW, Kagwiria R, et al. Inactivation of autophagy ameliorates glucocorticoid-  
1552 induced and ovariectomy-induced bone loss. *Ann Rheum Dis* 2016;75:1203-10.
- 1553 83. Shi J, Wang L, Zhang H, et al. Glucocorticoids: Dose-related effects on osteoclast formation  
1554 and function via reactive oxygen species and autophagy. *Bone* 2015;79:222-32.
- 1555 84. Dallas SL, Prideaux M, Bonewald LF. The osteocyte: an endocrine cell ... and more. *Endocrine*  
1556 *reviews* 2013;34:658-90.
- 1557 85. Sato AY, Cregor M, Delgado-Calle J, et al. Protection From Glucocorticoid-Induced  
1558 Osteoporosis by Anti-Catabolic Signaling in the Absence of Sost/Sclerostin. *Journal of bone and*  
1559 *mineral research : the official journal of the American Society for Bone and Mineral Research*  
1560 2016;31:1791-802.
- 1561 86. Piemontese M, Xiong J, Fujiwara Y, Thostenson JD, O'Brien CA. Cortical bone loss caused by  
1562 glucocorticoid excess requires RANKL production by osteocytes and is associated with reduced OPG  
1563 expression in mice. *American journal of physiology Endocrinology and metabolism* 2016;311:E587-  
1564 93.
- 1565 87. Yao W, Dai W, Jiang L, et al. Sclerostin-antibody treatment of glucocorticoid-induced  
1566 osteoporosis maintained bone mass and strength. *Osteoporosis international : a journal established*  
1567 *as result of cooperation between the European Foundation for Osteoporosis and the National*  
1568 *Osteoporosis Foundation of the USA* 2016;27:283-94.
- 1569 88. Achiou Z, Toumi H, Touvier J, et al. Sclerostin antibody and interval treadmill training effects  
1570 in a rodent model of glucocorticoid-induced osteopenia. *Bone* 2015;81:691-701.

1571 89. Sivagurunathan S, Muir MM, Brennan TC, Seale JP, Mason RS. Influence of glucocorticoids on  
1572 human osteoclast generation and activity. *J Bone Miner Res* 2005;20:390-8.

1573 90. Conaway HH, Henning P, Lie A, Tuckermann J, Lerner UH. Activation of dimeric  
1574 glucocorticoid receptors in osteoclast progenitors potentiates RANKL induced mature osteoclast  
1575 bone resorbing activity. *Bone* 2016;93:43-54.

1576 91. Hirayama T, Sabokbar A, Athanasou NA. Effect of corticosteroids on human osteoclast  
1577 formation and activity. *JEndocrinol* 2002;175:155-63.

1578 92. Kim HJ, Zhao H, Kitaura H, et al. Glucocorticoids suppress bone formation via the osteoclast.  
1579 *JClinInvest* 2006;116:2152-60.

1580 93. Jia D, O'Brien CA, Stewart SA, Manolagas SC, Weinstein RS. Glucocorticoids act directly on  
1581 osteoclasts to increase their life span and reduce bone density. *Endocrinology* 2006;147:5592-9.

1582 94. Dempster DW, Moonga BS, Stein LS, Horbert WR, Antakly T. Glucocorticoids inhibit bone  
1583 resorption by isolated rat osteoclasts by enhancing apoptosis. *J Endocrinol* 1997;154:397-406.

1584 95. Swanson C, Lorentzon M, Conaway HH, Lerner UH. Glucocorticoid regulation of osteoclast  
1585 differentiation and expression of receptor activator of nuclear factor-kappaB (NF-kappaB) ligand,  
1586 osteoprotegerin, and receptor activator of NF-kappaB in mouse calvarial bones. *Endocrinology*  
1587 2006;147:3613-22.

1588 96. Weinstein RS, Chen JR, Powers CC, et al. Promotion of osteoclast survival and antagonism of  
1589 bisphosphonate-induced osteoclast apoptosis by glucocorticoids. *J Clin Invest* 2002;109:1041-8.

1590 97. Zhang Y, Wei L, Miron RJ, Shi B, Bian Z. Anabolic bone formation via a site-specific bone-  
1591 targeting delivery system by interfering with semaphorin 4D expression. *Journal of bone and mineral*  
1592 *research : the official journal of the American Society for Bone and Mineral Research* 2015;30:286-  
1593 96.

1594 98. Deb Roy A, Yin T, Choudhary S, Rodionov V, Pilbeam CC, Wu YI. Optogenetic activation of  
1595 Plexin-B1 reveals contact repulsion between osteoclasts and osteoblasts. *Nature communications*  
1596 2017;8:15831.

1597 99. Chrousos GP, Torpy DJ, Gold PW. Interactions between the hypothalamic-pituitary-adrenal  
1598 axis and the female reproductive system: clinical implications. *Annals of internal medicine*  
1599 1998;129:229-40.

1600 100. Straub RH, Cutolo M, Pacifici R. Evolutionary medicine and bone loss in chronic inflammatory  
1601 diseases--A theory of inflammation-related osteopenia. *Seminars in arthritis and rheumatism*  
1602 2015;45:220-8.

1603 101. Sambrook PN. Corticosteroid osteoporosis: practical implications of recent trials. *JBone*  
1604 *MinerRes* 2000;15:1645-9.

1605 102. Kung AW, Chan TM, Lau CS, Wong RW, Yeung SS. Osteopenia in young hypogonadal women  
1606 with systemic lupus erythematosus receiving chronic steroid therapy: a randomized controlled trial  
1607 comparing calcitriol and hormonal replacement therapy. *Rheumatology* 1999;38:1239-44.

1608 103. Hall GM, Daniels M, Doyle DV, Spector TD. Effect of hormone replacement therapy on bone  
1609 mass in rheumatoid arthritis patients treated with and without steroids. *Arthritis Rheum*  
1610 1994;37:1499-505.

1611 104. Crawford BA, Liu PY, Kean MT, Bleasel JF, Handelsman DJ. Randomized placebo-controlled  
1612 trial of androgen effects on muscle and bone in men requiring long-term systemic glucocorticoid  
1613 treatment. *The Journal of clinical endocrinology and metabolism* 2003;88:3167-76.

1614 105. Lane NE, Sanchez S, Modin GW, Genant HK, ini E, Arnaud CD. Parathyroid hormone  
1615 treatment can reverse corticosteroid-induced osteoporosis. Results of a randomized controlled  
1616 clinical trial. *JClinInvest* 1998;102:1627-33.

1617 106. Ritz E, Kreusser W, Rambauek M. Effects of glucocorticoids on calcium and phosphate  
1618 excretion. *Advances in experimental medicine and biology* 1984;171:381-97.

1619 107. Fucik RF, Kukreja SC, Hargis GK, Bowser EN, Henderson WJ, Williams GA. Effect of  
1620 glucocorticoids on function of the parathyroid glands in man. *The Journal of clinical endocrinology*  
1621 *and metabolism* 1975;40:152-5.

1622 108. Rubin MR, Bilezikian JP. Clinical review 151: The role of parathyroid hormone in the  
1623 pathogenesis of glucocorticoid-induced osteoporosis: a re-examination of the evidence.  
1624 J Clin Endocrinol Metab 2002;87:4033-41.

1625 109. Giustina A, Mazziotti G, Canalis E. Growth hormone, insulin-like growth factors, and the  
1626 skeleton. Endocrine reviews 2008;29:535-59.

1627 110. Mauras N, George D, Evans J, et al. Growth hormone has anabolic effects in  
1628 glucocorticosteroid-dependent children with inflammatory bowel disease: a pilot study. Metabolism  
1629 2002;51:127-35.

1630 111. Simon D, Prieur A, Czernichow P. Treatment of juvenile rheumatoid arthritis with growth  
1631 hormone. Hormone research 2000;53 Suppl 1:82-6.

1632 112. Snow-Harter C, Bouxsein M, Lewis B, Charette S, Weinstein P, Marcus R. Muscle strength as  
1633 a predictor of bone mineral density in young women. J Bone Miner Res 1990;5:589-95.

1634 113. Lofberg E, Gutierrez A, Wernerman J, et al. Effects of high doses of glucocorticoids on free  
1635 amino acids, ribosomes and protein turnover in human muscle. Eur J Clin Invest 2002;32:345-53.

1636 114. Schakman O, Gilson H, de Coninck V, et al. Insulin-like growth factor-I gene transfer by  
1637 electroporation prevents skeletal muscle atrophy in glucocorticoid-treated rats. Endocrinology  
1638 2005;146:1789-97.

1639 115. Tomas FM, Munro HN, Young VR. Effect of glucocorticoid administration on the rate of  
1640 muscle protein breakdown in vivo in rats, as measured by urinary excretion of N tau-methylhistidine.  
1641 Biochem J 1979;178:139-46.

1642 116. Ponyi A, Borgulya G, Constantin T, Vancsa A, Gergely L, Danko K. Functional outcome and  
1643 quality of life in adult patients with idiopathic inflammatory myositis. Rheumatology (Oxford)  
1644 2005;44:83-8.

1645 117. Hardy RS, Doig CL, Hussain Z, et al. 11beta-hydroxysteroid dehydrogenase type 1 within  
1646 muscle protects against the adverse effects of local inflammation. The Journal of pathology 2016.

1647 118. Rafacho A, Ortsater H, Nadal A, Quesada I. Glucocorticoid treatment and endocrine pancreas  
1648 function: implications for glucose homeostasis, insulin resistance and diabetes. *The Journal of*  
1649 *endocrinology* 2014;223:R49-62.

1650 119. Ozen G, Pedro S, Holmqvist ME, Avery M, Wolfe F, Michaud K. Risk of diabetes mellitus  
1651 associated with disease-modifying antirheumatic drugs and statins in rheumatoid arthritis. *Ann*  
1652 *Rheum Dis* 2016.

1653 120. Brennan-Speranza TC, Henneicke H, Gasparini SJ, et al. Osteoblasts mediate the adverse  
1654 effects of glucocorticoids on fuel metabolism. *JClinInvest* 2012;122:4172-89.

1655 121. Lee NK, Sowa H, Hinoi E, et al. Endocrine regulation of energy metabolism by the skeleton.  
1656 *Cell* 2007;130:456-69.

1657 122. Yeap BB, Alfonso H, Chubb SA, et al. Higher serum undercarboxylated osteocalcin and other  
1658 bone turnover markers are associated with reduced diabetes risk and lower estradiol concentrations  
1659 in older men. *The Journal of clinical endocrinology and metabolism* 2015;100:63-71.

1660 123. Pi M, Kapoor K, Ye R, et al. Evidence for Osteocalcin Binding and Activation of GPRC6A in  
1661 beta-Cells. *Endocrinology* 2016;157:1866-80.

1662 124. Mazziotti G, Maffezzoni F, Doga M, Hofbauer LC, Adler RA, Giustina A. Outcome of glucose  
1663 homeostasis in patients with glucocorticoid-induced osteoporosis undergoing treatment with bone  
1664 active-drugs. *Bone* 2014;67:175-80.

1665 125. Suchacki KJ, Roberts F, Lovdel A, et al. Skeletal energy homeostasis: a paradigm of endocrine  
1666 discovery. *The Journal of endocrinology* 2017;234:R67-R79.

1667 126. Dirckx N, Tower RJ, Mercken EM, et al. Vhl deletion in osteoblasts boosts cellular glycolysis  
1668 and improves global glucose metabolism. *The Journal of clinical investigation* 2018;128:1087-105.

1669 127. Morgan SA, McCabe EL, Gathercole LL, et al. 11beta-HSD1 is the major regulator of the  
1670 tissue-specific effects of circulating glucocorticoid excess. *Proceedings of the National Academy of*  
1671 *Sciences of the United States of America* 2014;111:E2482-91.

1672 128. Yang M, Trettel LB, Adams DJ, Harrison JR, Canalis E, Kream BE. Col3.6-HSD2 transgenic  
1673 mice: a glucocorticoid loss-of-function model spanning early and late osteoblast differentiation.  
1674 Bone 2010;47:573-82.

1675 129. Toth M, Grossman A. Glucocorticoid-induced osteoporosis: lessons from Cushing's  
1676 syndrome. Clinical endocrinology 2013;79:1-11.

1677 130. Vestergaard P, Lindholm J, Jorgensen JO, et al. Increased risk of osteoporotic fractures in  
1678 patients with Cushing's syndrome. EurJEndocrinol 2002;146:51-6.

1679 131. Kendler DL, Bauer DC, Davison KS, et al. Vertebral Fractures: Clinical Importance and  
1680 Management. The American journal of medicine 2016;129:221 e1-10.

1681 132. Bliuc D, Nguyen ND, Milch VE, Nguyen TV, Eisman JA, Center JR. Mortality risk associated  
1682 with low-trauma osteoporotic fracture and subsequent fracture in men and women. JAMA  
1683 2009;301:513-21.

1684 133. Genant HK, Wu CY, van Kuijk C, Nevitt MC. Vertebral fracture assessment using a  
1685 semiquantitative technique. Journal of bone and mineral research : the official journal of the  
1686 American Society for Bone and Mineral Research 1993;8:1137-48.

1687 134. Tauchmanova L, Pivonello R, Di SC, et al. Bone demineralization and vertebral fractures in  
1688 endogenous cortisol excess: role of disease etiology and gonadal status. JClinEndocrinolMetab  
1689 2006;91:1779-84.

1690 135. Belaya ZE, Hans D, Rozhinskaya LY, et al. The risk factors for fractures and trabecular bone-  
1691 score value in patients with endogenous Cushing's syndrome. Archives of osteoporosis 2015;10:44.

1692 136. Barahona MJ, Sucunza N, Resmini E, et al. Deleterious effects of glucocorticoid replacement  
1693 on bone in women after long-term remission of Cushing's syndrome. JBone MinerRes 2009;24:1841-  
1694 6.

1695 137. Chiodini I, Carnevale V, Torlontano M, et al. Alterations of bone turnover and bone mass at  
1696 different skeletal sites due to pure glucocorticoid excess: study in eumenorrheic patients with  
1697 Cushing's syndrome. JClinEndocrinolMetab 1998;83:1863-7.



1698 138. Szappanos A, Toke J, Lippai D, et al. Bone turnover in patients with endogenous Cushing's  
1699 syndrome before and after successful treatment. *OsteoporosInt* 2010;21:637-45.

1700 139. Belaya ZE, Iljin AV, Melnichenko GA, et al. Diagnostic performance of osteocalcin  
1701 measurements in patients with endogenous Cushing's syndrome. *BoneKEy reports* 2016;5:815.

1702 140. Futo L, Toke J, Patocs A, et al. Skeletal differences in bone mineral area and content before  
1703 and after cure of endogenous Cushing's syndrome. *OsteoporosInt* 2008;19:941-9.

1704 141. Kawamata A, Iihara M, Okamoto T, Obara T. Bone mineral density before and after surgical  
1705 cure of Cushing's syndrome due to adrenocortical adenoma: prospective study. *World JSurg*  
1706 2008;32:890-6.

1707 142. Manning PJ, Evans MC, Reid IR. Normal bone mineral density following cure of Cushing's  
1708 syndrome. *ClinEndocrinol(Oxf)* 1992;36:229-34.

1709 143. Hermus AR, Smals AG, Swinkels LM, et al. Bone mineral density and bone turnover before  
1710 and after surgical cure of Cushing's syndrome. *JClinEndocrinolMetab* 1995;80:2859-65.

1711 144. Fassnacht M, Arlt W, Bancos I, et al. Management of adrenal incidentalomas: European  
1712 Society of Endocrinology Clinical Practice Guideline in collaboration with the European Network for  
1713 the Study of Adrenal Tumors. *European journal of endocrinology / European Federation of Endocrine*  
1714 *Societies* 2016;175:G1-G34.

1715 145. Goddard GM, Ravikumar A, Levine AC. Adrenal mild hypercortisolism. *Endocrinol Metab Clin*  
1716 *North Am* 2015;44:371-9.

1717 146. Chiodini I, Morelli V, Masserini B, et al. Bone mineral density, prevalence of vertebral  
1718 fractures, and bone quality in patients with adrenal incidentalomas with and without subclinical  
1719 hypercortisolism: an Italian multicenter study. *JClinEndocrinolMetab* 2009;94:3207-14.

1720 147. Eller-Vainicher C, Morelli V, Olivieri FM, et al. Bone quality, as measured by trabecular bone  
1721 score in patients with adrenal incidentalomas with and without subclinical hypercortisolism. *Journal*  
1722 *of bone and mineral research : the official journal of the American Society for Bone and Mineral*  
1723 *Research* 2012;27:2223-30.

1724 148. Chiodini I, Torlontano M, Carnevale V, et al. Bone loss rate in adrenal incidentalomas: a  
1725 longitudinal study. *The Journal of clinical endocrinology and metabolism* 2001;86:5337-41.

1726 149. Chiodini I, Viti R, Coletti F, et al. Eugonadal male patients with adrenal incidentalomas and  
1727 subclinical hypercortisolism have increased rate of vertebral fractures. *ClinEndocrinol(Oxf)*  
1728 2009;70:208-13.

1729 150. Morelli V, Eller-Vainicher C, Palmieri S, et al. Prediction of Vertebral Fractures in Patients  
1730 With Monolateral Adrenal Incidentalomas. *The Journal of clinical endocrinology and metabolism*  
1731 2016;101:2768-75.

1732 151. Chiodini I, Mascia ML, Muscarella S, et al. Subclinical hypercortisolism among outpatients  
1733 referred for osteoporosis. *Ann InternMed* 2007;147:541-8.

1734 152. Eller-Vainicher C, Cairoli E, Zhukouskaya VV, et al. Prevalence of subclinical contributors to  
1735 low bone mineral density and/or fragility fracture. *European journal of endocrinology / European*  
1736 *Federation of Endocrine Societies* 2013;169:225-37.

1737 153. Salcuni AS, Morelli V, Eller Vainicher C, et al. Adrenalectomy reduces the risk of vertebral  
1738 fractures in patients with monolateral adrenal incidentalomas and subclinical hypercortisolism.  
1739 *European journal of endocrinology / European Federation of Endocrine Societies* 2016;174:261-9.

1740 154. Tauchmanova L, Guerra E, Pivonello R, et al. Weekly clodronate treatment prevents bone  
1741 loss and vertebral fractures in women with subclinical Cushing's syndrome. *JEndocrinolInvest*  
1742 2009;32:390-4.

1743 155. Heshmati HM, Riggs BL, Burritt MF, McAlister CA, Wollan PC, Khosla S. Effects of the  
1744 circadian variation in serum cortisol on markers of bone turnover and calcium homeostasis in normal  
1745 postmenopausal women. *The Journal of clinical endocrinology and metabolism* 1998;83:751-6.

1746 156. Raff H, Raff JL, Duthie EH, et al. Elevated salivary cortisol in the evening in healthy elderly  
1747 men and women: correlation with bone mineral density. *JGerontolA BiolSciMed Sci* 1999;54:M479-  
1748 M83.

1749 157. Dennison E, Hindmarsh P, Fall C, et al. Profiles of endogenous circulating cortisol and bone  
1750 mineral density in healthy elderly men. *JClinEndocrinolMetab* 1999;84:3058-63.

1751 158. van Schoor NM, Dennison E, Lips P, Uitterlinden AG, Cooper C. Serum fasting cortisol in  
1752 relation to bone, and the role of genetic variations in the glucocorticoid receptor.  
1753 *ClinEndocrinol(Oxf)* 2007;67:871-8.

1754 159. Cooper MS, Syddall HE, Fall CH, et al. Circulating cortisone levels are associated with  
1755 biochemical markers of bone formation and lumbar spine BMD: the Hertfordshire Cohort Study.  
1756 *ClinEndocrinol(Oxf)* 2005;62:692-7.

1757 160. Shi L, Sanchez-Guijo A, Hartmann MF, et al. Higher glucocorticoid secretion in the  
1758 physiological range is associated with lower bone strength at the proximal radius in healthy children:  
1759 importance of protein intake adjustment. *Journal of bone and mineral research : the official journal*  
1760 *of the American Society for Bone and Mineral Research* 2015;30:240-8.

1761 161. Greendale GA, Unger JB, Rowe JW, Seeman TE. The relation between cortisol excretion and  
1762 fractures in healthy older people: results from the MacArthur studies-Mac. *JAmGeriatrSoc*  
1763 1999;47:799-803.

1764 162. Gonzalez Rodriguez E, Lamy O, Stoll D, et al. High Evening Cortisol Level Is Associated With  
1765 Low TBS and Increased Prevalent Vertebral Fractures: OsteoLaus Study. *The Journal of clinical*  
1766 *endocrinology and metabolism* 2017;102:2628-36.

1767 163. Huizenga NA, Koper JW, De Lange P, et al. A polymorphism in the glucocorticoid receptor  
1768 gene may be associated with and increased sensitivity to glucocorticoids in vivo.  
1769 *JClinEndocrinolMetab* 1998;83:144-51.

1770 164. Szappanos A, Patocs A, Toke J, et al. BclI polymorphism of the glucocorticoid receptor gene is  
1771 associated with decreased bone mineral density in patients with endogenous hypercortisolism.  
1772 *ClinEndocrinol(Oxf)* 2009;71:636-43.

1773 165. Koetz KR, van Rossum EF, Venz M, Diederich S, Quinkler M. BclI polymorphism of the  
1774 glucocorticoid receptor gene is associated with increased bone resorption in patients on  
1775 glucocorticoid replacement therapy. *Clinical endocrinology* 2013;78:831-7.

1776 166. van Oosten MJ, Dolhain RJ, Koper JW, et al. Polymorphisms in the glucocorticoid receptor  
1777 gene that modulate glucocorticoid sensitivity are associated with rheumatoid arthritis. *Arthritis Res*  
1778 *Ther* 2010;12:R159.

1779 167. Quax RA, Koper JW, Huisman AM, et al. Polymorphisms in the glucocorticoid receptor gene  
1780 and in the glucocorticoid-induced transcript 1 gene are associated with disease activity and response  
1781 to glucocorticoid bridging therapy in rheumatoid arthritis. *Rheumatology international*  
1782 2015;35:1325-33.

1783 168. Meijer OC, de Lange EC, Breimer DD, de Boer AG, Workel JO, de Kloet ER. Penetration of  
1784 dexamethasone into brain glucocorticoid targets is enhanced in mdr1A P-glycoprotein knockout  
1785 mice. *Endocrinology* 1998;139:1789-93.

1786 169. Issa S, Schnabel D, Feix M, et al. Human osteoblast-like cells express predominantly steroid  
1787 5alpha-reductase type 1. *The Journal of clinical endocrinology and metabolism* 2002;87:5401-7.

1788 170. Chapman K, Holmes M, Seckl J. 11beta-hydroxysteroid dehydrogenases: intracellular gate-  
1789 keepers of tissue glucocorticoid action. *Physiological reviews* 2013;93:1139-206.

1790 171. Lavery GG, Walker EA, Draper N, et al. Hexose-6-phosphate dehydrogenase knock-out mice  
1791 lack 11 beta-hydroxysteroid dehydrogenase type 1-mediated glucocorticoid generation. *JBiolChem*  
1792 2006;281:6546-51.

1793 172. Bellows CG, Ciaccia A, Heersche JN. Osteoprogenitor cells in cell populations derived from  
1794 mouse and rat calvaria differ in their response to corticosterone, cortisol, and cortisone. *Bone*  
1795 1998;23:119-25.

1796 173. Bland R, Worker CA, Noble BS, et al. Characterization of 11 $\beta$ -hydroxysteroid dehydrogenase  
1797 activity and corticosteroid receptor expression in human osteosarcoma cell lines. *JEndocrinol*  
1798 1999;161:455-64.

1799 174. Patel P, Hardy R, Sumathi V, et al. Expression of 11beta-hydroxysteroid dehydrogenase  
1800 enzymes in human osteosarcoma: potential role in pathogenesis and as targets for treatments.  
1801 EndocrRelat Cancer 2012;19:589-98.

1802 175. Cooper MS, Walker EA, Bland R, Fraser WD, Hewison M, Stewart PM. Expression and  
1803 functional consequences of 11 $\beta$ -hydroxysteroid dehydrogenase activity in human bone. Bone  
1804 2000;27:375-81.

1805 176. Eijken M, Hewison M, Cooper MS, et al. 11 $\beta$ -Hydroxysteroid dehydrogenase expression and  
1806 glucocorticoid synthesis are directed by a molecular switch during osteoblast differentiation.  
1807 MolEndocrinol 2005;19:621-31.

1808 177. Eyre LJ, Rabbitt EH, Bland R, et al. Expression of 11 $\beta$ -hydroxysteroid dehydrogenase in rat  
1809 osteoblastic cells: Pre-receptor regulation of glucocorticoid responses in bone. JCell Biochem  
1810 2001;81:453-62.

1811 178. Cooper MS, Rabbitt EH, Goddard PE, Bartlett WA, Hewison M, Stewart PM. Osteoblastic 11 $\beta$ -  
1812 hydroxysteroid dehydrogenase type 1 activity increases with age and glucocorticoid exposure. JBone  
1813 MinerRes 2002;17:979-86.

1814 179. Cooper MS, Bujalska I, Rabbitt E, et al. Modulation of 11 $\beta$ -hydroxysteroid dehydrogenase  
1815 isozymes by proinflammatory cytokines in osteoblasts: an autocrine switch from glucocorticoid  
1816 inactivation to activation. JBone MinerRes 2001;16:1037-44.

1817 180. Ahasan MM, Hardy R, Jones C, et al. Inflammatory regulation of glucocorticoid metabolism  
1818 in mesenchymal stromal cells. Arthritis Rheum 2012;64:2404-013.

1819 181. Sai S, Esteves CL, Kelly V, et al. Glucocorticoid regulation of the promoter of 11beta-  
1820 hydroxysteroid dehydrogenase type 1 is indirect and requires CCAAT/enhancer-binding protein-beta.  
1821 Mol Endocrinol 2008;22:2049-60.

1822 182. Yang Z, Zhu X, Guo C, Sun K. Stimulation of 11beta-HSD1 expression by IL-1beta via a C/EBP  
1823 binding site in human fetal lung fibroblasts. Endocrine 2009;36:404-11.

1824 183. Kaur K, Hardy R, Ahasan MM, et al. Synergistic induction of local glucocorticoid generation  
1825 by inflammatory cytokines and glucocorticoids: implications for inflammation associated bone loss.  
1826 *AnnRheumDis* 2009;Jun 22. [Epub ahead of print].

1827 184. Hwang JY, Lee SH, Kim GS, et al. HSD11B1 polymorphisms predicted bone mineral density  
1828 and fracture risk in postmenopausal women without a clinically apparent hypercortisolemia. *Bone*  
1829 2009;45:1098-103.

1830 185. Szappanos A, Patocs A, Gergics P, et al. The 83,557insA variant of the gene coding 11beta-  
1831 hydroxysteroid dehydrogenase type 1 enzyme associates with serum osteocalcin in patients with  
1832 endogenous Cushing's syndrome. *The Journal of steroid biochemistry and molecular biology*  
1833 2011;123:79-84.

1834 186. Feldman K, Szappanos A, Butz H, et al. The rs4844880 polymorphism in the promoter region  
1835 of the HSD11B1 gene associates with bone mineral density in healthy and postmenopausal  
1836 osteoporotic women. *Steroids* 2012;77:1345-51.

1837 187. Siggelkow H, Etmanski M, Bozkurt S, et al. Genetic polymorphisms in 11beta-hydroxysteroid  
1838 dehydrogenase type 1 correlate with the postdexamethasone cortisol levels and bone mineral  
1839 density in patients evaluated for osteoporosis. *The Journal of clinical endocrinology and metabolism*  
1840 2014;99:E293-302.

1841 188. Justesen J, Mosekilde L, Holmes M, et al. Mice Deficient in 11 $\beta$ -Hydroxysteroid  
1842 dehydrogenase Type 1 Lack Bone Marrow Adipocytes but Maintain Normal Bone Formation.  
1843 *Endocrinology* 2004.

1844 189. Kotelevtsev Y, Holmes MC, Burchell A, et al. 11 $\beta$ -hydroxysteroid dehydrogenase type 1  
1845 knockout mice show attenuated glucocorticoid-inducible responses and resist hyperglycemia on  
1846 obesity or stress. *ProcNatlAcadSciUSA* 1997;94:14924-9.

1847 190. Cooper MS, Blumsohn A, Goddard PE, et al. 11 $\beta$ -hydroxysteroid dehydrogenase type 1  
1848 activity predicts the effects of glucocorticoids on bone. *JClinEndocrinolMetab* 2003;88:3874-7.

1849 191. Hardy RS, Seibel MJ, Cooper MS. Targeting 11beta-hydroxysteroid dehydrogenases: a novel  
1850 approach to manipulating local glucocorticoid levels with implications for rheumatic disease. *Curr*  
1851 *Opin Pharmacol* 2013;13:440-4.

1852 192. Cooper MS, Kriel H, Sayers A, et al. Can 11beta-hydroxysteroid dehydrogenase activity  
1853 predict the sensitivity of bone to therapeutic glucocorticoids in inflammatory bowel disease?  
1854 *Calcified tissue international* 2011;89:246-51.

1855 193. Hansen M, Florescu A, Stoltenberg M, et al. Bone loss in rheumatoid arthritis. Influence of  
1856 disease activity, duration of the disease, functional capacity, and corticosteroid treatment. *Scand J*  
1857 *Rheumatol* 1996;25:367-76.

1858 194. Kirwan JR. The effect of glucocorticoids on joint destruction in rheumatoid arthritis. The  
1859 Arthritis and Rheumatism Council Low-Dose Glucocorticoid Study Group. *N Engl J Med*  
1860 1995;333:142-6.

1861 195. Hardy RS, Doig CL, Hussain Z, et al. 11beta-Hydroxysteroid dehydrogenase type 1 within  
1862 muscle protects against the adverse effects of local inflammation. *The Journal of pathology*  
1863 2016;240:472-83.

1864 196. Landewe RB, Boers M, Verhoeven AC, et al. COBRA combination therapy in patients with  
1865 early rheumatoid arthritis: long-term structural benefits of a brief intervention. *Arthritis Rheum*  
1866 2002;46:347-56.

1867 197. Kirwan JR. The effect of glucocorticoids on joint destruction in rheumatoid arthritis. The  
1868 Arthritis and Rheumatism Council Low-Dose Glucocorticoid Study Group. *N Engl J Med* 1995;333:142-  
1869 6.

1870 198. Gough A, Sambrook P, Devlin J, et al. Osteoclastic activation is the principal mechanism  
1871 leading to secondary osteoporosis in rheumatoid arthritis. *J Rheumatol* 1998;25:1282-9.

1872 199. van Staa TP, Leufkens HG, Abenhaim L, Begaud B, Zhang B, Cooper C. Use of oral  
1873 corticosteroids in the United Kingdom. *QJM* 2000;93:105-11.

1874 200. Walsh LJ, Wong CA, Pringle M, Tattersfield AE. Use of oral corticosteroids in the community  
1875 and the prevention of secondary osteoporosis: a cross sectional study. *BMJ* 1996;313:344-6.

1876 201. Overman RA, Yeh JY, Deal CL. Prevalence of oral glucocorticoid usage in the United States: a  
1877 general population perspective. *Arthritis care & research* 2013;65:294-8.

1878 202. Silverman S, Curtis J, Saag K, et al. International management of bone health in  
1879 glucocorticoid-exposed individuals in the observational GLOW study. *Osteoporosis international : a*  
1880 *journal established as result of cooperation between the European Foundation for Osteoporosis and*  
1881 *the National Osteoporosis Foundation of the USA* 2015;26:419-20.

1882 203. Fardet L, Petersen I, Nazareth I. Prevalence of long-term oral glucocorticoid prescriptions in  
1883 the UK over the past 20 years. *Rheumatology* 2011;50:1982-90.

1884 204. Laugesen K, Jorgensen JOL, Sorensen HT, Petersen I. Systemic glucocorticoid use in  
1885 Denmark: a population-based prevalence study. *BMJ open* 2017;7:e015237.

1886 205. van Staa TP, Leufkens HG, Cooper C. Use of inhaled corticosteroids and risk of fractures.  
1887 *JBone MinerRes* 2001;16:581-8.

1888 206. van Staa TP, Leufkens HG, Abenhaim L, Zhang B, Cooper C. Oral corticosteroids and fracture  
1889 risk: relationship to daily and cumulative doses. *Rheumatology(Oxford)* 2000;39:1383-9.

1890 207. Balasubramanian A, Wade SW, Adler RA, et al. Glucocorticoid exposure and fracture risk in  
1891 patients with new-onset rheumatoid arthritis. *Osteoporosis international : a journal established as*  
1892 *result of cooperation between the European Foundation for Osteoporosis and the National*  
1893 *Osteoporosis Foundation of the USA* 2016;27:3239-49.

1894 208. Majumdar SR, Morin SN, Lix LM, Leslie WD. Influence of recency and duration of  
1895 glucocorticoid use on bone mineral density and risk of fractures: population-based cohort study.  
1896 *Osteoporosis international : a journal established as result of cooperation between the European*  
1897 *Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA* 2013;24:2493-8.

1898 209. De VF, Bracke M, Leufkens HG, Lammers JW, Cooper C, van Staa TP. Fracture risk with  
1899 intermittent high-dose oral glucocorticoid therapy. *Arthritis Rheum* 2007;56:208-14.



- 1900 210. Waljee AK, Rogers MA, Lin P, et al. Short term use of oral corticosteroids and related harms  
1901 among adults in the United States: population based cohort study. *BMJ* 2017;357:j1415.
- 1902 211. Cohen S, Levy RM, Keller M, et al. Risedronate therapy prevents corticosteroid-induced bone  
1903 loss: a twelve-month, multicenter, randomized, double-blind, placebo-controlled, parallel-group  
1904 study. *Arthritis Rheum* 1999;42:2309-18.
- 1905 212. Adachi JD, Bensen WG, Brown J, et al. Intermittent etidronate therapy to prevent  
1906 corticosteroid-induced osteoporosis. *NEngJMed* 1997;337:382-7.
- 1907 213. Saag KG, Emkey R, Schnitzer TJ, et al. Alendronate for the prevention and treatment of  
1908 glucocorticoid-induced osteoporosis. Glucocorticoid-Induced Osteoporosis Intervention Study  
1909 Group. *NEngJMed* 1998;339:292-9.
- 1910 214. Amiche MA, Albaum JM, Tadrous M, et al. Fracture risk in oral glucocorticoid users: a  
1911 Bayesian meta-regression leveraging control arms of osteoporosis clinical trials. *Osteoporosis  
1912 international : a journal established as result of cooperation between the European Foundation for  
1913 Osteoporosis and the National Osteoporosis Foundation of the USA* 2016;27:1709-18.
- 1914 215. Zhu TY, Griffith JF, Qin L, et al. Cortical thinning and progressive cortical porosity in female  
1915 patients with systemic lupus erythematosus on long-term glucocorticoids: a 2-year case-control  
1916 study. *Osteoporosis international : a journal established as result of cooperation between the  
1917 European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA*  
1918 2015;26:1759-71.
- 1919 216. Paggiosi MA, Peel NF, Eastell R. The impact of glucocorticoid therapy on trabecular bone  
1920 score in older women. *Osteoporosis international : a journal established as result of cooperation  
1921 between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of  
1922 the USA* 2015;26:1773-80.
- 1923 217. Olsson A, Oturai DB, Sorensen PS, Oturai PS, Oturai AB. Short-term, high-dose glucocorticoid  
1924 treatment does not contribute to reduced bone mineral density in patients with multiple sclerosis.  
1925 *Multiple sclerosis (Houndmills, Basingstoke, England)* 2015;21:1557-65.

1926 218. Chen SJ, Liao WC, Huang KH, et al. Chronic obstructive pulmonary disease and allied  
1927 conditions is a strong independent risk factor for osteoporosis and pathologic fractures: a  
1928 population-based cohort study. QJM : monthly journal of the Association of Physicians  
1929 2015;108:633-40.

1930 219. Kanis JA, Johansson H, Oden A, McCloskey EV. Guidance for the adjustment of FRAX  
1931 according to the dose of glucocorticoids. Osteoporosis international : a journal established as result  
1932 of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis  
1933 Foundation of the USA 2011;22:809-16.

1934 220. Nguyen ND, Frost SA, Center JR, Eisman JA, Nguyen TV. Development of prognostic  
1935 nomograms for individualizing 5-year and 10-year fracture risks. Osteoporosis international : a  
1936 journal established as result of cooperation between the European Foundation for Osteoporosis and  
1937 the National Osteoporosis Foundation of the USA 2008;19:1431-44.

1938 221. Buckley L, Guyatt G, Fink HA, et al. 2017 American College of Rheumatology Guideline for  
1939 the Prevention and Treatment of Glucocorticoid-Induced Osteoporosis. Arthritis & rheumatology  
1940 (Hoboken, NJ) 2017;69:1521-37.

1941 222. Lekamwasam S, Adachi JD, Agnusdei D, et al. A framework for the development of guidelines  
1942 for the management of glucocorticoid-induced osteoporosis. Osteoporosis international : a journal  
1943 established as result of cooperation between the European Foundation for Osteoporosis and the  
1944 National Osteoporosis Foundation of the USA 2012;23:2257-76.

1945 223. Compston J, Cooper A, Cooper C, et al. UK clinical guideline for the prevention and  
1946 treatment of osteoporosis. Archives of osteoporosis 2017;12:43.

1947 224. Reid DM, Devogelaer JP, Saag K, et al. Zoledronic acid and risedronate in the prevention and  
1948 treatment of glucocorticoid-induced osteoporosis (HORIZON): a multicentre, double-blind, double-  
1949 dummy, randomised controlled trial. Lancet 2009;373:1253-63.

1950 225. Saag KG, Shane E, Boonen S, et al. Teriparatide or alendronate in glucocorticoid-induced  
1951 osteoporosis. NEnglJMed 2007;357:2028-39.

1952 226. Adachi JD, Saag KG, Delmas PD, et al. Two-year effects of alendronate on bone mineral  
1953 density and vertebral fracture in patients receiving glucocorticoids: a randomized, double-blind,  
1954 placebo-controlled extension trial. *Arthritis Rheum* 2001;44:202-11.

1955 227. Wallach S, Cohen S, Reid DM, et al. Effects of risedronate treatment on bone density and  
1956 vertebral fracture in patients on corticosteroid therapy. *CalcifTissue Int* 2000;67:277-85.

1957 228. Reginster JY, Kuntz D, Verdickt W, et al. Prophylactic use of alfacalcidol in corticosteroid-  
1958 induced osteoporosis. *OsteoporosInt* 1999;9:75-81.

1959 229. Ringe JD, Coster A, Meng T, Schacht E, Umbach R. Treatment of glucocorticoid-induced  
1960 osteoporosis with alfacalcidol/calcium versus vitamin D/calcium. *CalcifTissue Int* 1999;65:337-40.

1961 230. de Nijs RN, Jacobs JW, Lems WF, et al. Alendronate or alfacalcidol in glucocorticoid-induced  
1962 osteoporosis. *NEnglJMed* 2006;355:675-84.

1963 231. Gluer CC, Marin F, Ringe JD, et al. Comparative effects of teriparatide and risedronate in  
1964 glucocorticoid-induced osteoporosis in men: 18-month results of the EuroGIOPs trial. *Journal of*  
1965 *bone and mineral research : the official journal of the American Society for Bone and Mineral*  
1966 *Research* 2013;28:1355-68.

1967 232. Saag KG, Agnusdei D, Hans D, et al. Trabecular Bone Score in Patients With Chronic  
1968 Glucocorticoid Therapy-Induced Osteoporosis Treated With Alendronate or Teriparatide. *Arthritis &*  
1969 *rheumatology (Hoboken, NJ)* 2016;68:2122-8.

1970 233. Overman RA, Gourlay ML, Deal CL, Farley JF, Brookhart MA, Layton JB. Fracture rate  
1971 associated with quality metric-based anti-osteoporosis treatment in glucocorticoid-induced  
1972 osteoporosis. *Osteoporosis international : a journal established as result of cooperation between the*  
1973 *European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA*  
1974 *2015;26:1515-24.*

1975 234. Axelsson KF, Nilsson AG, Wedel H, Lundh D, Lorentzon M. Association Between Alendronate  
1976 Use and Hip Fracture Risk in Older Patients Using Oral Prednisolone. *JAMA* 2017;318:146-55.

1977 235. Bergman J, Nordstrom A, Nordstrom P. Alendronate Use and the Risk of Nonvertebral  
1978 Fracture During Glucocorticoid Therapy: A Retrospective Cohort Study. *The Journal of clinical*  
1979 *endocrinology and metabolism* 2018;103:306-13.

1980 236. Mellibovsky L, Prieto-Alhambra D, Mellibovsky F, et al. Bone Tissue Properties Measurement  
1981 by Reference Point Indentation in Glucocorticoid-Induced Osteoporosis. *Journal of bone and mineral*  
1982 *research : the official journal of the American Society for Bone and Mineral Research* 2015;30:1651-  
1983 6.

1984 237. Girgis CM, Sher D, Seibel MJ. Atypical femoral fractures and bisphosphonate use. *The New*  
1985 *England journal of medicine* 2010;362:1848-9.

1986 238. Schilcher J, Koeppen V, Aspenberg P, Michaelsson K. Risk of atypical femoral fracture during  
1987 and after bisphosphonate use. *Acta orthopaedica* 2015;86:100-7.

1988 239. Ishiguro S, Ito K, Nakagawa S, Hataji O, Sudo A. The clinical benefits of denosumab for  
1989 prophylaxis of steroid-induced osteoporosis in patients with pulmonary disease. *Archives of*  
1990 *osteoporosis* 2017;12:44.

1991 240. Sawamura M, Komatsuda A, Togashi M, Wakui H, Takahashi N. Effects of Denosumab on  
1992 Bone Metabolic Markers and Bone Mineral Density in Patients Treated with Glucocorticoids. *Internal*  
1993 *medicine (Tokyo, Japan)* 2017;56:631-6.

1994 241. Mok CC, Ho LY, Ma KM. Switching of oral bisphosphonates to denosumab in chronic  
1995 glucocorticoid users: a 12-month randomized controlled trial. *Bone* 2015;75:222-8.

1996 242. Bjornsdottir S, Saaf M, Bensing S, Kampe O, Michaelsson K, Ludvigsson JF. Risk of hip fracture  
1997 in Addison's disease: a population-based cohort study. *Journal of internal medicine* 2011;270:187-  
1998 95.

1999 243. Mor F, Green P, Wysenbeek AJ. Myopathy in Addison's disease. *Ann Rheum Dis* 1987;46:81-  
2000 3.

2001 244. Lee P, Greenfield JR. What is the optimal bone-preserving strategy for patients with  
2002 Addison's disease? *Clinical endocrinology* 2015;83:157-61.

2003 245. Koetz KR, Vents M, Diederich S, Quinkler M. Bone mineral density is not significantly reduced  
2004 in adult patients on low-dose glucocorticoid replacement therapy. *The Journal of clinical*  
2005 *endocrinology and metabolism* 2012;97:85-92.

2006 246. Camozzi V, Betterle C, Frigo AC, et al. Vertebral fractures assessed with dual-energy X-ray  
2007 absorptiometry in patients with Addison's disease on glucocorticoid and mineralocorticoid  
2008 replacement therapy. *Endocrine* 2018;59:319-29.

2009 247. Schulz J, Frey KR, Cooper MS, et al. Reduction in daily hydrocortisone dose improves bone  
2010 health in primary adrenal insufficiency. *European journal of endocrinology / European Federation of*  
2011 *Endocrine Societies* 2016;174:531-8.

2012 248. Ceccato F, Barbot M, Albiger N, et al. Long-term glucocorticoid effect on bone mineral  
2013 density in patients with congenital adrenal hyperplasia due to 21-hydroxylase deficiency. *European*  
2014 *journal of endocrinology / European Federation of Endocrine Societies* 2016;175:101-6.

2015 249. Elnecave RH, Kopacek C, Rigatto M, Keller BJ, Sisson de Castro JA. Bone mineral density in  
2016 girls with classical congenital adrenal hyperplasia due to CYP21 deficiency. *JPediatrEndocrinolMetab*  
2017 *2008;21:1155-62.*

2018 250. El-Maouche D, Collier S, Prasad M, Reynolds JC, Merke DP. Cortical bone mineral density in  
2019 patients with congenital adrenal hyperplasia due to 21-hydroxylase deficiency. *Clinical*  
2020 *endocrinology* 2015;82:330-7.

2021 251. Falhammar H, Filipsson H, Holmdahl G, et al. Fractures and bone mineral density in adult  
2022 women with 21-hydroxylase deficiency. *JClinEndocrinolMetab* 2007;92:4643-9.

2023 252. Falhammar H, Filipsson Nystrom H, Wedell A, Brismar K, Thoren M. Bone mineral density,  
2024 bone markers, and fractures in adult males with congenital adrenal hyperplasia. *European journal of*  
2025 *endocrinology / European Federation of Endocrine Societies* 2013;168:331-41.

2026 253. Florio M, Gunasekaran K, Stolina M, et al. A bispecific antibody targeting sclerostin and DKK-  
2027 1 promotes bone mass accrual and fracture repair. *Nature communications* 2016;7:11505.

2028 254. Newton R, Holden NS. Separating transrepression and transactivation: a distressing divorce  
2029 for the glucocorticoid receptor? *Molecular pharmacology* 2007;72:799-809.

2030 255. Cooper MS, Zhou H, Seibel MJ. Selective glucocorticoid receptor agonists: glucocorticoid  
2031 therapy with no regrets? *Journal of bone and mineral research : the official journal of the American*  
2032 *Society for Bone and Mineral Research* 2012;27:2238-41.

2033 256. Thiele S, Ziegler N, Tsourdi E, et al. Selective glucocorticoid receptor modulation maintains  
2034 bone mineral density in mice. *Journal of bone and mineral research : the official journal of the*  
2035 *American Society for Bone and Mineral Research* 2012;27:2242-50.

2036

2037