

Selective recognition and electrochemical sensing of dopamine using a ferrocene-based heteroditopic receptor

Miyaji, Hidekazu; Komada, Haruka; Goto, Keisuke; Fujimoto, Junko; Kiriya, Naoaki; Tucker, James H. R.

DOI:

[10.1016/j.tetlet.2018.09.026](https://doi.org/10.1016/j.tetlet.2018.09.026)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Miyaji, H, Komada, H, Goto, K, Fujimoto, J, Kiriya, N & Tucker, JHR 2018, 'Selective recognition and electrochemical sensing of dopamine using a ferrocene-based heteroditopic receptor', *Tetrahedron Letters*, pp. 3853-3857. <https://doi.org/10.1016/j.tetlet.2018.09.026>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Selective recognition and electrochemical sensing of dopamine using a ferrocene-based heteroditopic receptor

Hidekazu Miyaji^{a,*}, Haruka Komada^a, Keisuke Goto^a, Junko Fujimoto^a, Naoaki Kiriya^a and James H. R. Tucker^b

^a Department of Chemistry and Biomolecular Science, Faculty of Engineering, Gifu University, 1-1 Yanagido, Gifu 501-1193, Japan

^b School of Chemistry, University of Birmingham, Edgbaston, Birmingham B15 2TT, U.K.

ARTICLE INFO

Article history:

Received

Received in revised form

Accepted

Available online

Keywords:

Electrochemical molecular sensor

Dopamine

Ferrocene

Boronic acid

Crown ether

ABSTRACT

A redox-active ferrocene-based heteroditopic receptor bearing a boronic acid (as a catechol recognition site) and a benzo-18-crown-6-ether unit (as an ammonium ion recognition site) was synthesized. A 1:1 ditopic complex with dopamine was evidenced by mass spectrometry and NMR spectroscopy. Cyclic voltammetry measurements on the receptor in the presence of a series of organic guest species demonstrated the successful electrochemical sensing of dopamine through a distinct change in the ferrocene-centred redox-couple upon complex formation.

2009 Elsevier Ltd. All rights reserved.

Dopamine (DA) is a neurotransmitter that plays an important role in the central nervous system, with DA levels significantly affecting brain function. Low levels of DA in vivo are associated with Parkinson's disease, while excessive amounts are associated with attention deficit hyperactivity disorder (ADHD) and motor dysfunction. Therefore, accurately determining DA levels is important in terms of the diagnosis of disease and monitoring its progression. Although techniques such as capillary electrophoresis¹ and high performance liquid chromatography (HPLC),² have been used to detect DA, electrochemical techniques³ have an advantage as the measurement can be performed with high sensitivity, in real time and at low cost. However, since there are several compounds in vivo that have almost the same oxidation potential as DA,⁴ it is very difficult to unambiguously identify and quantify DA levels in biological media through electrochemical means.^{5,6} One potential solution to this issue is to develop redox-active receptors that signal the presence of the analyte at a separate oxidation potential, as has previously been demonstrated for other organic guest molecules.⁷

In this study, we report the synthesis of a ferrocene-containing receptor **1** designed to selectively bind and electrochemically sense DA as its hydrochloride salt. Compound **1** is a heteroditopic receptor⁸⁻¹⁰ that simultaneously recognizes both the catechol moiety of the guest through its boronic acid¹¹⁻¹⁵ and the ammonium ion through its benzocrown ether (Figure 1).^{16,17} We show that electrochemical detection of DA¹⁸ is achieved by monitoring the change in the ferrocene-centred redox couple of **1** upon its binding in acetonitrile.





Figure 1. A ferrocene-based heteroditopic receptor **1** and its multipoint recognition of dopamine hydrochloride.



Scheme 1. Synthetic route to compound **1**.

Synthesis of compound **1** was carried out after mono-protection of ferrocene diboronic acid using pinacol,¹⁹ followed by the introduction of a benzo-18-crown-6-ether unit by Suzuki coupling.²⁰ The pinacol was then deprotected by polystyrene boronic acid²¹ to give the compound **1** in overall yield of 28% (Scheme 1, see SI for experimental details).

A series of experiments were carried out to establish the binding properties of compound **1** with DA. The molecular ion peaks $[1+H]^+$ = 541.2, $[1+Na]^+$ = 563.2 and $[1+K]^+$ = 579.1 were clearly present in its ESI mass spectrum. When 1.0 molar equivalent of dopamine hydrochloride (DA·HCl) was added to a solution of **1** in CH₃CN containing 5.0% H₂O, these signals decreased at the expense of two new peaks of higher mass corresponding to the 1:1 complex with DA: $[1-2H_2O+DA+H]^+$ = 658.2 and $[1-2H_2O+DA\cdot HCl+H]^+$ = 694.3. This result clearly demonstrates the binding of compound **1** to DA, with the loss of two water molecules indicating the formation of the expected boronate ester adduct. In addition, the removal of metal ion adducts in the presence of DA is consistent with the formation of a ditopic complex **2** (Fig. 1), that also has a three-point H-bonding interaction between the protonated amine and the crown ether.




Figure 2. (a) The ESI mass spectrum of compound **1** (5.0% H₂O in CH₃CN) in (a) the presence and (b) the absence of 1.0 mol. equiv. of dopamine hydrochloride, DA·HCl.

The structure of complex **2** was calculated by DFT (B3LYP/6-31G(d,p)) using Gaussian,^{22,23} which revealed that the receptor **1** was indeed able to accommodate DA·HCl through the formation of a ditopic complex, in which the catechol unit is recognized at the boronic acid site and the primary ammonium ion at the crown ether site (Figure 3).




Figure 3. Optimized structure of complex **2** by DFT calculation.

A series of ¹H NMR experiments were then undertaken to probe the binding process further. DA·HCl (1.5 mol. equiv.) was added to 0.5 mM solutions of compound **1** in CD₃CN/H₂O with varying percentage compositions of water (see supplementary information). When the ratio of water was 1.5%,²⁴ four new signals at 4.3 ppm, 4.5 ppm, 4.7 ppm and 4.7 ppm (two overlapping), corresponding to the ferrocene cyclopentadienyl (Cp) protons (2 H each), were observed in the ¹H NMR spectrum at significantly different values to those for uncomplexed **1**. Furthermore, the characteristic multiplet at 3.5 ppm corresponding to the crown ether protons shifted slightly downfield to 3.7 ppm. In control experiments, no significant changes to the Cp proton resonances of **1** were observed when 1.5 equivalents of catechol (CA) and 2-phenylethylamine hydrochloride (2-PA·HCl) were added under the same conditions. This gives further evidence for the formation of a ditopic complex **2** between DA·HCl involving multiple point interactions that is facilitated by the ball-bearing effect of the ferrocene unit. Interestingly, under these conditions the exchange rate between unbound and bound forms of **1** was slower than the NMR time scale as signals originating from both unbound compound **1** and complex **2** were observed. Similar changes to the Cp and crown ether proton resonances were observed when the ratio of water was increased to 5.0% although the peaks for unbound **1** were significantly greater compared to the case for the 1.5% spectrum. Nevertheless, this indicates the ability of compound **1** to form a ditopic complex with DA·HCl even under more competitive solvent conditions.

In order to investigate the stoichiometry of the complex formed between compound **1** and DA·HCl, a Job plot was formed from the ¹H NMR spectra (3% H₂O in CD₃CN) by monitoring changes to the integration ratio of the newly appeared ferrocene Cp resonance at 4.5 ppm for the complex and the original ferrocene Cp resonance at 4.6 ppm for **1**. This indicated the expected 1:1 complex, albeit with a small amount of 1:2 (host:guest) complex also present (see supplementary information). Titration experiments were undertaken to assess the strength of the binding interaction between **1** and various guests by the curve fitting method.²⁵ The binding constant for the 1:1 complex with DA·HCl was determined to be $K_a = 2,900 M^{-1}$ (1.5% H₂O in CD₃CN). As expected for a ditopic complex such as **2** displaying a positive cooperative binding effect, this value was considerably higher than those determined under the same conditions for catechol (CA) and 2-phenylethylamine hydrochloride (2-PA·HCl) which gave values of $K_a = 0.8 M^{-1}$ and $1,100 M^{-1}$ respectively (see Figure 4 and supplementary information). It was also interesting to note that in contrast to the titration with DA·HCl, for 2-PA·HCl, no new peaks were observed upon complex formation, rather a gradual shift in the crown ether protons. This indicates fast guest exchange on the NMR timescale, which would be expected for a weaker complex that is not held so tightly by multiple point interactions found in **2**.

The ability of compound **1** to act as an electrochemical molecular sensor⁷ for DA was then probed using cyclic voltammetry (CV). Compound **1** gave a characteristic redox couple at $E_{1/2} = 460$ mV vs. Ag/AgCl in acetonitrile (1.5% H₂O), corresponding to the reversible oxidation of the ferrocene unit, where $E_{1/2} = (E_{pa} + E_{pc})/2$. A series of compounds were then added to investigate the effect of complexation on this redox couple.




Figure 4. Guest molecules used in this study.

Figure 5 shows the change in the CV upon the addition of 1.5 molar equivalents of DA·HCl (dissolved in water) to compound **1** (acetonitrile, 0.5 mM). A significant change was observed, with the two main waves, best explained as corresponding to the oxidation and reduction of the 1:1 complex between **1** and DA, appearing at more positive potentials, giving a $E_{1/2}$ value of 510 mV and a $\Delta E_{1/2}$ value of +70 mV (see the supplementary information for more details). When the amount of DA·HCl was increased to 2.0 and 2.5 molar equivalents, no further significant changes were observed, which again supports the formation of the 1:1 ditopic complex.




Figure 5. Cyclic voltammograms recorded for compound **1** (1.5% H₂O in CH₃CN) in the absence and in the presence of 1.5 mol. eq. of DA·HCl (sweep rate: 50 mV/s, electrode: GC, counter electrode: Pt).

In order to assess how distinct these changes were in terms of sensing and identifying DA electrochemically, a series of similar experiments was next undertaken with other organic molecules (Fig. 4, see also supplementary information). The addition of CA and 2-PA·HCl gave much lower $\Delta E_{1/2}$ values of +4 mV and +14 mV respectively. The presence of an ammonium ion bound in close proximity to the ferrocene unit would be expected to make the ferrocene unit harder to oxidise and shift the redox couple slightly to more positive potentials.⁹ However the much larger value observed with DA·HCl gives further support for the formation of the ditopic complex **2**, in which the guest is strapped across the ferrocene unit, making a more marked effect on the

electrochemical properties. Six other neurotransmitters were then tested in the same way including L-noradrenaline hydrochloride (L-NA·HCl), DL-adrenaline hydrochloride (DL-AD·HCl) and γ -aminobutyric acid hydrochloride (GABA·HCl) (Fig. 4). In all cases, less marked shifts were observed, with the greatest $\Delta E_{1/2}$ value of +30 mV observed for L-NA·HCl, presumably since this is the only guest species that can also form a multiple point ditopic complex,²⁶ having both a primary ammonium group and a catechol unit. Those compounds containing only a primary ammonium group and no catechol unit gave similar broadly similar $\Delta E_{1/2}$ values to PA·HCl, which was consistent with these guest species forming just a three-point H-bonding interaction with the benzocrown ether moiety.

In conclusion, we have succeeded in the synthesis of ferrocene compound **1** bearing both a boronic acid moiety and a benzocrown ether. It was found that this supramolecular receptor can form a 1:1 ditopic complex **2** with dopamine hydrochloride (DA·HCl), with binding able to be read out electrochemically, allowing it to function as an electrochemical sensor for this neurotransmitter.²⁷ The work demonstrates the ability to sense a redox-active target, not by monitoring the effect on its own redox properties but on those of a redox-active receptor to which it binds, thus allowing a different potential window to be monitored.

Acknowledgments

H. Miyaji and J. H. R. Tucker acknowledge the EPSRC for the funding of this work through the research grant GR/M85722/01. H. Miyaji, J. Fujimoto and H. Komada are grateful to Prof. T. Udagawa (Gifu University) for guidance in making DFT calculations.

References and notes

- Shou, M.; Ferrario, C. R.; Schultz, K. N.; Robinson, T. E.; Kennedy, R. T., *Anal. Chem.* **2006**, *78*, 6717-6725.
- Muzzi, C.; Bertocci, E.; Terzuoli, L.; porcelli, B.; Ciari, I.; Pagani, R.; Guerranti, R., *Biomed. Pharmacother.* **2008**, *62*, 253-258.
- Qian, T.; Wu, S.; Jian, J., *Chem Commun.* **2013**, *49*, 4610-4612.
- Forzani, E. S.; Rivas, G. A.; Solis, V. M., *J. Electroanal. Chem.* **1995**, *382*, 33-40.
- Hong, S.; Lee, L. Y. S.; So, M.-H.; Wong, K.-Y., *Electroanalysis.* **2013**, *25*, 1085-1094.
- Wu, W.; Zhu, H.; Fan, L.; Liu, D.; Renneberg, R.; Yang, S., *Chem Commun.* **2007**, 2345-2347.
- (a) Beer, P. D.; Gale, P. A.; Chen, G. Z., *Coord. Chem. Rev.* **1999**, *185-186*, 3-36; (b) Mulas, A.; Willener, Y.; Carr-Smith, J.; Joly, K. M.; Male, L.; Moody, C. J.; Horswell, S. L.; Nguyen, H. V.; Tucker, J. H. R., *Dalton Trans.*, **2015**, *44*, 7268-7275 and references cited therein.
- Miyaji, H.; Collinson, S. R.; Prokes, I.; Tucker, J. H. R., *Chem. Commun.* **2003**, 64-65.
- Miyaji, H.; Gasser, G.; Green, S. J.; Molard, Y.; Strawbridge, S. M.; Tucker, J. H. R., *Chem. Commun.* **2005**, 5355-5357.
- Miyaji, H.; Kim, D.-S.; Chang, B.-Y.; Park, E.; Park, S.-M.; Ahn, K. H., *Chem. Commun.* **2008**, 753-755.
- Nishiyabu, R.; Kubo, Y.; James, T. D.; Fossey, J. S., *Chem. Commun.* **2011**, *47*, 1106-1123.
- Wu, Z.; Li, M.; Fang, H.; Wang, B., *Bioorg. Med. Chem. Lett.* **2012**, *22*, 7179-7182.
- Bull, S. D.; Davidson, M. G.; van den Elsen, J. M. H.; Fossey, J. S.; Jenkins, A. T. A.; Jiang, Y.-B.; Kubo, Y.; Marken, R.; Sakurai, Z.; Zhao, J.; James, T. D., *Acc. Chem. Res.* **2013**, *46*, 312-326.
- Li, Y.; Xie, Y.; Qin, Y., *Sensors and Actuators B* **2014**, *191*, 227-232.
- Li, M.; Zhu, W.; Marken, F.; James, T. D., *Chem. Commun.* **2015**, *51*, 14562-14573.
- Pietrzyk, A.; Suriyanarayanan, S.; Kutner, W.; Maligaspe, E.; Zandler, M. E.; D' Souza, F., *Bioelectrochemistry* **2010**, *80*, 62-72.

17. Hashimoto, T.; Oyaidu, S.; Hayashita, T., *Bunseki kagaku* **2012**, *3*, 213-219.
18. For a related approach to dopamine sensing based on boronate ester formation, see Strawbridge, S. M.; Green, S. J.; Tucker, J. H. R., *Chem. Commun.* **2000**, 2393-2394.
19. D'Hooge, F.; Rogalle, D.; Thatcher, M. J.; Perera, S. P.; van den Elsen, J. M. H.; Jenkins, A. T. A.; James, T. D.; Fossey, J. S. *Polymer* **2008**, *49*, 3362-3365.
20. Miyaara, N.; Yamada, K.; Suzuki, A. *Tetrahedron Lett.* **1979**, *20*, 3437-3440.
21. Pennington, T. E.; Kardiman, C.; Hutton, C. A. *Tetrahedron Lett.* **2004**, *45*, 6657-6660.
22. Frisch, M. J., Trucks, G. W. & Schlegel, H. B. et al., Gaussian09, Revision B.01, Gaussian, Inc., Wallingford CT, 2010.
23. Kiriya, N.; Ebihara, M.; Udagawa, T.; Miyaji, H., *RSC Adv.* **2016**, *6*, 19794-19796.
24. In neat CD₃CN or in the presence of 1.0% H₂O, the binding constant between compound **1** and DA·HCl was too large to obtain an accurate value from ¹H NMR spectra ($K_a > 10,000 \text{ M}^{-1}$). Therefore, to allow a comparison of binding constants, measurements were performed in 1.5% H₂O in CD₃CN.
25. The binding constants were determined by a non-linear curve fitting method using Origin (OriginLab co.).
26. The binding constants (K_a) between compound **1** and L-NA·HCl and DL-AD·HCl were $1,800 \text{ M}^{-1}$ and 130 M^{-1} respectively, as determined from ¹H NMR titration experiments (1.5% H₂O in CD₃CN).
27. In this study, measurements were performed under standard CV conditions in organic solvents at millimolar concentrations. However, given a suitably strong binding interaction and more sensitive instrumentation, this approach could be used to detect lower concentrations of dopamine and in more competitive solvent conditions.