

Measurement of Chern numbers through center-of-mass responses

Price, Hannah M; Zilberberg, O.; Ozawa, T.; Carusotto, I.; Goldman, N.

DOI:

[10.1103/PhysRevB.93.245113](https://doi.org/10.1103/PhysRevB.93.245113)

License:

None: All rights reserved

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Price, HM, Zilberberg, O, Ozawa, T, Carusotto, I & Goldman, N 2016, 'Measurement of Chern numbers through center-of-mass responses', *Physical Review B*, vol. 93, no. 24, 245113.

<https://doi.org/10.1103/PhysRevB.93.245113>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Checked for eligibility 22/10/2018

H. M. Price, O. Zilberberg, T. Ozawa, I. Carusotto, and N. Goldman
Phys. Rev. B 93, 245113 – Published 6 June 2016
<https://doi.org/10.1103/PhysRevB.93.245113>

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.



Measurement of Chern numbers through center-of-mass responses

H. M. Price,^{1,*} O. Zilberberg,² T. Ozawa,¹ I. Carusotto,¹ and N. Goldman^{3,†}

¹*INO-CNR BEC Center and Dipartimento di Fisica, Università di Trento, I-38123 Povo, Italy*

²*Institute for Theoretical Physics, ETH Zurich, 8093 Zürich, Switzerland*

³*CENOLI, Faculté des Sciences, Université Libre de Bruxelles (U.L.B.), B-1050 Brussels, Belgium*

(Received 29 February 2016; published 6 June 2016)

Probing the center-of-mass of an ultracold atomic cloud can be used to measure Chern numbers, the topological invariants underlying the quantum Hall effects. In this work, we show how such center-of-mass observables can have a much richer dependence on topological invariants than previously discussed. In fact, the response of the center of mass depends not only on the current density, typically measured in a solid-state system, but also on the particle density, which itself can be sensitive to the topology of the band structure. We apply a semiclassical approach, supported by numerical simulations, to highlight the key differences between center-of-mass responses and more standard conductivity measurements. We illustrate this by analyzing both the two- and four-dimensional quantum Hall effects. These results have important implications for experiments in engineered topological systems, such as ultracold gases and photonics.

DOI: [10.1103/PhysRevB.93.245113](https://doi.org/10.1103/PhysRevB.93.245113)

I. INTRODUCTION

Over recent decades, there has been great interest in studying topological phases of matter [1,2]. In these systems, energy bands can be characterized by topological invariants, which have direct physical consequences in quantized bulk responses and robust edge physics. In the famous two-dimensional (2D) quantum Hall (QH) effect, for example, the Hall conductance is quantized in multiples of the integer first Chern number (1CN), an important topological invariant of 2D energy bands [3].

Although the 2D QH effect was first studied for electrons in solid-state materials subject to a perpendicular magnetic field, there has been much progress also in exploring this physics for other analog systems, such as ultracold atomic gases [4–9] and photonics [10,11]. In these systems, the particles are uncharged, and the effects of the magnetic field must be engineered artificially using other means [12–14]. These engineered platforms offer a variety of new opportunities, such as the recent proposal for engineering the four-dimensional (4D) QH effect for the first time [15,16]. This may be achieved by combining a three-dimensional system of atoms or photons with a “synthetic” dimension, where internal degrees of freedom are coupled to simulate an additional spatial dimension [8,9,17,18].

In atomic or photonic systems, different physical concepts are required, not only to access quantum Hall physics, but also to measure and probe its signatures. Usually in a solid-state material, the QH effect is observed in the electrical conductivity, namely, through voltage or current measurements. However, such measurements are difficult to perform in analog quantum Hall systems, and instead much work has gone into finding new tools for probing the topological and geometrical properties of energy bands [19–37]. In particular, QH responses can be measured in the center-of-mass drift of an ultracold atomic cloud [6,15,21,22], or in the displacement of the center of mass of the photon steady state [16,30].

Measurements of observables related to the center-of-mass (c.m.) motion typically depend not just on the current, which is usually measured in a solid-state system, but also on the particle density. For example, the c.m. velocity for a cloud of N_{tot} atoms with total velocity \mathbf{v}_{tot} is defined as

$$\mathbf{v}_{\text{c.m.}} = \frac{\mathbf{v}_{\text{tot}}}{N_{\text{tot}}} = \frac{\mathbf{j}}{n}, \quad (1)$$

where the current density is $\mathbf{j} = \mathbf{v}_{\text{tot}}/L^d$ and the particle density is $n = N_{\text{tot}}/L^d$ for a system of length L and dimension d . For a QH system, the current density is proportional to the topological invariants (Chern numbers) of filled energy bands [see, e.g., Eq. (2) below]. This behavior is captured directly by c.m. observables for the cases previously studied [6,16,21,22,30], in which the particle density n only contributes an unimportant constant factor. However, the particle density can itself become a function of the geometrical and topological properties of the energy bands in the presence of external (possibly artificial) magnetic perturbations [15,19,38–40]. As we show in this paper, center-of-mass responses can therefore exhibit a much richer dependence on topological invariants than previously discussed.

In this work, we use a semiclassical approach to explore how c.m. responses depend on the topological invariants of filled energy bands. We illustrate these results for the 2D quantum Hall effect, where we show that center-of-mass observables can exhibit nonlinear topological responses, which would not be observed in conductivity measurements, as routinely performed in solid-state systems. Our findings are directly relevant for current experiments in ultracold gases [4–9], photonics [10,11], and even classical mechanical systems [41]. As far as ultracold gases are concerned, we emphasize that our semiclassical treatment is valid for uniformly filled bands of either bosons or fermions, as can be achieved through fermionic statistics (for Fermi gases) or through thermal effects. We also build on our recent experimental proposals for the 4D quantum Hall effect [15,16] and show that in such 4D systems, our results have important implications for experimental design and detection. We demonstrate, for

*hannah.price@unitn.it

†ngoldman@ulb.ac.be

example, pathological cases in which 4D topological invariants would vanish from center-of-mass responses, while persisting in conductivity measurements.

A. Main implications for the 2D quantum Hall effect

Before proceeding, we emphasize the key implications of our results for the 2D quantum Hall effect. For a filled band of electrons, the transverse (Hall) current density follows the linear relation [3]

$$j^x = \frac{e^2}{h} E_y \nu_1^{xy}, \quad (2)$$

where E_y is an external electric field aligned along the y direction, e is the elementary charge, h is Planck's constant, and ν_1^{xy} is the topological 1CN of the populated band [defined in Eq. (4) below]; in the 2D QH effect, a nonzero 1CN, and hence a nonzero Hall current, is due to an applied perpendicular magnetic field. As the 1CN is topological, the Hall response in Eq. (2) is remarkably robust; for example, it is insensitive to small changes in the magnetic field provided that the energy gap of the system remains open. This leads to characteristic plateaus in the Hall conductivity $\sigma_{xy} = j^x/E_y$ plotted as a function of the applied magnetic field, where the height of a given plateau is proportional to the 1CN (summed over filled energy bands) [3].

In contrast, c.m. observables can have a much more complicated dependence on the topological index of an energy band. We find that the c.m. velocity (1) of an ultracold cloud, for example, can include nonlinear topological responses, such as a contribution proportional to $(\nu_1^{xy})^2$ under a small perturbation of the applied (artificial) magnetic field [see Eq. (28) below]. Such terms, which have no analog in current responses, stem from an interplay between the topological invariants appearing in both the current density and particle density in Eq. (1). These effects could be observed in state-of-the-art experiments with ultracold atoms and in photonics [4–11], where magnetic perturbations can both arise naturally from experimental uncertainties in the (artificial) magnetic flux imposed, and be engineered deliberately using current experimental techniques. Our theory allows one to clearly identify these topological effects and to understand the important differences between probing QH physics through c.m. observables versus conductivity measurements.

B. Outline

The structure of this paper is as follows: We begin by reviewing in Sec. II how the semiclassical equations of motion can be used to calculate the quantum Hall current response and other relevant observables. In Sec. III, we study a 2D quantum Hall system, emphasizing the effects of a magnetic field perturbation on the particle density and hence on any c.m. observables. In particular, we illustrate the effects of nonlinear topological responses in the c.m. transverse velocity. In Sec. IV, as a further example, we consider a 4D quantum Hall system, where the distinction between c.m. observables and current measurements has very striking implications for the design of future experiments; for example, we identify pathological configurations of the perturbing electromagnetic fields for which 4D topological invariants can be extracted

from a current response, and yet not from a center-of-mass response. Finally, we present additional experimental remarks in Sec. V and draw conclusions in Sec. VI.

II. SEMICLASSICAL APPROACH TO THE QUANTUM HALL EFFECT

In this section, we develop a semiclassical description of center-of-mass responses. In Sec. II A, we introduce the geometrical and topological properties of eigenstates in an energy band of a quantum Hall system, before reviewing the semiclassical equations of motion for a wave packet constructed out of these eigenstates in Sec. II B [42–46]. From these equations, we review how to derive the modified density of states, the quantum Hall current density, and relevant center-of-mass observables in Secs. II C–II E, respectively. By keeping terms up to second order in the external perturbing fields, this semiclassical framework can be used to describe both current experiments on 2D quantum Hall physics in ultracold gases and photonics [4–11] (see Sec. III) as well as proposed schemes for realizing the 4D quantum Hall effect [15,16] (see Sec. IV).

A. Berry curvature, the first Chern number, and the second Chern number

We begin from a particle in a periodic potential, where the eigenstates can be expressed through Bloch's theorem as $|\chi_{n,k}\rangle = e^{ik \cdot r} |u_{n,k}\rangle$, where $|u_{n,k}\rangle$ are the periodic Bloch functions and \mathbf{k} is the corresponding quasimomentum. In the Brillouin zone (BZ), the Bloch functions $|u_{n,k}\rangle$ form bands where the energy dispersion $\mathcal{E}_n(\mathbf{k})$ is labeled by the band index n . In this paper, we focus on the physics of an energetically isolated nondegenerate energy band, and so hereafter we drop the label n .

The eigenstates that make up the energy band can have nontrivial local geometrical properties as encoded, for example, in the Berry curvature [38]. In this paper, we deal with systems of various dimensionalities and so we express the Berry curvature as a general differential 2-form

$$\Omega = \frac{1}{2} \Omega^{\mu\nu}(\mathbf{k}) dk_\mu \wedge dk_\nu, \quad \Omega^{\mu\nu}(\mathbf{k}) = \partial_{k_\mu} \mathcal{A}_{k_\nu} - \partial_{k_\nu} \mathcal{A}_{k_\mu}, \quad (3)$$

where \wedge is the antisymmetric wedge product, $\mathcal{A}_{k_\mu} = i \langle u_{\mathbf{k}} | \partial_{k_\mu} | u_{\mathbf{k}} \rangle$ is the Berry connection, and where the indices μ, ν run over all spatial coordinates with Einstein summation convention. As can be seen from this definition, the Berry curvature components are antisymmetric under an exchange of indices $\Omega^{\mu\nu}(\mathbf{k}) = -\Omega^{\nu\mu}(\mathbf{k})$. Importantly, geometrical properties, such as the Berry curvature, are also closely related to key topological properties of the energy bands.

In general, topological phases of matter can be classified according to the symmetries and dimensionality of the system [48]. In this paper, we focus on noninteracting systems without any symmetries, where the energy bands are topologically trivial in odd dimensions but can be characterized by nonzero topological integers in even dimensions [3,47,48]. We consider, in particular, systems with two and four dimensions, where the relevant topological invariants are known as the first and second Chern numbers, respectively. For a 2D system

in, e.g. the x - y plane, the 1CN is calculated from the Berry curvature as

$$v_1^{xy} = \frac{1}{2\pi} \int_{\mathbb{T}^2} \Omega = \frac{1}{2\pi} \int_{\mathbb{T}^2} \Omega^{xy} dk_x dk_y \in \mathbb{Z}, \quad (4)$$

where the integral is over the first (magnetic) two-dimensional BZ, which is denoted here by \mathbb{T}^2 to emphasize that it is topologically equivalent to a 2-torus. We note that $v_1^{xy} = -v_1^{yx}$ by the antisymmetry of the Berry curvature.

In 4D, the second Chern number (2CN) can also be calculated from the Berry curvature as [49–53]

$$\begin{aligned} v_2 &= \frac{1}{8\pi^2} \int_{\mathbb{T}^4} \Omega \wedge \Omega \in \mathbb{Z} \\ &= \frac{1}{32\pi^2} \int_{\mathbb{T}^4} \epsilon_{\alpha\beta\gamma\delta} \Omega^{\alpha\beta} \Omega^{\gamma\delta} d^4k \\ &= \frac{1}{4\pi^2} \int_{\mathbb{T}^4} \Omega^{xy} \Omega^{zw} + \Omega^{wx} \Omega^{zy} + \Omega^{zx} \Omega^{yw} d^4k, \end{aligned} \quad (5)$$

where $\epsilon_{\alpha\beta\gamma\delta}$ is the 4D Levi-Civita symbol and where we have written out the antisymmetric wedge product $\Omega \wedge \Omega$ in components for clarity. Now, the integral is taken over the first (magnetic) BZ in 4D, which we denote by \mathbb{T}^4 . As can be seen, the 2CN is a genuine four-dimensional invariant, which vanishes in lower-dimensional systems due to the 4D Levi-Civita symbol. Physically, the 2CN underlies the quantization of current in the 4D quantum Hall effect as will be introduced below.

We emphasize that in the above definitions we have explicitly restricted ourselves to a single isolated energy band. More generally, there may be an isolated set of bands among which there are degeneracies, such as for a spin- $\frac{1}{2}$ particle in the presence of time-reversal symmetry. Then, the components of the Berry curvature $\Omega^{\mu\nu}$ are themselves matrices, with indices running over the set of bands. The definitions of the 1CN (4) and the 2CN (5) can then be generalized to the integral of a matrix trace over Ω and $\Omega \wedge \Omega$, respectively. We note that while the 1CN so defined always vanishes without time-reversal symmetry breaking, there can be nonzero 2CNs also in a time-reversal-invariant system in the presence of an SU(2) gauge field; indeed, this was the context in which the 4D quantum Hall effect was originally proposed [49] and studied [50,54–58].

B. Semiclassical equations of motion

Having introduced the geometrical and topological properties of the underlying energy bands, we now review the motion of a wave packet prepared in a given Bloch band $\mathcal{E}(\mathbf{k})$ and subject to perturbing electromagnetic fields [42–46].

1. Introducing the perturbing fields

We consider particles of charge $-e$ moving in the presence of two families of fields: (1) the “intrinsic” fields $\mathcal{B}_{\mu\nu}$ generating the band structure $\mathcal{E}(\mathbf{k})$ under scrutiny, and (2) the perturbing “extrinsic” fields in response to which transport is analyzed. While the intrinsic fields $\mathcal{B}_{\mu\nu}$ do not need to be specified at this stage [their effects are entirely captured by the dispersion $\mathcal{E}(\mathbf{k})$ and Berry curvature $\Omega^{\mu\nu}(\mathbf{k})$ of the band], the perturbing fields will be taken in the form of a

weak electric field $\mathbf{E} = E_\mu \mathbf{e}^\mu$ and a weak magnetic field strength $B_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$, where $\mathbf{A} = A_\mu \mathbf{e}^\mu$ denotes the electromagnetic vector potential. We assume that these weak external electromagnetic fields are both time independent and spatially uniform. In the following, we also set Planck’s constant $\hbar = 1$ and the elementary charge $e = 1$, such that $\hbar/e^2 = 2\pi$. We note that the discussion that follows is general and also directly applies to neutral particles subject to synthetic gauge fields [12,13].

Finally, we point out that, while the field $B_{\mu\nu}$ should be weak enough for the following perturbative analysis to be valid, the partition of external magnetic fields into intrinsic ($\mathcal{B}_{\mu\nu}$) and perturbing ($B_{\mu\nu}$) components is somewhat arbitrary; this aspect will be illustrated in Secs. III and IV.

2. Equations of motion

In a semiclassical approach, motion is assumed to be adiabatic with respect to a manifold of states, such that a wave packet can be constructed out of this manifold at all times. The wave packet is chosen to have a well-defined center of mass at position $\mathbf{r}_c = r_c^\mu \mathbf{e}_\mu$ and quasimomentum $\mathbf{k}^c = k_\mu^c \mathbf{e}^\mu$. The appropriate manifold to use for the wave-packet construction depends on the strength of the applied electromagnetic fields. To see this, the full quantum Hamiltonian, including the perturbing electromagnetic fields, can be expanded around \mathbf{r}_c as [38,42–46]

$$\hat{H} \approx \hat{H}_c + \hat{H}' + \hat{H}'' + \dots, \quad (6)$$

where \hat{H}_c is the full Hamiltonian evaluated at the center-of-mass position, and \hat{H}' (\hat{H}'') are first- (second-) order gradient corrections in the electromagnetic fields.

When the external fields are weak, the wave packet moves adiabatically with respect to the eigenstates of an isolated energy band $\mathcal{E}(\mathbf{k})$ of \hat{H}_c , corresponding to the original Bloch states $|u_k\rangle$ introduced above, up to a phase [42,43]. The resulting semiclassical equations of motion are then consistent up to first order in the perturbing electromagnetic fields; this is sufficient to capture the physics of the 2D quantum Hall effect for a filled band. To extend this validity up to second order, so that we may also capture the (nonlinear) 4D quantum Hall response, the wave packet should be constructed out of the perturbed states $|\tilde{u}_k\rangle = |u_k\rangle + |\tilde{u}'_k\rangle$ where $|\tilde{u}'_k\rangle$ are the first-order band-mixing corrections from \hat{H}' [44–46]. These perturbed states have geometrical properties encoded in modified Berry curvature components [44]

$$\begin{aligned} \tilde{\Omega}^{\mu\nu} &= \Omega^{\mu\nu} + \Omega_1^{\mu\nu}, \\ \Omega_1^{\mu\nu} &= \partial_{k_\mu} \mathcal{A}'_{k_\nu} - \partial_{k_\nu} \mathcal{A}'_{k_\mu}, \end{aligned} \quad (7)$$

where $\mathcal{A}'_{k_\mu} = i\langle u | \partial_{k_\mu} | u' \rangle + \text{c.c.}$ is the first-order correction to the Berry connection. As derived in Ref. [44], the semiclassical equations valid up to second order in dimensions $d \geq 2$ are

$$\dot{\mathbf{r}}^\mu(\mathbf{k}) = \frac{\partial \tilde{\mathcal{E}}(\mathbf{k})}{\partial k_\mu} - \dot{k}_\nu \tilde{\Omega}^{\mu\nu}(\mathbf{k}), \quad (8)$$

$$\dot{k}_\mu = -E_\mu - \dot{\mathbf{r}}^\nu B_{\mu\nu}, \quad (9)$$

where we have dropped the subscript c from the center-of-mass position and quasimomentum. Here, the second-order

wave-packet energy $\tilde{\mathcal{E}}(\mathbf{k})$ contains the unperturbed Bloch band $\mathcal{E}(\mathbf{k})$ plus corrections from the perturbing terms $\hat{H}' + \hat{H}''$ [44,46]. These equations can be combined repeatedly to give the mean velocity of the wave packet [15]

$$\dot{r}^\mu \approx \frac{\partial \tilde{\mathcal{E}}}{\partial k_\mu} + E_v \tilde{\Omega}^{\mu\nu} + \left(\frac{\partial \tilde{\mathcal{E}}}{\partial k_\gamma} + E_\delta \tilde{\Omega}^{\gamma\delta} + \frac{\partial \tilde{\mathcal{E}}}{\partial k_\alpha} B_{\delta\alpha} \tilde{\Omega}^{\gamma\delta} \right) B_{\nu\gamma} \tilde{\Omega}^{\mu\nu}, \quad (10)$$

where terms above second order in the perturbing electromagnetic fields are neglected. As can be seen, there are many terms in this expression; these will lead both to Bloch oscillations and unquantized anomalous Hall effects for a wave packet [44,46].

In order to reveal the topological quantum Hall effects, one has to compute the total current density j^μ associated with an entirely populated band. This can be obtained using the mean velocity in Eq. (10) and summing over all the states located in the band,

$$j^\mu = \frac{1}{L^d} \sum_{\mathbf{k}} \rho(\mathbf{k}) \dot{r}^\mu(\mathbf{k}), \quad (11)$$

where L^d is the volume of the system of dimension d , and where $\rho(\mathbf{k})$ is the distribution function of particles within the band. In general, converting the sum over momentum states in Eq. (11) into an integral over the BZ is subtle [38]: in the semiclassical limit, this operation can be performed through the so-called modified density of states $D(\mathbf{r}, \mathbf{k})$:

$$\frac{1}{L^d} \sum_{\mathbf{k}} \rho(\mathbf{k}) \rightarrow \int_{\mathbb{T}^d} d^d k D(\mathbf{r}, \mathbf{k}) \rho(\mathbf{k}), \quad (12)$$

where the integration is performed over the first (magnetic) Brillouin zone (\mathbb{T}^d). Notably, the explicit expression for the modified density of states relies on an interesting interplay between the perturbing magnetic field $B_{\mu\nu}$ and the Berry curvature of the band $\Omega^{\mu\nu}$.

C. Modified density of states

When all magnetic field effects are included intrinsically into the band structure, the phase-space density of states $D(\mathbf{r}, \mathbf{k})$ is a simple constant factor of $1/h^d = 1/(2\pi)^d$. This is also the case when all magnetic fields are treated extrinsically as perturbations acting on systems with zero Berry curvature. The fact that the density of states remains constant is guaranteed classically by Liouville's theorem, which states that the phase-space volume element is conserved under time evolution [39]. However, Liouville's theorem holds for the volume element $\Delta V = \Delta \mathbf{R} \Delta \mathbf{K}$ associated with the *canonical* position \mathbf{R} and momentum \mathbf{K} , while the semiclassical treatment above is for the *physical* position \mathbf{r} and momentum \mathbf{k} [39,40,59].

To see how canonical and physical variables are related, we consider three cases in turn. First, if all magnetic effects are treated extrinsically, i.e., are not incorporated into the band structure, and if the unperturbed band structure has a trivial geometry (vanishing Berry curvature), the particle moves in the presence of a magnetic vector potential but a trivial Berry connection. Then, the physical momentum is modified by the

minimal (or Peierls) substitution $\mathbf{k} = \mathbf{K} - \mathbf{A}(\mathbf{r})$, while the physical and canonical positions are equivalent [60].

Second, if all magnetic effects are treated intrinsically, i.e., are included in the band structure, the particle moves in the presence of a nontrivial Berry connection but no (additional) magnetic vector potential. In this case, the physical position is $\mathbf{r} = \mathbf{R} + \mathcal{A}(\mathbf{k})$, while the physical and canonical momenta are equivalent [38,61,62]. The symmetry between these substitutions can be understood as the Berry curvature acting like a magnetic field in momentum space [62–64], i.e., the Berry connection \mathcal{A} and magnetic vector potential \mathbf{A} are dual.

Third, if, as we consider here, some magnetic fields are treated intrinsically and some extrinsically, the particle experiences both a Berry connection and a magnetic vector potential. Then, neither the physical position nor momentum remain equal to their canonical counterparts, and generalized Peierls substitutions are required [38,44,59,65]. The modified density of states $D(\mathbf{r}, \mathbf{k})$ can then be understood as the usual phase-space density of states $1/(2\pi)^d$ multiplied by the Jacobian of the transformation from the canonical to physical variables [40,59,65,66]. For dimensions up to $d = 4$, the modified density of states, valid to second order in the external fields, is given by [15]

$$D(\mathbf{r}, \mathbf{k}) = \frac{1}{(2\pi)^d} \left[1 + \frac{1}{2} B_{\mu\nu} \tilde{\Omega}^{\mu\nu} + \frac{1}{64} (\varepsilon^{\alpha\beta\gamma\delta} B_{\alpha\beta} B_{\gamma\delta}) \times (\varepsilon_{\mu\nu\lambda\rho} \Omega^{\mu\nu} \Omega^{\lambda\rho}) \right], \quad (13)$$

where, thanks to the Levi-Civita symbol, the last term vanishes in fewer than four dimensions [39,40,59,65]. Since our analysis is restricted to second order, the last term in Eq. (13) only involves the zeroth-order components of the Berry curvature $\Omega^{\mu\nu}$. In contrast, the first correction to the density of states, which is linear in the perturbing magnetic field $B_{\mu\nu}$, involves the first-order corrections to the curvature through $\tilde{\Omega}^{\mu\nu}$. Importantly, the correction $\Omega_1^{\mu\nu}$, introduced in Eq. (7), necessarily vanishes upon integration over the BZ as it is the curl of \mathcal{A}_{k_μ}' , which is gauge invariant and periodic in the BZ [44]. Hence, one can safely substitute $\tilde{\Omega}^{\mu\nu} \rightarrow \Omega^{\mu\nu}$ in Eq. (13) whenever one considers a uniformly filled band.

D. Semiclassical current density

The modified density of states (13) can be combined with the mean velocity (10) to evaluate the semiclassical current density of a band filled with a given distribution of particles $\rho(\mathbf{k})$. In standard quantum Hall systems, one typically considers an isolated band that is completely filled with spinless fermions, in which case $\rho(\mathbf{k}) = 1$. The more general uniformly populated-band situation [i.e., $\rho(\mathbf{k}) = \rho$] is also particularly relevant to cold-atom and photonics experiments (see Secs. II E 1 and II E 2). However, as the current density j^μ for this configuration can be simply obtained from the completely filled-band result through the substitution $j^\mu(\rho) \rightarrow \rho j^\mu(\rho = 1)$ [see Eq. (11)], we focus without loss of generality on the completely filled-band case.

Then, the semiclassical current density is

$$j^\mu \approx \int_{\mathbb{T}^d} \frac{d^d k}{(2\pi)^d} \left[E_v \tilde{\Omega}^{\mu\nu} + E_\delta \Omega^{\gamma\delta} B_{\nu\gamma} \Omega^{\mu\nu} + \frac{1}{2} E_v \Omega^{\mu\nu} B_{\delta\gamma} \Omega^{\delta\gamma} \right. \\ \left. + \frac{\partial \tilde{\mathcal{E}}}{\partial k_\mu} + \frac{\partial \tilde{\mathcal{E}}}{\partial k_\gamma} B_{\nu\gamma} \tilde{\Omega}^{\mu\nu} + \frac{1}{2} \frac{\partial \tilde{\mathcal{E}}}{\partial k_\mu} B_{\gamma\nu} \tilde{\Omega}^{\gamma\nu} \right. \\ \left. + \left(\frac{\partial \mathcal{E}}{\partial k_\alpha} B_{\delta\alpha} \Omega^{\gamma\delta} + \frac{1}{2} \frac{\partial \mathcal{E}}{\partial k_\gamma} B_{\delta\alpha} \Omega^{\delta\alpha} \right) B_{\nu\gamma} \Omega^{\mu\nu} \right. \\ \left. + \frac{1}{64} \frac{\partial \mathcal{E}}{\partial k_\mu} (\varepsilon^{\alpha\beta\gamma\delta} B_{\alpha\beta} B_{\gamma\delta}) (\varepsilon_{\xi\eta\lambda\rho} \Omega^{\xi\eta} \Omega^{\lambda\rho}) \right], \quad (14)$$

where whenever needed Ω replaces $\tilde{\Omega}$ and \mathcal{E} replaces $\tilde{\mathcal{E}}$ to keep only terms up to second order in the external fields. Expression (14) can be substantially simplified; first, terms on the third and fourth lines sum to zero due to the antisymmetry of the magnetic field strength and Berry curvature [15]. As this cancellation is by symmetry, it holds also for a band with arbitrary filling. Second, it can be shown that terms on the second line vanish upon integration over the BZ, using, as needed, the periodicity of the corrected energy dispersion $\tilde{\mathcal{E}}$ and the Bianchi identity for the antisymmetric Berry curvature [45]. This leaves only the first line which we rewrite as

$$j^\mu = E_v \int_{\mathbb{T}^d} \Omega^{\mu\nu} \frac{d^d k}{(2\pi)^d} \\ + \varepsilon^{\mu\alpha\beta\nu} \frac{1}{8} E_v B_{\alpha\beta} \int_{\mathbb{T}^d} \varepsilon_{\gamma\delta\lambda\rho} \Omega^{\gamma\delta} \Omega^{\lambda\rho} \frac{d^d k}{(2\pi)^d}, \quad (15)$$

where we have used that the Berry curvature correction $\Omega_1(\mathbf{k})$ vanishes upon integration over the BZ as commented above [44]. We note that this expression does not depend on the corrections to the energy and Berry curvature appearing in the full second-order semiclassical equations (8) and (9); these would, however, play a role in the dynamics of bands with nonuniform partial fillings, which we do not discuss further here.

It is important to note, additionally, that the inclusion of the modified density of states in calculating the current density (14) implies that the band is filled in the presence of the perturbing magnetic field $B_{\mu\nu}$. The experimental implications of this will be discussed further in Sec. V. We also point out that the perturbing magnetic field can potentially split the unperturbed Bloch band under scrutiny $\mathcal{E}(\mathbf{k})$ into a set of subbands, in which case the filling condition discussed above should apply to the set of subbands.

1. Linear vs nonlinear responses

The first term in Eq. (15) can be nonzero for a system with two or more dimensions; this is the famous quantum Hall current response which is linear in the applied electric field. This is independent of any magnetic perturbations and, in 2D, is directly proportional to the 1CN in Eq. (4). The second term, conversely, may be nonzero only for systems with four or more dimensions due to the Levi-Civita symbols; it corresponds to a nonlinear current response as it depends on both the applied electric field and the magnetic perturbing field. In 4D, the nonlinear term is directly proportional to the 2CN in Eq. (5). While we have truncated the perturbative treatment

at second order in the applied fields, at each higher order there can be an additional quantum Hall response for systems with increasing even dimensions (potentially involving higher-dimensional topological invariants).

2. External field partitionment and the current response

We observe that the current response j^μ in Eq. (15) should not depend on the arbitrary partition of external magnetic fields into intrinsic ($\mathcal{B}_{\mu\nu}$) and perturbing ($B_{\mu\nu}$) components (see discussion in Sec. II B 1). In particular, one is formally allowed to include all external magnetic fields into the intrinsic component, which in our framework is directly realized through the substitutions

$$\mathcal{B}_{\mu\nu} \rightarrow \mathcal{B}_{\mu\nu} + B_{\mu\nu}, \quad B_{\mu\nu} \rightarrow 0. \quad (16)$$

In this singular picture, the intrinsic Berry curvature now depends on the included weak field components $\Omega \rightarrow \Omega(B_{\alpha\beta})$, and the transport equation in Eq. (15) becomes

$$j^\mu = E_v \int_{\mathbb{T}^d} \Omega^{\mu\nu}(\mathbf{k}; B_{\alpha\beta}) \frac{d^d k}{(2\pi)^d}, \quad (17)$$

where the linear and nonlinear responses are now mingled. We note that the area of the magnetic Brillouin zone \mathbb{T}^d over which the integration is performed now also depends on the included magnetic perturbation $A_{\text{MBZ}} = A_{\text{MBZ}}(B_{\alpha\beta})$. Moreover, including the perturbation $B_{\alpha\beta}$ within the band structure potentially leads to a band splitting (see, e.g., Fig. 1), in which case a trace should be performed over the matrix-valued Berry curvature in Eq. (17).

In this picture, the relation between the current responses and the Chern numbers of the underlying band structure is obscured, and so is the quantization of (Hall) transport coefficients. This drawback is particularly well illustrated in the case of time-reversal-invariant 4D systems subject to time-reversal-breaking perturbations [49,50,54–58]: the connection between the 4D-QH current response and the 2CN of the bands is only made clear when treating all external U(1) magnetic fields as perturbing components (i.e., by working in a picture where the unperturbed system is time-reversal invariant).

These observations highlight the fact that, when all magnetic fields are present at all times, the external field partition, although arbitrary and formal, could be chosen based on theoretical convenience (see Secs. III and IV for illustrations). However, in some experimental systems, as we discuss in Sec. V, it is possible to ramp up magnetic perturbations after the initial preparation of the energy bands, in which case the partitionment of the field into intrinsic and extrinsic components follows naturally.

E. Center-of-mass observables

While current or voltage measurements have long been used to study the quantum Hall effect in solid-state systems, other measurements are easier to make in the analog quantum Hall systems which are currently of great experimental interest [4–9,11,31]. In particular, key experimental observables in ultracold atomic gases and photonics can be related to center-of-mass (c.m.) behavior, e.g., the center-of-mass motion of an atomic cloud (1). Such c.m. observables depend generally not just on the quantum Hall current (15), but also on the particle




FIG. 1. Energy spectrum $\mathcal{E}(k_x, k_y)$ of the 2D Harper-Hofstadter model [67] for (a) $\Phi = \frac{1}{3}$ and (b) $\Phi = \frac{101}{300}$. For $\Phi = \frac{1}{3}$, the spectrum contains three well-separated bands, where the 1CN of the lowest band is indicated. Increasing the flux to $\Phi = \frac{101}{300}$ splits the lowest band into 101 subbands, while preserving the sum of 1CNs as shown. Due to the large number of bands in (b) compared to (a) it can be useful to instead view a system with total flux $\Phi = \frac{101}{300}$ as a system with a strong flux $\Phi = \frac{1}{3}$ and a weak perturbing flux $\tilde{\Phi} = \frac{101}{300}$. (c) Hofstadter butterfly (energy as a function of the magnetic flux Φ) in the vicinity of the value $\Phi = \frac{1}{3}$. The lowest band at exactly $\Phi = \frac{1}{3}$ is shown to split into many subbands as the flux deviates from this ideal value, leading to complex (fractal) patterns in the \mathcal{E} - Φ plane.

density n , which is calculated semiclassically as

$$n = \int_{\mathbb{T}^d} d^d k D(\mathbf{r}, \mathbf{k}) \rho(\mathbf{k}). \quad (18)$$

As the modified density of states $D(\mathbf{r}, \mathbf{k})$ directly depends on the Berry curvature Ω [see Eq. (13)], the particle density of a filled band potentially contains information about the topology of the filled band (see Secs. III B 2 and IV B 3). This has important implications for experiments that extract Chern numbers from the measurements of c.m. observables.

1. Center-of-mass drift in cold atoms

In ultracold atoms, as introduced above, a natural observable is the center-of-mass motion of a cloud, which can be extracted from *in situ* density measurements as in the experiment of Ref. [6]. As shown in Eq. (1), the c.m. velocity for a cloud of atoms is given by $v_{\text{c.m.}} = \mathbf{j}/n$, and so is influenced by both the (quantum Hall) current response (15) and the particle density n (18), with important consequences discussed below.

We also point out that the above semiclassical analysis equally applies to systems of noninteracting fermions and bosons, as it only relies on the distribution function of particles within the band $\rho(\mathbf{k})$. Importantly, when a Bloch band is uniformly populated, $\rho(\mathbf{k}) = \rho$, the c.m. observables such as in Eq. (1) become independent of the band-filling factor ρ . This is because both the current density j^μ from Eq. (11) and the particle density n in Eq. (18) are directly proportional to the band-filling factor ρ .

The uniformly populated-band situation [$\rho(\mathbf{k}) = \rho$] is particularly relevant to cold-atom experiments, whenever the temperature T is large compared to the bandwidth W of the lowest-energy band, but small (or of the order) of the band gap $W \ll k_B T \ll \Delta$. This typically occurs when the lowest Bloch band displays a large flatness ratio $\Delta/W \gg 1$, as was recently demonstrated in the Munich experiment [6] through band mapping. In particular, this indicates that atomic transport experiments based on center-of-mass responses could be equally performed using (thermal) Bose or Fermi gases; this is in sharp contrast to measurements based on current densities, where the filling factor ρ should be independently measured.

2. Center-of-mass displacement in photonics

In photonics, an optical analog of the quantum Hall effect could be measured in the displacement of the center of mass of the photon steady state in a driven-dissipative system of coupled photonic cavities [30]. In such a system, a continuous-wave laser can pump light resonantly with a given isolated energy band, while the photon loss rate γ in the lattice is chosen such that $W \ll \gamma \ll \Delta$, where W is the bandwidth of the chosen energy band, and Δ is the band gap to the nearest energy band. In this regime, for sufficiently long times, the light reaches a nonequilibrium steady state in which the losses lead to an approximately uniform population of the single energy band. Hence, the center-of-mass displacement of the photon steady state for a square (hypercubic) lattice can be expressed as [16,30]

$$\langle \mathbf{r}_{\text{photon}} \rangle = \frac{\sum_{\mathbf{r}} \mathbf{r} |a_{\mathbf{r}}|^2}{\sum_{\mathbf{r}} |a_{\mathbf{r}}|^2} \approx \frac{\mathbf{j}}{\gamma n}, \quad (19)$$

where $a_{\mathbf{r}}$ is the expectation value of the photon field in the cavity labeled by \mathbf{r} , the d -dimensional position index, and where n is again the particle density of the filled band (18). This idea is not restricted to optics, but an analogous effect could also be observed in any driven-dissipative system of coupled-classical harmonic oscillators simulating quantum Hall physics [68].

Due to the similarities between Eqs. (1) and (19), our discussion in the following sections focuses on the center-of-mass velocity for ultracold atoms but has important experimental implications also in photonics or even in classical mechanical systems. However, we note that the displacement of the steady state in Eq. (19) is the leading-order term to which there are corrections that, for example, do not depend on the loss rate, and which should be included when modeling realistic experiments [16,30].

III. MAGNETIC PERTURBATIONS AND THE 2D QUANTUM HALL EFFECT

To emphasize the differences between the current density and center-of-mass response, we first discuss the important case of a two-dimensional quantum Hall system in the x - y plane. As defined in Eq. (4), the integral of the Berry curvature

over the whole 2D BZ gives the topological first Chern number ν_1^{xy} of the band. In the following, we focus on the 2D Harper-Hofstadter model, introduced below, as a concrete example of a system with energy bands that have nontrivial ICNs. However, we note that it is straightforward to extend our discussion to other topologically nontrivial 2D models as required.

A. 2D Harper-Hofstadter model

The 2D Harper-Hofstadter (HH) model is a seminal lattice model for studying the quantum Hall effect that was originally developed to describe a charged particle hopping on a 2D tight-binding square lattice in the presence of a uniform perpendicular magnetic field $\mathbf{B} = B\mathbf{e}_z$ [67]. The Hamiltonian is given by

$$\hat{H} = -J \sum_{\mathbf{r}} (c_{\mathbf{r}+a\mathbf{e}_x}^\dagger c_{\mathbf{r}} + e^{i2\pi\Phi x/a} c_{\mathbf{r}+a\mathbf{e}_y}^\dagger c_{\mathbf{r}} + \text{H.c.}), \quad (20)$$

where $c_{\mathbf{r}}^\dagger$ creates a fermion at lattice site $\mathbf{r}=(x,y)$, a is the lattice spacing, J is the hopping amplitude, and $\Phi = -a^2 B/2\pi$ is the magnetic flux per plaquette in units of the flux quantum. Here, we have chosen the magnetic vector potential in the Landau gauge such that the hopping along \hat{y} is modified by complex spatially dependent Peierls phase factors, while the hopping along \hat{x} is unaffected by them.

A rational magnetic flux per plaquette $\Phi = p/q$, where p and q are coprime integers, can be directly incorporated into magnetic Bloch states [51]. These are arranged into q energy bands described by a band structure $\mathcal{E}_n(k_x, k_y)$ in the so-called magnetic Brillouin zone [see Fig. 1(a)]. The magnetic Brillouin zone is defined by the magnetic translational symmetry of the HH model, and is a factor of q smaller than the original BZ, having an area $A_{\text{MBZ}} = (2\pi)^2/q a^2$. Due to the incorporated magnetic flux, the eigenstates in the bands have nontrivial Berry curvatures and nonzero ICNs. For suitable values of the flux (e.g., $\Phi = 1/q$), the energy spectrum has a nondegenerate lowest band which is well separated from other bands, to which the above semiclassical approach can be directly applied.

1. Physical realizations of the HH model

The HH model is a directly experimentally relevant model in materials, where it has been realized for electronic transport in graphene placed on boron nitride substrates [69,70]. Furthermore, the HH model has recently been generated in a wide variety of analog systems with neutral particles, where the spatially dependent complex (Peierls) phase factors in the tunneling matrix elements are carefully engineered. In ultracold gases, for example, the HH model has been realized by trapping atoms in a 2D optical lattice, and then combining a superlattice (or a Wannier-Stark ladder) along one direction with a resonant time modulation of the optical-lattice potential [4–6]. In an alternative approach, atoms were trapped in a one-dimensional optical lattice while two-photon Raman couplings induced transitions between different internal atomic states [8,9]. In this setup, the internal states could then be viewed as sites along an extra “synthetic” dimension [17,18], meaning that the atoms moved in an effective 2D lattice. By controlling the spatial dependence of the Raman wave vector, the experiments were able to implement complex hopping phase factors along this synthetic dimension, and hence realize HH physics.

For photons, the HH model has been experimentally simulated in an array of silicon ring resonators [11], where link resonators were used to introduce artificial Peierls phase factors. The concept of synthetic dimensions can also be extended to photonics [71]; it has been proposed to realize a 2D HH model in either a one-dimensional array of optical cavities where different angular momentum modes are coupled by spatial light modulators [72] or in a one-dimensional array of ring resonators, where the modes are coupled via an external time-dependent modulation [16]. The HH model has also been implemented by controlling intersite couplings in 2D arrays of circuit elements [73,74] and classical pendula [41]. Additionally, proposals exist for systems of periodically modulated classical coupled harmonic oscillators [68]. Hence, a full understanding of center-of-mass observables in this model can have important and direct applications in many current experiments.

B. 2D quantum Hall response

To study the quantum Hall response in two dimensions, we consider an electric field applied along the y direction $\mathbf{E} = E_y \mathbf{e}_y$ and a perturbing magnetic field $B_{xy} \ll B$. In experiments with neutral particles, such as those introduced above, these perturbing fields can also be imposed artificially. An electric field could correspond, for example, in ultracold gases to a linear gradient created either magnetically [4,5] or optically [6]. In a 2D array of coupled photonic cavities, it can be generated by a spatial gradient in the cavity size or temperature; see also Ref. [16] for synthetic electric fields acting along synthetic dimensions.

The weak magnetic field B_{xy} may also arise naturally in experiments whenever the realized flux $\Phi_{\text{expt}} = \Phi + \tilde{\Phi}$ slightly deviates from the desired (rational) value $\Phi = p/q$. In the following, we write the perturbing flux as

$$\tilde{\Phi} = -a^2 B_{xy}/2\pi. \quad (21)$$

In recent cold-atom experiments, for example, the flux was estimated to be $\Phi_{\text{expt}} \approx (1/4) \times 0.73(5)$ in [4], $\Phi_{\text{expt}} \approx 0.185$ in [8], $\Phi_{\text{expt}} \approx 4/3$ in [9], and $\Phi_{\text{expt}} \approx 1/4$ in [6]. In photonic lattices, the flux realized in Ref. [11] was $\Phi_{\text{expt}} \approx 0.15$, although there was also an additional random variation in the flux over the lattice.

1. Current density

Under the perturbing fields \mathbf{E} and B_{xy} , the current density of the filled lowest band from Eq. (15) simply leads to Eq. (2), written out again here, now with our choice of units $e = \hbar = 1$:

$$j^x = \frac{E_y}{2\pi} \nu_1^{xy}. \quad (22)$$

This is the 2D quantum Hall current response [3], for which a weak perturbing magnetic field has no direct effect.

Let us comment further on why it may be convenient to separate the flux penetrating a 2D lattice into strong and weak components, i.e., $\Phi_{\text{tot}} = \Phi + \tilde{\Phi}$. Indeed, imagine that a synthetic flux $\Phi_{\text{expt}} = 101/300 \approx 0.34$ has been realized; the corresponding energy spectrum splits into three isolated sets of bands, each set being associated with numerous extremely flat subbands as shown in Figs. 1(b) and 1(c) (e.g., the lowest




FIG. 2. (a) The Hofstadter butterfly: energy spectrum as a function of the magnetic flux Φ . The main gaps are labeled by two integer (t_r, s_r) , which satisfy the Diophantine equation: $r = qs_r - pt_r$, where the integer r denotes the r th gap and where $\Phi = p/q$. The integer t_r is given by the sum of Chern numbers associated with the bands below the r th gap, and for the HH model was shown to satisfy $-q/2 < t_r < q/2$ [51]. Focusing on the first gap ($r = 1$), this integer is simply given by the Chern number of the lowest band $t_1 = v_1^{xy}$ [see Eq. (26)]. (b) The Wannier diagram associated with the main gaps of the butterfly [78]. In the r th gap, the reduced particle density satisfies the equation $n/n_0 = s_r - \Phi t_r$, where the integers (t_r, s_r) satisfy the aforementioned Diophantine equation.

set is constituted of 101 subbands). The same system can be seen as a lattice pierced by a main flux $\Phi = \frac{1}{3}$, leading to three nondegenerate isolated bands [Fig. 1(a)], which is then slightly perturbed by a very weak flux $\tilde{\Phi} = \frac{1}{300}$. This latter picture, which involves a small set of nondegenerate bands, significantly simplifies the analysis of the Hall current, which remains immune to weak perturbing fields [see Eq. (22)].

2. Semiclassical particle density and the Diophantine equation

Unlike the current response, the density of particles in a filled band is highly sensitive to perturbing magnetic fields. Semiclassically, this can be seen from Eq. (18) calculated for a 2D system

$$n = \int_{\mathbb{T}^2} \frac{d^2k}{(2\pi)^2} (1 + B_{xy} \Omega^{xy}) = \frac{A_{MBZ}}{(2\pi)^2} + \frac{B_{xy}}{2\pi} v_1^{xy}, \quad (23)$$

where we considered a filled lowest band, and where we used that first-order corrections to the Berry curvature Ω_1 vanish upon integration [see Eq. (13) and discussion below]. The particle density therefore varies smoothly with a weak magnetic perturbing field in proportion to the 1CN of the filled band. This is as expected from the Streda-Widom formula [75–77], which relates the Hall conductance to the variation of the particle density with respect to the magnetic field at fixed temperature and chemical potential.

As a concrete example of this physics, we consider the HH model introduced above, for which $A_{MBZ} = (2\pi)^2/qa^2$. Introducing the normalized particle density, we find [Eq. (23)]

$$\frac{n}{n_0} = \frac{1}{q} - \tilde{\Phi} v_1^{xy}, \quad (24)$$

where $n_0 = 1/a^2$ is the inverse of the unit-cell area and $\tilde{\Phi}$ is the perturbing magnetic flux defined in Eq. (21). This can be understood as the semiclassical derivation of the Wannier diagram [78]: as illustrated in Fig. 2, each gap in the energy spectrum of the HH model (the so-called Hofstadter butterfly [67]) can be described by a straight line on a plot of density versus applied magnetic flux [78]. The slope of each line

is given by minus the sum of first Chern numbers of the bands lying below the gap [79,80]. For the lowest gap, i.e., considering the lowest band, this line is simply given by [69,80]

$$\frac{n}{n_0} = s - \Phi_{\text{tot}} v_1^{xy}, \quad (25)$$

where $\Phi_{\text{tot}} = \Phi + \tilde{\Phi}$ is the total magnetic flux per plaquette, and s is an integer. For rational flux $\Phi_{\text{tot}} = \Phi = p/q$, the normalized particle density is simply $n/n_0 = 1/q$ as only one of q bands is filled; this leads to the well-known Diophantine equation for the HH model [3,78]

$$1 = qs - p v_1^{xy}. \quad (26)$$

Rearranging this as an equation for the integer s and substituting this back into Eq. (25) then recovers the semiclassical result of Eq. (24). While we have focused here on the HH model, we note that results such as Eqs. (25) and (26) can be derived relying only on magnetic translational symmetry in a two-dimensional periodic potential [79], while the semiclassical result (23) was derived for any system with a nondegenerate isolated energy band.

3. Center-of-mass observables

As introduced above, center-of-mass observables can be important for experiments in analog quantum Hall systems. Inputting the 2D current density (22) and particle density (23) into the c.m. velocity for an atomic cloud (1), we find

$$v_{\text{c.m.}}^x = \frac{j^x}{n} = \frac{E_y}{\frac{A_{MBZ}}{2\pi} + B_{xy} v_1^{xy}} v_1^{xy}. \quad (27)$$

Neglecting any perturbing magnetic field B_{xy} , the c.m. velocity is directly proportional to the quantum Hall current response up to a simple multiplicative factor [22]. In such a configuration, the first Chern number has recently been experimentally extracted from a measurement of the center-of-mass drift of an ultracold cloud of atoms [6]. However, when a perturbing magnetic field is present, the dependence on the first Chern

number in Eq. (27) is more involved. For a sufficiently weak additional magnetic field (i.e., $|2\pi B_{xy} v_1^{xy}|/A_{\text{MBZ}} \ll 1$), we can perform a Taylor expansion to write

$$v_{\text{c.m.}}^x \approx \frac{2\pi}{A_{\text{MBZ}}} E_y v_1^{xy} - \left(\frac{2\pi}{A_{\text{MBZ}}} \right)^2 E_y B_{xy} (v_1^{xy})^2. \quad (28)$$

As we see, even in 2D, an experiment could therefore measure a nonlinear response term in the center-of-mass drift, quantized in units of $(v_1^{xy})^2$. This second term can be isolated by performing differential measurements, reversing the sign of the perturbing magnetic field, as discussed in Example II below and shown in Fig. 3. We shall now detail two examples where this sensitivity to perturbing magnetic fields is distinctly manifested:

Example I. Let us first illustrate the result in Eq. (27) on a simple example. Consider a 2D lattice that is exactly pierced by a uniform flux $\Phi = \frac{1}{5}$. In this case, the lowest band of the spectrum is associated with a Chern number $v_1^{xy} = -1$, the area of the magnetic Brillouin zone is $A_{\text{MBZ}} = (2\pi/a)^2/5$, and the perturbing magnetic field is $B_{xy} = 0$. According to Eq. (27), the c.m. velocity is thus given by $v_{\text{c.m.}}^x = (-5a^2/2\pi)E_y$.

Now, one can reinterpret this system as being a 2D lattice pierced by an intrinsic uniform flux $\Phi = \frac{1}{4}$, which is perturbed by an extrinsic flux $\tilde{\Phi} = -\frac{1}{20}$. The total flux is then $\Phi_{\text{tot}} = \Phi + \tilde{\Phi} = \frac{1}{5}$, as defined above. The unperturbed spectrum, with flux $\Phi = \frac{1}{4}$, is still characterized by a lowest band with Chern number $v_1^{xy} = -1$, but now the area of the magnetic Brillouin zone is $A_{\text{MBZ}} = (2\pi/a)^2/4$ and the perturbing magnetic field is $B_{xy} = -2\pi\tilde{\Phi}/a^2 = \pi/10a^2$. One readily verifies that Eq. (27) yields $v_{\text{c.m.}}^x = (-5a^2/2\pi)E_y$, in agreement with the complementary picture above. Interestingly, this simple equivalence could not have been demonstrated without invoking the modified density of states in Eq. (23), and thus, its impact on c.m. observables.

Example II. Motivated by recent experiments in ultracold gases [4,6], we consider a 2D lattice pierced by a uniform flux $\Phi = \frac{1}{4}$ with an additional uniform uncertainty in the flux of $\tilde{\Phi} = \pm 10\% \times \Phi$. To validate the above semiclassical results, we have numerically simulated the c.m. displacement $x_{\text{c.m.}}(\pm B_{xy}; t)$ of an ultracold cloud as shown in Fig. 3. Numerically, the cloud is initially confined in the presence of all fluxes $\Phi_{\text{tot}} = \Phi + \tilde{\Phi}$ before the confinement is removed and the electric field is ramped up; see Appendix for details on the numerical method.

As can be seen in Fig. 3(a), we find excellent agreement between the c.m. trajectories from numerical simulations and semiclassical analytics [Eq. (28)], as shown here by dark-blue dots and light-blue solid lines, respectively. We note that there is a clear quantitative deviation between these trajectories and the semiclassical result when the perturbing flux is neglected, as indicated by the solid red line. In Fig. 3(b), we also verify that a differential measurement of the c.m. trajectories under positive and negative perturbing magnetic flux $x_{\text{c.m.}}(B_{xy}; t) - x_{\text{c.m.}}(-B_{xy}; t)$ could be used to extract an approximate $(v_1^{xy})_{\text{est}}^2$, as predicted by Eq. (28). We note that for these parameters the differential drift would be of the order of a lattice spacing over typical experimental times around $50\hbar/J$. As the c.m. velocity scales $\propto q^2 B_{xy}$, from the second term of Eq. (28), this effect




FIG. 3. (a) Center-of-mass trajectories $x_{\text{c.m.}}(t)$ after ramping up the electric field to $E_y = 0.2J/a$, in the presence of a strong flux $\Phi = \frac{1}{4}$ and a perturbing flux $\tilde{\Phi} = \pm 10\% \times \Phi$. The dark-blue dots are numerical simulations performed for a 200×200 HH lattice. The red dashed curve is the semiclassical analytical result in Eq. (28) with $v_1^{xy} = -1$ when the nonlinear response due to the perturbing magnetic flux is neglected (i.e., when disregarding the effects of the modified density of states). The light-blue solid curves are the corresponding analytical results when this response is also included. (b) The difference between the center-of-mass trajectories for positive and negative perturbing magnetic fluxes shown above in (a). The light-blue solid line is the analytical nonlinear response calculated from (28). The dark-blue dots are numerical simulations, from which we extract an approximate $(v_1^{xy})_{\text{est}}^2$ as indicated on the plot. The small deviation between the analytical and numerical results is due to approximation of Eq. (27) by the Taylor series expansion in Eq. (28). Corrections to this approximation scale with the dimensionless parameter $\delta = (2\pi |B_{xy} v_1^{xy}|)/A_{\text{MBZ}}$; for the parameters shown here $\delta = 0.10$. Residual deviations in the numerical results can be attributed to weak interband transitions and small inhomogeneities in the wave packet.

would of course be larger if the perturbing magnetic flux is increased and/or the unperturbed flux is reduced.

IV. 4D QUANTUM HALL RESPONSES AND MAGNETIC PERTURBATIONS

As a further example, we turn to a 4D quantum Hall system, for which magnetic perturbations enter both into the current and the particle density. A four-dimensional system is also richer than a two-dimensional one as the Berry curvature (3) can now have up to six independent components $\Omega^{\mu\nu}$: one for each possible 2D plane. To simplify the following discussion, we therefore first introduce a minimal topological lattice model, noting that to have a nonzero 2CN (5) it is sufficient to have only two nonzero Berry curvature components provided they are in “disconnected” planes. For instance, one can take the nonzero components to be $\Omega^{xz} = -\Omega^{zx}$ and $\Omega^{yw} = -\Omega^{wy}$, without loss of generality.

A. Minimal 4D topological model

As we have previously proposed in Refs. [15,16], a minimal 4D lattice model that has energy bands with nonzero 2CNs consists of two copies of the Harper-Hofstadter model [67] defined in the disconnected planes x - z and y - w . The corresponding tight-binding Hamiltonian is

$$\hat{H} = -J \sum_{\mathbf{r}} c_{\mathbf{r}+a\mathbf{e}_x}^\dagger c_{\mathbf{r}} + c_{\mathbf{r}+a\mathbf{e}_y}^\dagger c_{\mathbf{r}} + e^{i2\pi\Phi_1 x/a} c_{\mathbf{r}+a\mathbf{e}_z}^\dagger c_{\mathbf{r}} + e^{i2\pi\Phi_2 y/a} c_{\mathbf{r}+a\mathbf{e}_w}^\dagger c_{\mathbf{r}} + \text{H.c.}, \quad (29)$$

where $c_{\mathbf{r}}^\dagger$ creates a fermion at lattice site $\mathbf{r} = (x, y, z, w)$, and the x - z and y - w planes are penetrated by uniform magnetic fluxes $\Phi_{1,2}$, respectively. This model could be realized experimentally by exploiting the concept of synthetic dimensions [15,16,18], already introduced in Sec. III A. By combining a synthetic dimension with either a 3D optical lattice of ultracold atoms [15] or with a 3D coupled-cavity array for photons [16], experiments would be able to build and explore the physics of an effective 4D lattice. Standard techniques, such as those outlined in Sec. III, could then be used to generate the necessary complex Peierls’ phase factors for the x - z and y - w planes.

The energy spectrum of this 4D model is given by a Minkowski sum of the energy bands of the two constituent Harper-Hofstadter models [15,58], i.e.,

$$\mathcal{E}(\mathbf{k}) = \{E_1 + E_2 | E_1 \in \mathcal{E}_{xz}(k_x, k_z), E_2 \in \mathcal{E}_{yw}(k_y, k_w)\}.$$

In particular, we will focus on the lowest 4D band, which for appropriate choices of the uniform magnetic fluxes $\Phi_{1,2}$ is nondegenerate and well isolated from the higher bands. In this minimal lattice model, the eigenstates making up the 4D energy bands are also characterized by only two nonzero Berry curvature components

$$\Omega^{zx} = \Omega^{zx}(k_x, k_z) \neq 0, \quad \Omega^{yw} = \Omega^{yw}(k_y, k_w) \neq 0, \quad (30)$$

that only depend on the components of momentum along the considered plane. Consequently, the integral for the 2CN of the lowest band (5) can be performed to find $\nu_2 = \nu_1^{zx} \times \nu_1^{yw}$ [15,58], where we have introduced the 1CNs associated with the x - z and y - w planes, respectively. For instance, the 1CN

characterizing the x - z plane is defined as

$$\nu_1^{zx} = \frac{1}{2\pi} \int_{\mathbb{T}^2} \Omega^{zx}(k_x, k_z) dk_x dk_z = -\nu_1^{xz}. \quad (31)$$

B. 4D quantum Hall response

The semiclassical current density (15) for a filled lowest (nondegenerate) band in four dimensions is [15]

$$\mathbf{j}^\mu = E_\nu \frac{1}{(2\pi)^4} \int_{\mathbb{T}^4} \Omega^{\mu\nu} d^4k + \frac{\nu_2}{4\pi^2} \varepsilon^{\mu\alpha\beta\nu} E_\nu B_{\alpha\beta}, \quad (32)$$

where the second term now explicitly contains the second Chern number of the lowest band (5). In the following, we will choose with no loss of generality the perturbing (synthetic) electric field to be along the y direction $\mathbf{E} = E_y \mathbf{e}^y$. However, in 4D, there are various choices for the orientation of the perturbing magnetic field, which will lead to dramatic differences in center-of-mass observables, as we now discuss.

1. Perturbing magnetic flux through a 2D plane without Berry curvature

In the simplest case, all nonzero extrinsic perturbing field components $B_{\mu\nu}$ are set in planes for which there is no Berry curvature $\Omega^{\mu\nu}$ from the underlying lattice. For example, for the minimal model introduced above [Eqs. (29) and (30)], this could be $B_{zw} \neq 0$ since $\Omega^{zw} = 0$ by construction. For this choice of perturbing magnetic field, the modified density of states reduces to the standard form $D(\mathbf{r}, \mathbf{k}) = \frac{1}{(2\pi)^4}$, as can be seen from Eq. (13). Therefore, the density of particles for a filled lowest band is simply [Eq. (18)]

$$n = \frac{A_{\text{MBZ}}^{zx} A_{\text{MBZ}}^{yw}}{(2\pi)^4} = \frac{1}{V_{\text{cell}}}, \quad (33)$$

where $A_{\text{MBZ}}^{\mu\nu}$ is the area of the magnetic BZ in the μ - ν plane, and where V_{cell} is the four-dimensional magnetic unit-cell volume. For the lattice introduced in Eq. (29), with rational fluxes $\Phi_1 = p_1/q_1$ and $\Phi_2 = p_2/q_2$, we have the following expressions:

$$A_{\text{MBZ}}^{zx} = \frac{(2\pi)^2}{a^2 q_1}, \quad A_{\text{MBZ}}^{yw} = \frac{(2\pi)^2}{a^2 q_2}, \quad V_{\text{cell}} = q_1 q_2 a^4. \quad (34)$$

Importantly, from Eqs. (1) and (33), we obtain that the c.m. velocity is directly proportional to the current density up to a constant factor [15]

$$\mathbf{v}_{\text{c.m.}} = \frac{\mathbf{j}}{n} = \mathbf{j} V_{\text{cell}}. \quad (35)$$

Here, the current density \mathbf{j} is explicitly given by [Eq. (32)]

$$j^x = \frac{\nu_2}{4\pi^2} E_y B_{zw}, \quad (36)$$

$$j^w = E_y \frac{1}{(2\pi)^4} \int_{\mathbb{T}^4} \Omega^{wy} d^4k = -\frac{\nu_1^{yw} A_{\text{MBZ}}^{zx}}{(2\pi)^3} E_y, \quad (37)$$

$$j^y = j^z = 0, \quad (38)$$

where we have used that Ω^{wy} is a function only of the momenta in the y - w plane in order to perform the integral in Eq. (37). The current response in Eq. (36) is a nonlinear 4D QH response along the x direction, while that in Eq. (37) is similar to a 2D

QH effect taking place in the y - w plane. However, unlike the usual 2D QH response (22), the current j^w is reduced by a factor $1/a^2 q_1$, as it depends also on the area of the MBZ in the z - x plane [15].

Combining Eqs. (35) and (36), one finds that the center-of-mass displacement along the x direction is directly proportional to the 2CN:

$$x_{\text{c.m.}}(t) = v_2 (V_{\text{cell}} t / 4\pi^2) E_y B_{zx}. \quad (39)$$

Hence, in this configuration, measurements of center-of-mass observables [Eqs. (1) and (19)] in atomic or photonics systems can be used to directly extract the 2CN response, as proposed in Refs. [15,16]. The semiclassical predictions in Eqs. (36)–(39) have already been validated through numerical simulations in Ref. [15].

2. Perturbing magnetic flux through a 2D plane with Berry curvature

In this case, there is only one nonzero extrinsic perturbing field component $B_{\mu\nu}$, and this is in the same plane as a nonzero Berry curvature $\Omega^{\mu\nu}$ from the underlying lattice. For the minimal model proposed above [Eq. (30)], this would be the case, for example, when $B_{zx} \neq 0$ as $\Omega_{zx} \neq 0$. For such a configuration, the density of states is strongly modified [Eq. (13)], and the particle density for a filled band becomes [Eq. (18)]

$$\begin{aligned} n &= \int_{\mathbb{T}^4} \frac{d^4 k}{(2\pi)^4} (1 + B_{zx} \Omega^{zx}) \\ &= \frac{A_{\text{MBZ}}^{zx} A_{\text{MBZ}}^{yw}}{(2\pi)^4} + \frac{A_{\text{MBZ}}^{yw}}{(2\pi)^3} B_{zx} v_1^{zx}, \\ &= \frac{A_{\text{MBZ}}^{yw}}{(2\pi)^2} \left[\frac{A_{\text{MBZ}}^{zx}}{(2\pi)^2} + \frac{B_{zx}}{2\pi} v_1^{zx} \right], \end{aligned} \quad (40)$$

where we again used that first-order corrections to the Berry curvature vanish upon integration. Up to the overall factor $A_{\text{MBZ}}^{yw}/(2\pi)^2$, this is the same modification of the density as found in a 2D system [see Eq. (23)]; this was expected because the Berry curvature component $\Omega^{zx} = \Omega^{zx}(k_x, k_z)$ only depends on the momenta along the considered x - z 2D plane when performing the integral (see Ref. [15]).

In this case, the current density becomes [Eq. (32)]

$$\begin{aligned} j^x &= j^y = j^z = 0, \\ j^w &= -\frac{v_1^{yw} A_{\text{MBZ}}^{zx}}{(2\pi)^3} E_y - \frac{v_2}{4\pi^2} E_y B_{zx}, \end{aligned} \quad (41)$$

where now both the 2D-like quantum Hall effect (1CN response) and the 4D quantum Hall effect (2CN response) occur along the w direction. These responses could be separated by a differential current measurement, where the sign of the perturbing magnetic field is flipped. For instance, the 2CN can still be extracted from the differential current

$$\delta j^w = j^w(-B_{zx}) - j^w(B_{zx}) = \frac{v_2}{2\pi^2} E_y B_{zx}. \quad (42)$$

Surprisingly, if we now turn to the c.m. velocity, we find that the nonzero velocity component $v_{\text{c.m.}}^w$ simplifies to the

expression [Eqs. (1), (40), and (41)]

$$v_{\text{c.m.}}^w = \frac{j^w}{n} = -\frac{2\pi}{A_{\text{MBZ}}^{yw}} E_y v_1^{yw}. \quad (43)$$

Importantly, the c.m. velocity in Eq. (43) contains only the linear 1CN response; all effects from the perturbing magnetic field and 2CN have canceled out. Hence, in contrast to the current densities in Eqs. (41) and (42), the c.m. displacement cannot be exploited to extract the 2CN of the populated band. Similar to Examples I and II above, we emphasize these differences by presenting two examples:

Example III. Let us first illustrate Eqs. (41)–(43) on a simple example: a 4D lattice pierced by a uniform flux $\Phi_1 = \frac{1}{5}$ in the x - z plane and $\Phi_2 = \frac{1}{4}$ in the y - w plane. For this system, the lowest band of the spectrum has a second Chern number $v_2 = -1$ and first Chern numbers $v_1^{yw} = -1$ and $v_1^{zx} = 1$ in individual planes [15], while the magnetic Brillouin zone areas are $A_{\text{MBZ}}^{yw} = (2\pi/a)^2/4$ and $A_{\text{MBZ}}^{zx} = (2\pi/a)^2/5$. Let us now assume that there is no perturbing magnetic field $B_{zx} = 0$, in which case the current density from Eq. (41) is $j^w = (1/10\pi a^2) E_y$, and the c.m. velocity from Eq. (43) is $v_{\text{c.m.}}^w = (4a^2/\pi) E_y$.

As in Example I in Sec. III, we can reinterpret the flux through the x - z plane as a uniform flux $\Phi_1 = \frac{1}{4}$ perturbed by a weak flux $\tilde{\Phi} = -\frac{1}{20}$, which corresponds to $B_{zx} = -B_{xz} = 2\pi\tilde{\Phi}/a^2 = -\pi/10a^2$. The unperturbed spectrum of this system is still characterized by the properties detailed above, except that now $A_{\text{MBZ}}^{zx} = (2\pi/a)^2/4$. One then readily verifies that Eq. (41) again leads to $j^w = (1/10\pi a^2) E_y$, while Eq. (43) leads trivially to the same c.m. velocity as it is independent of any perturbing magnetic field in the x - z plane.

Example IV. We have explored the results in Eqs. (41)–(43) further by performing numerical simulations on a small 4D lattice. Here, the strong fluxes are $\Phi_{1,2} = \frac{1}{4}$ in disconnected planes [15], and we take the perturbing flux to be $\tilde{\Phi} = a^2 B_{zx}/2\pi = \pm 10\% \times \Phi_1$ in the x - z plane. As for the 2D case discussed in Sec. III B 3, an electric field is ramped up to the final value $E_y = 0.2J/a$, and the time-evolving particle density is obtained. The resulting center-of-mass trajectories $w_{\text{c.m.}}(t)$ are shown in Fig. 4, which demonstrates good agreement with the prediction in Eq. (43); in particular, one finds that the trajectories show no significant dependence on the perturbing flux. From our numerical data, we use Eq. (43) to extract $(v_1^{yw})_{\text{est}} = -0.99$ for $\tilde{\Phi} > 0$ and $(v_1^{yw})_{\text{est}} = -1.00$ for $\tilde{\Phi} < 0$, in excellent agreement with the expected 1CN $v_1^{yw} = -1$ of the y - w plane. The figure also compares the numerical results with the wrong prediction $v_{\text{c.m.}}^w = j^w V_{\text{cell}}$, which corresponds to neglecting the effects of the modified density of states on the particle density (40). We have also numerically verified the expression for the differential current in Eq. (42), which indicates that such measurements could equally be exploited to give an approximate value for the 2CN of the band.

Summarizing the results of Secs. IV B 1 and IV B 2: Even though a 4D-QH current response appears for both of the above choices of perturbing magnetic fields [Eqs. (36) and (42)], only the first of these is appropriate for extracting the 2CN from center-of-mass observables. This result highlights the importance of evaluating the modified density of states when considering density-dependent observables [Eqs. (1), (13), and (18)].




FIG. 4. Center-of-mass trajectories $w_{\text{c.m.}}(t)$ after ramping up the electric field to $E_y = 0.2J/a$, in the presence of strong fluxes $\Phi_{1,2} = \frac{1}{4}$ in disconnected x - z and y - w planes and a perturbing flux $\tilde{\Phi} = a^2 B_{zx}/2\pi = \pm 10\% \times \Phi_1$ in the x - z plane. The filled dark-blue dots (resp. empty dots) are numerical simulations performed for a small $(4 \times 41 \times 4 \times 41)$ 4D lattice with positive (resp. negative) perturbing flux $\tilde{\Phi}$. The light-blue solid curve is the semiclassical analytical result in Eq. (43). The two red dashed curves correspond to the wrong prediction $v_{\text{c.m.}}^w = j^w V_{\text{cell}}$ combined with Eq. (41), which corresponds to neglecting the effects of the modified density of states on the particle density (40). Combining the numerical data together with Eq. (43) allows one to extract approximate values for the first Chern number associated with the y - w plane: we find $(v_1^{yw})_{\text{est}} = -0.99$ for $\tilde{\Phi} > 0$ and $(v_1^{yw})_{\text{est}} = -1.00$ for $\tilde{\Phi} < 0$.

3. Perturbing magnetic fluxes through two 2D planes with Berry curvature

As a final example, we consider perturbing fields in two planes with nonzero Berry curvature, e.g., B_{zx} and B_{yw} for the above model [Eq. (30)]. In this case, the particle density for a filled band is strongly modified as [cf. Eq. (18)]

$$\begin{aligned} n &= \int_{\mathbb{T}^4} \frac{d^4 k}{(2\pi)^4} (1 + B_{zx} \Omega^{zx}) (1 + B_{yw} \Omega^{yw}) \\ &= \frac{A_{\text{MBZ}}^{zx} A_{\text{MBZ}}^{yw}}{(2\pi)^4} + \frac{A_{\text{MBZ}}^{yw} B_{zx} v_1^{zx}}{(2\pi)^3} \\ &\quad + \frac{A_{\text{MBZ}}^{zx} B_{yw} v_1^{yw}}{(2\pi)^3} + \frac{B_{zx} B_{yw} v_2}{(2\pi)^2} \end{aligned} \quad (44)$$

up to second order in the perturbing fields, where we used that each Berry curvature component $\Omega^{\mu\nu}$ is only a function of momenta in the μ - ν plane. Interestingly, for this configuration of perturbing fields and in contrast to Eq. (40), the particle density now explicitly depends on the topological 2CN of the lowest filled band. This suggests an extension of the Streda-Widom formula relating the Hall conductance to density variations with respect to magnetic fields, from 2D [75–77] to 4D (see also Supplemental Material of Ref. [15]).

Let us now investigate the current and c.m. responses for this third configuration. Since the perturbing electric field is aligned along the y direction, $\mathbf{E} = E_y \mathbf{e}_y$, the component B_{yw}

does not affect the current density [see Eq. (32)]. Hence, the transport equations obtained in the previous case,

$$\begin{aligned} j^x &= j^y = j^z = 0, \\ j^w &= -\frac{v_1^{yw} A_{\text{MBZ}}^{zx}}{(2\pi)^3} E_y - \frac{v_2}{4\pi^2} E_y B_{zx}, \end{aligned} \quad (45)$$

are still valid for this configuration. However, using the modified density in Eq. (44), the center-of-mass velocity $v_{\text{c.m.}}^w = j^w/n$ is now

$$v_{\text{c.m.}}^w = \frac{-v_1^{yw} A_{\text{MBZ}}^{zx} E_y - 2\pi v_2 E_y B_{zx}}{\frac{A_{\text{MBZ}}^{zx} A_{\text{MBZ}}^{yw}}{2\pi} + A_{\text{MBZ}}^{yw} B_{zx} v_1^{zx} + A_{\text{MBZ}}^{zx} B_{yw} v_1^{yw} + 2\pi B_{zx} B_{yw} v_2},$$

which depends on all three topological invariants v_1^{zx} , v_1^{yw} , and v_2 . To extract the 4D quantum Hall response and 2CN from such center-of-mass motion would therefore require a multistep protocol to separate all the different effects.

4. Microscopic interpretation

Even though the current density displays a 4D quantum Hall response in all three configurations (Secs. IV B 1–IV B 3), there can be striking differences in the topological properties revealed by center-of-mass observables, such as the c.m. velocity of a cloud of ultracold atoms calculated above. We now discuss how this can be understood through a microscopic interpretation of the 4D quantum Hall effect for a filled band of spinless particles.

From our semiclassical derivation of the current density [Eqs. (11)–(15)], we can see that the nonlinear 2CN response arises from combinations of terms in the mean velocity (10) and in the modified density of states (13). In the first configuration (Sec. IV B 1), as noted above, the density of states (13) is not affected by the perturbing magnetic field. Instead, the 2CN response in Eq. (36) stems entirely from the mean velocity, where it appears from the interplay of the Lorentz force in Eq. (9) with the anomalous velocity in Eq. (8), that can itself be interpreted as the analog of a Lorentz force acting in momentum space. Hence, we refer to this as a “Lorentz-type” 2CN current response.

In the second configuration (Sec. IV B 2), on the other hand, the situation is very different: the perturbing magnetic field strongly modifies the density of states [see Eq. (40)], which then combines with the anomalous velocity in Eq. (10) to give the 2CN response. In this case, the 4D quantum Hall effect arises from the change in particle density of a band due to a perturbing extrinsic magnetic field, and not from the Lorentz force. We refer to this a “density-type” 2CN current response. Consequently, when we take into account the particle density for center-of-mass observables, this type of 4D quantum Hall effect vanishes from the latter [Eq. (43)].

In the third configuration (Sec. IV B 3), the 2CN current again arises from the interplay of a change in particle density, induced by an extrinsic field component B_{zx} , and the anomalous velocity in Eq. (10), i.e., a “density-type” 2CN response. However, in this case, the second extrinsic magnetic field component B_{yw} , while also further changing the particle density, does not couple to the anomalous velocity. When we calculate center-of-mass observables, both changes to the

particle density must be taken into account but as only the first leads to a 2CN response, the resulting c.m. observables have a nontrivial dependence on the topological invariants.

All these configurations could be further combined to yield other more complicated responses, which include both “Lorentz-type” and “density-type” microscopic mechanisms. Also, we have illustrated our discussion by assuming the Berry curvature 2-form has only two nonzero components, as for the minimal lattice model presented in Sec. IV A. However, the extension to other lattice models would be straightforward.

V. EXPERIMENTAL REMARKS

In this section, we briefly comment on various experimental aspects related to the detection of the quantized responses identified in this work. The platforms that we consider include bosonic and fermionic cold-atomic gases, electrons in solid-state materials, and driven-dissipative systems such as coupled photonic cavity arrays [11], where we note that the discussion of the latter can also be extended to coupled electric circuits [73] or mechanical systems [41].

A. Preparation of filled energy bands in the presence of perturbing magnetic fields

As highlighted in Sec. II, we have assumed that the particle density corresponds to that of a system where the lowest band is filled in the presence of any perturbing magnetic fields. In discussing the preparation of such bands, we shall distinguish between three physical scenarios: closed systems in which the particle number is fixed, open systems in which particles can be exchanged with a reservoir, and finally driven-dissipative systems in which an analog quantum Hall effect can be observed in the long-time nonequilibrium steady state [30].

In closed systems, our assumption of filled bands requires that the initial preparation of the system should be carried out in the presence of all synthetic magnetic fields. A standard example of a closed system is an ultracold atomic gas (which is typically not connected to a reservoir). In this case, the assumption of filled bands then means that all magnetic fields should be present during the adiabatic loading of ultracold atoms into the band structure. For ultracold fermions, the energy bands can be filled through fermionic statistics by ensuring that, after loading the atoms, the Fermi energy lies in the middle of an energy band gap. If instead the perturbing magnetic field were to be ramped up after the atoms were loaded, the Fermi level would end up within an energy band, leading to unquantized anomalous Hall effects [38,44]. For thermal Bose gases, an energy band can be uniformly filled when the temperature is large compared with the combined bandwidth of the bands to be populated, as introduced also in Sec. II E 1.

In open systems, when particles can be exchanged with a reservoir, perturbing magnetic fields can be turned on during the experiment. However, to ensure that the bands are filled in the presence of all magnetic fields, the rate at which these perturbations are ramped up should be slow compared to the time scale over which particles are exchanged with the reservoirs to ensure equilibration of the particle density. Open systems with reservoirs include solid-state materials, in which electrons are

exchanged with the connecting leads, but also cold-atom setups involving constriction potentials (see Ref. [81]). Moreover, we point out that harmonically trapped ultracold fermionic gases potentially present incompressible density plateaus (in a local-density-approximation picture, these correspond to regions with filled bands) surrounded by compressible regions, which can act as an internal reservoir within the cloud [19]. Note, however, that such inhomogeneities in the density can also complicate the motion of such a cloud, making it difficult to cleanly observe a quantum Hall response.

Finally, in driven-dissipative systems, such as coupled photonic cavity arrays, we assume the system is coherently and continuously pumped with sufficient losses to cover the energy bands. Provided that measurements always take place on a sufficiently long time after any variation of the perturbing fields, the photon distribution is in the steady state and depends only on the current values of these applied perturbations [30]. In this case, therefore, perturbing magnetic fields do not need to be ramped up but can be switched on suddenly during the experiment.

B. Switching on of the perturbing electric field and the validity of the semiclassical approach

After the filled energy bands have been prepared, the electric field can be turned on to measure the quantum Hall response. Again, we can distinguish between the different physical scenarios introduced above.

On the one hand, for closed systems and open systems with reservoirs, the electric field should be ramped up adiabatically such that the speed of the ramp is sufficiently slow and that the final value of the fields is sufficiently small compared to the energy band gap such that diabatic interband transitions can be neglected. If these conditions are not fulfilled, then there can be additional contributions to the transport from excited bands, and the current density or c.m. responses will in general no longer be quantized in terms of topological invariants [38,44]. For systems with reservoirs, we also require that the time scale over which the electric field is ramped up is fast compared with the rate with which particles are exchanged, otherwise the equilibration of the system will suppress the quantum Hall response. On the other hand, for driven-dissipative systems, the perturbing electric field can be switched on suddenly, provided that afterwards we wait a sufficiently long time for the steady state to be reached as discussed above.

Finally, we emphasize that the semiclassical approach presented in this work can only capture dynamics after any external fields are fully switched on as we have assumed throughout this work that all fields are time independent. We note that the agreement between the semiclassics and the dynamics after an adiabatic ramping of the electric field is illustrated in Figs. 3 and 4.

C. Density measurements and the Streda-Widom formula

While we have focused in this paper on center-of-mass transport, the dependence of the density on perturbing magnetic fields can itself be used as an experimental tool to measure topological Chern numbers of an energy band, as indicated by the Streda-Widom formula. In ultracold atomic gases, this

was first proposed in Ref. [19], where it was demonstrated numerically that the 2D Hall conductance could be extracted by comparing the real-space density profile of a cloud for two different values of the total synthetic magnetic field. This protocol could be extended to a 4D system by measuring the density profile of a cloud, for example, for different values of the magnetic fields of the minimal model introduced in Sec. IV A [see Eq. (44)]. As one of the four dimensions is now synthetic, the real-space density imaging should be extended with an optical Stern-Gerlach measurement to also determine the distribution of atoms in the different internal states [8,9].

VI. CONCLUSIONS

We have discussed how center-of-mass responses can be used to measure Chern numbers and so to directly probe the topology of energy bands. Center-of-mass observables depend not only on the quantum Hall current density, but also on the particle density and so can have a more involved dependence on topological invariants than previously considered. In particular, the particle density is itself sensitive to band topology in the presence of extrinsic magnetic perturbations, enriching the quantum Hall physics that may be explored in ultracold atomic gases and photonic systems.

In the 2D quantum Hall effect, c.m. observables can depend nonlinearly on the first Chern number of a filled energy band, in striking contrast to the linear dependence expected for electrical conductivity measurements. Such effects may already be observable experimentally, as there are inherent uncertainties in the precise value of magnetic flux imposed [4–9,11]. Additionally, new experiments with larger magnetic perturbations could also be engineered straightforwardly to directly probe the physics discussed here.

Finally, as a further example of these effects, we have considered recent proposals for the realization of the 4D quantum Hall effect [15,16] in realistic experimental systems. As we have seen, the particular configuration of perturbing fields chosen can have a dramatic impact, even leading to a cancellation of all 4D QH effects in center-of-mass observables, despite the persistence of a clear signature in the current density. A clear understanding of the differences between c.m. observables and current density measurements is therefore crucial to the proper design of future experiments.

ACKNOWLEDGMENTS

The authors thank M. Aidelsburger, I. Bloch, J. Dalibard, M. Dalmonte, A. Dauphin, M. Lohse, S. Nascimbene, C. Schweizer, and D.-T. Tran for fruitful discussions. H.M.P., T.O., and I.C. are supported by the ERC through the QGBE grant, by the EU-FET Proactive grant AQuS, Project No. 640800, and by the Autonomous Province of Trento, partially through the project “On silicon chip quantum optics for quantum computing and secure communications” (“SiQuero”). H.M.P. was also supported by the EC through the H2020 Marie Skłodowska-Curie Action, Individual Fellowship Grant No. 656093 “SynOptic”. O.Z. acknowledges the Swiss National Foundation for financial support. N.G. is financed by the FRS-FNRS Belgium and by the BSPO under PAI Project No. P7/18 DYGEST.

APPENDIX: NUMERICAL METHOD FOR THE DYNAMICS OF INCOHERENT WAVE PACKETS

In this Appendix, we present the numerical method used in Secs. III and IV to simulate the full-time dynamics, and hence confirm the semiclassical predictions.

The present method aims to simulate the dynamics of a noninteracting gas that is initially located in a region of space, and which uniformly populates an isolated Bloch band $\mathcal{E}(\mathbf{k})$ in an incoherent manner. This “incoherent wave-packet” configuration typically describes an atomic gas whose temperature T is large compared to the bandwidth W of the lowest-energy band $\mathcal{E}(\mathbf{k})$, but smaller than the band gap, $W \ll k_B T \ll \Delta$, as discussed in Sec. II E 1. This is different with respect to the numerical method used in Refs. [15,22,82], which describes a Bloch band completely filled with noninteracting fermions at zero temperature. As already pointed out in Sec. II E 1, these two different band-filling configurations lead to the same semiclassical equations of motion, up to a constant band-filling factor ρ in the current density; we have verified this equivalence numerically by comparing both methods on several examples.

Let us start by building a state from an incoherent superposition of all the Bloch states in a given band,

$$|\psi_{\text{super}}\rangle = \sum_{\mathcal{E}_\lambda \in \text{band}} |\phi_\lambda\rangle \exp[i\theta_\lambda], \quad (\text{A1})$$

where $|\phi_\lambda\rangle$ (resp. \mathcal{E}_λ) are the single-particle eigenstates (resp. eigenenergies) of the Hamiltonian \hat{H} in the absence of the perturbing electric field E_μ , and where θ_λ is a random phase associated with the state λ . As our lattice models are treated within the tight-binding approximation, we introduce the Wannier basis $\{|j\rangle\}$, which are states localized around the




FIG. 5. Time-evolving particle density $n(\mathbf{r},t)$ for the 2D system of Sec. III subject to a constant electric field $E_y = 0.2J/a$ along the y direction. The c.m. drift along the x direction is highlighted by the static circles. Here, the density has been averaged over $N=8$ realizations. Time is measured in units of \hbar/J .

lattice sites j . We then define a closed region in the lattice \mathcal{S} , made of a set of lattice sites, and we project the state ψ_{super} unto this small region

$$|\psi_{\text{packet}}\rangle = (1/\mathcal{N}) \sum_{j \in \mathcal{S}} \sum_{\lambda \in \text{band}} |j\rangle \langle j|\phi_{\lambda}\rangle \exp[i\theta_{\lambda}], \quad (\text{A2})$$

where \mathcal{N} is a normalization factor. While this projection procedure can also weakly excite particles to higher bands, we have verified that the resulting lowest-band population is between 99% and 99.9% in our numerics and so we can safely neglect the contribution from particles in higher bands. The “wave packet” in Eq. (A2) defines the initial state for our simulations. We then act on this state with the time-evolution

operator associated with the full Hamiltonian (including the electric field), from which we compute the time-evolving particle density $n(\mathbf{r}, t)$, extract the center-of-mass trajectory $\mathbf{r}_{\text{c.m.}}^{\mu}(t)$, and determine the corresponding velocity $v_{\text{c.m.}}^{\mu}$. We finally average the results over $N \approx 10$ realizations in which the phases θ_{λ} are randomly generated. The (mean) particle density n and center-of-mass velocities can then be combined to extract the current density, through Eq. (1). Finally, we point out that the band-filling factor is simply evaluated as $\rho = 1/N_{\text{states}}$, where N_{states} is the number of states in the band, i.e., the number of states λ included in the sum (A2) since the wave packet is normalized. An example of a 2D time-evolving cloud is illustrated in Fig. 5.

-
- [1] M. Z. Hasan and C. L. Kane, *Rev. Mod. Phys.* **82**, 3045 (2010).
 - [2] X.-L. Qi and S.-C. Zhang, *Rev. Mod. Phys.* **83**, 1057 (2011).
 - [3] D. J. Thouless, M. Kohmoto, M. P. Nightingale, and M. den Nijs, *Phys. Rev. Lett.* **49**, 405 (1982).
 - [4] M. Aidelsburger, M. Atala, M. Lohse, J. T. Barreiro, B. Paredes, and I. Bloch, *Phys. Rev. Lett.* **111**, 185301 (2013).
 - [5] H. Miyake, G. A. Siviloglou, C. J. Kennedy, W. C. Burton, and W. Ketterle, *Phys. Rev. Lett.* **111**, 185302 (2013).
 - [6] M. Aidelsburger, M. Lohse, C. Schweizer, M. Atala, J. T. Barreiro, S. Nascimbène, N. R. Cooper, I. Bloch, and N. Goldman, *Nat. Phys.* **11**, 162 (2015).
 - [7] G. Jotzu, M. Messer, R. Desbuquois, M. Lebrat, T. Uehlinger, D. Greif, and T. Esslinger, *Nature (London)* **515**, 237 (2014).
 - [8] M. Mancini, G. Pagano, G. Cappellini, L. Livi, M. Rider, J. Catani, C. Sias, P. Zoller, M. Inguscio, M. Dalmonte, and L. Fallani, *Science* **349**, 1510 (2015).
 - [9] B. K. Stuhl, H. I. Lu, L. M. Ayccock, D. Genkina, and I. B. Spielman, *Science* **349**, 1514 (2015).
 - [10] M. C. Rechtsman, J. M. Zeuner, Y. Plotnik, Y. Lumer, D. Podolsky, F. Dreisow, S. Nolte, M. Segev, and A. Szameit, *Nature (London)* **496**, 196 (2013).
 - [11] M. Hafezi, S. Mittal, J. Fan, A. Migdall, and J. Taylor, *Nat. Photonics* **7**, 1001 (2013).
 - [12] J. Dalibard, F. Gerbier, G. Juzeliūnas, and P. Öhberg, *Rev. Mod. Phys.* **83**, 1523 (2011).
 - [13] N. Goldman, G. Juzeliūnas, P. Öhberg, and I. B. Spielman, *Rep. Prog. Phys.* **77**, 126401 (2014).
 - [14] M. Hafezi, *Int. J. Mod. Phys. B* **28**, 1441002 (2014).
 - [15] H. M. Price, O. Zilberberg, T. Ozawa, I. Carusotto, and N. Goldman, *Phys. Rev. Lett.* **115**, 195303 (2015).
 - [16] T. Ozawa, H. M. Price, N. Goldman, O. Zilberberg, and I. Carusotto, *Phys. Rev. A* **93**, 043827 (2016).
 - [17] O. Boada, A. Celi, J. I. Latorre, and M. Lewenstein, *Phys. Rev. Lett.* **108**, 133001 (2012).
 - [18] A. Celi, P. Massignan, J. Ruseckas, N. Goldman, I. B. Spielman, G. Juzeliūnas, and M. Lewenstein, *Phys. Rev. Lett.* **112**, 043001 (2014).
 - [19] R. O. Umucalilar, H. Zhai, and M. Ö. Oktel, *Phys. Rev. Lett.* **100**, 070402 (2008).
 - [20] E. Alba, X. Fernandez-Gonzalvo, J. Mur-Petit, J. K. Pachos, and J. J. Garcia-Ripoll, *Phys. Rev. Lett.* **107**, 235301 (2011).
 - [21] H. M. Price and N. R. Cooper, *Phys. Rev. A* **85**, 033620 (2012).
 - [22] A. Dauphin and N. Goldman, *Phys. Rev. Lett.* **111**, 135302 (2013).
 - [23] Dmitry A. Abanin, Takuya Kitagawa, Immanuel Bloch, and Eugene Demler, *Phys. Rev. Lett.* **110**, 165304 (2013).
 - [24] Marcos Atala, Monika Aidelsburger, Julio T. Barreiro, Dmitry Abanin, Takuya Kitagawa, Eugene Demler, and Immanuel Bloch, *Nat. Phys.* **9**, 795 (2013).
 - [25] H. M. Price and N. R. Cooper, *Phys. Rev. Lett.* **111**, 220407 (2013).
 - [26] X.-J. Liu, K. T. Law, T. K. Ng, and P. A. Lee, *Phys. Rev. Lett.* **111**, 120402 (2013).
 - [27] P. Hauke, M. Lewenstein, and A. Eckardt, *Phys. Rev. Lett.* **113**, 045303 (2014).
 - [28] D.-L. Deng, S.-T. Wang, and L.-M. Duan, *Phys. Rev. A* **90**, 041601(R) (2014).
 - [29] Marcos Atala, Monika Aidelsburger, Michael Lohse, Julio T. Barreiro, Belén Paredes, and Immanuel Bloch, *Nat. Phys.* **10**, 588 (2014).
 - [30] T. Ozawa and I. Carusotto, *Phys. Rev. Lett.* **112**, 133902 (2014).
 - [31] M. Hafezi, *Phys. Rev. Lett.* **112**, 210405 (2014); S. Mittal *et al.*, *Nat. Photonics* **10**, 180 (2016).
 - [32] C.-E. Bardyn, S. D. Huber, and O. Zilberberg, *New J. Phys.* **16**, 123013 (2014).
 - [33] T. Li, L. Duca, M. Reitter, F. Grusdt, E. Demler, M. Endres, M. Schleier-Smith, I. Bloch, and U. Schneider, *Science* **352**, 1094 (2016).
 - [34] Fabian Grusdt, Norman Y. Yao, Dmitry A. Abanin, Michael Fleischhauer, and Eugene A. Demler, *arXiv:1512.03407*.
 - [35] Z. Wu *et al.*, *arXiv:1511.08170*.
 - [36] L. Duca, T. Li, M. Reitter, I. Bloch, M. Schleier-Smith, and U. Schneider, *Science* **347**, 288 (2015).
 - [37] N. Fläschner, B. S. Rem, M. Tarnowski, D. Vogel, D.-S. Lühmann, K. Sengstock, and C. Weitenberg, *Science* **352**, 1091 (2016).
 - [38] D. Xiao, M.-C. Chang, and Q. Niu, *Rev. Mod. Phys.* **82**, 1959 (2010).
 - [39] D. Xiao, J. Shi, and Q. Niu, *Phys. Rev. Lett.* **95**, 137204 (2005).
 - [40] C. Duval, Z. Horváth, P. A. Horváthy, L. Martina, and P. C. Stichel, *Mod. Phys. Lett. B* **20**, 373 (2006).
 - [41] R. Süssstrunk and S. Huber, *Science* **349**, 47 (2015).
 - [42] M.-C. Chang and Q. Niu, *Phys. Rev. B* **53**, 7010 (1996).
 - [43] M.-C. Chang and Q. Niu, *Phys. Rev. Lett.* **75**, 1348 (1995).

- [44] Y. Gao, S. A. Yang, and Q. Niu, *Phys. Rev. Lett.* **112**, 166601 (2014).
- [45] D. Xiao, J. Shi, D. P. Clougherty, and Q. Niu, *Phys. Rev. Lett.* **102**, 087602 (2009).
- [46] Y. Gao, S. A. Yang, and Q. Niu, *Phys. Rev. B* **91**, 214405 (2015).
- [47] M. Nakahara, *Geometry, Topology and Physics* (IOP Publishing, Bristol, 2003).
- [48] S. Ryu, A. P. Schnyder, A. Furusaki, and A. W. W. Ludwig, *New J. Phys.* **12**, 065010 (2010).
- [49] S.-C. Zhang and J. Hu, *Science* **294**, 823 (2001).
- [50] X.-L. Qi, T. L. Hughes, and S.-C. Zhang, *Phys. Rev. B* **78**, 195424 (2008).
- [51] J. E. Avron, L. Sadun, J. Segert, and B. Simon, *Phys. Rev. Lett.* **61**, 1329 (1988).
- [52] T. Fukui, T. Fujiwara, and Y. Hatsugai, *J. Phys. Soc. Jpn.* **77**, 123705 (2008).
- [53] B. Lian and S.-C. Zhang, [arXiv:1604.07459](https://arxiv.org/abs/1604.07459).
- [54] Y. Li, S.-C. Zhang, and C. Wu, *Phys. Rev. Lett.* **111**, 186803 (2013).
- [55] Y. Li, K. Intriligator, Y. Yu, and C. Wu, *Phys. Rev. B* **85**, 085132 (2012).
- [56] Y. Li and C. Wu, *Phys. Rev. Lett.* **110**, 216802 (2013).
- [57] J. M. Edge, J. Tworzydło, and C. W. J. Beenakker, *Phys. Rev. Lett.* **109**, 135701 (2012).
- [58] Y. E. Kraus, Z. Ringel, and O. Zeitler, *Phys. Rev. Lett.* **111**, 226401 (2013).
- [59] K. Yu. Bliokh, *Phys. Lett. A* **351**, 123 (2006).
- [60] L. D. Landau and L. M. Lifshitz, *Quantum Mechanics*, Vol. 3, 3rd ed. (Butterworth Heinemann, Oxford, 1958).
- [61] E. N. Adams and E. I. Blount, *J. Phys. Chem. Solids* **10**, 286 (1959).
- [62] N. Nagaosa, J. Sinova, S. Onoda, A. H. MacDonald, and N. P. Ong, *Rev. Mod. Phys.* **82**, 1539 (2010).
- [63] K. Y. Bliokh and Y. P. Bliokh, *Ann. Phys. (NY)* **319**, 13 (2005).
- [64] H. M. Price, T. Ozawa, and I. Carusotto, *Phys. Rev. Lett.* **113**, 190403 (2014).
- [65] P. Gosselin, F. Ménas, A. Bérard, and H. Mohrbach, *Europhys. Lett.* **76**, 651 (2006).
- [66] P. Gosselin, A. Bérard, and H. Mohrbach, *Eur. Phys. J. B* **58**, 137 (2007).
- [67] D. R. Hofstadter, *Phys. Rev. B* **14**, 2239 (1976).
- [68] G. Salerno, T. Ozawa, H. M. Price, and I. Carusotto, *Phys. Rev. B* **93**, 085105 (2016).
- [69] C. R. Dean *et al.*, *Nature (London)* **497**, 598 (2013).
- [70] L. A. Ponomarenko *et al.*, *Nature (London)* **497**, 594 (2013).
- [71] V. Peano, C. Brendel, M. Schmidt, and F. Marquardt, *Phys. Rev. X* **5**, 031011 (2015).
- [72] X.-W. Luo, X. Zhou, C.-F. Li, J.-S. Xu, G.-C. Guo, and Z.-W. Zhou, *Nat. Commun.* **6**, 7704 (2015).
- [73] J. Ningyuan, C. Owens, A. Sommer, D. Schuster, and J. Simon, *Phys. Rev. X* **5**, 021031 (2015).
- [74] V. V. Albert, L. I. Glazman, and L. Jiang, *Phys. Rev. Lett.* **114**, 173902 (2015).
- [75] A. Widom, *Phys. Lett. A* **90**, 474 (1982).
- [76] P. Streda, *J. Phys. C: Solid State Phys.* **15**, L717 (1982).
- [77] P. Streda and L. Smrcka, *J. Phys. C: Solid State Phys.* **16**, L895 (1983).
- [78] G. H. Wannier, *Phys. Status Solidi B* **88**, 757 (1978).
- [79] I. Dana, Y. Avron, and J. Zak, *J. Phys. C: Solid State Phys.* **18**, L679 (1985).
- [80] M. Kohmoto, *Phys. Rev. B* **39**, 11943 (1989).
- [81] J.-P. Brantut, J. Meineke, D. Stadler, S. Krinner, and T. Esslinger, *Science* **337**, 1069 (2012).
- [82] N. Goldman, J. Dalibard, A. Dauphin, F. Gerbier, M. Lewenstein, P. Zoller, and I. B. Spielman, *Proc. Natl. Acad. Sci. USA* **110**, 6736 (2013).