## UNIVERSITY<sup>OF</sup> BIRMINGHAM University of Birmingham Research at Birmingham

# Quantifying the beliefs of key players in the UK sheep industry on the efficacy of two treatments for footrot

Winter, Joanne R.; Green, Laura E.

DOI: 10.1016/j.tvjl.2018.07.009

*License:* Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version Publisher's PDF, also known as Version of record

*Citation for published version (Harvard):* Winter, JR & Green, LE 2018, 'Quantifying the beliefs of key players in the UK sheep industry on the efficacy of two treatments for footrot', *The Veterinary Journal*, vol. 239, pp. 15-20. https://doi.org/10.1016/j.tvjl.2018.07.009

Link to publication on Research at Birmingham portal

Publisher Rights Statement: Checked for eligibility 25/10/2018

https://doi.org/10.1016/j.tvjl.2018.07.009

#### **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

•Users may freely distribute the URL that is used to identify this publication.

•Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.

•User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?) •Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

#### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Contents lists available at ScienceDirect

## The Veterinary Journal

journal homepage: www.elsevier.com/locate/tvjl

#### **Original Article**

## Quantifying the beliefs of key players in the UK sheep industry on the efficacy of two treatments for footrot

ABSTRACT

### J.R. Winter<sup>1</sup>, L.E. Green<sup>2,\*</sup>

School of Life Sciences, University of Warwick, Gibbet Hill Road, Coventry CV4 7AL, UK

#### ARTICLE INFO

Keywords: Expert elicitation Evidence base Footrot treatment efficacy Key players Sheep Clinical trials have demonstrated that sheep with footrot treated with parenteral and topical antibiotics without foot trimming (treatment A), cure faster than sheep treated with foot trimming and topical antibiotics (treatment B). We investigated how key players in the UK sheep industry recommended treating footrot, and tested whether reviewing the evidence surrounding treatment of footrot changed their beliefs. Eight key players attended a workshop to investigate their current practices, and their perceived efficacy of treatments, using probabilistic elicitation.

At the start of the study, all participants recommended use of antibiotic injection but only four recommended not foot trimming feet with footrot. Initial beliefs in the difference in cure rate within five days of treatment ranged from 30 to 97% in favour of treatment A (true difference 60%); this heterogeneity reduced after reviewing the evidence. Participants who believed the cure rate differed by >60% over-estimated the cure rate of treatment A whilst participants who believed the difference was <60% over-estimated the efficacy of treatment B. During discussions, participants stated that parenteral antibiotics had always been recommended as a treatment for footrot but that the new research clarified when to use them. In contrast, it was highly novel to hear that foot trimming was detrimental to recovery, and key players and farmers are taking longer to accept this evidence. Three months after the workshop, two participants stated that they now placed greater emphasis on rapid individual antibiotic treatment of lame sheep and one was no longer recommending trimming feet.

© 2018 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND licenses (http://creativecommons.org/licenses/by-nc-nd/4.0/).

#### Introduction

There is robust evidence from a clinical trial that compared antibiotic injection and foot trimming in a factorial design, with a control of topical treatment, that there is a difference of 60% in the cure rate of sheep with footrot five days after treatment with parenteral and topical antibiotics with no foot trimming ('treatment A') vs. foot trimming and topical antibiotics ('treatment B'). Cure rates five days after treatment are approximately 70% with treatment A and 11% with treatment B (Kaler et al., 2010a). Treatment A also prevents loss in body condition and productivity; and is therefore economically beneficial (Wassink et al., 2010).

The proportion of farmers using treatment A increased between 2004 (Kaler and Green, 2009) and 2013 (Winter et al., 2015), but

the majority of farmers were still not following the recommended treatment protocol; 24% of farmers 'always' treated ewes lame with footrot with antibiotic injection and only 3% of farmers 'never' trimmed the feet of ewes with footrot (Winter et al., 2015). Increasing the proportion of farmers using treatment A would improve the health, welfare and productivity of sheep.

Probabilistic elicitation enables an individual's knowledge and uncertainty regarding a topic to be captured as a probability distribution (Garthwaite et al., 2005). Reviewing the evidence for the efficacy of treatments among veterinarians with and without expertise in sheep changed their beliefs nearer to the reported efficacy of treatment A (Higgins et al., 2013). However, it is not known whether, or how, key players, such as consultants, influential farmers, political bodies, or knowledge exchange providers, involved in technology transfer in the UK sheep industry, have incorporated the evidence from Kaler et al. (2010a) into their advice on managing footrot. The current study was designed to investigate how key players in the UK sheep industry currently recommend managing footrot, and to test whether a review of the evidence would change their beliefs.

https://doi.org/10.1016/j.tvjl.2018.07.009

1090-0233/© 2018 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/).





<sup>\*</sup> Corresponding author.

*E-mail addresses:* laura.green@warwick.ac.uk, l.e.green.1@bham.ac.uk (L.E. Green).

<sup>&</sup>lt;sup>1</sup> Current address: Department of Primary Care and Public Health Sciences, King's College London, SE1 1UL, UK.

<sup>&</sup>lt;sup>2</sup> University of Birmingham, Edgbaston, Birmingham, B15 2TT.

#### Materials and methods

#### Ethical approval and data collection

A population of sixteen key players was identified based on having a relevant degree and more than ten years experience earning some or all of their income from consultancy within the English sheep industry. They were invited to participate in a 1-day workshop, held at the University of Warwick on 17th–18th June 2014. The study was approved by the Biomedical and Scientific Research Ethics Committee at the University of Warwick on 13th June 2014 (study reference REGO-2014-795). All participants were provided with an information sheet detailing the purpose and methods of the study and that all data would be anonymised prior to publication. Informed signed consent was obtained from all participants. Travel expenses and an honorarium were paid.

Participants completed a questionnaire on their role in the sheep industry, how they currently advised farmers to treat footrot and whether this was different from their perceived 'reference standard' care, the number of farmers they advised per year and the media used in knowledge exchange.

A probabilistic elicitation exercise was then used to capture their beliefs regarding the difference in cure rates between two treatments for footrot (A and B) as a probability distribution. The method used was the same as that in Higgins et al. (Garthwaite et al., 2005; Johnson et al., 2010; Higgins et al., 2013). The elicitation was explained using a standard script (available on request) that included the clinical context and the question of interest. The clinical context was a ewe lame on a single foot with footrot, separation of the hoof horn that started at the interdigital skin and grey oozing pus with a distinctive foul smell, caused by Dichelobacter nodosus (Beveridge, 1941); no other foot conditions were complicating the clinical presentation. The outcome was cure from footrot defined as 'ewes had no observable limp and were not head-flicking', within five days of treatment (Kaler et al., 2010a). The question was: 'Which treatment is more effective at curing footrot within five days?' Treatment A was an intramuscular injection of long-acting oxytetracycline, correctly dosed for the sheep's weight, combined with topical oxytetracycline spray and no trimming of hoof horn. Treatment B was foot trimming by someone experienced and proficient at trimming, and application of oxytetracycline spray to the foot. Any other factors which might influence cure rates were consistent across treatments.

Probabilistic elicitation was done using the roulette ('chip and bins') method (Johnson et al., 2010). Participants were given six example distributions (Appendix: Supplementary Figs. 1–6), demonstrating how they could make a distribution of different beliefs and certainty in those beliefs. The central tendency of the probability distributions is the participant's 'best guess' and the range of the distribution is a measure of 'certainty'. Participants were given 20, five pence coins, each representing a probability of 0.05, total probability = 1, which they used to create a probability distribution indicating the weight of their belief for  $\theta_d$  (the difference in cure rate between treatments) by placing the coins on a blank laminated sheet laid out as in Appendix: Supplementary Fig. 1. The interpretation of the probability distribution was described to the participant by a trained assistant and the participant swere also asked their estimate of the actual cure rate for each treatment, and the upper and lower boundaries they believed for these values.

#### Evidence presentation and follow up

Participants then listened to a 30-minute presentation delivered by LEG summarising the current peer-reviewed evidence regarding treatment of footrot, now published in Green and Clifton (2018). Information sheets were then provided to each participant. After the presentation, participants were asked to consider their initial distribution, and to make a second distribution, whether or not their opinions had changed and to complete a second questionnaire. This was followed by a group discussion on how the evidence produced from recent research into footrot had changed the recommendations they gave to farmers. Three months after the study, participants were asked whether the workshop had changed their beliefs regarding how best to treat footrot, or the advice they gave farmers.

#### Data analysis

Data were entered into Microsoft Excel (2010) and data analysis was done using R (R Development Core Team, 2008). There were two unknown parameters,  $\theta_1$  the probability of cure within five days when treated with treatment A ( $\theta_1 \in [0,1]$ ), and  $\theta_2$  the probability of cure within five days when treated with treatment B ( $\theta_2 \in [0,1]$ ). Participants' beliefs were quantified by eliciting a probability distribution for the difference in cure rates,

#### $\theta_d = \theta_1 - \theta_2$ , where $\theta_d \in [-1,+1]$

because this was a clinically intuitive scale for the participants to use.

Normal distributions were fitted to the elicited data. The best fitting hyperparameters (mean and standard deviation) were selected using numerical optimisation (Nelder and Mead, 1965) to minimise the sum of the squared differences between the distributions of the observed cumulative probabilities and the fitted cumulative probabilities.

Participants' change in belief was calculated from the differences in the fitted hyperparameters. The fitted distributions were used to calculate 95% Bayesian credible intervals, which identified the 95% probability interval for difference in cure rates.

#### Results

#### Response rates and participant characteristics

Ten (63%) of the 16 people identified as the most influential agreed to participate in the study; two cancelled at short notice due to unexpected events. Participants typically had several roles in the sheep industry (Table 1), and advised several hundred farmers, in a mixture of one-to-one and group settings. They did this through advisory committees, one-to-one consultancy, farm open days, meetings and workshops, press articles, newsletters, and technical publications, and reports for pharmaceutical companies. Some delivered continuing professional development (CPD) to other consultants, two of these organised experts to deliver CPD but did not deliver advice themselves. One participant had affiliations with several breed societies.

#### Recommendations for treating sheep lame with footrot

The participants listed 13 pieces of advice they gave as 'current' or 'reference standard care' (Table 2). The only advice all eight participants gave was to use parenteral antibiotics and only four participants routinely recommended no foot trimming. Not trimming feet, together with vaccination, segregating sheep with footrot and culling repeatedly lame ewes were considered as reference standard care; not always possible, practical or economical. Lack of farmer motivation was reported as a reason why reference standard treatment may not occur in practice.

#### Elicited probability distributions and estimates of cure rates

Initially, participants believed treatment A cured 75–95%, and treatment B 10–40%, of sheep within five days (Fig. 1). Values for  $\theta_d$  ranged from 38 to 90%, with Bayesian credible intervals ranging from 30 to 97%, in favour of treatment A. The credible intervals of some participants did not overlap, indicating a wide range of beliefs.

The range of estimates for  $\theta_d$  narrowed considerably to 60–90% although the mean  $\theta_d$  change was negligible, 0.70 before to 0.69 after the evidence was presented. However, the mean absolute change in  $\theta_d$  was 11.2% with all except one participant moving their estimate towards the 'true' value of 0.6. The mean standard deviation of  $\theta_d$  decreased from 5.9 to 4.3, indicating that reviewing the evidence increased participants confidence in their beliefs (Fig. 1, Table 3), although there were still participants whose credible intervals did not overlap. The estimates of cure rates for each treatment also moved nearer to published evidence but were still overestimates (Table 4).

 Table 1

 The roles of each participant in the sheep industry.

Participant	Role	Farmed sheep
1	Knowledge transfer	Yes
2	Consultant	Yes
3	Vet and consultant	No
4	Consultant	No
5	Knowledge transfer	No
6	Consultant	Yes
7	Vet and consultant	Yes
8	Consultant and breeder	Yes

#### Table 2

The number of participants in our study who currently gave each piece of advice to all clients, and the number of participants who would consider each piece of advice to be part of 'reference standard' care.

Advice	Number of participants currently giving this advice to all clients	Number of participants who considered this advice part of 'reference standard' care
Antibiotic injection	8	8
Vaccinate with Footvax	5	6
Treat promptly	4	4
Foot spray	4	4
Do not trim feet	4	5
Segregate sheep with footrot	4	5
Cull repeatedly lame ewes	4	6
Mark ewes	3	3
Record ear tag number	2	2
Check cause of lameness	2	2
Re-treat sheep later if necessary	2	2
Inspect sheep after treating	1	1
Cull sheep with chronically infected or misshapen feet	1	1
Quarantine new sheep	2	4
Footbath new sheep	1	3
Check the feet of all new sheep	1	1
Footbath after gathering	1	2
Avoid gathering sheep	1	3
Inspect the feet of the entire flock	0	1
Avoid buying replacement ewes with poor foot conformation	0	1



**Fig. 1.** Fitted Gaussian probability density functions of eight participants, before and after a review of the evidence on treatments for footrot. Positive responses favoured 'injection and spray' over 'trim and spray'. Values for the fitted hyperparameters (mean and standard deviation) are shown in Table 3. The published literature currently supports a difference in cure rates of approximately 60%.

#### Participants' discussions on recommended treatment of footrot

Some participants indicated that their opinion on how to treat footrot had not changed as result of the evidence presented, most commonly because they were already aware, at least to some extent, of the evidence-base surrounding treatments A and B. Five participants said they were already giving the evidence in the presentation. Two indicated that the actual cure rates were different from what they had thought, but that this would not change their advice. The eighth participant said that whilst they were already highlighting the negative impact of foot trimming, they would stress this more in future. The group were asked how their beliefs on treatment of footrot had changed since 2003, when the first paper hypothesising that the recommended managements for treatment of footrot at that time were ineffective or incorrect (Wassink et al., 2003) was published, and then as further papers were published on the benefits of antibiotic treatment and the detrimental effects of foot trimming. The consensus was that whilst it was believed that antibiotics were effective in 2003, when to use them was not clear and the 2003 paper had clarified that immediate treatment of all cases rather than only treatment of chronic cases of footrot was best practice. The new, and most challenging information, was that foot trimming as a routine procedure (Wassink et al., 2003; Kaler and Green, 2009; Winter et al., 2015) and as a treatment

I I was a was an a store	for the	C++ a d	Causaian	man hahilitur	مسما نعبيها نسعه ال
Hyperparameters	for the	mea	Gaussian	DIODADIIIIV	distributions
i i y per parameter o	101 0110	meeee	ounoblai	probability	anoundationio

Participant	Before evidence review		After e	vidence revie	Change in fitted parameters $(\theta_{d(after)} - \theta_{d(before)})$			
	$\theta_d$ cure rate	SD	95% CI	$\theta_d$ cure rate	SD	95% CI	Mean	SD
1	90.0	3.8	83-97	62.5	4.6	53-72	-27.5	+0.9
2	76.9	3.5	70-84	77.5	3.6	70-85	+0.6	+0.1
3	90.0	3.8	83-97	90.0	3.8	83-97	0.0	0.0
4	68.8	9.6	50-88	62.5	3.9	55-70	-6.3	-5.6
5	58.9	6.2	47-71	72.1	6.8	59-85	+13.3	+0.6
6	81.7	6.6	69-95	67.5	2.4	63-72	-14.2	-4.2
7	38.1	4.1	30-46	60.0	1.2	58-62	+21.9	-2.9
8	57.5	9.8	38-77	63.1	7.8	48-78	+5.6	-2.0

 $\theta_d$ , difference in cure rates; SD, standard deviation; CI, credible interval.

#### Table 4

Elicited values for the cure rate by treatments A (inject and spray) and B (trim and spray) and the difference between the two estimated cure rates, before and after the review of the evidence, and the change in estimates after the review of the evidence.

Part		e evidence revie		After evidence review				Change in diff in cure rates			
	Treatme	nt A	Treatmer	nt B	Diff $(\theta_1 - \theta_2)$	Treatment A		Treatment B		Diff $(\theta_1 - \theta_2)$	
	Cure rate $\theta_1$	95% CI	Cure rate $\theta_2$	95% CI		$\theta_1$ Cure rate	95% CI	$\theta_2$ Cure rate	95% CI		$(\theta_1 - \theta_2)_{after} - (\theta_1 - \theta_2)_{before}$
1	95	80-100	40	30-45	55	70	60-75	10	5-15	60	+5
2	80	60-90	10	5-20	70	80	60-90	10	5-20	70	0
3	75	60-90	20	0-40	55	75	60-90	15	0-30	60	+5
4	75	40-90	10	0-50	65	70	60-80	10	0-20	60	-5
5	80	60-95	30	25-45	50	90	70-95	25	15-30	65	+15
6	90	60-100	10	0-30	80	75	60-100	10	0-30	65	-15
7	75	70-85	30	15-35	45	72	70-75	11	10-20	61	+16
8	93	85-100	35	20-40	58	90	70-99	35	20-50	55	-3

Part, participant;  $\theta_1$ , injection and spray;  $\theta_2$ , trim and spray; Diff, difference; CI, credible interval.

(Kaler et al., 2010a) was not beneficial. By 2014 (when the workshop was run and results from all these studies were presented) all key players believed this evidence, although two participants were less convinced that trimming did not have a role in treatment.

#### Follow up

Seven of the eight participants responded to the follow up questions. All stated that the event had not changed their beliefs; five reiterated that this was because their beliefs were already in



Fig. 2. Eight participants probability density functions fitted to the raw elicited data from each participant. Positive responses favoured 'injection and spray' over 'trim and spray'. Values for the fitted hyperparameters (mean and standard deviation) are shown in Table 3.

line with the evidence presented. Despite this, three participants had changed the advice they gave to their clients as a result of the event; two were placing greater emphasis on rapid individual antibiotic treatment and a third participant stated that they now placed greater emphasis on reducing foot trimming.

#### Discussion

The key findings were that all the key players believed that treatment A, parenteral and topical antibiotic treatment was superior to treatment B, foot trimming and topical antibiotic treatment. However, despite convergence in estimates of  $\theta_d$  after reviewing the evidence, participants continued to overestimate the difference in cure rates. The credible intervals of three participants (2, 3, 6) did not contain the actual value for  $\theta_d$ (Table 3) suggesting strongly held beliefs that did not alter, indeed two of these made negligible changes to their distributions after the evidence was presented (Fig. 2) and stated that they were already aware of the evidence. A study on the strength and variation of veterinarians' clinical beliefs regarding dry cow therapy found that some veterinarians were also very confident in their prior beliefs (Higgins et al., 2012) and that even results from a very large clinical trial would be insufficient to change those veterinarians' opinions.

Overestimates of  $\theta_d$  came from participants overestimating the value of  $\theta_1$ , whilst underestimates came from overestimating  $\theta_2$  (Table 4). Participants who overestimated  $\theta_d$  may have been the participants' way of demonstrating their strong belief in the superiority of antibiotic injection over foot trimming, or they may have misremembered the evidence, possibly because of conscious or unconscious motivational bias (Spetzler and Staelvonholstein, 1975) when influencing others. It is also possible that participants recalled cure rates 10 days after treatment, which were still about 60% difference but with an absolute recovery of approximately 90% for treatment A and 30% for treatment B (Kaler et al., 2010a). Inaccurate estimations might also have occurred if participants did not fully understand/engage with the task. For example, prior to the evidence review, participant one was 100% certain that the difference in cure rates was between 80-100% (Table 3, Fig. 2) despite believing the cure rate for treatment A was 95% and the cure rate for treatment B was 40%; which is a difference of only 55% (Table 4). This inconsistency was highlighted to the participant, but they did not change their estimates.

In 2012 there was a similar study to the current study where 11 veterinary practitioners were elicited on their beliefs of treatments for footrot (Higgins et al., 2013). Opinions on the difference in cure rates between treatments A and B were narrower for participants in the current study than in Higgins et al. (2013) and in contrast to Higgins et al. (2013), all participants in the current study believed that treatment A was superior to treatment B at the start of the study. These differences may be because the key players in our study were more actively involved in knowledge exchange than the veterinarians in Higgins et al. (2013), and consequently may have been more aware of current research. Additionally, promotion of the efficacy of treatment A continued from 2012 to 2014 through knowledge exchange and new publications (Winter et al., 2015), which may have led to wider awareness of its efficacy. Both studies may be limited in their generalisability due to the small number of participants and potential participation bias (i.e. individuals willing to participate in research may be more open to adopting new evidence). However, in the current study we aimed to investigate the beliefs of key players in the industry, only 16 individuals were identified, a small number of individuals with large influence. All were invited to participate in the study, and 50% finally did.

The group discussion highlighted that the detrimental effect of routine foot trimming was highly novel when first reported in 2003 (Wassink et al., 2003) because it had been a recommended management for footrot for many years (Morgan, 1987; Winter, 2004). The 2003 results were challenged initially (Abbott et al., 2003). Stakeholders in the UK started changing advice and practice as further evidence on the negative effects of routine foot trimming arose (Green et al., 2007; Kaler and Green, 2009; Kaler et al., 2010b; Winter et al., 2015). The participants commented that the belief in the beneficial effect of trimming feet with footrot was to expose the wound to air to 'kill' the anaerobic bacteria, some then reflected that this information could encourage severe trimming of feet with the expectation of 'getting in more air'. The one clinical trial where therapeutic foot trimming was tested in controlled conditions indicated that foot trimming diseased feet more than halved the recovery rate, both when sheep were treated with parenteral and topical antibiotic or just topical antibiotic (Kaler et al., 2010a). Recently, Winter et al. (2015) reported a lower prevalence of lameness in flocks where sheep with footrot were never foot trimmed. In addition, there is no published evidence that therapeutic foot trimming is beneficial, although many trials include foot trimming as a baseline across all treatment groups (this is described in detail in Kaler et al., 2010a). Avoiding trimming has not been adopted across the globe; for example, an article from Switzerland promoted foot trimming in the conclusions without testing this in the study (Locher et al., 2015).

There were few differences between the participants' current recommendations to farmers and their perceptions of reference standard care (Table 2). This was in contrast to veterinarians in Higgins et al. (2013) who tailored advice to farmers according to what they believed farmers would realistically act upon. Knowledge exchange providers are often not directly dependent on individual farmers for their livelihood and this independence might help in providing impartial advice. Group advice is also likely to be reference standard advice because it is not possible to vary advice for individual farmer circumstances. One might hypothesise that consultants are similar to first opinion veterinarians and vary advice, however, only a small proportion of sheep farmers use specialist consultants, the majority believing such input is not beneficial or economical (Kaler and Green, 2013). Consequently, farmers who use specialist consultants may be more motivated, proactive and willing to change. In addition, consultants are less reliant on local clients than veterinarians and so might offer more challenging advice; because their reputation is based on making improvements and they might prefer to lose clients not prepared to make changes, rather than compromise their reputation.

Finally, the follow up questionnaire results suggested that change in beliefs continued after the workshop because more participants were recommending avoiding foot trimming and rapid treatment than indicated at the event. It might be that the discussion with others at the workshop gave participants more confidence to use the evidence that they discussed at the workshop with others, or that they tried the treatments on their own sheep. This delayed adoption of evidence has not been reported from elicitation studies before and it is an intriguing finding.

#### Conclusions

We demonstrated that key players in the UK sheep industry were aware that prompt parenteral and topical antibiotic was more effective than foot trimming and topical antibiotic to treat footrot. All participants had incorporated parenteral antibiotic in the advice they gave before 2014, but not all recommended avoiding foot trimming. Some participants overestimated the difference in cure rates in favour of treatment with topical and parenteral antibiotic and for some this belief did not change even after they had been presented with the evidence. Three participants who stated that the evidence had not changed their beliefs had changed the advice they gave farmers three months after the workshop, some were recommending rapid treatment and others avoiding foot trimming.

#### **Conflict of interest statement**

None of the authors of this paper has a financial or personal relationship with other people or organisations that could inappropriately influence or bias the content of the paper.

#### Acknowledgements

This study was funded by the Warwick Impact Fund. Laura Green is funded by BBSRC (BB/M012980/1). We thank the participants, Ron Crump for help with graphics and Corinna Clark, Rachel Clifton, Clare Grant, Daniel Franklin, Katharina Giebel, Emma Monaghan and Edward Smith who assisted participants with the elicitation.

#### **Appendix: Supplementary material**

Supplementary data associated with this article can be found, in the online version, at https://doi.org/10.1016/j.tvjl.2018.07.009.

#### References

- Abbott, K., McGowan, M., Pfeiffer, D., Sargison, N., 2003. Footrot in sheep. Veterinary Record 152, 510–511.
- Beveridge, W.I.B., 1941. Foot-rot in sheep: a transmissible disease due to infection with *Fusiformis nodosus* (n. sp.): studies on its cause, epidemiology and control. Commonwealth Scientific and Industrial Research Organisation (CSIRO) Australian Bulletin 140. 1–56.
- Garthwaite, P.H., Kadane, J.B., O'Hagan, A., 2005. Statistical methods for eliciting probability distributions. Journal of the American Statistical Association 100, 680–700.
- Green, L.E., Clifton, R., 2018. Diagnosing and managing footrot in sheep: an update. In Practice 40, 17–26.
- Green, L.E., Wassink, G.J., Grogono-Thomas, R., Moore, L.J., Medley, G.F., 2007. Looking after the individual to reduce disease in the flock: a binomial mixed

effects model investigating the impact of individual sheep management of footrot and interdigital dermatitis in a prospective longitudinal study on one farm. Preventive Veterinary Medicine 78, 172–178.

- Higgins, H.M., Dryden, I.L., Green, M.J., 2012. A Bayesian elicitation of veterinary beliefs regarding systemic dry cow therapy: variation and importance for clinical trial design. Preventive Veterinary Medicine 106, 87–96.
- Higgins, H.M., Green, L.E., Green, M.J., Kaler, J., 2013. How does reviewing the evidence change veterinary surgeons' beliefs regarding the treatment of ovine footrot? A quantitative and qualitative study. PLoS One 8.
- Johnson, S.R., Tomlinson, G.A., Hawker, G.A., Granton, J.T., Grosbein, H.A., Feldman, B. M., 2010. A valid and reliable belief elicitation method for Bayesian priors. Journal of Clinical Epidemiology 63, 370–383.
- Kaler, J., Daniels, S.L.S., Wright, J.L., Green, L.E., 2010a. Randomized clinical trial of long-acting oxytetracycline, foot trimming, and flunixine meglumine on time to recovery in sheep with footrot. Journal of Veterinary Internal Medicine 24, 420– 425.
- Kaler, J., Medley, G.F., Grogono-Thomas, R., Wellington, E.M.H., Calvo-Bado, L.A., Wassink, G.J., King, E.M., Moore, L.J., Russell, C., Green, L.E., 2010b. Factors associated with changes of state of foot conformation and lameness in a flock of sheep. Preventive Veterinary Medicine 97, 237–244.
- Kaler, J., Green, L.E., 2009. Farmers' practices and factors associated with the prevalence of all lameness and lameness attributed to interdigital dermatitis and footrot in sheep flocks in England in 2004. Preventive Veterinary Medicine 92, 52–59.
- Kaler, J., Green, L.E., 2013. Sheep farmer opinions on the current and future role of veterinarians in flock health management on sheep farms: a qualitative study. Preventive Veterinary Medicine 112, 370–377.
- Locher, I., Greber, D., Holdener, K., Luchinger, R., Haerdi-Landerer, C., Schupbach-Regula, G., Frey, J., Steiner, A., 2015. Longitudinal *Dichelobacter nodosus* status in 9 sheep flocks free from clinical footrot. Small Ruminant Research 132, 128–132.
   Morgan, K.L., 1987. Footrot. In Practice 9, 124–129.
- Nelder, J.A., Mead, R., 1965. A simplex method for function minimization. Computer
- Journal 7, 308–313.
- R Development Core Team, 2008. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria.
- Spetzler, C.S., Staelvonholstein, C.A.S., 1975. Probability encoding in decision analysis. Management Science 22, 340–358.
- Wassink, G.J., Grogono-Thomas, R., Moore, L.J., Green, L.E., 2003. Risk factors associated with the prevalence of footrot in sheep from 1999 to 2000. Veterinary Record 152, 351–358.
- Wassink, G.J., King, E.M., Grogono-Thomas, R., Brown, J.C., Moore, L.J., Green, L.E., 2010. A within farm clinical trial to compare two treatments (parenteral antibacterials and hoof trimming) for sheep lame with footrot. Preventive Veterinary Medicine 96, 93–103.
- Winter, A., 2004. Lameness in sheep 2. Treatment and control. In Practice 26, 130-139.
- Winter, J.R., Kaler, J., Ferguson, E., KilBride, A.L., Green, L.E., 2015. Changes in prevalence of, and risk factors for, lameness in random samples of English sheep flocks: 2004–2013. Preventive Veterinary Medicine 122, 121–128.