

## Safety of primary anastomosis following emergency left sided colorectal resection

The 2017 European Society of Coloproctology (ESCP) collaborating group; Pinkney, Thomas

DOI:

[10.1111/codi.14373](https://doi.org/10.1111/codi.14373)

License:

Other (please specify with Rights Statement)

Document Version

Peer reviewed version

Citation for published version (Harvard):

The 2017 European Society of Coloproctology (ESCP) collaborating group & Pinkney, T 2018, 'Safety of primary anastomosis following emergency left sided colorectal resection: an international, multi-centre prospective audit', *Colorectal Disease*, vol. 20, no. S6, pp. 47-57. <https://doi.org/10.1111/codi.14373>

[Link to publication on Research at Birmingham portal](#)

**Publisher Rights Statement:**

Checked for eligibility: 20/11/2018

This is the peer reviewed version of the following article: The 2017 European Society of Coloproctology (ESCP) collaborating group (2018), Safety of primary anastomosis following emergency left sided colorectal resection: an international, multi-centre prospective audit. *Colorectal Dis*, 20: 47-57., which has been published in final form at doi:10.1111/codi.14373. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

**General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

**Take down policy**

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

**Safety of primary anastomosis following emergency left sided colorectal resection: An international, multi-centre prospective audit**

The 2017 European Society of Coloproctology (ESCP) collaborating group\*

\*Collaborating authors listed at end of manuscript

**Correspondence to:**

Luis Sánchez-Guillén

European Society of Coloproctology (ESCP) Cohort Studies Committee

Hospital Universitari i Politècnic la Fe

Avinguda de Fernando Abril Martorell, 106,

46026 València,

Spain

Email: [drsanchezguillen@gmail.com](mailto:drsanchezguillen@gmail.com)

**Keywords:** Surgery; Emergency surgery; Colon cancer; Rectal cancer; Gastrointestinal surgery; Anastomotic leak; Surgical complications; Surgical outcomes.

**Conflicts of interest:** None to declare

**Funding:** None

**Word count:** 1854

**Abstract word count:** 237

**Acknowledgements:** Supported by the European Society of Coloproctology (ESCP). REDCap and infrastructural support was received from the Birmingham Surgical Trials Institute (BiSTC) at the Birmingham Clinical Trials Unit (BCTU).

## **Abstract**

*Introduction:* Some evidence suggests that primary anastomosis following left sided colorectal resection in the emergency setting may be safe in selected patients, and confer favourable outcomes to permanent enterostomy. The aim of this study was to compare the major postoperative complication rate in patients undergoing end stoma versus primary anastomosis following emergency left sided colorectal resection.

*Methods:* A pre-planned analysis of the European Society of Coloproctology 2017 audit. Adult patients (>16 years) who underwent emergency (unplanned, within 24 hours of hospital admission) left sided colonic or rectal resection were included. The primary endpoint was the 30-day major complication rate (Clavien-Dindo grade 3 to 5).

*Results:* From 591 patients, 455 (77%) received an end stoma, 103 a primary anastomosis (17%) and 33 primary anastomosis with defunctioning stoma (6%). In multivariable models, anastomosis was associated with a similar major complication rate to end stoma (adjusted odds ratio for end stoma 1.52, 95%CI 0.83-2.79. p=0.173). Although a defunctioning stoma was not associated with reduced anastomotic leak (12% defunctioned [4/33] versus 13% not defunctioned [13/97], adjusted odds ratio 2.19, 95%CI 0.43-11.02, p=0.343), it was associated with less severe complications (75% [3/4] with defunctioning stoma, 86.7% anastomosis only [13/15]), a lower mortality rate (0% [0/4] versus 20% [3/15]), and fewer reoperations (50% [2/4] versus 73% [11/15]) when a leak did occur.

*Conclusions:* Primary anastomosis in selected patients appears safe after left sided emergency colorectal resection. A defunctioning stoma might mitigate against risk of subsequent complications.

### **What this paper adds**

Anastomosis after emergency left sided colorectal resection is performed in up to one in five patients. In these highly selected patients, this study suggests that it is safe practice. A defunctioning stoma may mitigate against risk if an anastomotic leak subsequently occurs.

## **Introduction**

In patients undergoing emergency left sided colorectal surgery, resection with end colostomy is a commonly described procedure. Concerns about the safety of any anastomosis in the emergency setting are particularly high in the presence of contamination or an unstable patient [1, 2]. Although a stoma avoids the risk of anastomotic leak, it carries with its own morbidity and mortality profile (27-55% and 4-27% respectively) [2]. For patients that undergo end stoma formation, the reversal rate is as low as 44% in published series [3], with a significant impact on long-term quality of life and a risk of stoma-related complications.

Many studies have evaluated primary anastomosis in the emergency setting with generally favourable results. Multiple single-centre, retrospective, observational studies have demonstrated that anastomosis can be safely performed in selected patients within the emergency setting, even in presence of peritonitis [4-6]. However, ~~it remains fairly uncommon practice and often only undertaken by more specialised or experienced colorectal surgeons.~~ The number of supporting randomised trials in the literature is low and those that exist are mainly related to peritonitis secondary to perforated diverticulitis, with primary anastomosis often only undertaken by specialised colorectal surgeons [4, 7, 8].

Decision-making about whether to create a primary anastomosis in selected, stable patients in an emergency setting remains a challenge for the individual surgeon. The decision must take into account patient comorbidities, intraoperative findings, underlying colorectal pathology, clinical status of the patient and expertise of the surgeon [9]. The aim of this multi-centre international study was to ~~provide examine whether more evidence to guide whether~~ current decision-making in real-world settings supports primary anastomosis as a safe technique in selected patients after emergency left sided colorectal resection.



## **Methods**

### *Protocol and centres*

This prospective, observational, multi-centre study was conducted in line with a pre-specified protocol (<http://www.escp.eu.com/research/cohort-studies>). All participating centres were responsible for compliance to local approval requirements for ethics approval or indemnity as required. In the UK, the National Research Ethics Service tool recommended that this project was not classified as research, and the protocol was registered as clinical audit in all participating centres. Any unit performing gastrointestinal surgery was eligible to register to enter patients into the study. No minimum case volume, or centre-specific limitations were applied. The study protocol was disseminated to registered members of the European Society of Coloproctology (ESCP), and through national surgical or colorectal societies.. This study represents planned analysis of the European Society of Coloproctology 2017 audit database.

### *Patient eligibility*

Adult patients (>16 years) undergoing left side colectomy or rectal resection, via any operative approach in emergency (~~within 24 hours~~) settings (within 24 hours of hospital admission) were extracted from the main audit database. Any indication for surgery (benign or malignant) were eligible. Patients undergoing planned elective surgery were excluded, as were those undergoing left colorectal resection as part of a more extensive resection (e.g. subtotal colectomy, panproctocolectomy).

### *Data capture*

Consecutive sampling was performed of eligible patients over an 8-week study period in each included centre. Local investigators commenced data collection on any date between the 1 January 2017 and 15 March 2017, with the last eligible patient being enrolled on 10 May 2017. Small teams of up to five surgeons or surgical trainees worked together to collect prospective

data on all eligible patients at each centre. Quality assurance was provided by at least one consultant or attending-level surgeon. Data was recorded contemporaneously and stored on a secure, user-encrypted online platform (REDCap) without using patient identifiable information. Centres were asked to validate that all eligible patients during the study period had been entered, and to attain >95% completeness of data field entry prior to final submission.

#### *Outcome measure*

The primary outcome measure was the 30-day postoperative major complication rate other, defined as Clavien-Dindo classification grade 3 to 5 (other than anastomotic leak including reoperation, reintervention, unplanned admission to critical care, organ support requirement or death). The secondary outcome measure was anastomotic leak, pre-defined as either i) gross anastomotic leakage proven radiologically or clinically, or ii) the presence of an intraperitoneal (abdominal or pelvic) fluid collection on post-operative imaging.

#### *Statistical analysis*

This report has been prepared in accordance to guidelines set by the STROBE (strengthening the reporting of observational studies in epidemiology) statement for observational studies [10]. Patient, disease and operative characteristics were compared using Student's t-test for normal, continuous data, Mann-Whitney U test for non-normal continuous data or Chi-squared test for categorical data. To test the association between the outcome measures and the main explanatory variables of interest (expedited versus emergency, end stoma versus primary anastomosis), a mixed-effects logistic regression model was fitted. Clinically plausible patient, disease and operation-specific factors were entered into the model for risk-adjustment, treated as fixed effects. These were defined *a priori* within the study protocol, and included irrespective of their significance on univariate analysis. Hospitals were entered into the model as a random-effect, to adjust for hospital-level variation in outcome. Effect estimates are presented as odds

ratios (OR) with 95% confidence intervals (95% CI) and two-tailed p-values. Model discrimination was ~~tested quantified~~ using C-statistic, ~~or~~, ~~analogous to~~ the area under the receiver operating characteristic curve curve (AUC) of the model. An alpha level of 0.05 was used throughout. Data analysis was undertaken using R Studio V3.1.1 (R Foundation, Boston, USA).

## Results

### *Patients*

This study included 591 patients undergoing emergency surgery from 43 countries ([Figure 1](#)).

The mean age of patients was 67.4 years (ranging from 18-96). 51.8% were male and 57.4% had a high anaesthetic risk class (ASA 3-5). Differences in demographics between patients with anastomosis and end stoma are shown in Table 1. Primary anastomosis was performed in 136 patients (23%) with 33 of these patients receiving a defunctioning stoma. This stoma was a loop ileostomy in 84.8% (28/33), an end/double-barreled ileostomy in 6.1% (2/33) and a loop colostomy in 3.0% (3/33). 30.1% (178/591) of included operations were done for malignancy, with end stoma being most common operative strategy (73.0%, 130/178). Of these, 20.2% were undergoing neoadjuvant therapy prior to their presentation for emergency surgery (short course radiotherapy, 7/36; long course chemoradiotherapy, 18/36; chemotherapy only: 11/36). Primary anastomosis with or without defunctioning stoma was performed less frequently than end stoma in disease affecting the rectum (14.9% and 7.2% versus 77.9% respectively). An anastomosis was attempted in 27% (87/326) of patients operated upon by a colorectal surgeon and 18% (49/265) by a general surgeon ( $p=0.059$ , [table 4](#)[Figure 2](#)).

### *Major complications (Clavien-Dindo 3 to 5)*

Results of analysis for factors associated with the occurrence of major complications are shown in Table [23](#). An end stoma was significantly associated with increased major postoperative complications at upon the univariable analysis (OR 1.98 95% CI 1.17–3.54,  $p=0.015$ ), but this association was not seen following risk adjustment (adjusted odds ratio for end stoma in mixed effects model 1.52, 95%CI 0.83-2.79,  $p=0.173$ ). In the multilevel model significant predictors for major complications were high ASA risk (grade 3-5) (OR 2.54, 95% CI 1.59–4.07,  $p<0.001$ ), overweight or obese (OR 1.88, 95% CI 1.17-3,  $p=0.009$ ) and male gender (OR 1.66, 95% CI 1.10–2.51,  $p=0.016$ ). Overweight BMI was associated with a lower major complication rate

than a normal BMI (OR 0.53, 95% CI 0.33-0.85, p=0.009), however the location of resection (involving rectum or colonic only) demonstrated no association. Anastomosis was associated with a similar major complication rate to end stoma (adjusted odds ratio for end stoma 1.52, 95%CI 0.83-2.79, p=0.173). The model demonstrated fair discrimination (AUC: 0.71).

#### *Anastomotic Leak*

Unadjusted outcomes according the anastomotic strategy, stratified by presence of leak, are shown in Table 3. Although a defunctioning stoma was not associated with reduced anastomotic leak (12% defunctioned [4/33] versus 13% not defunctioned [13/97], adjusted odds ratio 2.19, 95%CI 0.43-11.02, p=0.343), it was associated with fewer major complications (75% [3/4] with defunctioning stoma, 86.7% anastomosis only [13/15]), lower mortality (0% [0/4] versus 20% [3/15]), and reoperation 50% [2/4] versus 73% [11/15] when a leak did occur (Figure 3). The minor complication rate was similar between groups where the anastomosis successfully healed without leak (41.4% defunctioned [12/27] versus 34.1% not defunctioned [30/88]) and where an end stoma was formed (32.7% [149/455]). There was no difference in the anastomotic leak rate between anastomosis with and without defunctioning stoma (p=0.343). On the univariable analysis, age over 70 years old (OR 10, 95% CI 1.61-100, p=0.037) and (Table 4) previous history of IHD/CVA (OR 5.06, 95% CI 1.50-16.27, p=0.007) were was associated with an increased risk of leak, whilst being of middle age was protective (age 55 to 70 years old; OR 0.10, 95% CI 1.61-100, p=0.037). When a multilevel model was fitted (Figure 4), a significant association was found with age over than 70 years old (OR 20, 95% CI 0.00-1.51, p=0.023) a history ofnd diabetes also conveyed an increased risk of leak (OR 8.56, 95% CI 1.16-63.38, p=0.035). The model demonstrated good discrimination (AUC: 0.87).

|

## Discussion

This study showed that primary anastomosis was performed in up to one in five patients and appears safe in this highly selected group after emergency emergencyleft sided colorectal resection (unplanned, within 24 hours of hospital admission). A defunctioning stoma was only used in 24% of patients with a primary anastomosis. The exploratory findings of this study, limited by small numbers, suggested that a defunctioning stoma may mitigate against risk if an anastomotic leak occurs. Other patient-related risk characteristics (male gender, high ASA classgrade and overweight or obese BMI) and an open approach were identified as independent risk factors for major postoperative complications. Furthermore, young and elderly age and or a history of diabetes were shown as risk factors for anastomotic leak in emergency procedures.

Previously, the simple formation of an end colostomy after resection of the pathology (ubiquitously known as a ‘Hartmann’s procedure’) has been advocated as the gold standard treatment in emergency left colonic resection, to eliminate risk of anastomotic leak [11-13]. In the last fifteen years, several studies have questioned this strategy [14, 15]. A primary anastomosis is not only feasible, it may even be associated with better postoperative outcomes, both in terms of complications and mortality [16, 17]. Given that more than 40 % of temporary stomas become permanent, selecting patients correctly for a primary anastomosis is attractive [18-20]. In addition, Hartmann reversal can be a technically demanding operation resulting in further morbidity and mortality [21]. These findings support a recent consensus statements and prospective multi-centre randomized trials that suggest primary anastomosis with proximal diversion as an optimal strategy for sigmoid diverticulitis in selected patients with Hinchey 3 or 4 disease [4, 22, 23]. This current study gives credence to the current situation and confirms that surgeons are making appropriate decisions on a case-by-case level, thereby

effectively stratifying patients for primary anastomosis or end stoma. It is known that a defunctioning stoma in elective surgery has utility in mitigating the clinical impact of anastomotic leak [24, 25]. Loop ileostomies and their closure are not complication-free and several studies have shown that temporary loop ileostomies can become permanent in up to 25% of patients [24-28]. However this study comparably suggests that a defunctioning stoma may mitigate some risk when a leak occurs. This must be interpreted with caution since numbers in this study were low; for example, only four patients with an anastomosis and defunctioning stoma suffered a leak.

There were slightly more primary anastomotic attempts by colorectal versus general surgeons in this study. Even though there is no homogenous definition of colorectal surgeon internationally, results of multiple studies confirm the importance of colorectal specialisation in the emergency setting [29, 30]. An individual surgeon's personality and their response to perceived operative risk may also influence choice of anastomotic strategy [31]. Further research is needed to determine whether the grade and surgical specialism of the operating (or senior) surgeon, and specialisation and experience of included centres affect both the decision to-for anastomosis and the subsequent clinical outcome.

There are inherent limitations to the 'snapshot' observational study reported here in this study which we have attempted to overcome in the study design, statistical analysis and interpretation. There is an obvious selection bias in this study, although we planned the analysis around this *a priori*. We aimed to analyse safety of current practice; this study showed that end stoma was more frequently used in older patients, with poor general status, in smokers, and in those with arteriopathy and benign disease. However this paper defines outcomes in the highly selected group of patients undergoing anastomosis, and thus supports surgical decision making in specific cases, rather than in recommending a general change in approach. The low numbers

of anastomotic leak and major complication within secondary analyses of the subgroup undergoing anastomosis (<25% of included patients) makes estimation of effect sizes inaccurate here (reflected by broad confidence intervals). Therefore, this should be seen as exploratory only; the analysis would likely be underpowered to detect a small to moderate effect size. We are also unable to comment on the appropriateness of decision making and have not collected detailed information on parameters that may effect this (for example: contamination (Mannheim Peritonitis Index [32]), previous surgery, intraoperative physiological instability). Most of the literature available on this topic is based on retrospective or single centres data which lacks sufficient detail to allow case-mix adjustment in multivariable models. This study therefore adds to the literature in providing a contemporary perspective using a prospective international observational study design, with a pre-specified protocol and analysis plan. In addition, the variety of centers included (in terms of number of patients, facilities and different technologies available) in this study delivers a realistic picture of the current management of emergency left colorectal resections, avoiding reducing selection bias and increasing the external validity of the findings. The different countries and even continents involved ensured the result's validity resolving the demographic differences in diverticulitis and cancer across countries. Finally, the study is limited by short-term follow up to 30 days only; we have not collected data on stoma reversal rates, quality of life or stoma-related complications following surgery. An alternative complication categorisation system such as the Comprehensive Complications Index [33] may also give increased fidelity in comparisons between intermediate term outcomes. Further evaluation of these important parameters following emergency left sided colorectal surgery is warranted.

The data from this study supports current international practice of primary anastomosis following emergency left sided colorectal resection in a highly selected group of patients, demonstrating satisfactory safety and an acceptable morbidity profile. Where an anastomosis is formed, a

defunctioning stoma does not appear to reduce the risk of leak, but may mitigate the severity of resultant complications.

## References

1. Hartmann H. Note sur un procédé nouveau d'extirpation des cancers de la partie du côlon. *Bull Mem Soc Chir Paris* 1923;49:1474—7.
2. J. Barbieux, F. Plumerias, A. Hammy. Current indications for Hartmann procedure. *J Visc Surg.* 2016 Feb;153(1):31-8.
3. van de Wall BJ, Draaisma WA, Schouten ES, Broeders IA, Consten EC. Conventional and laparoscopic reversal of the Hartmann procedure: a review of literature. *J Gastrointest Surg* 2010; **14:** 743-52.
4. Bridoux V, Regimbeau JM, Ouaissi M, Mathonnet M, Mauvais F, Houivet E, et al. Hartmann's Procedure or Primary Anastomosis for Generalized Peritonitis due to Perforated Diverticulitis: A Prospective Multicenter Randomized Trial (DIVERTI). *J Am Coll Surg* 2017; **225:** 798-805.
5. Salem L, Flum DR. Primary anastomosis or Hartmann's procedure for patients with diverticular peritonitis? A systematic review. *Dis Colon Rectum* 2004; **47:** 1953-64.
6. Trenti L, Biondo S, Golda T, Monica M, Kreisler E, Fraccalvieri D, et al. Generalized peritonitis due to perforated diverticulitis: Hartmann's procedure or primary anastomosis? *Int J Colorectal Dis* 2011; **26:** 377-84.
7. Oberkofer CE, Rickenbacher A, Raptis DA, Lehmann K, Villiger P, Buchli C, et al. A multicenter randomized clinical trial of primary anastomosis or Hartmann's procedure for perforated left colonic diverticulitis with purulent or fecal peritonitis. *Ann Surg* 2012; **256:** 819-26; discussion 26-7.
8. Binda GA, Karas JR, Serventi A, Sokmen S, Amato A, Hydo L, et al. Primary anastomosis vs nonrestorative resection for perforated diverticulitis with peritonitis: a prematurely terminated randomized controlled trial. *Colorectal Dis* 2012; **14:** 1403-10.

9. Gibbons G, Tan CJ, Bartolo DC, Filgate R, Makin G, Barwood N, et al. Emergency left colonic resections on an acute surgical unit: does subspecialization improve outcomes? *ANZ J Surg* 2015; **85**: 739-43.
10. von Elm E, Altman DG, Egger M, Pocock SJ, Gotzsche PC, Vandebroucke JP, et al. Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *BMJ* 2007; **335**: 806-8.
11. Myers E, Hurley M, O'Sullivan GC, Kavanagh D, Wilson I, Winter DC. Laparoscopic peritoneal lavage for generalized peritonitis due to perforated diverticulitis. *Br J Surg* 2008; **95**: 97-101.
12. Sanderson ER. Henri Hartmann and the Hartmann operation. *Arch Surg* 1980; **115**: 792-3.
13. Walklett CL, Yeomans NP. A retrospective case note review of laparoscopic versus open reversal of Hartmann's procedure. *Ann R Coll Surg Engl* 2014; **96**: 539-42.
14. Constantinides VA, Tekkis PP, Athanasiou T, Aziz O, Purkayastha S, Remzi FH, et al. Primary resection with anastomosis vs. Hartmann's procedure in nonelective surgery for acute colonic diverticulitis: a systematic review. *Dis Colon Rectum* 2006; **49**: 966-81.
15. Zorcolo L, Covotta L, Carlomagno N, Bartolo DC. Safety of primary anastomosis in emergency colo-rectal surgery. *Colorectal Dis* 2003; **5**: 262-9.
16. Gachabayov M, Oberkofer CE, Tuech JJ, Hahnloser D, Bergamaschi R. Resection with primary anastomosis vs nonrestorative resection for perforated diverticulitis with peritonitis: a systematic review and meta-analysis. *Colorectal Dis* 2018
17. Schilling MK, Maurer CA, Kollmar O, Buchler MW. Primary vs. secondary anastomosis after sigmoid colon resection for perforated diverticulitis (Hinchey Stage III and IV): a prospective outcome and cost analysis. *Dis Colon Rectum* 2001; **44**: 699-703; discussion -5.

18. Fleming FJ, Gillen P. Reversal of Hartmann's procedure following acute diverticulitis: is timing everything? *Int J Colorectal Dis* 2009; **24**: 1219-25.
19. Hallam S, Mothe BS, Tirumulaju R. Hartmann's procedure, reversal and rate of stoma-free survival. *Ann R Coll Surg Engl* 2018; **100**: 301-7.
20. Roig JV, Cantos M, Balciscueta Z, Uribe N, Espinosa J, Rosello V, et al. Hartmann's operation: how often is it reversed and at what cost? A multicentre study. *Colorectal Dis* 2011; **13**: e396-402.
21. Garber A, Hyman N, Osler T. Complications of Hartmann takedown in a decade of preferred primary anastomosis. *Am J Surg* 2014; **207**: 60-4.
22. Feingold D, Steele SR, Lee S, Kaiser A, Boushey R, Buie WD, et al. Practice parameters for the treatment of sigmoid diverticulitis. *Dis Colon Rectum* 2014; **57**: 284-94.
23. Roig JV, Sanchez-Guillen L, Garcia-Armengol JJ. Acute diverticulitis and surgical treatment. *Minerva Chir* 2018; **73**: 163-78.
24. Chow A, Tilney HS, Paraskeva P, Jeyarajah S, Zacharakis E, Purkayastha S. The morbidity surrounding reversal of defunctioning ileostomies: a systematic review of 48 studies including 6,107 cases. *Int J Colorectal Dis* 2009; **24**: 711-23.
25. Garcia-Botello SA, Garcia-Armengol J, Garcia-Granero E, Espi A, Juan C, Lopez-Mozos F, et al. A prospective audit of the complications of loop ileostomy construction and takedown. *Dig Surg* 2004; **21**: 440-6.
26. Wong NY, Eu KW. A defunctioning ileostomy does not prevent clinical anastomotic leak after a low anterior resection: a prospective, comparative study. *Dis Colon Rectum* 2005; **48**: 2076-9.
27. Kong AP, Kim J, Holt A, Konyalian V, Huynh R, Udani SM, et al. Selective treatment of rectal cancer with single-stage coloanal or ultralow colorectal anastomosis does not adversely affect morbidity and mortality. *Int J Colorectal Dis* 2007; **22**: 897-901.

28. Chun LJ, Haigh PI, Tam MS, Abbas MA. Defunctioning loop ileostomy for pelvic anastomoses: predictors of morbidity and nonclosure. *Dis Colon Rectum* 2012; **55**: 167-74.
29. Biondo S, Kreisler E, Millan M, Fraccalvieri D, Golda T, Frago R, et al. Impact of surgical specialization on emergency colorectal surgery outcomes. *Arch Surg* 2010; **145**: 79-86.
30. Patel S, Senagore A. General surgeons vs. colorectal surgeons: Who should be doing what to whom?. *Seminars Colon Rectal Surgery* 2013; **24**: 191-194.
31. Moug SJ, Henderson N, Tiernan J, Bisset CN, Ferguson E, Harji D, Maxwell-Armstrong C, MacDermid E, Acheson AG, Steele RJC, Fearnhead NS; Edinburgh Delphi Collaborative Group. The colorectal surgeon's personality may influence the rectal anastomotic decision. *Colorectal Dis.* 2018 Jun 14. doi: 10.1111/codi.14293.
32. Wacha H, Linder MM, Feldman U, Wesch G, Gundlach E, Steifensand RA. Mannheim peritonitis index – prediction of risk of death from peritonitis: construction of a statistical and validation of an empirically based index. *Theoretical Surg.* 1987;1:169–77.
33. Clavien PA, Vetter D, Staiger RD, Slankamenac K, Mehra T, Graf R, Puhan MA. The Comprehensive Complication Index (CCI®): Added Value and Clinical Perspectives 3 Years "Down the Line". *Ann Surg.* 2017 Jun;265(6):1045-1050.

**Authorship list (PubMed citable)****Writing Group**

Luis Sánchez-Guillén (Chair), Dmitri Nepogodiev, Emre Sivrikoz, Tim van Elst, Ana Minaya, Francesco Pata, James Glasbey, Thomas Pinkney, Aneel Bhangu

**ESCP Cohort Studies and Audits Committee**

Alaa El-Hussuna (2017 Audit Lead), Aneel Bhangu, Nicholas Buchs, Christianne Buskens, Sanjay Chaudri, Matteo Frasson, Gaetano Gallo, James Glasbey, Ana Minaya, Dion Morton, Ionut Negoi, Dmitri Nepogodiev, Francesco Pata, Luis Sánchez-Guillén, Baljit Singh, Oded Zmora, Thomas Pinkney (Chair)

**Statistical Analysis and Data Management**

James Glasbey, Dmitri Nepogodiev, Rita Perry, Laura Magill, Aneel Bhangu (Guarantor)

**ESCP Research Committee**

Dion Morton (Chair), Donato Altomare, Willem Bemelman, Steven Brown, Christianne Buskens, Quentin Denost, Charles Knowles, Søren Laurberg, Jérémie Lefèvre, Gabriela Möeslein, Tom Pinkney, Carolynne Vaizey, Oded Zmora

**Collaborators**

Albania: S. Bilali, V. Bilali (University Hospital Center Mother Teresa).

Argentina: M. Salomon, M. Cillo, D. Estefania, J. Patron Uriburu, H. Ruiz (Buenos Aires British Hospital); P. Farina, F. Carballo, S. Guckenheimer (Hospital Pirovano).

Australia: D. Proud, R. Brouwer, A. Bui, B. Nguyen, P. Smart (Austin Hospital); A. Warwick, J. Theodore (Redcliffe Hospital).

Austria: F. Herbst, T. Birsan, B. Dauser, S. Ghaffari, N. Hartig (Barmherzige Brueder, Wien); A. Stift, S. Argeny, L. Unger (Medical University of Vienna); R. Strouhal, A. Heuberger (Oberndorf b. Salzburg).

Belarus: A. Varabei, N. Lahodzich, A. Makhmudov, L. Selniahina (Minsk Regional Clinical Hospital).

Belgium: T. Feryn, T. Leupe, L. Maes, E. Reynvoet, K. Van Langenhove (AZ Sint-Jan Brugge); M. Nachtergaele (AZ St Jozef); B. Monami, D. Francart, C. Jehaes, S. Markiewicz, J. Weerts (Clinique St Joseph, Liege); K. Van Belle, B. Bomans, V. Cavenaile, Y. Nijs, M. Vertruyen (Europe Hospitals Brussels); P. Pletinckx, D. Claeys, B. Defoort, F. Muysoms, S. Van Cleven (Maria Middelares Gent); C. Lange, K. Vindevoghel (OLV van Lourdes Hospital Waregem); A. Wolthuis (University Hospital Leuven).

Bosnia and Heregovina: M. Todorovic, S. Dabic, B. Kenjic, S. Lovric, J. Vidovic (JZU Hospital Sveti Vračevi); S. Delibegovic, Z. Mehmedovic (University Clinic Center Tuzla).

Brazil: A. Christiano, B. Lombardi, M. Marchiori Jr, V. Terciotti Jr (Hospital Centro Médico de Campinas).

Bulgaria: D. Dardanov, P. Petkov, L. Simonova, A. Yonkov, E. Zhivkov (Alexandrovska Hospital - First Surgery); S. Maslyankov, V. Pavlov, M. Sokolov, G. Todorov (Alexandrovska Hospital, Second Surgery Clinic); V. Stoyanov, I. Batashki, N. Iarumov, I. Lozev, B. Moshev (Medical

Institute - Ministry of Interior); M. Slavchev, B. Atanasov, N. Belev, P. Krstev, R. Penkov (University Hospital - Eurohospital).

Croatia: G. Šantak, J. Čosić, A. Previšić, L. Vukušić, G. Zukanović (County Hospital Požega); M. Zelić, D. Kršul, V. Lekić Vitlov, D. Mendrila (University Hospital Rijeka).

Czech Republic: J. Orhalmi, T. Dusek, O. Maly, J. Paral, O. Sotona (Charles University Hospital). M. Skrovina, V. Bencurik, M. Machackova (Complex Oncology Center Novy Jicin, Surgical Department); Z. Kala, M. Farkašová, T. Grolich, V. Procházka (Surgical Department, University Hospital Brno); J. Hoch, P. Kocian, L. Martinek (University Hospital Motol, Prague); F. Antos, V. Pruchova (University Hospital Prague Bulovka).

Denmark: A. El-Hussuna, A. Ceccotti, T. Madsbøll, D. Straarup, A. Uth Ovesen (Aalborg University Hospital); P. Christensen, P. Bondeven, P. Edling, H. Elfeki, V. Alexandrovich Gameza, S. Michelsen Bach, I. Zheltiakova (Aarhus University Hospital/Randers Regional Hospital); PM. Krarup, A. Krogh, H-C. Rolff (Bispebjerg); J. Lykke, A. F. Juvik, H. H. K. Lóven, M. Marckman, J. T. F. Osterkamp (Herlev Hospital); A. H. Madsen, J. Worsøe (Hospital Unit West); A. Ugianskis (North Denmark Regional Hospital); M. D. Kjær, B. Youn Cho Lee (Odense University Hospital); A. Khalid, M. H. Kristensen (Regional Hospital Viborg).

Egypt: M. El Sorogy, A. Elgeidie, M. Elhemaly, A. ElNakeeb, M. Elrefai (Gastrointestinal Surgery Center, Mansoura University); M. Shalaby, S. Emile, W. Omar, A. Sakr, W. Thabet (Mansoura University Hospital); S. Awny, I. Metwally, B. Refky, N. Shams, M. Zuhdy (Oncology Center Mansoura University).

Finland: A. Lepistö, I. Keränen, A. Kivelä, T. Lehtonen, P. Siironen (Helsinki University Hospital); T. Rautio, M. Ahonen-Siirtola, K. Klintrup, K. Paarnio, H. Takala (Oulu University

Hospital); M. Hyöty, E. Haukijärvi, S-M. Kotaluoto, K. Lehto, T. Tomminen (Tampere University Hospital); H. Huhtinen, A. Carpelan, J. Karvonen, A. Rantala, P. Varpe (Turku University Hospital).

France: E. Cotte, Y. Francois, O. Glehen, G. Passot (Centre Hospitalier Lyon Sud); A. d'Alessandro, E. Chouillard, J. C. Etienne, E. Ghilles, B. Vinson-Bonnet (CHIPS); A. Germain, A. Ayav, L. Bresler (CHU Nancy-Brabois); R. Chevalier, Q. Denost, R. Didailleur, E. Rullier (Hôpital Haut Leveque); E. Tiret, N. Chafai, J. Lefevre, Y. Parc (Hôpital Saint-Antoine); I. Sielezneff, D. Mege (Timone Hospital); Z. Lakkis (University Hospital of Besançon); M. Barussaud (University Hospital of Poitiers).

Germany: C. Krones, B. Bock, R. Webler (Marienhospital Aachen); J. Baral, T. Lang, S. Münch, F. Pullig, M. Schön (Städtisches Klinikum Karlsruhe); S. Hinz, T. Becker, T. Möller, F. Richter, C. Schafmayer (University Hospital Schleswig-Holstein, Kiel); J. Hardt, P. Kienle (University Medical Center Mannheim); F. Crescenti, M. Ahmad, Y. Soleiman (Verden KRH).

Greece: I. Papaconstantinou , A. Gklavas, K. Nastos, T. Theodosopoulos, A. Vezakis (Areteion Hospital); K. Stamou, A. Saridaki (Athens Bioclinic); E. Xynos, S. Paraskakis, N. Zervakis (Creta-InterClinic Hospital); G. Skroubis, T. Amanatidis, S. Germanos, I. Maroulis, G. Papadopoulos (General University Hospital of Patras); N. Dimitriou, A. Alexandrou, E. Felekouras, J. Griniatsos, I. Karavokyros (Laiko Hospital); A. Papadopoulos, C. Chouliaras, P. Ioannidis, D. Katsounis, E. Kefalou (Nikaia General Hospital); I. Katsoulis, D. Balalis, D. Manatakis (St. Savvas Cancer Hospital); G. Tzovaras, I. Baloyiannis, I. Mamaloudis (University Hospital of Larissa).

Hungary: G. Lázár, S. Ábraham, A. Paszt, Z. Simonka (Department of Surgery, University of Szeged); A. Zaránd, Z. Baranyai, G. Ferreira, L. Harsányi, P. Ónody (Semmelweis University,

1st Clinic of Surgery); B. Banky, Á. Burány, M. Lakatos, J. Marton, A. Solymosi (St. Borbala Hospital); I. Besznyák, A. Bursics, G. Papp, G. Saftics, I. Svastics (Uzsoki Hospital);

Iceland: E. Valsdottir, J. Atladottir, T. Jonsson, P. Moller, H. Sigurdsson (University Hospital of Iceland).

India: S. K. Gupta, S. Gupta, N. Kaul, S. Mohan, G. Sharma (Government Medical College, Jammu, Jammu and Kashmir, India); R. Wani, N. Chowdri, M. Khan, A. Mehraj, F. Paray (Sher-i-Kashmir Institute of Medical Sciences).

Ireland: A. Coveney, J. Burke, J. Deasy, S. El-Masry, D. McNamara (Beaumont Hospital); M. F. Khan, R. Cahill, E. Faul, J. Mulsow, C. Shields (Mater Misericordiae University Hospital); M. E. Kelly, G. Bass, S. T. Martin, R. O'Connell, E. Ryan (St Vincent's Private Hospital); T. Connelly, G. Ahmad, W. Bukhari, F. Cooke (University Hospital Waterford).

Israel: O. Zmora, R. Gold Deutch, N. Haim, R. Lavy, A. Moscovici (Assaf Harofe Medical Center); N. Shussman, R. Gefen, G. Marom, A. Pikarsky, D. Weiss (Hadassah Hebrew University Medical Center); S. Avital, N. Hermann, B. Raguan, M. Slavin, I. White (Meir Medical Center); N. Wasserberg, H. Arieli, N. Gurevich (RMC, Beilinson Campus); M. R. Freund, S. Dorot, G. Halfteck, P. Reissman, E. Yair (Shaare Zedek Mount Sinai); Y. Eden, R. Pery (Sheba Medical Center); H. Tulchinsky, A. Weizman (Sourasky Medical Center).

Italy: F. Agresta, R. Curinga, E. Finotti, G. Savino, L. A. Verza (Adria Hospital); C. R. Asteria, L. Boccia, A. Pascariello (ASST - Mantua); N. Tamini, A. Bugatti, L. Gianotti, M. Totis (Asst-Monza, Ospedale San Gerardo); L. Vincenti, V. Andriola, I. Giannini, E. Travaglio (Azienda Ospedaliero Universitaria Consorziale Policlinico di Bari); R. Balestri, P. Buccianti, N. Roffi, E.

Rossi, L. Urbani (Azienda Ospedaliero Universitaria Pisana); A. Mellano, A. Cinquegrana (Candiolo Cancer Institute IRCCS); A. Lauretta, C. Belluco (Centro di Riferimento Oncologico, IRCCS, Aviano); M. Mistrangelo, M. E. Allaix, S. Arolfo, M. Morino, V. Testa (Città della Salute e della Scienza di Torino); P. Delrio, U. Pace, D. Rega, D. Scala (Colorectal Surgical Oncology Istituto Nazionale per lo Studio e la Cura dei Tumori); G. Gallo, G. Clerico, S. Cornaglia, A. Realis Luc, M. Trompetto (Department of Colorectal Surgery, S. Rita Clinic); G. Ugolini, N. Antonacci, S. Fabbri, I. Montroni, D. Zattoni (Faenza Hospital); C. D'Urbano, A. Cornelli, M. Viti (G. Salvini); M. Inama, M. Bacchion, A. Casaril, H. Impellizzeri, G. Moretto (Hospital Dott. Pederzoli); A. Spinelli, M. Carvello, G. David, F. Di Candido, M. Sacchi (Humanitas Research Hospital); A. Frontali, V. Ceriani, M. Molteni (IRCCS MultiMedica); R. Rosati, F. Aleotti, U. Elmore, M. Lemma, A. Vignali (IRCCS Ospedale San Raffaele); S. Scabini, G. Casoni Pattacini, A. Luzzi, E. Romairone (IRCCS San Martino IST); F. Marino, D. Lorusso, F. Pezzolla (IRCCS 'Saverio de Bellis'); F. Colombo, C. Baldi, D. Foschi, G. Sampietro, L. Sorrentino (L. Sacco University Hospital); S. Di Saverio, A. Birindelli, E. Segalini, D. Spacca (Maggiore Hospital); G. M. Romano, A. Belli, F. Bianco, S. De franciscis, A. Falato (National Cancer Institute Naples); A. Muratore, P. Marsanic (Ospedale Agnelli Pinerolo); S. Grimaldi, N. Castaldo, M. G. Ciolli, P. Picarella, R. Porfidia (Ospedale Convenzionato Villa Dei Fiori); S. Di Saverio, A. Birindelli, G. Tugnoli (Ospedale Maggiore); A. Bondurri, D. Cavallo, A. Maffioli, A. Pertusati (Ospedale Sacco Italy); F. Pulighe, F. Balestra, C. De Nisco, M. Podda (Ospedale San Francesco); E. Opocher, M. Longhi, N. M. Mariani, N. Maroni, A. Pisani Ceretti (Ospedale San Paolo); R. Galleano, P. Aonzo, G. Curletti, L. Reggiani (Ospedale Santa Corona); M. Marconi, L. Del Prete, M. Oldani, R. Pappalardo, S. Zaccione I (Ospedale Santa Maria delle Stelle); M. Scatizzi, M. Baraghini, S. Cantafio, F. Feroci, I. Giani (Ospedale Santo Stefano, Prato); R. Tutino, G. Cocorullo, G. Gulotta, L. Licari, G. Salamone (Policlinico 'P. Giaccone'); P. Sileri, F. Saraceno (Policlinico Tor Vergata); F. La Torre, P. Chirletti, D. Coletta, G. De Toma, A. Mingoli (Policlinico Umberto I 'Sapienza University'); M. Papandrea, E. De Luca, R. Sacco, G. Sammarco, G. Vescio

(Policlinico Universitario di Catanzaro); V. Tonini, S. Bianchini, M. Cervellera, S. Vaccari (Policlinico universitario Sant'Orsola-Malpighi, Universita degli Studi di Bologna); N. Cracco, G. Barugola, E. Bertocchi, R. Rossini, G. Ruffo (Sacro Cuore Don Calabria Hospital); A. Sartori, N. Clemente, M. De Luca, A. De Luca, G. Scaffidi (San Valentino Hospital); L. Lorenzon, G. Balducci, T. Bocchetti, M. Ferri, P. Mercantini (Sant'Andrea Hospital); F. Pata, S. Bause, A. Benevento, C. Bottini, P. R. Crapa (Sant' Antonio Abate Hospital, Gallarate); M. Rubbini, G. Anania, P. Carcoforo, G. Cavallesco, C. Feo (University Hospital of Ferrara).

Japan: T. Yamamoto (Yokkaichi Hazu Medical Centre).

Latvia: A. Sivins, G. Ancans, S. Gerkis, R. Lunis, A. Pcolkins (Latvia Oncology Center).

Lithuania: D. Venskutonis, S. Bradulskis, E. Dainius, A. Subocius, J. Vencius (Department of General Surgery, LSMU, Kaunas Clinical Hospital); P. Zeromskas, V. Eismontas, V. Nutautiene, D. Simcikas, A. Tamosiunas (Klaipeda University Hospital); S. Svagzdys, T. Latkauskas, P. Lizdenis, Z. Saladzinskas, A. Tamelis (Lithuanian University of Health Sciences Hospital Kauno Klinikos); A. Dulskas, J. Kulavas, N. E. Samalavicius (National Cancer Institute, Lithuania); T. Poskus, V. Jotautas, S. Mikalauskas, E. Poskus, K. Strupas (Vilnius University).

Malaysia: A. Zakaria, N. N.Lah, M. Wong, Z. Zain, Z. Zakaria (Hospital Universiti Sains Malaysia); L. Mazlan, Z. A. Mohd Azman, I. Sagap (UKM Medical Centre).

Malta: J. Psaila, P. Andrejevic, C. Cini, S. Ellul, K. Pace (Mater Dei Hospital). Morocco: M. Ahallat, M. Hamid, A. Hrora, M. A. Majbar, M. Raiss (Ibn Sina University Hospital).

Netherlands: E. Westerduin, W. Bemelman, C. Buskens, P. Tanis (Academic Medical Centre); P.C. van der Sluis, P.H. Davids, A. Pronk, A.H.W. Schiphorst, N. Smakman (Diakonessenhuis); D. Zimmerman, T. Koeter, J. Stijns, Y-T. van Loon (Elisabeth TweeSteden Hospital); M. Vermaas, E. de Graaf, P. Doornbosch, P. van Hagen, O. van Ruler (IJsstelland Ziekenhuis); B. Toorenvliet, J. Nonner, I. van den Berg, L. van Steensel, W. Vles (Ikazia); J. Melenhorst, R. Orsini, R. Visschers (Maastricht University Medical Centre); C. Hoff (Medical Center Leeuwarden); R. Blom, H. Marsman (Onze Lieve Vrouwe Gasthuis); I. Mulder, H. Cense, S. de Castro, A. Demirkiran, M. Hunfeld (Rode Kruis Ziekenhuis Beverwijk); A. van Geloven, J. de Groof, E. Hendriks, M. Leeuwenburg, N. van Oorschot (Tergooi); F. Wit, C. Rupert, P. Veldman (Tjongerschans ziekenhuis); M. Keijzers, J. Konsten (VieCuri Medisch Centrum voor Noord Limburg); F. Den Boer, M. Corver (Zaans Medical Center); E-J. Boerma, L. Koolen, M. Martens, K. Van Wijck (Zuyderland Medical Center).

Norway: D. Ignjatovic, R. Breuer, B. Gurpreet, T. Oresland, T. Tetens Moe (Akershus University Hospital); A. Nesbakken, I. Flateh Backe, T-A. Wik (Oslo University Hospital); K. Radiya, T. Dehli, P. Gjessing, S. Norderval, K. Woll (University Hospital of North Norway).

Pakistan: M. Anwer, M. S. Qureshi (JPMC WARD 2); A. U. Qureshi, M. Billah, M. Y. Jawad, A. Raza, N. Urooj (King Edward Medical University/Mayo Hospital, Lahore).

People's Republic of China: X. Wang, L. Li (West China Hospital). Poland: D. Jajtner, B. Gasinski, W. Kabiesz (Beskidian Oncology Center); P. Walega, M. Romaniszyn (Third Department of General Surgery, Jagiellonian University Medical College); M. Zawadzki, R. Czarnecki, Z. Obuszko, M. Rzaca, M. Sitarska (Wojewódzki Szpital Specjalistyczny).

Portugal: P. Silva, A. Duarte, D. Gonçalves, M. Morais (Centro Hospitalar de S. João); N. Rama, J. Nobre, I. Sales (Centro Hospitalar Leiria, EPE); J. Costa Pereira, S. Costa, C. Costa Pereira, C. Insua, I. Romero (Centro Hospitalar Tâmega e Sousa); N. Figueiredo, J. Cunha, H. Domingos, P. Vieira (Champalimaud Foundation); M. Cunha, M. Americano, E. Amorim, J. Rachadell (Cirurgia 2 - CHA - Unidade Portimão); J. Carvas, I. Armas, P. Fernandes, C. Pires, R. Reis (Hospital de Bragança); R. Martins, M. Dos Santos, P. Henriques (Hospital de Faro, Centro Hospitalar do Algarve); O. Oliveira, M. Duarte, L. Ferreira, J. Miranda, N. Vilela (Hospital Distrital de Santarém, E.P.E.); J. Corte Real, S. Carlos, M. Frois Borges, P. Moniz Pereira, J. Simões (Hospital Garcia de Orta); P. Silva-vaz, V. Bettencourt, A. Gouveia, H. Perez, R. Rainho (Unidade Local de Saúde de Castelo Branco).

Romania: V. Bintintan, C. Ciuce, G. Dindelegan, R. Scurtu, R. Seicean (Clinica Chirurgie I); D. Cristian, T. Burcos, F. Grama, D. M. Mandi, G. Richiteanu (Coltea Clinical Hospital); A. Miron, V. Calu, O. Enciu, M. Nadragea, R. Parvuletu (Elias Emergency Hospital); S. S. Mogoanta, A. Crafcicu, S. Paitici (Emergency County Hospital of Craiova); I. Negoi, M. Beuran, C. Ciubotaru, A. Prodan, M. Vartic (Emergency Hospital of Bucharest); V. Tomulescu, C. Copăescu (Ponderas Academic Hospital).

Russia: D. Popov, A. Sednev, A. Klimenko, A. Semenov, S. Vasiliyev (City Hospital 9); A. Pozdnyakov, D. Cherdancev, D. Mahotin, A. Nesytykh, V. Samsonyuk (Krasnoyarsk Regional Clinical Hospital); I. Pravosudov, D. Ivlev, A. Karachun, K. Lebedev, D. Samsonov (N.N. Petrov Research Institute of Oncology); R. Aiupov, D. Feoktistov, M. Garipov, S. Nail, N. Tarasov (National Republic Oncology Center); A. Yanishev, A. Abelevich, A. Kokobelyan, M. Lebedeva, R. Luzan (Nizhny Novgorod Regional Clinical Hospital); A. Rasulov, H. Dzhamabaev, Z. Mamedli (Russian Cancer Research Center); A. Bedzhanyan (Russian Reserch Center of Surgery named after B.V.Petrovsky); A. Khazov, M. Khanovich, G. Khrykov (Saint-Petersburg Clinical Oncological Health Center); S. Katorkin, P. Andreev, A. Chernov, O. Davidova, A.

Zhuravlev (Samara State Medical University); S. Achkasov, D. Shakhmatov, Y. Shelygin, O. Sushkov, A. Vardanyan (State Scientific Centre of Coloproctology); A. Ilkanich, N. Barbashinov, V. Darvin, S. Onishchenko, Y. Voronin (Surgut District Hospital).

Serbia: Z. Krivokapić, G. Barišić, I. Dimitrijević, V. Marković, A. Sekulić (Clinic for Digestive Surgery-I Surgical Clinic, Clinical Ceter of Serbia); G. Stanojevic, B. Brankovic, M. Nestorovic, V. Pecic, D. Petrovic (Clinic for General Surgery, Clinical Center Nis); I. Kostic, A. Aleksic, D. Dabic, B. Maric, V. Perunicic (General Hospital Cacak); Z. Radovanovic, M. Djuric, D. Lukic, D. Radovanovic (Oncology Institute of Vojvodina); V. Cuk, V. Cuk, J. Juloski, M. Kenic, I. Krdzic (Surgical Clinic KBC Zvezdara).

Singapore: J. C. Ngu, Y. Y. Ng, N. Teo (Changi General Hospital).

Slovak Republic: J. Korcek, A. Lazorisak, (Faculty Hospital Nitra).

Slovenia: M. Rems, Š. Ramovš Trampuš (General Hospital Jesenice); A. Tomazic, J. Grosek, J. Kosir, G. Norcic (University Medical Centre Ljubljana).

Spain: V. Vigorita, N. Caceres, E. Casal, A. Ruano, I. Trostchansky (Alvaro Cunqueiro Hospital); T. Golda, A. Galvez Saldaña, E. Kreisler Moreno, J. Lopez Dominguez, M. Vila Tura (Bellvitge University Hospital); F. Labarga, P. Galvez, V. Maderuelo, C. Suero (Complejo Asistencial Universitario de Palencia); J. Bargallo, L. Cayetano, S. Lamas, M. C. Silva (Consorci Sanitari de Terrassa - Hospital de Terrassa); J. C. Bernal-Sprekelsen, R. Gómez, S. Jareño, A. Ríos, D. Vercher (Consorcio Hospital General Universitario); J-M. García-González, J. Cervera-Aldama, J. Ramos-Prada, M. Santamaría-Olabarrieta (Cruces University Hospital); N. Borda, J. M. Enríquez-Navascués, Y. Saralegui (Donostia University Hospital); A. Calero-Lillo, S. Aznar-

Puig, M. A. López-Lara, S. Muñoz-Collado, J. Valverde-Sintas (Fundacio Hospital Esperit Sant); P. Menendez, C. Leon (Gutierrez Ortega Hospital); N. Truan, R. Baldonedo, D. Martínez, J. Otero, L. Solar (Hospital Central de Asturias); V. Turrado-Rodriguez, F. de Lacy Oliver, A. M. Lacy Fortuny, B. Martin Perez, A. M. Otero Piñeiro (Hospital Clinic Barcelona); J. Paredes, F. Fernandez, M. J. Ladra, A. Paulos, D. Prieto (Hospital Clinico Universitario de Santiago de Compostela); J. P. Beltrán de Heredia, F. Blanco Antona, B. de Andrés Asenjo, C. Ferreras García, A. Romero de Diego (Hospital Clínico Universitario de Valladolid); E. Cordoba Diaz de Laspra, E. Echazarreta Gallego, M. Elia Guedea (Hospital Clinico Universitario de Zaragoza); D. Escola, S. Martinez (Hospital Comarcal Alt Penedes); V. Primo Romaguera, R. Parreño, L. Pastor, E. Rosell (Hospital de Dénia); R. Lozoya Trujillo, R. Alós Company, M. D. Ruiz Carmona, A. Solana Bueno (Hospital de Sagunto); S. Salvans Ruiz, S. Alonso Gonçalves, M. Jiménez- Toscano, M. Pascual Damieta, M. Pera Roman (Hospital Del Mar); E. M. Pellicer-Franco, J. A. Garcia-Marin, M. Mengual-Ballester, V. Soria-Aledo, G. Valero-Navarro (Hospital Morales Meseguer); M. Vicente-Ruiz, C. Garcia-Zamora, A. Gonzalez-Gil, M. J. Montoya-Tabares, M. Paredes-Quiles (Hospital Rafael Mendez); J. Die Trill, P. Abadia, I. Moreno, J. D. Pina, D. Ramos Rubio (Hospital Ramon y Cajal); J. Escartin, J. L. Blas, J. Fernando, R. Ferrer, J. Garcia Egea (Hospital Royo Villanova); I. Pros, W. Martinez, J. Rius, M. Socías (Hospital Sant Joan de Deu de Martorell); D. Sabia, J. Castellvi Valls, V. Gonzalez Santin, S. Mompart Garcia, L. Viso Pons (Hospital Sant Joan Despí Moises Broggi); D. Julià, A. Codina-Cazador, R. Farrés, N. Gómez, P. Planellas (Hospital Universitari de Girona Doctor Josep Trueta); M. Cuadrado, I. Camps (Hospital Universitari Germans Trias I Pujol); M. Rufas, J. Escoll, A. Fermiñán, P. Muriel, E. Sierra (Hospital Universitario Arnau de Vilanova de Lerida); C. Alvarez-Laso, P. Lora, H. Padín (Hospital Universitario de Cabueñes); J. Garcia-Septiem, C. Bustamante, V. Jimenez, J. Jimenez-Miramon, J. L. Ramos (Hospital Universitario de Getafe); A. B. Gallardo, P. Benito, L. Colao, P. Galindo, C. Garcia (Hospital Universitario de Torrejón de Ardoz); A. Forero-Torres, A. Alonso Poza, B. Dieguez Fernandez, C. Gilsanz Martin, M.

Hernandez Garcia (Hospital Universitario del Sureste); J. A. Rojo López, J. M. Gil López, M. González Zunzárren, J. Martínez Alegre, J. P. Zorrilla Matilla (Hospital Universitario Infanta Sofia); A. Navarro-Sánchez, F. J. Alcalá Serrano, J. López-Fernández, D. Montesdeoca Cabrera (Hospital Universitario Insular de Gran Canaria); M. Alvarez-Gallego, J. Guevara, I. Pascual Miguelañez, I. Rubio-Perez (Hospital Universitario La Paz); M. Gomez Ruiz, J. Alonso Martín, C. Cagigas Fernández, J. Castillo Diego (Hospital Universitario Marques de Valdecilla); J. A. Pando, C. Maristany, A. Muñoz-Duyos, A. Rada-Palomino, H. Vargas-Pierola (Hospital Universitario Mutua Terrassa); E. Peña Ros, J. A. Benavides Buleje, J. M. Muñoz Camarena, P. A. Parra Baños, M. Ramirez Faraco (Hospital Universitario Reina Sofía); J. J. Arenal, M. A. Cidores, J. L. Marcos, J. Sánchez, C. Tinoco (Hospital Universitario Río Hortega); L. J. García Flórez, R. D. Arias Pacheco, G. Mínguez Ruiz, N. Rodríguez Corral, A. Rodríguez Infante (Hospital Universitario San Agustín); M. J. Carrillo López, M. M. Carrasco Prats, A. Lage Laredo, Á. Martínez Manzano, P. Rodríguez García (Hospital Universitario Santa Lucia); J. J. Segura-Sampedro, N. Alonso-Hernández, M. Fernandez Isart, M. Gamundi Cuesta, A. Ochogavia Segui (Hospital Universitario Son Espases); N. Ibañez, J. Abrisqueta, J. Lujan (Hospital Universitario Virgen de la Arrixaca); R. Gómez Pérez, E. Corrales Valero, C. Monje Salazar, E. Sanchiz Cardenas, R. Soler Humanes (Hospital Universitario Virgen de la Victoria); R. M. Jimenez-Rodriguez, F. De la Portilla, J. M. Diaz Pavon, A. M. Garcia Cabrera, M. L. Reyes Diaz (Hospital Universitario Virgen del Rocio); E. Espin, F. Marinello, M. Martí, J. L. Sanchez, F. Vallribera (Hospital Valle de Hebron); F. J. Orts Mico, M. Ortin Navarro, M. Perez Climent, C. Serra Diaz (Hospital Virgen de los Lirios); M. Millan, A. Caro, J. Escuder, B. Espina, F. Feliu (Joan XXIII University Hospital); A. Climent Aira, A. Estévez Diz, M. T. Moreno Asencio, A. Varela Mato, R. Vázquez Bouzán (POVISA Hospital); A. M. Minaya-Bravo, M.M. Diez-Alonso, R. Villetta-Plaza (Principe de Asturias Hospital); H. Guadalajara, D. Alías, D. García Olmo, C. Pastor, I. Valverde (Quironsalud Publicos); A. Sanchez Romero, A. Gardea, M. Gil Santos, T. Nimmersgern, P. Serrano Paz (Unidad de Coloproctología, Hospital Vinalopó-

Torrevieja); M. Romero-Simó, T. Blasco-Segura, I. Caravaca-García, D. Costa-Navarro, A. Zarco-Pleguezuelos (University General Hospital of Alicante); L. Sánchez-Guillén, B. Flor-Lorente, M. Frasson, Á. García-Granero, E. García-Granero (University Hospital La Fe Valencia); B. Arencibia, J. Alonso, G. Febles, E. M. Nogués, C. Roque (University Hospital of Gran Canaria Dr. Negrín).

Sweden: J. Segelman, J. Nygren (Ersta Hospital); G. Nestler (Falu lasarett); M. Abraham-Nordling, M. Egenvall (Karolinska University Hospital); P. Myrelid, B. Jung, P. Loftås (Linköping University Hospital); M-L. Lydrup, N. Azahr, P. Buchwald, P. Mangell, I. Syk (Skane University Hospital); M. Nikberg, J. Carlander, A. Chabok, K. Smedh, C. Tiselius (Västmanlands Hospital Västerås); S. Haapaniemi, A. Benckert (Vrinnevi Hospital Norrkoping).

Switzerland: M. Adamina, C. Freil-Lanter, C. Gingert, P. Müller, J. Schäfli (Kantonsspital Winterthur); L. Regusci, M. Brenna, F. Fasolini (Regional Hospital Mendrisio); H. Misteli, P. Kirchhoff, D. Oertli (University Hospital Basel, Switzerland); D. Hahnloser, D. Clerc, M. Hübner (University Hospital of Lausanne, CHUV); F. Ris, N. C. Buchs, M. Chevallay, P. Morel, B. Schiltz (University Hospitals Geneva).

Taiwan: J. Y. Wang, W-C. Su, C-W. Huang, C-J. Ma, H-L. Tsai (Kaohsiung Medical University Hospital).

Turkey: G. S. Özbalcı, B. B. Özkan, U. Karabacak (19 Mayıs University Faculty of Medicine); D. Bugra (American Hospital); F. Agalar, H. Baloglu, I. Basoglu (Anadolu Medical Center [in aff with Johns Hopkins Med]); N. Okkabaz, E. Binboga, A. Biricik, A. Celik, E. Yavuz (Bagcilar Training and Research Hospital); A. E. Canda, C. Agalar, M. Fuzun, S. Sokmen, C. Terzi (Dokuz Eylul University); A. Isik (Erzincan University, Mengucek Gazi Training and Research Hospital); B. Karip, A. C. Bilgili (Fatih Sultan Mehmet Training and Research Hospital); S.

Leventoglu, B. Aytac, A. Yıldız, O. Yuksel (Gazi University Medical School); H. Sinan, O. Hancerliogullari, S. Kaymak, O. Kozak, M. T. Ozer (Gulhane Training and Research Hospital); I. S. Sarici, O. Akca, M. U. Kalayci, Y. Kara (Kanuni Sultan Suleyman Training and Research Hospital); D. Bugra, O. Agcaoglu, E. Balik, O. Bayram (Koc University School of Medicine); U. Sungurtekin, U. Ozgen (Pamukkale University School of Medicine); S. Demirbas (TOBB-ETU University Hospital); E. Öztürk, O. Isik, T. Yilmazlar (Uludag University School of Medicine); E. Colak, S. Karagul, V. Kinias (University of Health Sciences, Samsun Training and Research Hospital).

UK: N. Fearnhead, I. Lord, P. Stewart (Addenbrooke's [Cambridge University] Hospital); M. Zammit (Basildon Hospital); S. Arnold, N. Battersby, J. Broadhurst, A. Mehta, F. Seretis (Basingstoke and North Hampshire Hospital); J. Shabbir, C. Jones, J. Kynaston (Bristol Royal Infirmary); D. Vimalachandran, E. Blower, C. McFaul, D. McWhirter, J. Pilkington (Countess of Chester Hospital); T. Wilson, M. Chowdhary (Doncaster Royal Infirmary); B. Stubbs, M. Abdalkoddus, C. Lai, N. Thavanesan, C. Yao (Dorset County Hospital); T. Agarwal, S. Dindyal, R. M. C. Hill, S. Reade, A. Slesser (Ealing Hospital); H. Paterson, A. Balfour, M. Boland, A. Geraghty, J. O'Kelly (Edinburgh Western General Hospital); P. Patel, S. Tezas (Furness General Hospital); S. Yahia, V. Jadhav, K. Marimuthu, A. Narayanan, B. Piramanayagam (George Eliot Hospital); N. Bradley, F. Buchanan, K. Paul, J. Singh, K. Thomson (Glasgow Royal Infirmary); S. Korsgen, M. Bedford, K. Lee, K. Leong (Good Hope Hospital); D. McArthur, A. Bhangu, S. Malik, I. Mohamed (Heartlands Hospital); P. Cunha, A. Pilavas (Homerton University Hospital NHS Trust); A. Reddy, S. Ahmed, A. Ahmed, J. Voll (James Cook University Hospital); V. Velchuru, R. Lal, B. Mirshekar-Syahkal (James Paget Hospital); M. Kassai, M. Aleem, S. Keogh-Bootland (Jersey General Hospital); P. Sarmah, S. Brown, R. Keegan, A. Kelkar, P. Sen (Kettering General Hospital); M. Oliveira-Cunha, S. Chaudhri, R. Fares, B. Singh, W. M. Thomas (Leicester General Hospital); M. I. Aslam, K. Boyle, D. Hemingway, A. Miller, M.

Norwood (Leicester Royal Infirmary); S. Gurjar, M. Al-Saeedi, L. Anandan, A. Sudlow, N. Zampitis (Luton & Dunstable Hospital); K. Malik, M. Bogdan, C. Smart (Macclesfield District General Hospital); M. R. Iqbal, S. Bailey, D. Lawes, G. Omar, R. Tamhane (Maidstone and Tunbridge Wells NHS Trust); M. Evans, S. Ather, J. Lim, H. Nageswaran, G. Taylor (Morriston Hospital); L. Hunt, J. Nicholls (Musgrove Park Hospital); I. Shaikh, F. Muscara, J. O'Brien, E. Photi, A. Stearns (Norfolk and Norwich University Hospital); D. Meylemans, C. Cunningham, R. Hompes (Oxford University Hospitals); A. Tennakoon, N. Kumarasinghe, M. Rao, I. Upanishad (Pilgrim Hospital); S. Smolarek, E. Platt, B. Rossi, J. C. Tham (Plymouth Hospital NHS Trust); J. Khan, N. Ahmad, Z. Shweejawee, S. Stefan (Queen Alexandra Hospital); N. Smart, I. Daniels, T. Gregoir, L. Longstaff, F. McDermott (Royal Devon & Exeter Hospital); M. Varcada, I. Drami, T. Gala, E. Moggia, K. Ratnatunga (Royal Free Hospital NHS Trust Hampstead); R. Harries, J. Hayes, G. Williams (Royal Gwent Hospital); T. Raymond, C. Bronder, E. Davies, P. Hawkin, O. Ryska (Royal Lancaster Infirmary); K. Ayral, A. Beveridge, A. Bhowmik, M. Gill, R. Simpson (Royal Preston Hospital); A. Schofield, K. McArdle, M. Parmar (Royal Shrewsbury Hospital); M. Williamson, H. Burton, E. Courtney, C. Grant, A. Saracino (Royal United Hospital Bath); K. Newton, J. Epstein (Salford Royal NHS Foundation Trust); G. Branagan, M. Bignell, M. Symankewicz (Salisbury District Hospital); S. Zaman, R. Mankotia, Z. Siddiqui, A. Torrance (Sandwell General Hospital); D. Artioukh, M. Eggleston, K. Gokul, D. Selwyn (Southport and Ormskirk Hospitals); J. Warusavitarne, P. Chandrasinghe, J. Grainger , C. A. Leo, C. J. Vaizey (St Mark's Hospital); G. Harris, B. Levy, A. Skull (St Richard's Hospital); M. Thaha, S. Ahmed, A. Garg, H. Patel, A. Ramsanahie (The Royal London Hospital, Barts Health NHS Trust); M. Mondragon-Pritchard, K. Cuinas Leon, G. Williams (The Royal Wolverhampton NHS Trust); A. Shukla, H. Brewer, J. Fitzgerald, H. Kho (United Lincolnshire Hospitals NHS Trust); J. Torkington, S. Tate, J. Wheat (University Hospital of Wales); J. Knight, J. Richardson, A. Tzivanakis (University Hospital Southampton); M. Gregori, M. A. Ashraf, M. Atif, A. Birindelli, J. Santos (University Hospitals Birmingham NHS FT); N. Saffaf, M. I. Aslam, L. Canning (Warwick

Hospital); N. Chandratreya, M. Bowen, B. Graham, Y. Hamad, M. Kaubrys (Weston General Hospital at Weston super Mare); Z. U. Chaudhry, C. Bhan, H. Mukhtar, A. Oshowo, J. Wilson (Whittington Hospital NHS Trust); J. Richardson, N. Gouvas, D. Nicol, S. Pandey, M. Zilvetti (Worcestershire Royal Hospital); A. Sharma, T. Fatayer, S. Mothe, M. Rahman (Wythenshawe Hospital, UHSM); N. Curtis, A. Allison, R. Dalton, N. Francis, J. Ockrim (Yeovil District Hospital). Ukraine: G. Psaras, H. Dudarovaska, T. Marharint, E. Mostovoy, S. Voloshin (Mariupol Cancer Center); O. Kolesnik, D. Makhmudov (National Cancer Institute, Ukraine).

United States: Y. Altinel (Cleveland Clinic); A. Iqbal, L. Cunningham, K. Go, S. Tan (University of Florida).