

Salmonella-induced thrombi in mice develop asynchronously in the spleen and liver and are not effective bacterial traps

Beristain-Covarrubias, Nonantzin; Perez-Toledo, Marisol; Flores-langarica, Adriana; Zuidscherwoude, Malou; Hitchcock, Jessica; Channell, William; King, Lloyd David William; Thomas, Mark R.; Henderson, Ian; Rayes, Julie; Watson, Steve; Cunningham, Adam

DOI:

[10.1182/blood-2018-08-867267](https://doi.org/10.1182/blood-2018-08-867267)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Beristain-Covarrubias, N, Perez-Toledo, M, Flores-langarica, A, Zuidscherwoude, M, Hitchcock, J, Channell, W, King, LDW, Thomas, MR, Henderson, I, Rayes, J, Watson, S & Cunningham, A 2019, 'Salmonella-induced thrombi in mice develop asynchronously in the spleen and liver and are not effective bacterial traps', *Blood*, vol. 133, no. 6, pp. 600–604. <https://doi.org/10.1182/blood-2018-08-867267>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Final Version published in *Blood* as detailed above and is available at: <https://doi.org/10.1182/blood-2018-08-867267>

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

***Salmonella*-induced thrombi in mice develop asynchronously in the spleen and liver and are not effective bacterial traps**

Short title:

***Salmonella*-induced thrombi do not trap bacteria**

Nonantzin Beristain-Covarrubias^{1*}, Marisol Perez-Toledo^{1*}, Adriana Flores-Langarica¹, Malou Zuidschewoude², Jessica R Hitchcock¹, Will M. Channell¹, Lloyd D.W. King¹, Mark R. Thomas², Ian R. Henderson³, Julie Rayes², Steve P. Watson^{2,4}, and Adam F. Cunningham¹

¹Institute of Immunology and Immunotherapy, College of Medical and Dental Sciences, University of Birmingham, UK; ²Institute of Cardiovascular Sciences, College of Medical and Dental Sciences, University of Birmingham, UK; ³Institute for Microbiology and Infection, College of Medical and Dental Sciences, University of Birmingham, UK; and ⁴Centre of Membrane Proteins and Receptors, Universities of Birmingham and Nottingham, The Midlands, UK

* N.B.C. and M.P.T. contributed equally to this work.

Corresponding authors:

Adam F. Cunningham, ¹Institute of Immunology and Immunotherapy, College of Medical and Dental Sciences, University of Birmingham, B15 2TT, UK. A.f.cunningham@bham.ac.uk. Phone: +441214144068, Fax: +441214143599

Steve P. Watson, ²Institute of Cardiovascular Sciences, College of Medical and Dental Sciences, University of Birmingham, B15 2TT, UK. S.p.watson@bham.ac.uk. Tel: +441214146514

Text word count: 1192

Abstract word count: 195

Figures: 2

Tables: 0

References: 19

Key words: Immunothrombosis, platelets, monocytes, inflammation, podoplanin, *Salmonella*

Key points:

- Thrombosis develops in the spleen and liver with distinct kinetics following *Salmonella* infection
- Thrombi in the spleen and liver are not major sites of bacterial localisation

Abstract

Thrombosis is a frequent, life-threatening complication of systemic infection, associated with multiple organ damage. We have previously described a novel mechanism of inflammation-driven thrombosis induced by *Salmonella* Typhimurium infection of mice. Thrombosis in the liver develops 7 days post-infection persisting after the infection resolves, and is monocytic cell-dependent. Unexpectedly, thrombosis was not prominent in the spleen at this time, despite carrying a similar bacterial burden as the liver. In this study, we show that thrombosis does occur in the spleen but with strikingly accelerated kinetics compared to the liver, being evident by 24 h and resolving rapidly thereafter. The distinct kinetics of thrombosis and bacterial burden provide a test of the hypothesis that thrombi form in healthy vessels to trap or remove bacteria from the circulation, often termed immunothrombosis. Remarkably, despite bacteria being detected throughout infected spleens and livers in the early days of infection, immunohistological analysis of tissue sections show that thrombi contain very low numbers of bacteria. In contrast, bacteria are present throughout platelet aggregates induced by *Salmonella* *in vitro*. Therefore, we show that thrombosis develops with organ-specific kinetics and challenge the universality of immunothrombosis as a mechanism to capture bacteria *in vivo*.

Introduction

The consequences of thrombosis are the leading cause of death worldwide¹. Thrombosis is common after infection and can lead to organ failure and poor outcome²⁻⁵. There are however significant gaps in our understanding of blood-borne infection-associated thrombosis, including whether it occurs at multiple sites through distinct mechanisms and/or kinetics². Immune-driven thrombosis, broadly-termed “immunothrombosis”, can occur in the presence or absence of infection. Nevertheless, when

triggered by infection, it is still unclear whether the induced thrombi capture and contain blood-borne pathogens within the vasculature as proposed⁶⁻⁸.

We recently reported on a novel pathway of thrombosis in the liver after infection with *Salmonella* Typhimurium (STm), involving inflammation-driven upregulation of podoplanin on monocytic cells and activation of platelets⁹. A striking feature of this thrombosis is that it takes a week to develop and then persists as the bacterial burden declines. Furthermore, thrombi are largely undetectable in the spleen at this time, despite this organ being a major site of bacterial colonisation^{10,11}. In this paper, we show that extensive thrombosis does occur in the spleen but is rapid in onset and transient, with distinct kinetics to liver. Furthermore, we show that thrombi present in either organ contain surprisingly few bacteria, despite the high bacterial burdens in the organs themselves, indicating that bacterial entrapment is not a major consequence of thrombosis after infection with STm.

Study design

Full details are provided in the Supplemental Material

Mice and infection with STm

Wild-type (WT), C57BL/6 mice (Home Office Licenses 3028/50 and P2E63AE7B) were infected intraperitoneally (i.p.) or intravenously (i.v.) with $1-5 \times 10^5$ attenuated SL3261 or virulent SL1344 STm^{12,13}.

Immunohistology and fluorescent microscopy

Cryosections were stained for immunohistochemistry (IHC) or immunofluorescence (IF)^{13,14} to detect CD41, CD31, fibrin/fibrinogen, Ly6G, Ly6C, F4/80, *Salmonella* and nuclei (DAPI; Supplemental Table 1)⁹.

Clodronate treatment

Mice were treated i.p. with either 200 μ l (5 mg/ml) of clodronate or PBS liposomes 24 h before STm infection^{15,16}.

Results and Discussion

Thrombosis develops with distinct kinetics in the spleen and liver

Thrombosis in the liver becomes established 7 days after infection, whereas few thrombi are detectable in the spleen at this time⁹. In the liver, thrombosis is driven by the up-regulation of podoplanin on monocytes/macrophages, triggering activation of CLEC-2 on platelets⁹. The spleen is a reservoir of monocytic cells, with high numbers of these cells present pre-infection compared to the liver, and 24 hours post-infection there were increased numbers of inflammatory splenic monocytes (Supplemental Figure S1), suggesting that thrombosis may occur at a much earlier stage in the infection. Consistent with this, we found numerous, large platelet-rich thrombi within the spleen at 24 h post-infection (Figure 1A), independent of whether mice were infected i.v. or i.p. or of the virulence of the infecting strain (Figure 1B-C). Thrombi typically stained positive for citrullinated-histone H3, Ly6G⁺ cells and myeloperoxidase (Supplemental Figure S2). Ly6C⁺ and F4/80⁺ cells were located at the periphery of thrombi (Supplemental Figure S3). Splenic thrombosis resolved rapidly after day 1, with few thrombi detected thereafter (Figure 1D,E), often leaving the remnants of a fibrin core (Figure 1 and 2A). In contrast, at these early times, thrombosis was undetectable in the liver (Figure 1B,C,D,F). Moreover, thrombosis was absent in the spleens of clodronate-liposome treated mice (Figure 1G) suggesting that, like in the liver, monocytic cells are important in this process⁹. Therefore, systemic infection with STm can induce thrombosis in distinct sites and with distinct kinetics, likely due to the levels of tissue-resident macrophages present at the time of infection.

Most thrombi induced in the spleen and liver contain limited numbers of bacteria

It has been proposed that thrombus formation can trap and remove bacteria, a process sometimes known as immunothrombosis⁶. After platelet activation induced by STm *in vitro*, bacteria are present throughout the aggregate as shown in Supplemental Figure S4 and video 1. This demonstrates that bacteria can closely associate with platelets in aggregates formed *in vitro*. We used IHC and IF microscopy to identify the relationship between bacterial localisation and thrombi *in vivo* at the peak times of thrombosis in the spleen (day 1) and liver (day 7) (Figure 2A-B and Supplemental Figure

S5). A 3-dimensional reconstruction of a thrombus and proximal bacteria is shown in Supplemental Figure S6 and Supplemental Video 2. Collectively, these approaches all showed that thrombi contained a surprisingly low number of bacteria, despite their relative abundance in the surrounding tissues. Quantification of the bacteria within sections of splenic thrombi (>200 thrombi from 37 mice, 3 time points evaluated) showed that no bacteria were detected in 38% of thrombus sections at day 1, and that 33% of sectioned thrombi contained 1-2 bacteria (Figure 2C). At later times, bacteria were detected at an even lower frequency within sections of splenic thrombi, with >90% of thrombi containing 0-2 bacteria at day 7 and day 21 (Figure 2C). In the liver, only ~20% of thrombi sections (>400 thrombi counted from 23 mice) contained bacteria at day 7 and this proportion was even lower (<5%) at day 21 (Figure 2D). Analysis of serial sections from the same thrombi confirmed a paucity of bacteria within individual thrombi (Supplemental Figure S7). When the bacterial burdens per organ were compared with the levels of thrombosis at days 1 to 21 post-infection, no direct relationship was found between the two, other than the necessity for infection to induce thrombosis. In the spleen, thrombosis peaks before bacterial numbers peak (Figure 2E) and falls whilst bacterial numbers are still rising, whereas in the liver thrombosis develops later and peaks when the bacterial levels are beginning to decline (Figure 2F)^{9,13,17}. Therefore, thrombi induced during this infection do not trap significant numbers of bacteria, regardless of the bacterial loads in the organs. This contrasts with other models of infection^{18,19}, which used 1000-fold higher numbers of bacteria compared to here⁹. Thus, although STm infection can drive thrombosis, thrombi do not necessarily contribute to bacterial containment and moreover they form in different organs with distinct kinetics. These findings are important for our understanding of the consequences of infection on the haematological system since they show that the presence of equivalent levels of bacteria is not enough to induce thrombosis in an organ.

Although these data show that during a single infection thrombosis can occur sequentially in multiple tissues, further work is needed to evaluate whether other systemic bacterial infections induce thrombosis with similar kinetics. Thrombi are induced by many different pathogens and although the role of thrombosis after infection remains unclear, the presumption must be that they are pathological

in some circumstances, particularly if they are large and/or embolise. Since we have shown that thrombi develop in the venous system, they may form due to differences in local infection-associated changes in blood flow and as part of altered vessel homeostasis. Perhaps the bigger clinical question is about what controls their ultimate size and what triggers thrombus resolution as this may influence whether thrombosis becomes clinically problematic. Therefore, for a known infection it may be possible to target therapeutically those organs at greatest risk of developing thrombosis at particular stages of infection. These findings deepen our understanding of the concept of immunothrombosis and shows thrombi can form as a non-canonical haemostatic response to infection-driven inflammation but not to capture bacteria.

Acknowledgements

The authors wish to thank the staff of the BMSU for their assistance. This work was supported by grants from the Medical Research Council (MR/N023706/1) and the British Heart Foundation (BHF) (RG/13/18/30563). N.B.C. was supported by National Council of Science and Technology (CONACYT)-Mexico overseas postdoctoral fellowship (CVU 332390). S.P.W. is a BHF Chair (CH/03/003).

Authorship

Contribution: N.B.C. and M.P.T. designed and performed the experiments, analysed the data and wrote the manuscript; A.F.L., M.Z., J.R.H., L.D.W.K., W.M.C. performed the experiments and analysed the data; M.R.T., J.R. contributed vital reagents, experimental design and proofread the manuscript; I.R.H. experimental design, novel reagents and proofread the manuscript; A.F.C. and S.P.W. supervised the research, analysed the data, and wrote the manuscript.

Conflict-of-interest disclosure:

The authors declare no competing financial interests.

References

1. World Health Organization. Fact Sheet The Top Ten Causes of Death. Vol. 2015; 2015.
2. Furie B, Furie BC. Mechanisms of thrombus formation. *N Engl J Med*. 2008;359(9):938-949.
3. Pawlinski R, Pedersen B, Schabbauer G, et al. Role of tissue factor and protease-activated receptors in a mouse model of endotoxemia. *Blood*. 2004;103(4):1342-1347.
4. Pawlinski R, Wang JG, Owens AP, 3rd, et al. Hematopoietic and nonhematopoietic cell tissue factor activates the coagulation cascade in endotoxemic mice. *Blood*. 2010;116(5):806-814.
5. Angus DC, van der Poll T. Severe sepsis and septic shock. *N Engl J Med*. 2013;369(9):840-851.
6. Engelmann B, Massberg S. Thrombosis as an intravascular effector of innate immunity. *Nature Reviews Immunology*. 2013;13(1):34-45.
7. van der Poll T, Herwald H. The coagulation system and its function in early immune defense. *Thromb Haemost*. 2014;112(4):640-648.
8. Davis RP, Miller-Dorey S, Jenne CN. Platelets and coagulation in infection. *Clin Transl Immunology*. 2016;5(7):e89.
9. Hitchcock JR, Cook CN, Bobat S, et al. Inflammation drives thrombosis after Salmonella infection via CLEC-2 on platelets. *Journal of Clinical Investigation*. 2015;125(12):4429-4446.
10. Monack DM, Mueller A, Falkow S. Persistent bacterial infections: The interface of the pathogen and the host immune system. *Nature Reviews Microbiology*. 2004;2(9):747-765.
11. Mastroeni P, Grant A, Restif O, Maskell D. A dynamic view of the spread and intracellular distribution of Salmonella enterica. *Nat Rev Microbiol*. 2009;7(1):73-80.
12. Cunningham AF, Khan M, Ball J, et al. Responses to the soluble flagellar protein FlhC are Th2, while those to FlhC on Salmonella are Th1. *Eur J Immunol*. 2004;34(11):2986-2995.
13. Cunningham AF, Gaspal F, Serre K, et al. Salmonella induces a switched antibody response without germinal centers that impedes the extracellular spread of infection. *Journal of Immunology*. 2007;178(10):6200-6207.
14. Flores-Langarica A, Marshall JL, Bobat S, et al. T-zone localized monocyte-derived dendritic cells promote Th1 priming to Salmonella. *European Journal of Immunology*. 2011;41(9):2654-2665.
15. Buiting AM, Van Rooijen N. Liposome mediated depletion of macrophages: an approach for fundamental studies. *J Drug Target*. 1994;2(5):357-362.
16. Van Rooijen N. The liposome-mediated macrophage 'suicide' technique. *J Immunol Methods*. 1989;124(1):1-6.
17. Gil-Cruz C, Bobat S, Marshall JL, et al. The porin OmpD from nontyphoidal Salmonella is a key target for a protective B1b cell antibody response. *Proc Natl Acad Sci U S A*. 2009;106(24):9803-9808.
18. Massberg S, Grahl L, von Bruehl ML, et al. Reciprocal coupling of coagulation and innate immunity via neutrophil serine proteases. *Nature Medicine*. 2010;16(8):887-U887.
19. Gaertner F, Ahmad Z, Rosenberger G, et al. Migrating Platelets Are Mechano-scavengers that Collect and Bundle Bacteria. *Cell*. 2017;171(6):1368-1382 e1323.

Figure legends

Figure 1. Thrombosis in the spleen and liver follows different kinetics after STm infection. (A)

Frozen spleens from WT mice infected with 5×10^5 STm i.p were sectioned longitudinally to the hilum ($\geq 1200 \mu\text{m}$ deep) and $5\text{-}\mu\text{m}$ sections were stained by IHC. Scans of stained spleen sections from day 0, 1, 7 and 21-infected mice show blood vessels identified with anti-CD31 in blue and thrombi with anti-fibrin/fibrinogen in brown. Arrows identify individual thrombi. (B) Representative low power

images of spleen and liver sections stained for CD41 (platelets; blue) and fibrin/fibrinogen (brown) from mice infected for 24 hours via the i.v. route with 5×10^5 STm SL3261. The left hand graph shows the number of thrombi per spleen section for mice infected i.p. or i.v. The right hand graph shows the number of thrombi per section in the spleen and liver from these i.v. infected mice. (C) Representative low power images of spleen and liver sections stained for CD41 (platelets; blue) and fibrin/fibrinogen (brown) from mice infected for 24 hours with the virulent 10^5 STm SL1344 strain. The left hand graph shows the number of thrombi per spleen section for mice infected with SL3261 (attenuated) or SL1344 (virulent). The right hand graph shows the number of thrombi per section in the spleen and liver from mice infected with SL1344. (D) Representative scans at higher magnification of spleen sections (upper panels) and liver sections (lower panels) from WT mice at day 0, 1, 7 and 21 post-infection with 5×10^5 STm SL3261. Sections are stained for fibrin/fibrinogen (brown) and CD41 (blue). V=Vein, RP=Red Pulp, WP=White Pulp, T=Thrombus. (E) Quantification of numbers of thrombi per spleen section (left graph) and the proportion of section area covered by thrombi (right graph) at days 0, 1, 2, 7 and 21 after infection with 5×10^5 STm SL3261. Each point represents a single mouse (Data are combined from 3 independent experiments); * $p < 0.05$, 1 way ANOVA. (F) Line graph showing the level of thrombosis in the spleen (black line) and the liver (grey line) over the first 3 weeks of infection with 5×10^5 STm SL3261. The data are expressed as mean \pm S.E.M. from at least 4 mice per group combined from 3 independent experiments. (G) Quantification of thrombi in spleen sections from PBS liposomes or clodronate liposome pre-treated mice, infected for 24 hours with 5×10^5 STm SL3261. Combined data from 2 experiments with a total of 8 mice in each group. * $p < 0.05$. 2-tailed non-parametric t test. N.D.=Not detected.

Figure 2. Detection of bacteria within thrombi. Representative immunofluorescence photomicrographs of (A) spleens and (B) livers from WT mice infected with 5×10^5 STm SL3261 for 0, 1, 7 and 21 days. (V=Vein, RP=Red Pulp, WP=White Pulp, T=Thrombus, MK=Megakaryocyte). Fibrin, blue; CD31, white; CD41, red and STm, green indicated with white arrows. For both (A) and (B) the second row shows a higher magnification image of the area identified by the white box. (C) and (D) Frequency of detecting 0, 1, 2 or ≥ 3 bacteria in thrombi in sections from spleens and livers

232 respectively, infected with 5×10^5 STm SL3261 for 0, 1, 7 or 21 days. (E) and (F) Line graphs showing
233 the kinetics of thrombosis (black) and bacterial colonisation (gray) in spleens and livers respectively,
234 from mice infected with 5×10^5 STm SL3261 for 0, 1, 2, 7 or 21 days. Data are expressed as mean \pm
235 S.E.M. from 152 thrombi counted in spleens from day 1; 40 from day 7; and 18 from day 21-infected
236 mice. In liver, 411 thrombi were counted for day 7 and 23 for day 21 after infection. In each case
237 thrombi were counted from at least 4 mice per group and are combined from 3 independent
238 experiments. CFU= Colony-forming unit.